

6 Literaturverzeichnis

- Annuk M, Lind L, Linde T, Fellstrom B. Impaired endothelium-dependent vasodilatation in renal failure in humans. *Nephrol Dial Transplant*. 2001 Feb;16(2):302-6.
- Annuk M, Zilmer M, Lind L, Linde T, Fellstrom B. Oxidative stress and endothelial function in chronic renal failure. *J Am Soc Nephrol*. 2001 Dec;12(12):2747-52.
- Asmar R, Rudnichi A, Blacher J, London GM, Safar ME. Pulse pressure and aortic pulse wave are markers of cardiovascular risk in hypertensive populations. *Am J Hypertens*. 2001 Feb;14(2):91-7.
- Bachmann J, Tepel M, Raidt H et al. Hyperhomocysteinemia and the risk for vascular disease in hemodialysis patients. *J Am Soc Nephrol*. 1995 Jul;6(1):121-5.
- Barenbrock M, Hausberg M, Kosch M, Kisters K, Hoeks AP, Rahn KH. Effect of hyperparathyroidism on arterial distensibility in renal transplant recipients. *Kidney Int*. 1998 Jul;54(1):210-5.
- Benetos A, Safar M, Rudnichi A et al. Pulse pressure: a predictor of long-term cardiovascular mortality in a French male population. *Hypertension*. 1997 Dec;30(6):1410-5.
- Blacher J, Asmar R, Djane S, London GM, Safar ME. Aortic pulse wave velocity as a marker of cardiovascular risk in hypertensive patients. *Hypertension*. 1999 May;33(5):1111-7.
- Blacher J, Guerin AP, Pannier B, Marchais SJ, London GM. Arterial calcifications, arterial stiffness, and cardiovascular risk in end-stage renal disease. *Hypertension*. 2001 Oct;38(4):938-42.
- Blacher J, Safar ME, Guerin AP, Pannier B, Marchais SJ, London GM. Aortic pulse wave velocity index and mortality in end-stage renal disease. *Kidney Int*. 2003 May;63(5):1852-60.
- Boaz M, Weinstein T, Matas Z, Green, Smetana S. Peripheral vascular disease and serum phosphorus in hemodialysis: a nested case-control study. *Clin Nephrol*. 2005 Feb;63(2):98-105.
- Bowes WA 3rd, Corke BC, Hulka J. Pulse oximetry: a review of the theory, accuracy, and clinical applications. *Obstet Gynecol*. 1989 Sep;74(3 Pt 2):541-6.
- Briganti EM, Kerr PG, Shaw JE, Zimmet PZ, Atkins RC. Prevalence and treatment of cardiovascular disease and traditional risk factors in Australian adults with renal insufficiency. *Nephrology (Carlton)*. 2005 Feb;10(1):40-7.
- Broome IJ, Mills GH, Spiers P, Reilly CS. An evaluation of the effect of vasodilatation on oxygen saturations measured by pulse oximetry and venous blood gas analysis. *Anaesthesia* 1993 May;48(5):415-6.
- Buccianti G, Baragetti I, Bamonti F, Furiani S, Dorighet V, Patrosso C. Plasma homocysteine levels and cardiovascular mortality in patients with end-stage renal disease. *J Nephrol*. 2004 May-Jun;17(3):405-10.
- Calabro P, Willerson JT, Yeh ET. Inflammatory cytokines stimulate C-reactive protein production by human coronary artery smooth muscle cells. *Circulation*. 2003 Oct 21;108(16):1930-2. Epub 2003 Oct 6.
- Charra B, Chazot C. Volume control, blood pressure and cardiovascular function. Lessons from hemodialysis treatment. *Nephron Physiol*. 2003;93(4):p94-101.
- Chow KM, Szeto CC, Li PK. Parathyroid hormone and mineral metabolism do not have significant impact on pulse pressure in patients undergoing peritoneal dialysis. *Clin Nephrol*. 2003 Oct;60(4):266-9.

- Chowienczyk PJ, Kelly RP, MacCallum H et al. Photoplethysmographic assessment of pulse wave reflection: blunted response to endothelium-dependent beta2-adrenergic vasodilation in type II diabetes mellitus. *J Am Coll Cardiol.* 1999 Dec;34(7):2007-14.
- Clarke R, Daly L, Robinson K et al. Hyperhomocysteinemia: an independent risk factor for vascular disease. *N Engl J Med.* 1991 Apr 25;324(17):1149-55.
- Cockcroft DW, Gault MH. Prediction of creatinine clearance from serum creatinine. *Nephron.* 1976;16(1):31-41.
- Cooke JP. Does ADMA Cause Endothelial Dysfunction? *Arterioscler Thromb Vasc Biol.* 2000 Sep;20(9):2032-7.
- Corretti MC, Anderson TJ, Benjamin EJ et al. International Brachial Artery Reactivity Task Force. Guidelines for the ultrasound assessment of endothelial-dependent flow-mediated vasodilation of the brachial artery: a report of the International Brachial Artery Reactivity Task Force. *J Am Coll Cardiol.* 2002 Jan 16;39(2):257-65.
- Cressmann MD, Heyka RJ, Paganini EP, O'Neil J, Skibinski CI, Hoff HF. Lipoprotein(a) is an independent risk factor for cardiovascular disease in hemodialysis patients. *Circulation.* 1992 Aug;86(2):475-82.
- Fellner SK, Lang RM, Neumann A, Bushinski DA, Borow KM. Parathyroid hormone and myocardial performance in dialysis patients. *Am J Kidney Dis.* 1991 Sep;18(3):320-5.
- Fitchett DH. Forearm arterial compliance: a new measure of arterial compliance? *Cardiovasc Res.* 1984 Nov;18(11):651-6.
- Foley RN, Herzog CA, Collins AJ. Smoking and cardiovascular outcomes in dialysis patients: the United States Renal Data System Wave 2 study. *Kidney Int.* 2003 Apr;63(4):1462-7.
- Foley RN, Parfrey PS, Harnett JD et al. Hypocalcemia, morbidity, and mortality in end-stage renal disease. *Am J Nephrol.* 1996;16(5):386-93.
- Foley RN, Parfrey PS, Harnett JD, Kent GM, Murray DC, Barre PE. The impact of anemia on cardiomyopathy, morbidity, and mortality in end-stage renal disease. *Am J Kidney Dis.* 1996 Jul;28(1):53-61.
- Foley RN, Parfrey PS, Harnett JD, Kent GM, Murray DC, Barre PE. Hypoalbuminemia, cardiac morbidity, and mortality in end-stage renal disease. *J Am Soc Nephrol.* 1996 May;7(5):728-36.
- Foley RN, Parfrey PS, Sarnak MJ. Epidemiology of cardiovascular disease in chronic renal disease. *J Am Soc Nephrol.* 1998 Dec;9(12 Suppl):S16-23.
- Frei U, Schober-Halstenberg HJ. Nierenersatztherapie in Deutschland. *QuaSi-Niere Jahresbericht 2003/2004*, Berlin, Deutschland.
- Furumoto T, Saito N, Dong J, Mikami T, Fujii S, Kitabatake A. Association of cardiovascular risk factors and endothelial dysfunction in Japanese hypertensive patients: implications for early atherosclerosis. *Hypertens Res.* 2002 May;25(3):475-80.
- Gerstein HC, Pogue J, Mann JF et al. The relationship between dysglycaemia and cardiovascular and renal risk in diabetic and non-diabetic participants in the HOPE study: a prospective epidemiological analysis. *Diabetologia.* 2005 Sep;48(9):1749-55. Epub 2005
- Hansell J, Henareh L, Agewall S, Norman M. Non-invasive assessment of endothelial function - relation between vasodilatory responses in skin microcirculation and brachial artery. *Clin Physiol Funct Imaging.* 2004 Nov;24(6):317-22.
- Hsu CY, McCulloch CE, Darbinian J, Go AS, Iribarren C. Elevated blood pressure and risk of end-stage renal disease in subjects without baseline kidney disease. *Arch Intern Med.* 2005 Apr 25;165(8):923-8.
- Iseki K, Tozawa M, Yoshi S, Fukiyama K. Serum C-reactive protein (CRP) and risk of death in chronic dialysis patients. *Nephrol Dial Transplant.* 1999 Aug;14(8):1956-60.
- Jialal I, Devaraj S, Venugopal SK. C-reactive protein: risk marker or mediator in atherosclerosis? *Hypertension.* 2004 Jul;44(1):6-11. Epub 2004 May 17.

- Jiang JP, Hou FF, Gao LZ et al. Improvement of left ventricular hypertrophy by anemic-correcting erythropoietin therapy in chronic renal insufficiency *Zhonghua Nei Ke Za Zhi*. 2005 Jan;44(1):25-9.
- Joannides R, Bakkali EH, Le Roy F et al. Altered flow-dependent vasodilatation of conduit arteries in maintenance haemodialysis. *Nephrol Dial Transplant*. 1997 Dec;12(12):2623-8.
- Joannides R, Haefeli WE, Linder L et al. Nitric oxide is responsible for flow-dependent dilatation of human peripheral conduit arteries in vivo. *Circulation*. 1995 Mar 1;91(5):1314-9.
- Jungers P, Chauveau P, Bandin O et al. Hyperhomocysteinemia is associated with atherosclerotic occlusive arterial accidents in predialysis chronic renal failure patients. *Miner Electrolyte Metab*. 1997;23(3-6):170-3.
- Kalantar-Zadeh K, Stenvinkel P, Pillon L, Kopple JD. Inflammation and nutrition in renal insufficiency. *Adv Ren Replace Ther*. 2003 Jul;10(3):155-69.
- Kaysen GA, Rathore V, Shearer GC, Depner TA. Mechanisms of hypoalbuminemia in hemodialysis patients. *Kidney Int*. 1995 Aug;48(2):510-6.
- Kelly R, Hayward C, Avolio A, O'Rourke M. Noninvasive determination of age-related changes in the human arterial pulse. *Circulation*. 1989 Dec;80(6):1652-9.
- Kelm M. Flow-mediated dilatation in human circulation: diagnostic and therapeutic aspects. *Am J Physiol Heart Circ Physiol*. 2002 Jan;282(1):H1-5.
- Kimura Y, Matsumoto M, Den YB et al. Impaired endothelial function in hypertensive elderly patients evaluated by high resolution ultrasonography. *Can J Cardiol*. 1999 May;15(5):563-8.
- Kundhal K, Lok CE. Clinical Epidemiology of Cardiovascular Disease in Chronic Kidney Disease. *Nephron Clin Pract*. 2005;101(2):c47-52. Epub 2005 Jun 7.
- Laurent S, Boutouyrie P, Asmar R et al. Aortic stiffness is an independent predictor of all-cause and cardiovascular mortality in hypertensive patients. *Hypertension*. 2001 May;37(5):1236-41.
- Leskinen Y, Salenius JP, Lehtimaki T, Huhtala H, Saha H. The prevalence of peripheral arterial disease and medial arterial calcification in patients with chronic renal failure: requirements for diagnostics. *Am J Kidney Dis*. 2002 Sep;40(3):472-9.
- Levey AS, Bosch JP, Lewis JB, Greene T, Rogers N, Roth D. A more accurate method to estimate glomerular filtration rate from serum creatinine: a new prediction equation. Modification of Diet in Renal Disease Study Group. *Ann Intern Med*. 1999 Mar 16;130(6):461-70.
- Locatelli F, Canaud B, Eckardt KU, Stenvinkel P, Wanner C, Zocalli C. Oxidative stress in end-stage renal disease: an emerging threat to patient outcome. *Nephrol Dial Transplant*. 2003 Jul;18(7):1272-80.
- London GM, Blacher J, Pannier B, Guerin AP, Marchais SJ, Safar ME. Arterial wave reflections and survival in end-stage renal failure. *Hypertension*. 2001 Sep;38(3):434-8.
- London GM, Guerin AP, Marchais SJ et al. Cardiac and arterial interactions in end-stage renal disease. *Kidney Int*. 1996 Aug;50(2):600-8.
- London GM. Left ventricular alterations and end-stage renal disease. *Nephrol Dial Transplant*. 2002;17 Suppl 1:29-36.
- Low I, Grutzmacher P, Bergmann M, Schoeppe W. Echocardiographic findings in patients on maintenance hemodialysis substituted with recombinant human erythropoietin. *Clin Nephrol*. 1989 Jan;31(1):26-30.
- Machado AP, Alcantara P. Nitric oxide-dependent endogenous and exogenous vasodilators. The mechanism of action of nitrates. *Rev Port Cardiol*. 1995 Jan;14(1):61-71.

- Madhavan S, Ooi WL, Cohen H, Alderman MH. Relation of pulse pressure and blood pressure reduction to the incidence of myocardial infarction. *Hypertension*. 1994 Mar;23(3):395-401.
- Mallamaci F, Zocalli C, Tripepi G et al. CREED Investigators. Hyperhomocysteinemia predicts cardiovascular outcomes in hemodialysis patients. *Kidney Int*. 2002 Feb;61(2):609-14.
- Massy ZA, Chadefaux-Vekemans B, Chevalier A et al. Hyperhomocysteinaemia: a significant risk factor for cardiovascular disease in renal transplant recipients. *Nephrol Dial Transplant*. 1994;9(8):1103-8.
- McClellan W, Aronoff SL, Bolton WK et al. The prevalence of anemia in patients with chronic kidney disease. *Curr Med Res Opin*. 2004 Sep;20(9):1501-10.
- Meeus F, Kourilsky O, Guerin AP, Gaudry C, Marchais SJ, London GM. Pathophysiology of cardiovascular disease in hemodialysis patients. *Kidney Int Suppl*. 2000 Aug;76:S140-7.
- Migliacci R, Falcinelli F, Imperiali P, Floridi A, Nenci GG, Gresele P. Endothelial dysfunction in patients with kidney failure and vascular risk factors: acute effects of hemodialysis. *Ital Heart J*. 2004 May;5(5):371-7.
- Millasseau SC, Guigui FG, Kelly RP et al. Noninvasive assessment of the digital volume pulse. Comparison with the peripheral pressure pulse. *Hypertension*. 2000 Dec;36(6):952-6.
- Millasseau SC, Kelly RP, Ritter JM, Chowienczyk PJ. Determination of age-related increases in large artery stiffness by digital pulse contour analysis. *Clin Sci (Lond)*. 2002 Oct;103(4):371-7.
- Moustapha A, Naso A, Nahlawi M et al. Prospective study of hyperhomocysteinemia as an adverse cardiovascular risk factor in end-stage renal disease. *Circulation*. 1998 Jan 20;97(2):138-41.
- Nguyen-Khoa T, Massy ZA, De Bandt JP et al. Oxidative stress and haemodialysis: role of inflammation and duration of dialysis treatment. *Nephrol Dial Transplant*. 2001 Feb;16(2):335-40.
- Orth SR, Schroeder T, Ritz E, Ferrari P. Effects of smoking on renal function in patients with type 1 and type 2 diabetes mellitus. *Nephrol Dial Transplant*. 2005 Nov;20(11):2414-9.
- Palmer RM, Ferrige AG, Moncada S. Nitric oxide release accounts for the biological activity of endothelium-derived relaxing factor. *Nature*. 1987 Jun 11-17;327(6122):524-6.
- Pawlaczyk K, Oko A, Lindholm B, Czekalski S. Malnutrition -- inflammation -- atherosclerosis (MIA syndrome) in patients with renal failure *Pol Merkur Lek*. 2003 Oct;15(88):334-41; discussion 341-3.
- Perunicic-Pekovic G, Rasic-Milutinovic Z, Pljesa S. Predictors of mortality in dialysis patients-- association between malnutrition, inflammation and atherosclerosis (MIA syndrome) *Med Pregl*. 2004 Mar-Apr;57(3-4):149-52.
- Pohl U, Holtz J, Busse R, Bassenge E. Crucial role of endothelium in the vasodilator response to increased flow in vivo. *Hypertension*. 1986 Jan;8(1):37-44.
- Pozzoni P, Del Vecchio L, Locatelli F. Anemia treatment to reduce mortality risk and to improve quality of life in chronic uremic patients. *G Ital Nefrol*. 2005 Jan-Feb;22 Suppl 31:S41-6.
- Safar ME, Blacher J, Pannier B et al. Central pulse pressure and mortality in end-stage renal disease. *Hypertension*. 2002 Mar 1;39(3):735-8.
- Saka B, Oflaz H, Erten N et al. Non-invasive evaluation of endothelial function in hypertensive elderly patients. *Arch Gerontol Geriatr*. 2005 Jan-Feb;40(1):61-71.
- Santoro A, Canova C. Anemia and erythropoietin treatment in chronic kidney diseases. *Minerva Urol Nefrol*. 2005 Mar;57(1):23-31.
- Schachinger V, Britten MB, Zeiher AM. Prognostic impact of coronary vasodilator dysfunction on adverse long-term outcome of coronary heart disease. *Circulation*. 2000 Apr 25;101(16):1899-906.

- Scholze A, Rinder C, Beige J, Riezler R, Zidek W, Tepel M. Acetylcysteine reduces plasma homocysteine concentration and improves pulse pressure and endothelial function in patients with end-stage renal failure. *Circulation*. 2004 Jan 27;109(3):369-74. Epub 2004 Jan 19.
- Schramm WM, Bartunek A, Gilly H. Effect of local limb temperature on pulse oximetry and the plethysmographic pulse wave. *Int J Clin Monit Comput*. 1997 Feb;14(1):17-22.
- Silber HA, Bluemke DA, Ouyang P, Du YP, Post WS, Lima JA. The relationship between vascular wall shear stress and flow-mediated dilation: endothelial function assessed by phase-contrast magnetic resonance angiography. *J Am Coll Cardiol*. 2001 Dec;38(7):1859-65.
- Simmons EM, Langone A, Sezer MT et al. Effect of renal transplantation on biomarkers of inflammation and oxidative stress in end-stage renal disease patients. *Transplantation*. 2005 Apr 27;79(8):914-9.
- Stenvinkel P. Inflammation in end-stage renal disease--a fire that burns within. *Contrib Nephrol*. 2005;149:185-99.
- Suwaidei JA, Hamasaki S, Higano ST, Nishimura RA, Holmes DR Jr., Lerman A. Long-term follow-up of patients with mild coronary artery disease and endothelial dysfunction. *Circulation*. 2000 Mar 7;101(9):948-54.
- Taddei S, Virdis A, Ghiadoni L, Magagna A, Salvetti A. Vitamin C improves endothelium-dependent vasodilation by restoring nitric oxide activity in essential hypertension. *Circulation*. 1998 Jun 9;97(22):2222-9.
- Takazawa K, Tanaka N, Fujita M et al. Assessment of vasoactive agents and vascular aging by the second derivative of photoplethysmogram waveform. *Hypertension*. 1998 Aug;32(2):365-70.
- Tepel M, Echelmeyer M, Orië NN, Zidek W. Increased intracellular reactive oxygen species in patients with end-stage renal failure: effect of hemodialysis. *Kidney Int*. 2000 Aug;58(2):867-72.
- Tepel M, van der Giet M, Statz M, Jankowski J, Zidek W. The antioxidant acetylcysteine reduces cardiovascular events in patients with end-stage renal failure: a randomized, controlled trial. *Circulation*. 2003 Feb 25;107(7):992-5.
- Tepel M. Oxidative stress: does it play a role in the genesis of essential hypertension and hypertension of uraemia? *Nephrol Dial Transplant*. 2003 Aug;18(8):1439-42.
- Ting HH, Timimi FK, Haley EA, Roddy MA, Ganz P, Creager MA. Vitamin C improves endothelium-dependent vasodilation in forearm resistance vessels of humans with hypercholesterolemia. *Circulation*. 1997 Jun 17;95(12):2617-22.
- Tschöpe W, Koch M, Thomas B, Ritz E. Serum lipids predict cardiac death in diabetic patients on maintenance hemodialysis. Results of a prospective study. The German Study Group Diabetes and Uremia. *Nephron*. 1993;64(3):354-8.
- Wang AY, Woo J, Wang M et al. Association of inflammation and malnutrition with cardiac valve calcification in continuous ambulatory peritoneal dialysis patients. *J Am Soc Nephrol*. 2001 Sep;12(9):1927-36.
- White SL, Cass A, Atkins RC, Chadban SJ. Chronic kidney disease in the general population. *Adv Chronic Kidney Dis*. 2005 Jan;12(1):5-13.
- Yao Q, Axelsson J, Heimbürger O, Stenvinkel P, Lindholm B. Systemic inflammation in dialysis patients with end-stage renal disease: causes and consequences. *Minerva Urol Nefrol*. 2004 Sep;56(3):237-48.
- Zager PG, Nikolic J, Brown RH et al. "U" curve association of blood pressure and mortality in hemodialysis patients. Medical Directors of Dialysis Clinic, Inc. *Kidney Int*. 1998 Aug;54(2):561-9.

- Zarzecki M, Chudek J, Kukla M et al. Prevalence of anemia, calcium-phosphorus abnormalities and metabolic acidosis in different stages of chronic renal failure. *Pol Arch Med Wewn.* 2004 Oct;112(4):1211-9.
- Zimmermann J, Herrlinger S, Pruy A, Metzger T, Wanner C. Inflammation enhances cardiovascular risk and mortality in hemodialysis patients. *Kidney Int.* 1999 Feb;55(2):648-58.
- Zocalli C, Benedetto FA, Maas R et al. CREED Investigators. Asymmetric dimethylarginine, G reactive protein, and carotid intima-media thickness in end-stage renal disease. *J Am Soc Nephrol.* 2002 Feb;13(2):490-6.
- Zocalli C, Bode-Boger S, Mallamaci F et al. Plasma concentration of asymmetrical dimethylarginine and mortality in patients with end-stage renal disease: a prospective study. *Lancet.* 2001 Dec 22-29;358(9299):2113-7.