

9. Literaturverzeichnis

Achen M.G., Jeltsch M., Kukk E., Mäkinen T., Vitali A., Wilks A.F., Alitalo K., Stacker S.A. (1998).

Vascular endothelial growth factor D (VEGF-D) is a ligand for the tyrosine kinase VEGF receptor 2 (Flk1) and VEGF receptor 3 (Flt4).

Proc. Natl. Acad. Sci. USA **95**: 548-553

Aiello L.P., Avery R.L., Arrigg P.G., Keyt B.A., Jampel H.D., Sabera T., Shah S.T., Pasquale L.R., Thieme H., Iwamoto M.A., Park J.E., Nguyen H.V., Aiello L.M., Ferrara N., King G.L. (1994).

Vascular endothelial growth factor in ocular fluid of patients with diabetic retinopathy and other retinal disorders.

N. Engl. J. Med. **331**: 1480-1487

Algire G.H. (1943).

An adaptation of the transparent-chamber technique to the mouse.

J. Natl. Cancer Inst. USA **4**: 1-11

Algire G.H., Chalkley H.W. (1945).

Vascular reactions of normal and malignant tissues in vivo. I. Vascular reactions of mice to wounds and to normal and neoplastic transplants.

J. Natl. Cancer Inst. **6**: 73-85

Amselgruber W.M., Schäfer M., Sinowatz F. (1999).

Angiogenesis in the bovine corpus luteum: an immunocytochemical and ultrastructural study.

Anat. Histol. Embryol. **28**: 157-166

Andrade S.P., Fan T.P., Lewis G.P. (1987).

Quantitative in-vivo studies on angiogenesis in a rat sponge model.

Br. J. Exp. Pathol. **68**: 755-766

Arthur W.T., Vernon R.B., Sage E.H., Reed M.J. (1998).

Growth Factors reverse the impaired sprouting of microvessels from aged mice.

Microvasc. Res. **55**: 260-270

Asahara T., Bauters C., Zheng L.P., Satoshi T., Bunting S., Ferrara N., Symes J.F., Isner J.M. (1995).

Synergistic effect of vascular endothelial growth factor and basic fibroblast growth factor on angiogenesis in vivo.

Circulation **92**: II365-II371

Asahara T., Murohara T., Sullivan A., Silver M., van der Zee R., Li T., Witzenbichler B., Schatteman G., Isner J.M. (1997).

Isolation of putative progenitor endothelial cells for angiogenesis.

Science **275**: 964-967

Asahara T., Masuda H., Takahashi T., Kalka C., Pastore C., Silver M., Kearne M., Magner M., Isner J.M. (1999).

Bone marrow origin of endothelial progenitor cells responsible for postnatal vasculogenesis in physiological and pathological neovascularization.

Circ. Res. **85**: 221-228

Asahara T., Chen D., Takahashi T., Fujikawa K., Kearney M., Magner M., Yancopoulos G.D., Isner J.M. (1998).

Tie2 receptor ligands, angiopoietin-1 and angiopoietin-2 modulate VEGF-induced postnatal neovascularization.

Circ. Res. **83**: 233-240

Auerbach R., Kubai L., Knighton D., Folkman J. (1974).

A simple procedure for the long-term cultivation of chicken embryos.

Devl. Biol. **41**: 391-394

Auerbach R., Auerbach W., Polakowski I. (1991).

Assays for Angiogenesis: A Review.

Pharmac. Ther. **51**: 1-11

- Auerbach R., Plendl J., Barmak K. (1992).
Endothelial cell heterogeneity and differentiation. In: Angiogenesis in health and disease (M.E. Maragoudakis et al., eds.), 55-62. Plenum Press, New York.
- Auerbach R., Akhtar N., Lewis R. L., Shinnars B. L. (2000).
Angiogenesis assays: problems and pitfalls.
Cancer Metastasis Rev. **19**: 167-172
- Augustin H.G., Voss A.K., Pauli B.U. (1993).
Senescence of aortic endothelial cells in culture: effects of basic fibroblast growth factor expression on cell phenotype, migration, and proliferation.
J. Cell. Physiol. **157**: 279-288
- Augustin H.G., Braun K., Telemenakis I., Modlich U., Kuhn W. (1995).
Ovarian angiogenesis: Phenotypic characterization of endothelial cells in a physiological model of blood vessel growth and regression.
Am. J. Pathol. **147**: 339-351
- Ausprunk D.H., Knighton D.R., Folkman J. (1975).
Vascularization of normal and neoplastic tissues grafted to the chick chorioallantois. Role of host and preexisting graft blood vessels.
Am. J. Pathol. **79**: 597-628
- Ausprunk D.H., Folkman J. (1977).
Migration and proliferation of endothelial cells in preformed and newly formed blood vessels during tumor angiogenesis.
Microvasc. Res. **14**: 53-65
- Bacic M., Edward N.A., Merrill M.J. (1995).
Differential expression of vascular endothelial growth factor (vascular permeability factor) forms in rat tissues.
Growth Factors **12**: 11-15

Bagavandoss P., Wilks J.W. (1990).

Specific inhibition of endothelial cell proliferation by thrombospondin.

Biochem. Biophys. Res. Commun. **170**: 867-72

Bangsow T., Male D.A. (1999).

Polymerase-Kettenreaktion (PCR). In: Gentechnische Methoden (Gassen H.G. und Schrimpf G., Hrsg.), 333-346. Spektrum Akademischer Verlag, Heidelberg, Berlin.

Barleon B., Sozzani S., Zhou D., Weich H.A., Mantovani A., Marme D. (1996).

Migration of human monocytes in response to vascular endothelial growth factor (VEGF) is mediated via the VEGF receptor flt-1.

Blood **87**: 3336-3343

Barleon B., Siemeister G., Martiny-Baron G., Weindel K., Herzog C., Marme D. (1997).

Vascular endothelial growth factor up-regulates its receptor *fms*-like tyrosine kinase 1 (FLT-1) and a soluble variant of FLT-1 in human vascular endothelial cells.

Cancer Res. **57**: 5421-5425

Basilico C., Moscatelli D. (1992).

The FGF family of growth factors and oncogenes.

Adv. Cancer Res. **59**: 115-165

Baumann G., Stolar M.W., Amburn K., Barsano C.P., DeVries B.C. (1986).

A specific growth hormone-binding protein in human plasma: initial characterization.

J. Clin. Endocrinol. Metab. **62**: 134-141

Baumann G., Shaw M.A. (1990).

A second, lower affinity growth hormone-binding protein in human plasma.

J. Clin. Endocrinol. Metab. **70**: 680-686

Benjamin L.A., Keshet E. (1997).

Conditional switching of vascular endothelial growth factor (VEGF) expression in tumors: induction of endothelial cell shedding and regression of hemangio-blastoma-like vessels by VEGF withdrawal.

Proc. Natl. Acad. Sci. USA **94**: 8761-8766

Benjamin L.E., Hemo I., Keshet E. (1998).

A plasticity window for blood vessel remodelling is defined by pericyte coverage of the preformed endothelial network and is regulated by PDGF-B and VEGF.

Development **125**: 1591-1598

Breier G., Albrecht U., Sterrer S., Risau W. (1992).

Expression of vascular endothelial growth factor during embryonic angiogenesis and endothelial cell differentiation.

Development **114**: 521-532

Brown L.F., Berse B., Jackman R.W., Tognazzi K., Manseau E.J., Dvorak H.F., Senger D.R. (1993a).

Increased expression of vascular permeability factor (vascular endothelial growth factor) and its receptors in kidney and bladder carcinomas.

Am. J. Pathol. **143**: 1255-1262

Brown L.F., Berse B., Jackman R.W., Tognazzi K., Manseau E.J., Senger D.R., Dvorak H.F. (1993b).

Expression of vascular permeability factor (vascular endothelial growth factor) and its receptors in adenocarcinomas of the gastrointestinal tract.

Cancer Res. **53**: 4727-4735

Brown K.J., Haynes S.F., Bezos A., Maguire D.J., Ford M.D., Parish C.R. (1996).

A novel in vitro assay for human angiogenesis.

Lab. Invest. **75**: 539-555

Budde A. (1999).

Immunhistochemischer Nachweis der Rezeptoren für Wachstumshormon und Oxytocin im bovinen Ovar. Dissertation. Tierärztliche Fakultät der Ludwig-Maximilians-Universität München.

Burri P.H. (1992).

Intussusceptive microvascular growth, a new mechanism of capillary network formation.

EXS 62: 32-39

Bussolini F., Di Renzo M.F., Ziche M., Bocchietto E., Olivero M., Naldini L., Gaudino G., Tamagnone L., Coffer A., Comoglio P.M. (1992).

Hepatocyte growth factor is a potent angiogenic factor which stimulates endothelial cell motility and growth.

J. Cell Biol. **119:** 629-641

Cao Y., Ju W.-R., Rosin A., Cao Y. (1997).

Placenta growth factor: identification and characterization of a novel isoform generated by RNA alternative splicing.

Biochem. Biophys. Res. Commun. **235:** 493-498

Cao Y., Linden P., Farnebo J., Cao R., Eriksson A., Kumar V., Qi J.-H., Claesson-Welsh L., Alitalo K. (1998).

Vascular endothelial growth factor C induces angiogenesis in vivo.

Proc. Natl. Acad. Sci. USA **95:** 14389-14394

Carmeliet P., Ferreira V., Breier G., Pollefeyt S., Kieckens L., Gertsenstein M., Fahrig M., Vandenhoeck A., Harpal K., Eberhardt C., Declercq C., Pawling J., Moons L., Collen D., Risau W., Nagy A. (1996).

Abnormal blood vessel development and lethality in embryos lacking a single VEGF allele.

Nature **380:** 435-439

Carr D., Friesen H.G. (1976).

Growth hormone and insulin binding to human liver.

J. Clin Endocrinol. Metab. **42**: 484-493

Cavender J.L., Murdoch W.J. (1988).

Morphological studies of the microcirculatory system of periovulatory ovine follicles.

Biol. Reprod. **39**: 989-997

Charnock-Jones D.S., Sharkey A.M., Rajput-Williams J., Burch D., Schofield J.P.,

Fountain S.A., Boocock C.A., Smith S.K. (1993).

Identification and localization of alternately spliced mRNAs for vascular endothelial growth factor in human uterus and estrogen regulation in endometrial carcinoma cell lines.

Biol. Reprod. **48**: 1120-1128

Charnock-Jones D.S., Sharkey A.M., Boocock C.A., Ahmed A., Plevin R., Ferrara N.,

Smith S.K. (1994).

Vascular endothelial growth factor receptor localization and activation in human trophoblast and choriocarcinoma cells.

Biol. Reprod. **51**: 524-530

Chomczynski P., Sacchi N. (1987).

Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction.

Anal. Biochem. **162**: 156-159

Clark E.R., Clark E.L. (1932).

Observations on living preformed blood vessels as seen in a transparent chamber inserted into the rabbit's ear.

Am. J. Anat. **49**: 441-473

Cohen T., Gitay-Goren H., Sharon R., Shibuya M., Halaban R., Levi B., Neufeld G. (1995).

VEGF₁₂₁, a vascular endothelial growth factor (VEGF) isoform lacking heparin binding ability, requires cell surface heparan sulfates for efficient binding to the VEGF receptors of human melanoma cells.

J. Biol. Chem. **270**: 11322-11326

Colville-Nash P.R., Alam C.A.S., Appleton I., Brown J.R., Seed M.P., Willoughby D.A. (1995).

The pharmacological modulation of angiogenesis in chronic granulomatous inflammation.

J. Pharmacol. Exp. Ther. **274**: 1463-1472

Connolly D.T., Heuvelman D.M., Nelson R., Olander J.V., Eppley B.L., Delfino J.J., Siegel N.R., Leimgruber R.M., Feder J. (1989).

Tumor vascular permeability factor stimulates endothelial cell growth and angiogenesis.

J. Clin. Invest. **84**: 1470-1478

Davis S., Aldrich T.H., Jones P.F., Acherson A., Compton D.L., Jain F., Ryan T.E., Bruno J., Radziejewski C., Maisonpierre P.C., Yancopoulos G.D. (1996).

Isolation of angiopoietin-1, a ligand for the Tie-2 receptor, by secretion-trap expression cloning.

Cell **87**: 1161-1169

Davis-Smyth T., Chen H., Park J., Presta L.G., Ferrara N. (1996).

The second immunoglobulin-like domain of the VEGF tyrosine kinase receptor Flt-1 determines ligand binding and may initiate a signal transduction cascade.

EMBO J. **15**: 4919-4927

De Vries C., Escobedo J.A., Ueno H., Houck K., Ferrara N., Williams L.T. (1992).

The fms-like tyrosine kinase, a receptor for vascular endothelial growth factor.

Science **255**: 989-991

Dejana E. (1996).

Endothelial adherens junction: Implications in the control of vascular permeability and angiogenesis.

J. Clin. Invest. **98**: 1948-1953

DeLisser H.M., Newman P.J., Albelda S.M. (1994).

Molecular and functional aspects of PECAM-1/CD31.

Immunology Today **15**: 490-495

Dellian M., Witwer B.P., Salehi H.A., Yuan F., Jain R.K. (1996).

Quantitation and physiological characterization of angiogenic vessels in mice: effect of basic fibroblast growth factor, vascular endothelial growth factor/vascular permeability factor, and host microenvironment.

Am. J. Pathol. **149**: 59-71

Denekamp J. (1984).

Vascular endothelium as the vulnerable element in tumours.

Acta Radiol. Oncol. **4**: 217-225

Dibbens J.A., Miller D.L., Damert A., Risau W., Vadas M.A., Goodall G.J. (1999).

Hypoxic regulation of vascular endothelial growth factor mRNA stability requires the cooperation of multiple RNA elements.

Mol. Biol. Cell **10**: 907-919

Dilley R.J., Schwartz S.M. (1989).

Vascular remodeling in the growth hormone transgenic mouse.

Circ. Res. **65**:1233-1240

DiSalvo J., Bayne M.L., Conn G, Kwok P.W., Trivedi P.G., Soderman D.D., Palisi T.M., Sullivan K.A., Thomas K.A. (1995).

Purification and characterization of a naturally occurring vascular endothelial growth factor - placenta growth factor heterodimer.

J. Biol. Chem. **270**: 7717-7723

Djonov V., Schmid M., Tschanz S.A., Burri P.H. (2000).
Intussusceptive angiogenesis: its role in embryonic vascular network formation.
Circ. Res. **86**: 286-292

Djonov V., Andres A.C., Ziemiecki A. (2001).
Vascular remodelling during the normal and malignant life cycle of the mammary gland.
Microsc. Res. Tech. **52**: 182-189

Dor Y., Rinnat P., Keshet E. (2001).
Vascular endothelial growth factor and vascular adjustments to perturbations in oxygen homeostasis.
Am. J. Physiol. Cell Physiol. **280**: C1367-C1374

Drake C.J., Little C.D. (1995).
Exogenous vascular endothelial growth factor induces malformed and hyperperfused vessels during embryonic neovascularization.
Proc. Natl. Acad. Sci. **92**: 7657-7661

Dvorak H.F., Harvey V.S., Estrella P., Brown L.F., McDonagh J., Dvorak A.M. (1987).
Fibrin containing gels induce angiogenesis. Implications for tumor stroma generation and wound healing.
Lab. Invest. **57**: 673-686

Dvorak H.F., Brown L.F., Detmar M., Dvorak A.M. (1995).
Vascular permeability factor/vascular endothelial growth factor, microvascular hyperpermeability, and angiogenesis.
Am. J. Pathol. **146**: 1029-1039

Ergün S., Kilic N., Fiedler I.W., Mukhopadhyay A.K. (1997).
Vascular endothelial growth factor and its receptors in normal human testicular tissue.
Mol. Cell Endocrinol. **131**: 9-20

Espey L.L. (1980).

Ovulation as an inflammatory reaction - a hypothesis.

Biol. Reprod. **22**: 73-106

Esser S., Wolburg K., Wolburg H., Breier G., Kurzchalia T., Risau W. (1998).

Vascular endothelial growth factor induces endothelial fenestrations in vitro.

J. Cell Biol. **140**: 947-959

Evans H.M., Long J.A. (1921).

The effect of the anterior lobe administered intraperitoneally upon growth, maturity, and oestrus cycles of the rat.

Anat Rec. **21**: 62

Fajardo L.F., Kowalski J., Kwan H.H., Prionas S.D., Allison A.C. (1988).

The disc angiogenesis system.

Lab. Invest. **58**: 718-724

Fajardo L.F., Kwan H.H., Kowalski J., Prionas S.D., Allison A.C. (1992).

Dual role of tumor necrosis factor-alpha in angiogenesis.

Am. J. Pathol. **140**: 539-544

Fawcett D.W. (1977).

Temporäre Differenzierungen und Oberflächenaktivität. In: Atlas zur Elektronenmikroskopie der Zelle (Fawcett D.W., edt.), 394-414. Urban & Schwarzb erg, München, Berlin, Wien

Ferrara N., Henzel W.J. (1989).

Pituitary follicular cells secrete a novel heparin-binding growth factor specific for vascular endothelial cells.

Biochem. Biophys. Res. Commun. **161**: 851-858

Ferrara N., Carver-Moore K., Chen H., Dowd M., Lu L., O'Shea K.S., Powell-Braxton L., Hillan K.J., Moore M.W. (1996).

Heterozygous embryonic lethality induced by targeted inactivation of the VEGF gene.
Nature **380**: 439-442

Ferrara N., Chen H., Davis-Smyth T., Gerber H.-P., Nguyen T.-N., Peers D., Chisholm V., Hillian K.J., Schwall R.H. (1998).

Vascular endothelial growth factor is essential for corpus luteum angiogenesis.
Nat. Med. **4**: 336-340

Findlay J.K. (1986).

Angiogenesis in reproductive tissues.

J. Endocrinol. **111**: 357-366

Finkenzeller G., Marme D., Weich H.A., Hug H. (1992).

Platelet derived growth factor-induced transcription of the vascular endothelial growth factor gene is mediated by protein kinase C.

Cancer Res. **52**: 4821-4823

Foidart J.M., Bere E.W. Jr., Yaar M., Rennard S.I., Gullino M., Martin G.R., Katz S.I. (1980).

Distribution and immunoelectron microscopic localization of laminin, a noncollagenous basement membrane glycoprotein.

Lab. Invest. **42**: 336-342

Folkman J., Cole P., Zimmerman S. (1966).

Tumor behavior in isolated perfused organs.

Ann Surg. **164**: 491-502

Folkman J. (1971).

Tumor angiogenesis: therapeutic implications.

N. Engl. J. Med. **285**: 1182-1186

Folkman J., Merler E., Abernathy C., Williams G. (1971).
Isolation of a tumor factor responsible for angiogenesis.
J. Exp. Med. **133**: 275-288

Folkman J., Haudenschild C.C., Zetter B.R. (1979).
Long-term culture of capillary endothelial cells.
Proc. Natl. Acad. Sci. USA **76**: 5217-5221

Folkman J., Haudenschild C. (1980).
Angiogenesis by capillary endothelial cells in culture.
Trans. Ophthal. Soc. U.K. **100**: 346-353

Folkman J. (1984).
Angiogenesis. In: Biology of endothelial cells (Jaffe E.A., Hrsg.), 412-428. Martinus Nijhoff Publishers, Boston, The Hague, Dordrecht.

Fong G.H, Rossant J., Gertsenstein M., Breitman M.L. (1995).
Role of the Flt-1 receptor tyrosine kinase in regulating the assembly of vascular endothelium.
Nature **376**: 66-70

Frank S., Hübner G., Breier G., Longaker M.T., Greenhalgh D.G., Werner S. (1995).
Regulation of vascular endothelial growth factor expression in cultured keratinocytes - implication for normal and impaired wound healing.
J. Biol. Chem. **270**: 12607-12613

Frater-Schröder M., Risau W., Hallmann R., Gautschi R., Böhlen P. (1987).
Tumor necrosis factor type α , a potent inhibitor of endothelial cell growth *in vitro*, is angiogenic *in vivo*.
Proc. Natl. Acad. Sci. USA **84**: 5277-5281

Fuchs-Schönleber T. (1999).
Etablierung eines *in vitro*-Modells des bovinen Corpus luteum. Dissertation.
Tierärztliche Fakultät der Ludwig-Maximilians-Universität München.

Galland F., Karamysheva A., Mattei M.-G., Rosnet O., Marchetto S., Birnbaum D. (1992).

Chromosomal localization of FLT4, a novel receptor-type tyrosine kinase gene.
Genomics **13**: 475-478

Gaudric A., N'guyen T., Moenner M., Glacet-Bernard A., Barritault D. (1992).
Quantification of angiogenesis due to basic fibroblast growth factor in a modified rabbit corneal model.

Ophthalmic. Res. **24**: 181-188

Gerber H.-P., Condorelli F., Park J., Ferrara N. (1997).
Differential transcriptional regulation of the two vascular endothelial growth factor receptor genes. Flt-1, but not flk-1/KDR, is up-regulated by hypoxia.
J. Biol. Chem. **272**: 23659-23667

Gimbrone M.A. Jr., Cotran R.S., Folkman J. (1973).
Endothelial regeneration: studies with human endothelial cells in culture.
Ser. Haematol. **6**: 453-455

Gimbrone M.A., Cotran R.S., Leapman S.B., Folkman J. (1974).
Tumor growth and neovascularization: an experimental model using the rabbit cornea.
J. Natl. Cancer Inst. USA **52**: 413-419

Gitay-Goren H., Soker S., Vlodavsky I., Neufeld G. (1992).
The binding of vascular endothelial growth factor to its receptors is dependent on cell surface-associated heparin-like molecules.
J. Biol. Chem. **267**: 6093-6098

Gitay-Goren H., Halaban R., Neufeld G. (1993).
Human melanoma cells but not normal melanocytes express vascular endothelial growth factor receptors.
Biochem. Biophys. Res. Commun. **190**: 702-709

Gluzman-Poltorak Z, Cohen T, Herzog Y, Neufeld G. (2000).

Neuropilin-2 and neuropilin-1 are receptors for the 165-amino acid form of vascular endothelial growth factor (VEGF) and of placenta growth factor-2, but only neuropilin-2 functions as a receptor for the 145-amino acid form of VEGF.

J. Biol. Chem. **275**: 18040-18045

Gnarra J.R., Zhou S., Merrill M.J., Wagner J.R., Krumm A., Papavassiliou E., Oldfield E.H., Klausner R.D., Linehan W.M. (1996).

Post-transcriptional regulation of vascular endothelial growth factor mRNA by the product of the *VHL* tumor suppressor gene.

Proc. Natl. Acad. Sci. USA **93**: 10589-10594

Goad D.L., Rubin J., Wang H., Tashjian A.H., Patterson C. (1996).

Enhanced expression of vascular endothelial growth factor in human SaOS-2 osteoblast-like cells and murine osteoblasts induced by insulin-like growth factor I. Endocrinology **137**: 2262-2268

Goede V., Schmidt T., Kimmina S., Kozian D., Augustin H.G. (1998).

Analysis of blood vessel maturation processes during cyclic ovarian angiogenesis. Lab. Invest. **78**: 1385-1393

Gonzalez-Crussi F. (1971).

Vasculogenesis in the chick embryo. An ultrastructural study.

Am. J. Anat. **130**: 441-446

Gordon S.R., Essner E. (1985).

Plasma membrane-associated vesicles in retinal capillaries of the rat.

Am. J. Anat. **174**: 161-172

Gospodarowicz D., Abraham J.A., Schilling J. (1989).

Isolation and characterization of a vascular endothelial cell mitogen produced by pituitary-derived folliculo stellate cells.

Proc. Natl. Acad. Sci. **86**: 7311-7315

Gould J., Aramburo C., Capdevielle M., Scanes C.G. (1995).
Angiogenic activity of anterior pituitary tissue and growth hormone on the chick embryo chorioallantoic membrane: a novel action of GH.
Life Sci. **56**: 587-594

Grant M.B., Mames R.N., Fitzgerald C., Ellis E.A., Caballero S., Chegini N., Guy J. (1993).

Insulin-like growth factor I as an angiogenic agent.
Ann. N.Y. Acad. Sci. **692**: 230-240

Grant D.S., Kleinman H.K. (1997).

Regulation of capillary formation by laminin and other components of the extracellular matrix.
EXS **79**: 317-333

Greenblatt M., Shubik P. (1968).

Tumor angiogenesis: Transfilter diffusion studies in the hamster by the transparent chamber technique.

J. Natl. Cancer Inst. USA **41**: 111-124

Greene H.S.N. (1941).

Heterologous transplantation of mammalian tumors.
J. Exper. Med. **73**: 461-468

Grimmond S., Lagercrantz J., Drinkwater C., Silins G., Townson S., Pollock P., Gotley D., Carson E., Rakar S., Nordenskjöld M., Ward L., Hayward N., Weber G. (1996).

Cloning and characterization of a novel human gene related to vascular endothelial growth factor.
Genome Res. **6**: 124-131

Grosskreutz C.L., Anand-Apte B., Duplaa C., Quinn T.P., Terman B.I., Zetter B., D'Amore P.A. (1999).

Vascular endothelial growth factor-induced migration of vascular smooth muscle cells *in vitro*.

Microvasc. Res. **58**: 128-136

Guidi A.J., Abu-Jawdeh G., Berse B., Jackman R.W., Tognazzi K., Dvorak H.F., Brown L.F. (1995).

Vascular permeability factor (vascular endothelial growth factor) expression and angiogenesis in cervical neoplasia.

J. Natl. Cancer Inst. USA **87**: 1237-1245

Hauser S.D., McGrath M.F., Collier R.J., Krivi G.G. (1990).

Cloning and *in vivo* expression of bovine growth hormone receptor mRNA.

Mol. Cell Endocrinol. **72**: 187-200

Hauser S., Weich H. (1993).

A heparin-binding form of placenta growth factor (PIGF-2) is expressed in human umbilical vein endothelial cells and in placenta.

Growth Factors **9**: 259-268

Hees H. (2000).

Kreislaufsystem. In: Histologie (H. Hees, F. Sinowitz, Hrsg.), 191-204. Deutscher Ärzte-Verlag, Köln.

Hees H. (2000).

Weiblicher Geschlechtsapparat. In: Histologie (H. Hees, F. Sinowitz, Hrsg.), 303-321. Deutscher Ärzte-Verlag, Köln.

Hellström A., Svensson E., Carlsson B., Niklasson A., Albertsson-Wikland K. (1999).

Reduced retinal vascularization in children with growth hormone deficiency.

J. Clin. Endocrinol. Metab. **84**: 795-798

Henkind P. (1978).
Ocular neovascularization.
Am. J. Ophthal. **85**: 287-301

Herrmann G., Spanel Borowski K. (1998).
A sparsely vascularised zone in the cortex of the bovine ovary.
Anat. Histol. Embryol. **27**: 143-146

Hirschi K.K., D'Amore P.A. (1996).
Pericytes in the microvasculature.
Cardiovasc. Res. **32**: 687-698

Hirschi K.K., Rohovsky S.A., Beck L.H., Smith S.R., D'Amore P.A. (1999).
Endothelial cells modulate the proliferation of mural cell precursors via platelet-derived growth factor-BB and heterotypic cell contact.
Circ. Res. **84**: 298-305

Houck K.A., Ferrara N., Winer J., Cachianes G., Li B., Leung D.W. (1991).
The vascular endothelial growth factor family: Identification of a fourth molecular species and characterization of alternative splicing of RNA.
Mol. Endocrinol. **5**: 1806-1814

Houck K.A., Leung D.W., Rowland A.M., Winer J., Ferrara N. (1992).
Dual regulation of vascular endothelial growth factor bioavailability by genetic and proteolytic mechanisms.
J. Biol. Chem. **267**: 26031-26037

Hughes S., Chan-Ling T. (2000).
Roles of endothelial cell migration and apoptosis in vascular remodeling during development of the central nervous system.
Microcirculation **7**: 317-333

Hull K.L., Harvey S. (2001).

Growth hormone: roles in female reproduction.

J. Endocrinol. **168**: 1-23

Ikeo S., Yamauchi K., Shigematsu S., Nakajima K., Aizawa T., Hashizume K. (2001).

Differential effects of growth hormone and insulin-like growth factor I on human endothelial cell migration.

Am. J. Physiol. Cell Physiol. **280**: C1255-1261

Imaizumi T., Itaya H., Nasu S., Yoshida H., Matsubara Y., Fujimoto K., Matsumiya T., Kimura H., Satoh K. (2000).

Expression of vascular endothelial growth factor in human umbilical vein endothelial cells stimulated with interleukin-1alpha: an autocrine regulation of angiogenesis and inflammatory reactions.

Thromb. Haemostasis **83**: 949-955

Iwahana M., Nakayama Y., Tanaka N.G., Goryo M., Okada K. (1996).

Quantification of tumour-induced angiogenesis by image analysis.

Int. J. Exp. Pathol. **77**: 109-114

Jaffe E.A., Nachman R.L., Becker C.G., Minick C.R. (1973).

Culture of human endothelial cells derived from umbilical veins.

J. Clin. Invest. **52**: 2745-2756

Jain R., Schlenger K., Höckel M., Yuan F. (1997).

Quantitative angiogenesis assay: Progress and problems.

Nat. Med. **3**: 1203-1208

Jakeman L.B., Winer J., Bennett G.L., Altar C.A., Ferrara N. (1992).

Binding sites for vascular endothelial growth factor are localized on endothelial cells in adult rat tissues.

J. Clin. Invest. **89**: 244-253

Joukov V., Pajusola K., Kaipainen A., Chilov D., Lahtinen I., Kukk E., Saksela O., Kalkkinen N., Alitalo K. (1996).

A novel vascular endothelial growth factor, VEGF-C, is a ligand for the Flt4 (VEGFR-3) and KDR (VEGFR-2) receptor tyrosine kinases.

EMBO J. **15**: 290-298

Joukov V., Sorsa T., Kumar V., Jeltsch M., Claesson-Welsh L., Cao Y., Saksela O., Kalkkinen N., Alitalo K. (1997).

Proteolytic processing regulates receptor specificity and activity of VEGF-C.

EMBO J. **16**: 3898-3911

Kaipainen A., Korhonen J., Mustonen T., Van Hinsbergh V.W.M., Fang G.-H., Dumont D., Breitman M., Alitalo K. (1995).

Expression of the fms-like tyrosine kinase 4 gene becomes restricted to lymphatic endothelium during development.

Proc. Natl. Acad. Sci. USA **92**: 3566-3570

Kaiser G.G., Sinowatz F., Palma G.A. (2001).

Effects of growth hormone on female reproductive organs.

Anat. Histol. Embryol. **30**: 265-271

Kalashnik L., Bridgeman C.J., King A.R., Francis S.E., Mikhalovsky S., Wallis C., Denyer S.P., Crossman D., Faragher R.G. (2000)

A cell kinetic analysis of human umbilical vein endothelial cells.

Mech. Ageing Dev. **120**: 23-32.

Katoh O., Tauchi H., Kawaishi K., Kimura A., Satow Y. (1995).

Expression of the vascular endothelial growth factor (VEGF) receptor gene, *KDR*, in hematopoietic cells and inhibitory effect of VEGF on apoptotic cell death caused by ionizing radiation.

Cancer Res. **55**: 5687-5692

Keck P.J., Hauser S.D., Krivi G., Sanzo K., Warren T., Feder J., Connolly D.T. (1989).
Vascular permeability factor, an endothelial cell mitogen related to PDGF.
Science **246**: 1309-1312

Kelly P.A., Djiane J., Banville D., Ali S., Edery M., Rozakis M. (1991).
The growth hormone/prolactin receptor gene family.
Oxf. Surv. Eukaryot. Genes **7**: 29-50

Kendall R.L., Thomas K.A. (1993).
Inhibition of vascular endothelial cell growth factor activity by an endogenously
encoded soluble receptor.
Proc. Natl. Acad. Sci. USA **90**: 10705-10709

Kessel J., Fabian B. (1985).
Graded morphogenetic patterns during the development of the extraembryonic
blood system and coelom of the chick blastoderm: A scanning electron microscope
and light microscope study.
Am. J. Anat. **173**: 99-112

Keyt B.A., Nguyen H.V., Berleau L.T., Duarte C.M., Park J., Chen H., Ferrara N.
(1996).
Identification of vascular endothelial growth factor determinants for binding KDR
and FLT-1 receptors. Generation of receptor-selective VEGF variants by site-
directed mutagenesis.
J. Biol. Chem. **271**: 5638-5646

Kieser A., Weich H., Brandner G., Marme D., Kolch W. (1994).
Mutant p53 potentiates protein kinase C induction of vascular endothelial growth
factor expression.
Oncogene **9**: 963-969

- Kim K.J., Li B., Winer J., Armanini M., Gillett N., Phillips H.S., Ferrara N. (1993). Inhibition of vascular endothelial growth factor-induced angiogenesis suppresses tumor growth *in vivo*. *Nature* **362**: 841-844
- Klagsbrun M., D'Amore P.A. (1991). Regulators of angiogenesis. *Annu. Rev. Physiol.* **53**: 217-239
- Kleinman H.K., Weeks B.S., Schnaper H.W., Kibbey M.C., Yamamura K., Grant D.S. (1993). The laminins: a family of basement membrane glycoproteins important in cell differentiation and tumor metastases. *Vit. Hormon.* **47**: 161-186
- Koblizek T.I., Weiss C., Yancopoulos G.D., Deutsch U., Risau W. (1998). Angiopoietin-1 induces sprouting angiogenesis in vitro. *Curr. Biol.* **8**: 529-532
- Koch A.E., Harlow L.A., Haines G.K., Amento E.P., Unemori E.N., Wong W.L., Pope R.M., Ferrara N. (1994). Vascular endothelial growth factor: a cytokine modulating endothelial function in rheumatoid arthritis. *J. Immunol.* **152**: 4149-4156
- Kölle S., Sinowitz F., Boie G., Lincoln D. (1998). Developmental changes in the expression of the growth hormone receptor messenger ribonucleic acid and protein in the bovine ovary. *Biol. Reprod.* **59**: 836-842
- König H.E., Amselgruber W., Rüsse I. (1988). Zur Mikrozirkulation in Follikeln und Corpora lutea des Rinderovars - eine korrosionsanatomische Studie. *Tierärztl. Prax.* **16**: 25-31

Kopchick J.J., Andry J.M. (2000).

Growth hormone (GH), GH receptor and signal transduction.

Mol. Genet. Metab. **71**: 293-314

Kramer R.H., Bensch K.G., Davison P.M., Karsek M.A. (1984).

Basal lamina formation by cultured microvascular endothelial cells.

J. Cell Biol. **99**: 692-698

Kukk E., Lymboussaki A., Taira S., Kaipainen A., Jeltsch M., Joukov V., Alitalo K. (1996).

VEGF-C receptor binding and pattern of expression with VEGFR-3 suggests a role in lymphatic vascular development.

Development **122**: 3829-3837

Ladoux A., Frelin C. (1993).

Expression of vascular endothelial growth factor by cultured endothelial cells from brain microvessels.

Biochem. Biophys. Res. Commun. **194**: 799-803

Langer R., Siegel R., Brown L., Leong K., Kost J., Edelman E. (1985).

Controlled release and magnetically modulated systems for macromolecular drugs.

Ann. N.Y. Acad. Sci. **446**: 1-13

Lauer G., Sollberg S., Cole M., Flamme I., Stürzebecher J., Mann K., Krieg T., Eming S.A. (2000).

Expression and proteolysis of vascular endothelial growth factor is increased in chronic wounds.

J. Invest. Dermatol. **115**: 12-18

Lebovic D.I., Mueller M.D., Tylor R.N. (1999).

Vascular endothelial growth factor in reproductive biology.

Curr. Opinion Obst. Gyn. **11**: 255-260

- Lee J., Gray A., Yuan J., Luoh S.M., Avraham H., Wood W.I. (1996).
Vascular endothelial growth factor-related protein: A ligand and specific activator of the tyrosine kinase receptor Flt4.
Proc. Natl. Acad. Sci. USA **93**: 1988-1992
- Leibovich S.J., Polverini P.J., Shepard H.M., Wiseman D.M., Shively V., Nuseir N. (1987).
Macrophage-induced angiogenesis is mediated by tumor necrosis factor- α .
Nature **329**: 630-632
- Leiser R. (1990).
Weibliche Geschlechtsorgane: In: Zytologie, Histologie und mikroskopische Anatomie der Haussäugetiere (Mosimann W., Kohler T., Hrsg.), 232-248. Paul Parey, Berlin, Hamburg.
- Leiser R. (1999).
Weibliche Geschlechtsorgane. In: Lehrbuch der Anatomie der Haustiere; Band II: Eingeweide (Nickel R., Schummer A., Seiferle E., Hrsg.), 393-439. Blackwell Wissenschafts-Verlag, Berlin, Wien.
- Leung D.W., Spencer S.A., Cachianes G., Hammonds R.G., Collins C., Henzel W.J., Barnard R., Waters M.J., Wood W.I. (1987).
Growth hormone receptor and serum binding protein: purification, cloning and expression.
Nature **330**: 537-543
- Leung D.W., Cachianes G., Kuang W.-J., Goeddel D.V., Ferrara N. (1989).
Vascular endothelial growth factor is a secreted angiogenic mitogen.
Science **246**: 1306-1309
- Levasseur J.E., Wei E.P., Raper A.J., Kontos H.A., Patterson J.L. (1975).
Detailed description of a cranial window technique for acute and chronic experiments.
Stroke **6**: 308-317

Li C.H., Evans H.M. (1944).

The isolation of pituitary growth hormone.

Science **99**: 183-184

Li J., Brown L.F., Hibberd M.G., Grossman J.D., Morgan J.P., Simons M. (1996).

VEGF, flk-1, and flt-1 expression in a rat myocardial infarction model of angiogenesis.

Am. J. Physiol. **270**: H1803-H1811

Liebich H.-B. (1999).

Kreislaufsystem (Systema cardiovasculare et lymphovasculare). In: Funktionelle Histologie der Haussäugetiere (Liebich H.-G., Hrsg.), 117-131. F. K. Schattauer Verlagsgesellschaft mbH, Stuttgart.

Liebich H.-B. (1999).

Weibliche Geschlechtsorgane (Organa genitalia feminina). In: Funktionelle Histologie der Haussäugetiere (Liebich H.-G., Hrsg.), 284-302. F. K. Schattauer Verlagsgesellschaft mbH, Stuttgart.

Lindenbaum E.S., Langer N., Beach D. (1991).

Isolation and culture of human decidual capillary endothelial cells in serum-free medium supplemented with human uterine angiogenic factor.

Acta Anat. **140**: 273-279

Lindl T., Bauer J. (1987).

Räumliche und apparative Voraussetzungen und Sicherheitsvorschriften. In: Zell- und Gewebekultur (Lindl T., Bauer J., Hrsg.), 1-10. Gustav Fischer Verlag, Stuttgart, New York.

Lyttle D.J., Fraser K.M., Fleming S.B., Mercer A.A., Robinson A.J. (1994).

Homologs of vascular endothelial growth factor are encoded by the poxvirus orf virus.

J. Virol. **68**: 84-92

Madri J.A., Pratt B.M. (1986).

Endothelial cell-matrix interactions: in vitro models of angiogenesis.

J. Histochem. Cytochem. **34**: 85-91

Madri J. A., Pratt B.M., Tucker A.M. (1988).

Phenotypic modulation of endothelial cells by transforming growth factor- β depends on the composition and organization of the extracellular matrix.

J. Cell Biol. **106**: 1375-1384

Maglione D., Guerriero V., Viglietto G., Delli-Bovi P., Persico M.G. (1991).

Isolation of a human placenta cDNA coding for a protein related to the vascular permeability factor.

Proc. Natl. Acad. Sci. USA **88**: 9267-9271

Maisonpierre P.C., Suri C., Jones P.F., Bartunkova S., Wiegand S.J., Radziejewski C., Compton D., McClain J., Aldich TH., Papadopoulos N., Daly T.J., Davis S., Sato T.N., Yancopoulos G.D. (1997).

Angiopoietin-2, a natural antagonist for Tie2 that disrupts in vivo angiogenesis.

Science **277**: 55-60

Makinen T., Olofsson B., Karpanen T., Hellman U., Soker S., Klagsbrun M., Eriksson U., Alitalo K. (1999).

Differential binding of vascular endothelial growth factor B splice and proteolytic isoforms to neuropilin-1.

J. Biol. Chem. **274**: 21217-21222

Mandriota S.J., Menoud P.-A., Pepper M.S. (1996).

Transforming growth factor β 1 down-regulates vascular endothelial growth factor receptor 2/flk-1 expression in vascular endothelial cells.

J. Biol. Chem. **271**: 11500-11505

Marconcini L., Marchio S., Morbidelli L., Cartocci E., Albini A., Ziche M., Bussolino F., Oliviero S. (1999).

c-fos-induced growth factor/vascular endothelial growth factor D induces angiogenesis *in vivo* and *in vitro*.

Proc. Natl. Acad. Sci. USA **96**: 9671-9676

Matsumoto K., Ohi H., Kanmatsuse K. (1997).

Interleukin 10 and interleukin 13 synergize to inhibit vascular permeability factor release by peripheral blood mononuclear cells from patients with lipoid nephrosis.

Nephron **77**: 212-218

Matsumoto M., Nishinakagawa H., Kurohmaru M., Hayashi Y., Otsuka J. (1992).

Pregnancy and lactation affect the microvasculature of the mammary gland in mice.
J. Vet. Med. Sci. **54**: 937-943

Matthews W., Jordan C.T., Gavin M., Jenkins N.A., Copeland N.G., Lemischka I.R. (1991).

A receptor tyrosine kinase cDNA isolated from a population of enriched primitive hematopoietic cells and exhibiting close genetic linkage to *c-kit*.

Proc. Natl. Acad. Sci. USA **88**: 9026-9030

Meyer G.T., Matthias L.J., Noack L., Vadas M.A., Gamble J.R. (1997).

Lumen formation during angiogenesis *in vitro* involves phagocytic activity, formation and secretion of vacuoles, cell death and capillary tube remodelling by different populations of endothelial cells.

Anat. Rec. **249**: 327-340

Meyer M., Clauss M., Lepple-Wienhues A., Waltenberger J., Augustin H.G., Ziche M., Lanz C., Buttner M., Rziha H.-J., Dehio C. (1999).

A novel vascular endothelial growth factor encoded by Orf virus, VEGF-E, mediates angiogenesis via signalling through VEGFR-2 (KDR) but not VEGFR-1 (Flt-1) receptor tyrosine kinases.

EMBO J. **18**: 363-374

Migdal M., Huppertz B., Tessler S., Comforti A., Shibuya M., Reich R., Baumann H., Neufeld G. (1998).

Neuropilin-1 is a placenta growth factor-2 receptor.

J. Biol. Chem. **273**: 22272-22278

Millauer B., Wizigmann-Voos S., Schnürch H., Martinez R., Moller N.P.H., Risau W., Ullrich A. (1993).

High affinity VEGF binding and developmental expression suggest Flk-1 as a major regulator of vasculogenesis and angiogenesis.

Cell **72**: 835-846

Modlich U., Kaup F.-J., Augustin H. G. (1996).

Cyclic angiogenesis and blood vessel regression in the ovary: Blood vessel regression during luteolysis involves endothelial cell detachment and vessel occlusion.

Lab. Invest. **74**: 771-780

Monacci W.T., Merrill M.J., Oldfield E.H. (1993).

Expression of vascular permeability factor/vascular endothelial growth factor in normal rat tissues.

Am. J. Physiol. **264**: C995-C1002

Montesano R., Pepper M.S., Vassalli J.D., Orci L. (1992).

Modulation of angiogenesis in vitro.

EXS. **61**: 129-136

Moses M.A., Klagsbrun M., Shing Y. (1995).

The role of growth factors in vascular cell developmental differentiation.

Int. Rev. Cytol. **161**: 1-48

Nakao-Hayashi J., Ito H., Kanayasu T., Morita I., Murota S. (1992).

Stimulatory effects of insulin and insulin-like growth factor I on migration and tube formation by vascular endothelial cells.

Atherosclerosis **92**: 141-149

Namiki A., Brogi E., Kearney M., Kim E.A., Wu T., Couffinhal T., Varticovski L., Isner J.M. (1995).

Hypoxia induces vascular endothelial growth factor in cultured human endothelial cells.

J. Biol. Chem. **270**: 31189-31195

Nehls V., Denzer K., Drenckhahn D. (1992).

Pericyte involvement in capillary sprouting during angiogenesis in situ.

Cell Tissue Res. **270**: 469-474

Nguyen M., Shing Y., Folkman J. (1994).

Quantitation of angiogenesis and antiangiogenesis in the chick embryo chorioallantoic membrane.

Microvasc. Res. **47**: 31-40

Nickel R., Schummer A., Seiferle E. (1996).

Vena cava caudalis. In: Lehrbuch der Anatomie der Haustiere; Band III: Kreislaufsystem, Haut und Hautorgane (Nickel R., Schummer A., Seiferle E., Hrsg.), 240-275. Blackwell Wissenschafts-Verlag, Berlin, Wien.

Nicosia R.F., Tchao R., Leighton J. (1982).

Histotypic angiogenesis in vitro: light microscopic, ultrastructural, and radioautographic studies.

In Vitro **18**: 538-549

Nicosia R.F., Ottinetti A. (1990).

Growth of microvessels in serum-free matrix culture of rat aorta.

Lab. Invest. **63**: 115-122

Nicosia R.F., Nicosia S.V., Smith M. (1994).

Vascular endothelial growth factor, platelet-derived growth factor, and insulin-like growth factor-1 promote rat aortic angiogenesis in vitro.

Am. J. Pathol. **145**: 1023-1029

Nicosia R.F., Villaschi S. (1999).

Autoregulation of angiogenesis by cells of the vascular wall.

Int. Rev. Cytol. **185**: 1-43

Nomura M., Yamanashi S.I., Harada S.I., Hayashi Y., Yamashima T., Yamashita J., Yamamoto H. (1995).

Possible participation of autocrine and paracrine vascular endothelial growth factor in hypoxia-induced proliferation of endothelial cells and pericytes.

J. Biol. Chem. **270**: 28316-28324

Norrby K., Jakobsson A., Sörbo J. (1986).

Mast-cell-mediated angiogenesis: a novel experimental model using the rat mesentery.

Virchows Arch. B Cell. Pathol. Incl. Mol. Pathol. **52**: 195-206

Norrby K., Jakobsson A., Sörbo J. (1996).

Vascular endothelial growth factor and de novo mammalian angiogenesis.

Microvasc. Res. **51**: 153-163

Öberg C., Waltenberger J., Claesson-Welsh L., Welsh M. (1994).

Expression of protein tyrosine kinases in islet cells: possible role of the Flk-1 receptor for β -cell maturation from duct cells.

Growth Factors **10**: 115-126

Oh S.J., Jeltsch M.M., Birkenhage R., McCarthy J.E.G., Weich H.A., Christ B., Alitalo K., Wilting J. (1997).

VEGF and VEGF-C: specific induction of angiogenesis and lymphangiogenesis in the differentiated avian chorioallantoic membrane.

Dev. Biol. **188**: 96-109

Olofsson B., Pajusola K., Kaipainen A., von Euler G., Joukov V., Saksela O., Orpana A., Pettersson R.F., Alitalo K., Eriksson U. (1996a).

Vascular endothelial growth factor B, a novel growth factor for endothelial cells.

Proc. Natl. Acad. Sci. USA **93**: 2576-2581

Olofsson B., Pajusola K., von Euler G., Chilov D., Alitalo K., Eriksson U. (1996b). Genomic organization of the mouse and human genes for vascular endothelial growth factor B (VEGF-B) and characterization of a second splice isoform. *J. Biol. Chem.* **271**: 19310-19317

Olofsson B., Korpeleinen E., Pepper M.S., Mandriota S., Aase K., Kumar V., Gunji Y., Jeltsch M.M., Shibuya M., Alitalo K., Eriksson U. (1998). Vascular endothelial growth factor B (VEGF-B) binds VEGF receptor-1 and regulates plasminogen activator activity in endothelial cells. *Proc. Natl. Acad. Sci. USA* **95**: 11709-11714

O'Reilly M.S., Holmgren L., Shing Y., Chen C., Rosenthal R.A., Moses M., Lane W.S., Cao Y., Sage E.H., Folkman J. (1994). Angiostatin: a novel angiogenesis inhibitor that mediates the suppression of metastases by a Lewis lung carcinoma. *Cell* **79**: 315-328

O'Reilly M.S., Boehm T., Shing Y., Fukai N., Vasios G., Lane W.S., Flynn E., Birkhead J.R., Olsen B.R., Folkman J. (1997). Endostatin: an endogenous inhibitor of angiogenesis and tumor growth. *Cell* **88**: 277-285

Orlandini M., Marconcini L., Ferruzzi R., Oliviero S. (1996). Identification of a c-fos-induced gene that is related to the platelet-derived growth factor/vascular endothelial growth factor family. *Proc. Natl. Acad. Sci. USA* **93**: 11675-11680

O'Shea J.D., Cran D.G., Hay M.F. (1980). Fate of the theca interna following ovulation in the ewe. *Cell Tissue Res.* **210**: 305-319

O'Shea J.D. (1987). Heterogeneous cell types in the corpus luteum of sheep, goat and cattle. *J. Reprod. Fert. (Suppl)* **34**: 71-85

Pajusola K., Aprelikova O., Korhonen J., Kaipainen A., Alitalo R., Alitalo K. (1992). FLT4 receptor tyrosine kinase contains seven immuno-globulin-like loops and is expressed in multiple human tissues and cell lines.

Cancer Res. **52**: 5738-5743

Paku S., Paweletz N. (1991).

First steps of tumor-related angiogenesis.

Lab. Invest. **65**: 334-346

Paku S. (1998).

Current concepts of tumor-induced angiogenesis.

Pathol. Oncol. Res. **4**: 62-75

Papapetropoulos A., Garcia-Cardena G., Dengler T.J., Maisonpierre P.C., Yancopoulos G.D., Sessa W.C. (1999).

Direct actions of angiopoietin-1 on human endothelium: evidence for network stabilization, cell survival, and interaction with other angiogenic growth factors.

Lab. Invest. **79**: 213-223

Park J.E., Keller G.A., Ferrara N. (1993).

The vascular endothelial growth factor (VEGF) isoforms - differential deposition into the subepithelial extracellular matrix and bioactivity of extracellular matrix-bound VEGF.

Mol. Biol. Cell **4**: 1317-1326

Park J.E., Chen H.H., Winer J., Houck K.A., Ferrara N. (1994).

Placenta growth factor. Potentiation of vascular endothelial growth factor bioactivity, *in vitro* and *in vivo*, and high affinity binding to Flt-1 but not to Flk-1/KDR.

J. Biol. Chem. **269**: 25646-25654

Patan S., Haenni B., Burri P.H. (1996).

Implementation of intussusceptive microvascular growth in the chicken chorioallantoic membrane (CAM): 1. Pillar formation by folding of the capillary wall.
Microvasc. Res. **51**: 80-98

Pe'er J., Folberg R., Itin A., Gnessin H., Hemo I., Keshet E. (1996).

Upregulated expression of vascular endothelial growth factor in proliferative diabetic retinopathy.

Br. J. Ophthalmol. **80**: 241-245

Pepper M.S., Ferrara N., Orci L., Montesano R. (1992).

Potent synergism between vascular endothelial growth factor and basic fibroblast growth factor in the induction of angiogenesis in vitro.

Biochem. Biophys. Res. Commun. **189**: 824-831

Pepper J., Vassalli J.D., Orci L., Montesano R. (1993).

Biphasic effect of transforming growth factor- β 1 on *in vitro* angiogenesis.

Exp. Cell Res. **204**: 356-363

Pepper M.S., Mandriota S.J., Vassalli J.-D., Orci L., Montesano R. (1996).

Angiogenesis-regulating cytokines: activities and interactions.

Curr. Top. Microbiol. Immunol. **213**: 31-67

Pertovaara L., Kaipainen A., Mustonen T., Orpana A., Ferrara N., Saksela O., Alitalo K. (1994).

Vascular endothelial growth factor is induced in response to transforming growth factor- β in fibroblastic and epithelial cells.

J. Biol. Chem. **269**: 6271-6274

Peters K.G., De Vries C., Williams L.T. (1993).

Vascular endothelial growth factor receptor expression during embryogenesis and tissue repair suggests a role in endothelial differentiation and blood vessel growth.

Proc. Natl. Acad. Sci. USA **90**: 8915-8919

Petrides P.E. (1997).

Endokrine Gewebe II: Die schnelle Stoffwechselregulation. In: Biochemie und Pathobiochemie (Löffler G., Petrides P.E., Hrsg.), 787-812. Springer Verlag Berlin Heidelberg New York.

Phillips H.S., Hains J., Leung D.W., Ferrara N. (1990).

Vascular endothelial growth factor is expressed in rat corpus luteum.

Endocrinology **127**: 965-967

Picton H.M. (2001).

Activation of follicle development: the primordial follicle.

Theriogenology **55**: 1193-1210

Pipili-Synetos E., Sakkoula E., Maragoudakis M.E. (1993).

Nitric oxide is involved in the regulation of angiogenesis.

Br. J. Pharmacol. **108**: 855-857

Plate K.H., Breier G., Weich H.A., Risau W. (1992).

Vascular endothelial growth factor is a potential tumour angiogenesis factor in human gliomas *in vivo*.

Nature **359**: 845-848

Plendl J., Sinowitz F., Auerbach R. (1992).

Die Heterogenität des vaskulären Endothels.

Anat. Histol. Embryol. **21**: 256-262

Plendl J. (1997).

Das Gefäßendothel: organ- und entwicklungsphasenspezifische Heterogenität im *in vitro* Modell.

Habilitationsschrift. Tierärztliche Fakultät der Ludwig-Maximilians-Universität München.

Plendl J., Sinowitz F. (1999).

Angiogenese und Antiangiogenese: neue therapeutische Strategien.

Tierärztl. Prax. **27**: 348-353

Plendl J., Stierstorfer B., Fuchs T., Boie G., Sinowitz F. (1999).

Expression of vascular endothelial growth factor and its receptor in an in vitro model of the bovine corpus luteum.

Reprod. Dom. Anim. **43**: 22 (abstract)

Plendl J. (2000).

Angiogenesis and vascular regression in the ovary.

Anat. Histol. Embryol. **29**: 257-266

Plunkett M.L., Hailey J.A. (1990).

An in vivo quantitative angiogenesis model using tumor cells entrapped in alginate.

Lab. Invest. **62**: 510-517

Poltorak Z., Cohen T., Sivan R., Kanelis Y., Spira G., Vlodavsky I., Keshet E.,

Neufeld G. (1997).

VEGF₁₄₅, a secreted vascular endothelial growth factor isoform that binds to extracellular matrix.

J. Biol. Chem. **272**: 7151-7158

Poole T.J., Coffin J.D. (1988).

Developmental angiogenesis: quail embryonic vasculature.

Scanning Microsc. **2**: 443-448

Rak J., Mitsuhashi Y., Bayko L., Filmus J., Shirasawa S., Sasazuki T., Kerbel R.S.

(1995).

Mutant *ras* oncogenes upregulate VEGF/VPF expression: implications for induction and inhibition of tumor angiogenesis.

Cancer Res. **55**: 4575-4580

Ravindranath N., Little-Ihrig L., Phillips H.S., Ferrara N., Zeleznik A.J. (1992).
Vascular endothelial growth factor messenger ribonucleic acid expression in the primate ovary.
Endocrinology **131**: 254-260

Redmer D.A., Reynolds L.P. (1996).

Angiogenesis in the ovary.

Rev. Reprod. **1**: 182-192

Reynolds L.P., Redmer D.A. (1998).

Expression of the angiogenic factors, basic fibroblast growth factor and vascular endothelial growth factor in the ovary.

J. Anim. Sci. **76**: 1671-1681

Ricken A., Rahner C., Landmann L., Spanel-Borowski S. (1996).

Bovine endothelial-like cells increase intercellular junctions under treatment with interferon- γ . An in vitro study.

Anat. Anz. **178**: 321-330

Risau W., Sariola H., Zerwes H., Sasse J., Ekblom P., Kemler R., Doetschman T. (1988).

Vasculogenesis and angiogenesis in embryonic-stem-cell-derived embryoid bodies.

Development **102**: 471-478

Risau W. (1996).

What, if anything, is an angiogenic factor?

Cancer Metastasis Rev. **15**: 149-151

Roberts W.G., Palade G.E. (1995).

Increased microvascular permeability and endothelial fenestration induced by vascular endothelial growth factor.

J. Cell Sci. **108**: 2369-2379

Rodger F.E., Young F.M., Fraser H.M., Illingworth P.J. (1997).
Endothelial cell proliferation follows the mid-cycle luteinizing hormone surge, but
not human chorionic gonadotrophin rescue, in the human corpus luteum.
Hum. Reprod. **12**: 1723-1729

Rosenthal R.A., Megyesi J.F., Henzel W.J., Ferrara N., Folkman J. (1990).
Conditioned medium from mouse sarcoma 180 cells contains vascular endothelial
growth factor.
Growth Factors **4**: 53-59

Rüsse I. (1998).
Fruchthüllen des Vogels. In: Lehrbuch der Embryologie der Haustiere (Rüsse I.,
Sinowitz F., Hrsg.), 204-206. Paul Parey, Berlin, Hamburg.

Rüsse I. (1998).
Weiblicher Sexualzyklus. In: Lehrbuch der Embryologie der Haustiere (Rüsse I.,
Sinowitz F., Hrsg.), 93-116. Paul Parey, Berlin, Hamburg.

Rüsse I., Amselgruber W., Sinowitz F. (1987).
Gefäßarchitektonische Untersuchungen am ovulierenden Follikel des Rindes.
Fertilität **3**: 94-98

Rüsse I., Sinowitz F. (1998).
Gametogenese. In: Lehrbuch der Embryologie der Haustiere (Rüsse I., Sinowitz F.,
Hrsg.), 42-98. Paul Parey, Berlin, Hamburg.

Rymaszewski Z., Cohen R.M., Chomczynski P. (1991).
Human growth hormone stimulates proliferation of human retinal microvascular
endothelial cells *in vitro*.
Proc. Natl. Acad. Sci. USA **88**: 617-621

Ryuto M., Ono M., Izumi H., Yoshida S., Weich H.A., Kohno K., Kuwano M. (1996).
Induction of vascular endothelial growth factor by tumor necrosis factor alpha in
human glioma cells.

J. Biol. Chem. **271**: 28220-28228

Salmon W.D., Daughaday W.H. (1957).

A hormonal controled serum factor which stimulates sulfate incorporation by
cartilage in vitro.

J. Lab. Clin. Med. **49**: 825-836

Sandison J.C. (1924).

A new method for the microscopic study of living growing tissues by the introduction
of a transparent chamber in the rabbit's ear.

Anat. Rec. **28**: 281-287

Sandison J.C. (1932).

Contraction of blood vessels and observations on the circulation in the transparent
chamber in the rabbit's ear.

Anat. Rec. **54**: 105-127

Sato T.N., Tozawa Y., Deutsch U., Wolburg-Buchholz K., Fujiwara Y., Gendron-
Maguire M., Gridley T., Wolburg H., Risau W., Qin Y. (1995).

Distinct roles of the receptor tyrosine kinases Tie-1 and Tie-2 in blood vessel
formation.

Nature **376**: 70-74

Sato Y., Shimada T., Takaki R. (1991).

Autocrinological role of basic fibroblast growth factor on tube formation of vascular
endothelial cells in vitro.

Biochem. Biophys. Res. Commun. **180**: 1098-1102

Sawano A., Takahashi T., Yamaguchi S., Aonuma T., Shibuya M. (1996). Flt-1 but not KDR/Flk-1 tyrosine kinase is a receptor for placenta growth factor (PIGF), which is related to vascular endothelial growth factor (VEGF). *Cell Growth Differ.* **7:** 213-221

Schams D., Berisha B., Kosmann M., Einspanier R., Amselgruber W.M. (1999). Possible role of growth hormone, IGFs, and IGF-binding proteins in the regulation of ovarian function in large farm animals. *Dom. Anim. Endocrinol.* **17:** 279-285

Schröder M. (1999). Isolierung von RNA. In: Gentechnische Methoden (Gassen H.G. und Schrimpf G., Hrsg.), 225-242. Spektrum Akademischer Verlag, Heidelberg, Berlin.

Schweigerer L., Malerstein B., Gospodarowicz D. (1987). Tumor necrosis factor inhibits the proliferation of cultured capillary endothelial cells.

Biochem. Biophys. Res. Commun. **143:** 997-1004

Senger D.R., Galli S.J., Dvorak A.M., Perruzzi C.A., Harvey V.S., Dvorak H.F. (1983). Tumor cells secrete a vascular permeability factor that promotes accumulation of ascites fluid.

Science **25:** 983-985

Senger D.R., Perruzzi C.A., Feder J., Dvorak H.F. (1986). A highly conserved vascular permeability factor secreted by a variety of human and rodent tumor cell lines. *Cancer Res.* **46:** 5629-5632

Shalaby F., Rossant J., Yamaguchi T.P., Gertsenstein M., Wu X.F., Breitman M.L., Schuh A.C. (1995). Failure of blood-island formation and vasculogenesis in Flk-1-deficient mice. *Nature* **376:** 62-66

Shi Q., Rafii S., Wu M.H.-D., Wijelath E.S., Yu C., Ishida A., Fujita Y., Kothari S., Mohle R., Sauvage L.R., Moore M.A.S., Storb R.F., Hammond W.P. (1998). Evidence for circulating bone marrow-derived endothelial cells. *Blood* **92**: 362-367

Shibuya M., Yamaguchi S., Yamane A., Ikeda T., Tojo A., Matsushima H., Sato M. (1990).

Nucleotide sequence and expression of a novel human receptor-type kinase gene (*flt*) closely related to the *fms* family.

Oncogene **5**: 519-524

Shweiki D., Itin A., Soffer D., Keshet E. (1992).

Vascular endothelial growth factor induced by hypoxia may mediate hypoxia-initiated angiogenesis.

Nature **359**: 843-845

Shweiki D., Itin A., Neufeld G., Gitay-Goren H., Keshet E. (1993).

Patterns of expression of vascular endothelial growth factor (VEGF) and VEGF receptors in mice suggest a role in hormonally regulated angiogenesis.

J. Clin. Invest. **91**: 2235-2243

Sim B.K.L., MacDonald N.J., Gubish E.R. (2000).

Angiostatin and endostatin: endogenous inhibitors of tumor growth.

Cancer Metastasis Rev. **19**: 181-190

Sinowatz F. (2000).

Sinnesorgane. In: *Histologie* (H. Hees, F. Sinowatz, Hrsg.), 383-427. Deutscher Ärzte-Verlag, Köln.

Sinowatz F. (2000).

Die Zelle. In: *Histologie* (H. Hees, F. Sinowatz, Hrsg.), 35-93. Deutscher Ärzte-Verlag, Köln.

Sinowatz F., Kölle S., Töpfer-Petersen E. (2001).

Biosynthesis and expression of zona pellucida glycoproteins in mammals.

Cells Tissues Organs **168**: 24-35

Smith L.E.H., Kopchick J., Chen W., Knapp J., Kinose f., Daley D., Foley E., Smith R.G., Schaeffer J.M. (1997).

Essential role of growth hormone in ischemia-induced retinal neovascularization.

Science **276**: 1706-1709

Smith M.F., McIntosh E.W., Smith G.W. (1994).

Mechanisms associated with corpus luteum development.

J. Anim. Sci. **72**: 1857-1872

Soker S., Takashima S., Miao H.Q., Neufeld G., Klagsbrun M. (1998).

Neuropilin-1 is expressed by endothelial and tumor cells as an isoform-specific receptor for vascular endothelial growth factor.

Cell **92**: 735-745

Sorokin L., Girg W., Gopfert T., Hallmann R., Deutzmann R. (1994).

Expression of novel 400-kDa laminin chains by mouse and bovine endothelial cells.

Eur. J. Biochem. **223**: 603-610

Stone J., Chang-Ling T., Pe'er J., Itin A., Gnessin H., Keshet E. (1996).

Roles of vascular endothelial growth factor and astrocyte degeneration in the genesis of retinopathy of prematurity.

Invest. Ophthalmol. Vis. Sci. **37**: 290-299

Struman I., Bentzien F., Lee H., Mainfroid V., D'Angelo G., Goffin V., Weiner R.I., Martial J.A. (1999).

Opposing actions of intact and N-terminal fragments of the human prolactin/growth hormone family members on angiogenesis: an efficient mechanism for the regulation of angiogenesis.

Proc. Natl. Acad. Sci. USA **96**: 1246-1251

Stryer L. (1996).

Replikation und Expression von Genen. In: Biochemie (Stryer L., Hrsg.), 825-1058.
Akademischer Verlag GmbH Heidelberg, Berlin, Oxford.

Suri C., Jones P.F., Patan S., Bartunkova S., Maisonpierre P.C., Davis S., Sato T.N.,
Yancopoulos G.D. (1996).

Requisite role of angiopoietin-1, a ligand for the Tie2 receptor, during embryonic
angiogenesis.

Cell **87**: 1171-1180

Suri C., McClain J., Thurston G., McDonald D.M., Zhou H., Oldmixon E.H., Sato T.N.,
Yancopoulos G.D. (1998).

Increased vascularization in mice overexpressing angiopoietin-1.

Science **282**: 468-471

Takagi H., King G.L., Aiello L.P. (1996)

Identification and characterization of vascular endothelial growth factor receptor (flt)
in bovine retinal pericytes.

Diabetes **45**: 1016-1023

Takahashi T., Shirasawa T., Miyake K., Yahagi Y., Maruyama N., Kasahara N.,
Kawamura T., Matsumura O., Mitarai T., Sakai O. (1995).

Protein tyrosine kinases expressed in glomeruli and cultured glomerular cells: Flt-1
and VEGF expression in renal mesangial cells.

Biochem. Biophys. Res. Commun. **209**: 218-226

Terman B.I., Carrion M.E., Kovacs E., Rasmussen B.A., Eddy R.L., Shows T.B.
(1991).

Identification of a new endothelial cell growth factor receptor tyrosine kinase.
Oncogene **6**: 1677-1683

Terman B.I., Dougher-Vermazen M., Carrion M.E., Dimitrov D., Armellino D.C., Gospodarowicz D., Bohlen P. (1992).

Identification of the KDR tyrosine kinase as a receptor for vascular endothelial growth factor.

Biochem. Biophys. Res. Commun. **187**: 1579-1586

Terman B.I., Khandke L., Dougher-Vermazan M., Maglione D., Lassam N.J., Gospodarowicz D., Persico M.G., Böhlen P., Eisinger M. (1994).

VEGF receptor subtypes KDR and FLT1 show different sensitivities to heparin and placenta growth factor.

Growth Factors **11**: 187-195

Thurston G., Suri C., Smith K., McClain J., Sato T.N., Yancopoulos G.D., McDonald D.M. (1999).

Leakage-resistant blood vessels in mice transgenically overexpressing angiopoietin-1.

Science **286**: 2511-2514

Tischer E., Mitchell R., Hartman T., Silva M., Gospodarowicz D., Fiddes J.C., Abraham J.A. (1991).

The human gene for vascular endothelial growth factor: multiple protein forms are encoded through alternative exon splicing.

J. Biol. Chem. **266**: 11947-11954

Tolsma S.S., Volpert O.V., Good D.J., Frazier W.A., Polverini P.J., Bouck N. (1993).

Peptides derived from two separate domains of the matrix protein thrombospondin-1 have anti-angiogenic activity.

J. Cell Biol. **122**: 497-511

Tuder R.M., Flook B.E., Voelkel N.F. (1995).

Increased gene expression for VEGF and the VEGF receptors *KDR/flk* and *flt* in lungs exposed to acute or to chronic hypoxia. Modulation of gene expression by nitric oxide.

J. Clin. Invest. **95**: 1798-1807

- Ullrich A., Schlessinger J. (1990).
Signal transduction by receptors with tyrosine kinase activity.
Cell **61**: 203-212
- Vacca A., Ribatti D., Iurlaro M., Albini A., Minischetti M., Bussolino F., Pellegrino A., Ria R., Rusnati M., Presta M., Vincenti V., Persico M.G., Dammacco F. (1998).
Human lymphoblastoid cells produce extracellular matrix-degrading enzymes and induce endothelial cell proliferation, migration, morphogenesis, and angiogenesis.
Int. J. Clin. Lab. Res. **28**: 55-68
- Van Groningen J.P., Wenink A.C.G., Testers L.H.M. (1991).
Myocardial capillaries: increase in number by splitting of existing vessels.
Anat. Embryol. **184**: 65-70
- Vega-Diaz B., Herron S.G., Michel S. (2001).
An autocrine loop mediates expression of vascular endothelial growth factor in human dermal microvascular endothelial cells.
J. Invest. Dermatol. **116**: 525-530
- Voigt K. (1994).
Endokrines System. In: *Lehrbuch der Physiologie* (Klinke R., Silbernagel S., Hrsg.), 439-490. Georg Thieme Verlag, Stuttgart, New York.
- Volpert O.V., Tolsma S.S., Pellerin S., Feige J.J., Chen H., Mosher D.F., Bouck N. (1995).
Inhibition of angiogenesis by thrombospondin-2.
Biochem. Biophys. Res. Commun. **217**: 326-332
- Von den Driesch A. (1998).
Geschichte der Embryologie der Haustiere. In: *Lehrbuch der Embryologie der Haustiere* (Rüsse I., Sinowitz F., Hrsg.), 11-25. Paul Parey, Berlin, Hamburg.

Waltenberger J., Claesson-Welsh L., Siegbahn A., Shibuya M., Heldin C.H. (1994). Different signal transduction properties of KDR and Flt1, two receptors for vascular endothelial growth factor.

J. Biol. Chem. **269**: 26988-26995

Waltenberger J., Mayr U., Pentz S., Hombach V. (1996).

Functional upregulation of the vascular endothelial growth factor receptor KDR by hypoxia.

Circulation **94**: 1647-1654

Wen Y., Edelman J.L., Kang T., Zeng N.X. Sachs G (1998).

Two functional forms of vascular endothelial growth factor receptor-2/Flik-1 mRNA are expressed in normal rat retina.

J. Biol. Chem. **273**: 2090-2097

Xin X., Yang S., Ingle G., Zlot C., Rangell L., Kowalski J., Schwall R., Ferrara N., Gerritsen M.E. (2001).

Hepatocyte growth factor enhances vascular endothelial growth factor-induced angiogenesis *in vitro* and *in vivo*.

Am. J. Pathol. **158**: 1111-1120

Yamada Y., Nezu J., Shimane M., Hirata Y. (1997).

Molecular cloning of a new vascular endothelial growth factor, VEGF-D.

Genomics **42**: 483-488

Yamagami I. (1970).

Electron microscopic study on the cornea. 1. the mechanism of experimental new vessel formation.

Japan. J. Ophthalmol. **14**: 41-58

Yang X., Cepko C.L. (1996).

Flik-1, a receptor for vascular endothelial growth factor (VEGF), is expressed by retinal progenitor cells.

J. Neurosci. **16**: 6089-6099

Yasunaga C., Nakashima Y., Sueishi K. (1989).

A role of fibrinolytic activity in angiogenesis. Quantitative assay using *in vitro* method.

Lab. Invest. **61**: 698-704

Young P.E., Baumhueter S., Lasky L.A. (1995).

The sialomucin CD34 is expressed on hematopoietic cells and blood vessels during murine development.

Blood **85**: 96-105

Zheng J., Redmer D.A., Reynolds L.P. (1993).

Vascular development and heparin-binding growth factors in the bovine corpus luteum at several stages of the estrous cycle.

Biol. Reprod. **49**: 1177-1189

Zhu T., Goh E.L.K., Graichen R., Ling L., Lobie P.E. (2001).

Signal transduction via the growth hormone receptor.

Cell Signal. **13**: 599-616

Ziche M., Morbidelli L., Masini E., Amerin S., Granger H.J., Maggi C.A., Geppetti P., Ledda F. (1994).

Nitric oxide mediates angiogenesis *in vivo* and endothelial cell growth and migration *in vitro* promoted by substance P.

J. Clin. Invest. **94**: 2036-2044

Ziche M., Maglione D., Ribatti D., Morbidelli L., Lago C.T., Battisti M., Paoletti I., Barra A., Tucci M., Parise G., Vincenti V., Granger H.J., Viglietto G., Persico M.G. (1997).

Placenta growth factor-1 is chemotactic, mitogenic, and angiogenic.

Lab. Invest. **76**: 517-531