

6. Literaturverzeichnis

- Albin, R. L.; Young, A. B. und Penney, J. B. (1989): The functional anatomy of basal ganglia disorders, Trends Neurosci 12 [10], Seite 366-75.
- Alderson, H. L.; Latimer, M. P. und Winn, P. (2005): Involvement of the laterodorsal tegmental nucleus in the locomotor response to repeated nicotine administration, Neurosci Lett 380 [3], Seite 335-9.
- Alheid, G. F.; Heimer, L. und Schwitzer, R. C. (1990): Basal Ganglia. In: Paxinos, G. (ed.): The Human Nervous System Academic Press, San Diego - New York - Boston - London - Sydney - Tokyo - Toronto, Seite 483-582.
- Amat, J.; Sparks, P. D.; Matus-Amat, P.; Griggs, J.; Watkins, L. R. und Maier, S. F. (2001): The role of the habenular complex in the elevation of dorsal raphe nucleus serotonin and the changes in the behavioral responses produced by uncontrollable stress, Brain Res 917 [1], Seite 118-26.
- Andres, K. H.; von During, M. und Veh, R. W. (1999): Subnuclear organization of the rat habenular complexes, J Comp Neurol.
- Araki, M.; McGeer, P. L. und Kimura, H. (1988): The efferent projections of the rat lateral habenular nucleus revealed by the PHA-L anterograde tracing method, Brain Res.
- Beccari, N. (1943): Neurologia Comparata anatomo-funzionale dei vertebrati, compresso l'uomo, Firenze: Sansoni Edizioni Scientifiche.
- Behzadi, G.; Kalen, P.; Parvopassu, F. und Wiklund, L. (1990): Afferents to the median raphe nucleus of the rat: retrograde cholera toxin and wheat germ conjugated horseradish peroxidase tracing, and selective D-[3H]aspartate labelling of possible excitatory amino acid inputs, Neuroscience 37 [1], Seite 77-100.
- Benabid, A. L. und Jeaugey, L. (1989): Cells of the rat lateral habenula respond to high-threshold somatosensory inputs, Neurosci Lett 96 [3], Seite 289-94.
- Bielajew, C. H. (1991): Distribution of cytochrome oxidase in response to rewarding brain stimulation: effect of different pulse durations, Brain Res Bull 26 [3], Seite 379-84.
- Blaha, C. D.; Allen, L. F.; Das, S.; Inglis, W. L.; Latimer, M. P.; Vincent, S. R. und Winn, P. (1996): Modulation of dopamine efflux in the nucleus accumbens after cholinergic stimulation of the ventral tegmental area in intact, pedunculopontine tegmental nucleus-lesioned, and laterodorsal tegmental nucleus-lesioned rats, J Neurosci 16 [2], Seite 714-22.
- Blander, A. und Wise, R. A. (1989): Anatomical mapping of brain stimulation reward sites in the anterior hypothalamic area: special attention to the stria medullaris, Brain Res 483 [1], Seite 12-6.
- Bleier, R.P. ; Cohn, P. und Sigelkow, I.R. (1979): A cytoarchitectonic atlas of the hypothalamus and hypothalamic third ventricle of the rat. In P. J. Morgane and J. Panksepp (eds) Anatomy of the Hypothalamus 1, Marcel Dekker, New York.
- Bon, K.; Lanteri-Minet, M.; de Pommery, J.; Michiels, J. F. und Menetrey, D. (1997): Cyclophosphamide cystitis as a model of visceral pain in rats: minor effects at mesodiencephalic levels as revealed by the expression of c-fos, with a note on Krox-24, Exp Brain Res 113 [2], Seite 249-64.
- Brinschwitz, K.; Lommel, R.; Penkalla, A.; Goertzen, A.; Geisler, S. und Veh, R.W. (2005): GLUTAMATERGIC FIBERS FROM THE LATERAL HABENULA DO NOT TERMINATE ON DOPAMINERGIC NEURONS IN THE VENTRAL

- TEGMENTAL AREA Program No. 891.23. 2005 Abstract Viewer/Itinerary Planner. Washington, DC: Society for Neuroscience, 2005. Online.
- Briski, K. und Gillen, E. (2001): Differential distribution of Fos expression within the male rat preoptic area and hypothalamus in response to physical vs. psychological stress, *Brain Res Bull* 55 [3], Seite 401-8.
- Broadwell, R. D. und Bleier, R. (1976): A cytoarchitectonic atlas of the mouse hypothalamus, *J Comp Neurol* 167 [3], Seite 315-39.
- Cajal, S.R. (1904): *La Textura del Sistema Nervioso del Hombre y los Vertebrados*, Moya, Madrid.
- Caldecott-Hazard, S.; Mazziotta, J. und Phelps, M. (1988): Cerebral correlates of depressed behavior in rats, visualized using ¹⁴C-2-deoxyglucose autoradiography, *J Neurosci* 8 [6], Seite 1951-61.
- Cechetto, D. F. und Saper, C. B. (1988): Neurochemical organization of the hypothalamic projection to the spinal cord in the rat, *J Comp Neurol* 272 [4], Seite 579-604.
- Chastrette, N.; Pfaff, D. W. und Gibbs, R. B. (1991): Effects of daytime and nighttime stress on Fos-like immunoreactivity in the paraventricular nucleus of the hypothalamus, the habenula, and the posterior paraventricular nucleus of the thalamus, *Brain Res* 563 [1-2], Seite 339-44.
- Chen, J.; Nakamura, M.; Kawamura, T.; Takahashi, T. und Nakahara, D. (2006): Roles of pedunculopontine tegmental cholinergic receptors in brain stimulation reward in the rat, *Psychopharmacology (Berl)* 184 [3-4], Seite 514-22.
- Chen, S. und Aston-Jones, G. (1995): Evidence that cholera toxin B subunit (CTb) can be avidly taken up and transported by fibers of passage, *Brain Res* 674 [1], Seite 107-11.
- Christoph, G. R.; Leonzio, R. J. und Wilcox, K. S. (1986): Stimulation of the lateral habenula inhibits dopamine-containing neurons in the substantia nigra and ventral tegmental area of the rat, *J Neurosci*.
- Cohen, S. R. und Melzack, R. (1985): Morphine injected into the habenula and dorsal posteromedial thalamus produces analgesia in the formalin test, *Brain Res* 359 [1-2], Seite 131-9.
- Cohen, S. R. und Melzack, R. (1986): Habenular stimulation produces analgesia in the formalin test, *Neurosci Lett* 70 [1], Seite 165-9.
- Cohen, S. R. und Melzack, R. (1993): The habenula and pain: repeated electrical stimulation produces prolonged analgesia but lesions have no effect on formalin pain or morphine analgesia, *Behav Brain Res* 54 [2], Seite 171-8.
- Concha, M. L. und Wilson, S. W. (2001): Asymmetry in the epithalamus of vertebrates, *J Anat* 199 [Pt 1-2], Seite 63-84.
- Conrad, L. C. und Pfaff, D. W. (1976): Efferents from medial basal forebrain and hypothalamus in the rat. I. An autoradiographic study of the medial preoptic area, *J Comp Neurol* 169 [2], Seite 185-219.
- Conrad, L. C. und Pfaff, D. W. (1976): Efferents from medial basal forebrain and hypothalamus in the rat. II. An autoradiographic study of the anterior hypothalamus, *J Comp Neurol* 169 [2], Seite 221-61.
- Cornwall, J.; Cooper, J. D. und Phillipson, O. T. (1990): Afferent and efferent connections of the laterodorsal tegmental nucleus in the rat, *Brain Res Bull* 25 [2], Seite 271-84.
- Corodimas, K. P.; Rosenblatt, J. S.; Canfield, M. E. und Morrell, J. I. (1993): Neurons in the lateral subdivision of the habenular complex mediate the hormonal onset of maternal behavior in rats, *Behav Neurosci* 107 [5], Seite 827-43.

- Corodimas, K. P.; Rosenblatt, J. S. und Morrell, J. I. (1992): The habenular complex mediates hormonal stimulation of maternal behavior in rats, Behav Neurosci 106 [5], Seite 853-65.
- Crossman, A. R. (1987): Primate models of dyskinesia: the experimental approach to the study of the basal ganglia-related involuntary movement disorders, Neurosci 21, Seite 1-40.
- Dado, R. J.; Burstein, R.; Cliffer, K. D. und Giesler, G. J., Jr. (1990): Evidence that Fluoro-Gold can be transported avidly through fibers of passage, Brain Res 533 [2], Seite 329-33.
- DeLong, M. R. (1990): Primate models of movement disorders of basal ganglia origin, Trends Neurosci 13 [7], Seite 281-5.
- Donovick, P. J.; Burright, R. G. und Zuromski, E. (1970): Localization of quinine aversion within the septum, habenula, and interpeduncular nucleus of the rat, J Comp Physiol Psychol 71 [3], Seite 376-83.
- Dougherty, P. M.; Qiao, J. T.; Wiggins, R. C. und Dafny, N. (1990): Microiontophoresis of cocaine, desipramine, sulpiride, methysergide, and naloxone in habenula and parafasciculus, Exp Neurol 108 [3], Seite 241-6.
- Edinger, L. (1896): Vorlesungen über den Bau der nervösen Zentralorgane des Menschen und der Thiere 5, Vogel, Leipzig.
- Elisevich, K.; Rajakumar, N.; Hrycyshyn, A.W. und Flumerfelt, B.A. (1993): Topographical organization of the corticostriatal projection in the rat., SOCN. eurosci. Abstr. 19:1433.
- Ellison, G. (1992): Continuous amphetamine and cocaine have similar neurotoxic effects in lateral habenular nucleus and fasciculus retroflexus, Brain Res 598 [1-2], Seite 353-6.
- Ellison, G. (1994): Stimulant-induced psychosis, the dopamine theory of schizophrenia, and the habenula, Brain Res Brain Res Rev.
- Ellison, G.; Irwin, S.; Keys, A.; Noguchi, K. und Sulur, G. (1996): The neurotoxic effects of continuous cocaine and amphetamine in Habenula: implications for the substrates of psychosis, NIDA Res Monogr.
- Felton, T. M.; Linton, L.; Rosenblatt, J. S. und Morrell, J. I. (1998): Intact neurons of the lateral habenular nucleus are necessary for the nonhormonal, pup-mediated display of maternal behavior in sensitized virgin female rats, Behav Neurosci.
- Forster, G. L. und Blaha, C. D. (2000): Laterodorsal tegmental stimulation elicits dopamine efflux in the rat nucleus accumbens by activation of acetylcholine and glutamate receptors in the ventral tegmental area, Eur J Neurosci 12 [10], Seite 3596-604.
- Forster, G. L.; Falcon, A. J.; Miller, A. D.; Heruc, G. A. und Blaha, C. D. (2002): Effects of laterodorsal tegmentum excitotoxic lesions on behavioral and dopamine responses evoked by morphine and d-amphetamine, Neuroscience 114 [4], Seite 817-23.
- Gallistel, C. R.; Gomita, Y.; Yadin, E. und Campbell, K. A. (1985): Forebrain origins and terminations of the medial forebrain bundle metabolically activated by rewarding stimulation or by reward-blocking doses of pimozide, J Neurosci 5 [5], Seite 1246-61.
- Ganser, S. (1881): Vergleichend anatomische Studien ueber das Gehirn des Maulwurfs, Gegenbaurs Morphol Jahrb 7.
- Gao, D. M.; Hoffman, D. und Benabid, A. L. (1996): Simultaneous recording of spontaneous activities and nociceptive responses from neurons in the pars compacta of substantia nigra and in the lateral habenula, Eur J Neurosci.

- Gao, D. M.; Jeaugey, L.; Pollak, P. und Benabid, A. L. (1990): Intensity-dependent nociceptive responses from presumed dopaminergic neurons of the substantia nigra, pars compacta in the rat and their modification by lateral habenula inputs, *Brain Res* 529 [1-2], Seite 315-9.
- Geeraedts, L. M.; Nieuwenhuys, R. und Veening, J. G. (1990a): Medial forebrain bundle of the rat: III. Cytoarchitecture of the rostral (telencephalic) part of the medial forebrain bundle bed nucleus, *J Comp Neurol* 294 [4], Seite 507-36.
- Geeraedts, L. M.; Nieuwenhuys, R. und Veening, J. G. (1990b): Medial forebrain bundle of the rat: IV. Cytoarchitecture of the caudal (lateral hypothalamic) part of the medial forebrain bundle bed nucleus, *J Comp Neurol* 294 [4], Seite 537-68.
- Geisler, S.; Andres, K. H. und Veh, R. W. (2003): Morphologic and cytochemical criteria for the identification and delineation of individual subnuclei within the lateral habenular complex of the rat., *J Comp Neurol*.
- Geisler, S. und Zahm, D. S. (2005): Afferents of the ventral tegmental area in the rat-anatomical substratum for integrative functions, *J Comp Neurol* 490 [3], Seite 270-94.
- Gerfen, C. R. und Sawchenko, P. E. (1984): An anterograde neuroanatomical method that shows the detailed morphology of neurons, their axons and terminals: immunohistochemical localization of an axonally transported plant lectin, *Phaseolus vulgaris* leucoagglutinin (PHA-L), *Brain Res* 290, Seite 219-238.
- Goldstein, R. (1983): A GABAergic habenulo-raphe pathway mediation of the hypnotogenic effects of vasotocin in cat, *Neuroscience* 10 [3], Seite 941-5.
- Gomita, Y. und Gallistel, C. R. (1982): Effects of reinforcement-blocking doses of pimozide on neural systems driven by rewarding stimulation of the MFB: a ¹⁴C-2-deoxyglucose analysis, *Pharmacol Biochem Behav* 17 [4], Seite 841-5.
- Gonzalez-Lima, F.; Helmstetter, F. J. und Agudo, J. (1993): Functional mapping of the rat brain during drinking behavior: a fluorodeoxyglucose study, *Physiol Behav* 54 [3], Seite 605-12.
- Gonzalez-Lima, F. und Scheich, H. (1986): Classical conditioning of tone-signaled bradycardia modifies 2-deoxyglucose uptake patterns in cortex, thalamus, habenula, caudate-putamen and hippocampal formation, *Brain Res* 363 [2], Seite 239-56.
- Goto, M.; Swanson, L. W. und Canteras, N. S. (2001): Connections of the nucleus incertus, *J Comp Neurol*.
- Gurdjian, E.S. (1927): The diencephalon of the albino rat, *J Comp Physiol Psychol* 43, Seite 1-144.
- Harlan, R. E.; Shivers, B. D.; Romano, G. J.; Howells, R. D. und Pfaff, D. W. (1987): Localization of preproenkephalin mRNA in the rat brain and spinal cord by in situ hybridization, *J Comp Neurol* 258 [2], Seite 159-84.
- Haynes, A. C.; Jackson, B.; Overend, P.; Buckingham, R. E.; Wilson, S.; Tadayyon, M. und Arch, J. R. (1999): Effects of single and chronic intracerebroventricular administration of the orexins on feeding in the rat, *Peptides* 20 [9], Seite 1099-105.
- Heimer, L.; Alheid, G. F. und Zaborszky, L. (1985): Basal Ganglia. In G. Paxinos (ed): *The Rat Nervous System*, Academic Press, Sydney
- Herkenham, M. und Nauta, W. J. (1977): Afferent connections of the habenular nuclei in the rat. A horseradish peroxidase study, with a note on the fiber-of-passage problem, *J Comp Neurol*.
- Herkenham, M. und Nauta, W. J. (1979): Efferent connections of the habenular nuclei in the rat, *J Comp Neurol*.
- His, W. (1880): *Anatomie menschlicher Embryonen.*, S. Hirzel, Leipzig.

- His, W. (1895): Die anatomische Nomenclatur. Nomina anatomica, Verzeichnis der von der Anatomischen Gesellschaft auf ihrer IX. Versammlung in Basel angenommenen Namen IV, Veit, Leipzig.
- Hunt, G. E. und McGregor, I. S. (1998): Rewarding brain stimulation induces only sparse Fos-like immunoreactivity in dopaminergic neurons, *Neuroscience* 83 [2], Seite 501-15.
- Iwahori, N. (1977): A Golgi study on the habenular nucleus of the cat, *J Comp Neurol* 72 [3], Seite 319-44.
- Ji, H. und Shepard, P.D. (2005): MODELING A NEGATIVE TEMPORAL DIFFERENCE ERROR IN THE RAT Program No. 400.13. 2005 Abstract Viewer/Itinerary Planner. Washington, DC: Society for Neuroscience, 2005. Online.
- Kalen, P.; Karlson, M. und Wiklund, L. (1985): Possible excitatory amino acid afferents to nucleus raphe dorsalis of the rat investigated with retrograde wheat germ agglutinin and D-[3H]aspartate tracing, *Brain Res* 360 [1-2], Seite 285-97.
- Kalen, P.; Strecker, R. E.; Rosengren, E. und Bjorklund, A. (1989): Regulation of striatal serotonin release by the lateral habenula-dorsal raphe pathway in the rat as demonstrated by in vivo microdialysis: role of excitatory amino acids and GABA, *Brain Res* 492 [1-2], Seite 187-202.
- Kalinichev, M.; Rosenblatt, J. S.; Nakabeppu, Y. und Morrell, J. I. (2000): Induction of c-fos-like and fosB-like immunoreactivity reveals forebrain neuronal populations involved differentially in pup-mediated maternal behavior in juvenile and adult rats, *J Comp Neurol* 416 [1], Seite 45-78.
- Karreman, M.; Westerink, B. H. und Moghaddam, B. (1996): Excitatory amino acid receptors in the ventral tegmental area regulate dopamine release in the ventral striatum, *J Neurochem* 67 [2], Seite 601-7.
- Kelly, P. H. (1998): Defective inhibition of dream event memory formation: a hypothesized mechanism in the onset and progression of symptoms of schizophrenia, *Brain Res Bull* 46 [3], Seite 189-97.
- Kha, H. T.; Finkelstein, D. I.; Pow, D. V.; Lawrence, A. J. und Horne, M. K. (2000): Study of projections from the entopeduncular nucleus to the thalamus of the rat, *J Comp Neurol*.
- Kim, R.; Nakano, K.; Jayaraman, A. und Carpenter, M. B. (1976): Projections of the globus pallidus and adjacent structures: an autoradiographic study in the monkey, *J Comp Neurol* 169 [3], Seite 263-90.
- Kim, U. und Chang, S. Y. (2005): Dendritic morphology, local circuitry, and intrinsic electrophysiology of neurons in the rat medial and lateral habenular nuclei of the epithalamus, *J Comp Neurol*.
- Koelliker, A. (1861): Entwicklungsgeschichte des Menschen, S. Hirzel, Leipzig.
- Koelliker, A. (1896): Handbuch der Gewebelehre des Menschen 2, 6., umgearb. Aufl. . Auflage, Nervensystem des Menschen und der Thiere, Engelmann, Leipzig.
- Kohler, C.; Haglund, L. und Swanson, L. W. (1984): A diffuse alpha MSH-immunoreactive projection to the hippocampus and spinal cord from individual neurons in the lateral hypothalamic area and zona incerta, *J Comp Neurol* 223 [4], Seite 501-14.
- Krieg, W.J.S. (1932): The Hypothalamus Of The Albino Rat, *J Comp Neurol* 55 [1], Seite 19-89.
- Kuhlenbeck, H. (1977): The central nervous system of vertebrates Vol. 5, part I: Derivatives of the prosencephalon: Diencephalon and telencephalon, S Karger, Basel, München, Paris, London, New York, Sydney.

- Landis, C. A.; Collins, B. J.; Cribbs, L. L.; Sukhatme, V. P.; Bergmann, B. M.; Rechtschaffen, A. und Smalheiser, N. R. (1993): Expression of Egr-1 in the brain of sleep deprived rats, *Brain Res Mol Brain Res* 17 [3-4], Seite 300-6.
- Landry, M.; Trembleau, A.; Arai, R. und Calas, A. (1991): Evidence for a colocalization of oxytocin mRNA and galanin in magnocellular hypothalamic neurons: a study combining *in situ* hybridization and immunohistochemistry, *Brain Res Mol Brain Res* 10 [1], Seite 91-5.
- Larsen, K. D. und Sutin, J. (1978): Output organization of the feline entopeduncular and subthalamic nuclei, *Brain Res* 157 [1], Seite 21-31.
- Lecourtier, L. und Kelly, P. H. (2005): Bilateral lesions of the habenula induce attentional disturbances in rats, *Neuropsychopharmacology*.
- Lecourtier, L.; Neijt, H. C. und Kelly, P. H. (2004): Habenula lesions cause impaired cognitive performance in rats: implications for schizophrenia, *Eur J Neurosci*.
- Lee, E. H. und Huang, S. L. (1988): Role of lateral habenula in the regulation of exploratory behavior and its relationship to stress in rats, *Behav Brain Res* 30 [3], Seite 265-71.
- Lehmann, A. ; Geisler, S.; Schneider, K. und Veh, R.W. (2002): INDIVIDUAL SUBNUCLEI OF THE LATERAL HABENULAR COMPLEX DISTINCTLY PROJECT TO THE DORSAL AND MEDIAN RAPHE NUCLEI, Program No. 587.1. 2002 Abstract Viewer/Itinerary Planner. Washington, DC: Society for Neuroscience, 2002. Online.
- Lehmann, A.; Geisler, S. und Veh, R.W. (2003): MEDIAN AND PARAMEDIAN RAPHE NUCLEI SHOW DISTINCT PATTERNS OF INNERVATION BY THE SUBNUCLEI OF THE LATERAL HABENULAR COMPLEX IN RAT, Program No. 963.16. 2003 Abstract Viewer/Itinerary Planner. Washington, DC: Society for Neuroscience, 2003. Online.
- Levin, M. C.; Sawchenko, P. E.; Howe, P. R.; Bloom, S. R. und Polak, J. M. (1987): Organization of galanin-immunoreactive inputs to the paraventricular nucleus with special reference to their relationship to catecholaminergic afferents, *J Comp Neurol* 261 [4], Seite 562-82.
- Li, T.; Gao, W. und Rao, Z. R. (1998): Noxious somatic stimulation-induced expression of Fos-like immunoreactivity in catecholaminergic neurons with habenular nucleus projection in the medullary visceral zone of rat, *Brain Res*.
- Liu, G. J. und Wang, S. (1988): [Effects of nucleus raphe magnus and locus coeruleus in descending modulation of the habenula on pain threshold and acupuncture analgesia], *Zhongguo Yao Li Xue Bao* 9 [1], Seite 18-22.
- Ma, Q. P.; Shi, Y. S. und Han, J. S. (1992): Further studies on interactions between periaqueductal gray, nucleus accumbens and habenula in antinociception, *Brain Res* 583 [1-2], Seite 292-5.
- Marcus, J. N.; Aschkenasi, C. J.; Lee, C. E.; Chemelli, R. M.; Saper, C. B.; Yanagisawa, M. und Elmquist, J. K. (2001): Differential Expression of Orexin Receptors 1 and 2 in the Rat Brain, *J Comp Neurol* 435, Seite 6-25.
- Matsumoto, N.; Yahata, F.; Kawarada, K.; Kamata, K. und Suzuki, T. A. (1994): Tooth pulp stimulation induces c-fos expression in the lateral habenular nucleus of the cat, *Neuroreport* 5 [17], Seite 2397-400.
- Matthews-Felton, T.; Corodimas, K. P.; Rosenblatt, J. S. und Morrell, J. I. (1995): Lateral habenula neurons are necessary for the hormonal onset of maternal behavior and for the display of postpartum estrus in naturally parturient female rats, *Behav Neurosci* 109 [6], Seite 1172-88.

- Matthews-Felton, T.; Linton, L.; Rosenblatt, J. S. und Morell, J. I. (1999): First and second order maternal behavior related afferents of the lateral habenula, *Neuroreport*.
- McBride, R. L. (1981): Organization of afferent connections of the feline lateral habenular nucleus, *J Comp Neurol* 198 [1], Seite 89-99.
- McCulloch, J.; Savaki, H. E. und Sokoloff, L. (1980): Influence of dopaminergic systems on the lateral habenular nucleus of the rat, *Brain Res* 194 [1], Seite 117-24.
- McMullen, N. T. und Almli, C. R. (1981): Cell types within the medial forebrain bundle: a Golgi study of preoptic and hypothalamic neurons in the rat, *Am J Anat* 161 [3], Seite 323-40.
- Melander, T.; Hokfelt, T. und Rokaeus, A. (1986): Distribution of galaninlike immunoreactivity in the rat central nervous system, *J Comp Neurol* 248 [4], Seite 475-517.
- Melrose, P. A. und Knigge, K. M. (1988): Comparative topography of the immunoreactive alpha-melanocyte-stimulating hormone neuronal systems in the brains of horses and rats, *Brain Behav Evol* 32 [4], Seite 226-35.
- Meshul, C. K.; Noguchi, K.; Emre, N. und Ellison, G. (1998): Cocaine-induced changes in glutamate and GABA immunolabeling within rat habenula and nucleus accumbens, *Synapse*.
- Meynert, T. (1870): Beiträge zur Kenntniss der centralen Projection der Sinnesoberflächen, *Sitzungberichte der Kaiserlichen Akademie der Wissenschaften, Wien. Mathematish-Naturwissenschaftliche Classe* 60, Seite 547-562.
- Miller, M. A.; Kolb, P. E. und Raskind, M. A. (1993): Extra-hypothalamic vasopressin neurons coexpress galanin messenger RNA as shown by double in situ hybridization histochemistry, *J Comp Neurol* 329 [3], Seite 378-84.
- Milner, T. A. und Pickel, V. M. (1986): Ultrastructural localization and afferent sources of substance P in the rat parabrachial region, *Neuroscience* 17 [3], Seite 687-707.
- Morgan, M. J.; O'Donnell, J. M. und Oliver, R. F. (1973): Development of left-right asymmetry in the habenular nuclei of *Rana temporaria*, *J Comp Neurol* 149 [2], Seite 203-14.
- Morris, J. S.; Smith, K. A.; Cowen, P. J.; Friston, K. J. und Dolan, R. J. (1999): Covariation of activity in habenula and dorsal raphe nuclei following tryptophan depletion, *Neuroimage* 10 [2], Seite 163-72.
- Murphy, C. A.; Ghazi, L.; Kokabi, A. und Ellison, G. (1999): Prenatal cocaine produces signs of neurodegeneration in the lateral habenula, *Brain Res.*
- Nagao, M.; Kamo, H.; Akiguchi, I. und Kimura, J. (1993): Induction of c-Fos-like protein in the lateral habenular nucleus by persistent noxious peripheral stimulation, *Neurosci Lett* 151 [1], Seite 37-40.
- Nambu, T.; Sakurai, T.; Mizukami, K.; Hosoya, Y.; Yanagisawa, M. und Goto, K. (1999): Distribution of orexin neurons in the adult rat brain, *Brain Res* 827 [1-2], Seite 243-60.
- Nauta, H. J. (1974): Evidence of a pallidohabenular pathway in the cat, *J Comp Neurol* 156 [1], Seite 19-27.
- Neal, C. R., Jr.; Mansour, A.; Reinscheid, R.; Nothacker, H. P.; Civelli, O.; Akil, H. und Watson, S. J., Jr. (1999): Opioid receptor-like (ORL1) receptor distribution in the rat central nervous system: comparison of ORL1 receptor mRNA expression with (125)I-[(14)Tyr]-orphanin FQ binding, *J Comp Neurol* 412 [4], Seite 563-605.
- Nieuwenhuys, R.; Geeraedts, L. M. und Veening, J. G. (1982): The medial forebrain bundle of the rat. I. General introduction, *J Comp Neurol* 206 [1], Seite 49-81.

- Nissl, F. (1913): Die Großhirnanteile des Kanninchens, Arch f Psychiatr Nervenkr 52, Seite 867-953.
- Olszewski, P. K.; Billington, C. J. und Levine, A. S. (2000): Fos expression in feeding-related brain areas following intracerebroventricular administration of orphanin FQ in rats, Brain Res 855 [1], Seite 171-5.
- Olszewski, P. K.; Grace, M. K.; Billington, C. J. und Levine, A. S. (2000): The effect of [Phe(1)psi(CH(2)-NH)Gly(2)]-nociceptin(1-13)NH(2) on feeding and c-Fos immunoreactivity in selected brain sites, Brain Res 876 [1-2], Seite 95-102.
- Pain, Y. Z.; Wang, L. H.; Tang, Y. H.; Yin, X. M. und Wang, S. (1992): [Antagonistic effect of electro-acupuncture analgesia with Ca²⁺ injection into habenula could be reversed by gallamine triethiodide], Sheng Li Xue Bao 44 [4], Seite 326-32.
- Palkovits, M. und Van Cuc, H. (1980): Quantitative light and electron microscopic studies on the lateral hypothalamus in rat. Cell and synaptic densities, Brain Res Bull 5 [6], Seite 643-7.
- Panula, P.; Yang, H. Y. und Costa, E. (1984): Comparative distribution of bombesin/GRP- and substance-P-like immunoreactivities in rat hypothalamus, J Comp Neurol 224 [4], Seite 606-17.
- Parent, A. (1979): Identification of the pallidal and peripallidal cells projecting to the habenula in monkey, Neurosci Lett 15 [2-3], Seite 159-64.
- Parent, A. (1986): Comparative Neurobiology of the Basal Ganglia, Wiley-Interscience, New York.
- Parent, A.; Gravel, S. und Boucher, R. (1981): The origin of forebrain afferents to the habenula in rat, cat and monkey, Brain Res Bull 6 [1], Seite 23-38.
- Parent, M.; Levesque, M. und Parent, A. (2001): Two types of projection neurons in the internal pallidum of primates: single-axon tracing and three-dimensional reconstruction, J Comp Neurol.
- Pavel, S. und Eisner, C. (1984): A GABAergic habenulo-raphe pathway mediates both serotonergic and hypnogenic effects of vasotocin in cats, Brain Res Bull 13 [5], Seite 623-7.
- Paxinos, G. und Watson, C. (1986): The Rat Brain in Stereotaxic Coordinates, Academic Press, Sydney.
- Paxinos, G. und Watson, C. (1989): The Rat Brain in Stereotaxic Coordinates, Academic Press, Sydney.
- Phillipson, O. T. und Pycock, C. J. (1982): Dopamine neurones of the ventral tegmentum project to both medial and lateral habenula. Some implications for habenular function, Exp Brain Res.
- Planas, B.; Kolb, P. E.; Raskind, M. A. und Miller, M. A. (1995): Sex difference in coexpression by galanin neurons accounts for sexual dimorphism of vasopressin in the bed nucleus of the stria terminalis, Endocrinology 136 [2], Seite 727-33.
- Pomonis, J. D.; Billington, C. J. und Levine, A. S. (1996): Orphanin FQ, agonist of orphan opioid receptor ORL1, stimulates feeding in rats, Neuroreport 8 [1], Seite 369-71.
- Qu, D.; Ludwig, D. S.; Gammeltoft, S.; Piper, M.; Pelleymounter, M. A.; Cullen, M. J.; Mathes, W. F.; Przypek, R.; Kanarek, R. und Maratos-Flier, E. (1996): A role for melanin-concentrating hormone in the central regulation of feeding behaviour, Nature 380 [6571], Seite 243-7.
- Ragsdale, C. W. und Graybiel, A. M. (1988): Fibers from the basolateral nucleus of the amygdala selectively innervate striosomes in the caudate nucleus of the rat, J Comp Neurol 269, Seite 506-522.
- Rajakumar, N.; Elisevich, K. und Flumerfelt, B. A. (1993): Compartmental Origin of the Striato-Entopeduncular Projection in the Rat, J Comp Neurol 331.

- Rajakumar, N.; Elisevich, K. und Flumerfelt, B. A. (1994): Parvalbumin-containing GABAergic neurons in the basal ganglia output system of the rat, J Comp Neurol 350 [2], Seite 324-36.
- Ruggiero, D. A.; Ross, C. A.; Anwar, M.; Park, D. H.; Joh, T. H. und Reis, D. J. (1985): Distribution of neurons containing phenylethanolamine N-methyltransferase in medulla and hypothalamus of rat, J Comp Neurol 239 [2], Seite 127-54.
- Sakanaka, M.; Shibasaki, T. und Lederis, K. (1987): Corticotropin releasing factor-like immunoreactivity in the rat brain as revealed by a modified cobalt-glucose oxidase-diaminobenzidine method, J Comp Neurol 260 [2], Seite 256-98.
- Sakanaka, M.; Shiosaka, S.; Takatsuki, K.; Inagaki, S.; Hara, Y.; Kawai, Y.; Senba, E. und Tohyama, M. (1982): Origins of substance P-containing fibers in the lateral septal area of young rats: immunohistochemical analysis of experimental manipulations, J Comp Neurol 212 [3], Seite 268-77.
- Sakurai, T. (1999): Orexins and orexin receptors: implication in feeding behavior, Regul Pept 85 [1], Seite 25-30.
- Sandyk, R. (1992): Pineal and habenula calcification in schizophrenia, Int J Neurosci 67 [1-4], Seite 19-30.
- Saper, C. P.; Swanson, L. W. und Cowan, W. M. (1979): An autoradiographic study of the efferent connections of the lateral hypothalamic area in the rat, J Comp Neurol 183, Seite 689-706.
- Satoh, K. und Fibiger, H. C. (1986): Cholinergic neurons of the laterodorsal tegmental nucleus: efferent and afferent connections, J Comp Neurol 253 [3], Seite 277-302.
- Semba, K. und Fibiger, H. C. (1992): Afferent connections of the laterodorsal and the pedunculopontine tegmental nuclei in the rat: a retro- and antero-grade transport and immunohistochemical study, J Comp Neurol 323 [3], Seite 387-410.
- Serres, E.R.A. (1827): Anatomie comparée du cerveau, dans les quatre classes des animaux vertébrés, Gabon, Paris.
- Shepard, P. D.; Holcomb, H. H. und Gold, J. M. (2006): Schizophrenia in translation: the presence of absence: habenular regulation of dopamine neurons and the encoding of negative outcomes, Schizophr Bull 32 [3], Seite 417-21.
- Sherin, J. E.; Elmquist, J. K.; Torrealba, F. und Saper, C. B. (1998): Innervation of histaminergic tuberomammillary neurons by GABAergic and galaninergic neurons in the ventrolateral preoptic nucleus of the rat, J Neurosci 18, Seite 4705-4721.
- Sipe, J. C. und Moore, R. Y. (1977): The lateral hypothalamic area. An ultrastructural analysis, Cell Tissue Res 179 [2], Seite 177-96.
- Skagerberg, G.; Lindvall, O. und Björklund, A. (1984): Origin, course and termination of the mesohabenular dopamine pathway in the rat, Brain Res 307 [1-2], Seite 99-108.
- Sridhya, R.; Mallick, H. N. und Kumar, V. M. (2006): Differences in the effects of medial and lateral preoptic lesions on thermoregulation and sleep in rats, Neuroscience 139 [3], Seite 853-64.
- Stieda, L. (1868): Studien ueber das centrale Nervensystem der Knochenfische, Leipzig
- Stricker, E. M.; Swerdloff, A. F. und Zigmond, M. J. (1978): Intrahypothalamic injections of kainic acid produce feeding and drinking deficits in rats, Brain Res 158 [2], Seite 470-3.
- Sutherland, R. J. (1982): The dorsal diencephalic conduction system: a review of the anatomy and functions of the habenular complex, Neurosci Biobehav Rev 6 [1], Seite 1-13.

- Sutherland, R. J. und Nakajima, S. (1981): Self-stimulation of the habenular complex in the rat, *J Comp Physiol Psychol* 95 [5], Seite 781-91.
- Swanson, L. W. (1992): Brain Maps: Structure of the Rat Brain., Elsevier, Amsterdam.
- Swanson, L. W. (1996): Brain Maps: Structure of the Rat Brain., Elsevier, Amsterdam.
- Swanson, L. W. (2004): Brain Maps: Structure of the Rat Brain., Elsevier, Amsterdam.
- Swanson, L. W. und Cowan, W. M. (1979): The connections of the septal region in the rat, *J Comp Neurol* 186, Seite 621-656.
- Tavakoli-Nezhad, M. und Schwartz, W. J. (2005): c-Fos expression in the brains of behaviorally "split" hamsters in constant light: calling attention to a dorsolateral region of the suprachiasmatic nucleus and the medial division of the lateral habenula, *J Biol Rhythms*.
- Terenzi, M. G.; Guimaraes, F. S. und Prado, W. A. (1990): Antinociception induced by stimulation of the habenular complex of the rat, *Brain Res* 524 [2], Seite 213-8.
- Terenzi, M. G. und Prado, W. A. (1990): Antinociception elicited by electrical or chemical stimulation of the rat habenular complex and its sensitivity to systemic antagonists, *Brain Res* 535 [1], Seite 18-24.
- Tronel, S. und Sara, S. J. (2002): Mapping of olfactory memory circuits: region-specific c-fos activation after odor-reward associative learning or after its retrieval, *Learn Mem* 9 [3], Seite 105-11.
- Ullsperger, M. und von Cramon, D. Y. (2003): Error monitoring using external feedback: specific roles of the habenular complex, the reward system, and the cingulate motor area revealed by functional magnetic resonance imaging, *J Neurosci*.
- Vachon, M. P. und Miliaressis, E. (1992): Dorsal diencephalic self-stimulation: a movable electrode mapping study, *Behav Neurosci* 106 [6], Seite 981-91.
- van der Kooy, D. und Carter, D. A. (1981): The organization of the efferent projections and striatal afferents of the entopeduncular nucleus and adjacent areas in the rat, *Brain Res* 211 [1], Seite 15-36.
- Veening, J. G.; Swanson, L. W.; Cowan, W. M.; Nieuwenhuys, R. und Geeraedts, L. M. (1982): The medial forebrain bundle of the rat. II. An autoradiographic study of the topography of the major descending and ascending components, *J Comp Neurol* 206 [1], Seite 82-108.
- Vercelli, A.; Repici, M.; Garbossa, D. und Grimaldi, A. (2000): Recent techniques for tracing pathways in the central nervous system of developing and adult mammals, *Brain Res Bull* 51 [1], Seite 11-28.
- Villarreal, J. S.; Gonzalez-Lima, F.; Berndt, J. und Barea-Rodriguez, E. J. (2002): Water maze training in aged rats: effects on brain metabolic capacity and behavior, *Brain Res* 939 [1-2], Seite 43-51.
- Vincent, S. R. und Brown, J. C. (1986): Somatostatin immunoreactivity in the entopeduncular projection to the lateral habenula in the rat, *Neurosci Lett* 68 [2], Seite 160-4.
- Vincent, S. R.; Kimura, H. und McGeer, E. G. (1982): A histochemical study of GABA-transaminase in the efferents of the pallidum, *Brain Res* 241 [1], Seite 162-5.
- Vincent, S. R.; McIntosh, C. H.; Buchan, A. M. und Brown, J. C. (1985): Central somatostatin systems revealed with monoclonal antibodies, *J Comp Neurol* 238 [2], Seite 169-86.
- Wagner, C. K.; Silverman, A. J. und Morrell, J. I. (1998): Evidence for estrogen receptor in cell nuclei and axon terminals within the lateral habenula of the rat: regulation during pregnancy, *J Comp Neurol*.
- Wang, S.; Liu, G. J.; Gao, Y. L. und Tang, Y. H. (1987): [Effects of exciting the habenula by L-glutamic acid on the pain threshold and acupuncture analgesia], *Zhongguo Yao Li Xue Bao* 8 [3], Seite 200-3.

- Wernicke, C. (1881): Lehrbuch der Gehirnerkrankungen, Kassel.
- Wilcox, K. S.; Christoph, G. R.; Double, B. A. und Leonzio, R. J. (1986): Kainate and electrolytic lesions of the lateral habenula: effect on avoidance responses, *Physiol Behav* 36 [3], Seite 413-7.
- Winn, P. (2006): How best to consider the structure and function of the pedunculopontine tegmental nucleus: Evidence from animal studies, *J Neurol Sci.*
- Wirtshafter, D. (2005): Cholinergic involvement in the cortical and hippocampal Fos expression induced in the rat by placement in a novel environment, *Brain Res* 1051 [1-2], Seite 57-65.
- Wirtshafter, D.; Asin, K. E. und Pitzer, M. R. (1994): Dopamine agonists and stress produce different patterns of Fos-like immunoreactivity in the lateral habenula, *Brain Res* 633 [1-2], Seite 21-6.
- Wirtshafter, D. und Krebs, J. C. (1997): Interactive effects of stimulation of D1 and D2 dopamine receptors on Fos expression in the lateral habenula, *Brain Res.*
- Yanez, J. und Anadon, R. (1994): Afferent and efferent connections of the habenula in the larval sea lamprey (*Petromyzon marinus* L.): an experimental study, *J Comp Neurol* 345 [1], Seite 148-60.
- Yanez, J. und Anadon, R. (1996): Afferent and efferent connections of the habenula in the rainbow trout (*Oncorhynchus mykiss*): an indocarbocyanine dye (Dil) study, *J Comp Neurol.*
- Yeomans, J. S. (1995): Role of tegmental cholinergic neurons in dopaminergic activation, antimuscarinic psychosis and schizophrenia, *Neuropsychopharmacology* 12, Seite 3-16.
- Yu, L. C. und Han, J. S. (1990): Habenula as a relay in the descending pathway from nucleus accumbens to periaqueductal grey subserving antinociception, *Int J Neurosci* 54 [3-4], Seite 245-51.
- Yuan, P. Q.; Granas, C.; Kallstrom, L.; Yu, J.; Huhman, K.; Larhammar, D.; Albers, H. E. und Johnson, A. E. (1997): Differential distribution of glutamic acid decarboxylase-65 and glutamic acid decarboxylase-67 messenger RNAs in the entopeduncular nucleus of the rat, *Neuroscience*.
- Ziehen, G.T. (1901): Zoologische Forschungsreisen in Australien und dem Malayischen Archipel. Dritter Band: Monotremen und Marsupialier. Das Zentralnervensystem der Monotremen und Marsupialier. 2. Teil: Makroskopische Anatomie. 1. Abschnitt: Der Faserverlauf im Hirnstamm von *Pseudochirus peregrinus*., II. Denkschriften der Medizinisch-Naturwissenschaftlichen Gesellschaft zu Jena. Gustav Fischer Verlag, Jena, ISBN: 3334-605817

Publikationen

A. Kowski, S. Geisler and R.W. Veh. THE SUBNUCLEI OF THE LATERAL HABENULAR COMPLEX ARE INDIVIDUALLY TARGETED FROM LATERAL PREOPTIC SUBAREAS. 99th Annual Meeting der Anatomischen Gesellschaft, Würzburg, 2004.

A. Kowski, S. Geisler, and R.W. Veh. THE LATERAL PREOPTIC AREA PROJECTS TO INDIVIDUAL SUBNUCLEI OF THE LATERAL HABENULAR COMPLEX. 98th Annual Meeting der Anatomischen Gesellschaft, Würzburg 2003.

A. Kowski, S. Geisler, A. Goertzen, R.W. Veh. The LATERAL PREOPTIC AREA AND THE LATERAL HYPOTHALMUS PROJECT TO INDIVIDUAL SUBNUCLEI OF THE LATERAL HABENULAR COMPLEX. 33th Meeting of the Society of Neuroscience, New Orleans, USA, 2003.

A. Kowski, S. Geisler, R.W. Veh. THE OVAL SUBNUCLEUS OF THE LATERAL HABENULA IS A MAIN TARGET OF AFFERENTS FROM THE ENTOPEDUNCULAR NUCLEUS IN THE RAT. 34th Meeting of the Society of Neuroscience, San Diego, USA, 2004.

Kowski A et al., „DIFFERENTIAL PROJECTIONS FROM SUBFIELDS IN THE LATERAL PREOPTIC AREA TO THE LATERAL HABENULAR COMPLEX OF THE RAT“, J Comp Neurol., 2007.

Krauss M, Weiss T, Langnaese K, Richter K, Kowski A, Veh RW, Laube G., „Cellular and subcellular rat brain spermidine synthase expression patterns suggest region-specific roles for polyamines, including cerebellar pre-synaptic function.“ J Neurochem., 2007.

Gruber C, Kahl A, Lebenheim L, Kowski A, Dittgen A, Veh RW, „Dopaminergic projections from the VTA substantially contribute to the mesohabenular pathway in the rat.“, Neurosci Lett., 2007.

Kowski A et. al., „Dopaminergic activation excites rat lateral habenula neurons in vivo“ in Vorbereitung.