

6. Literatur

- Barrow, J.R., H.S. Stadler, and M.R. Capecchi. 2000. Roles of Hoxa1 and Hoxa2 in patterning the early hindbrain of the mouse. *Development* **127**: 933-44.
- Bell, E., R.J. Wingate, and A. Lumsden. 1999. Homeotic transformation of rhombomere identity after localized Hoxb1 misexpression. *Science* **284**: 2168-71.
- Birnboim, H.C. and J. Doly. 1979. A rapid alkaline extraction procedure for screening recombinant plasmid DNA. *Nucleic Acids Res* **7**: 1513-23.
- Briscoe, J. and J. Ericson. 2001. Specification of neuronal fates in the ventral neural tube. *Curr Opin Neurobiol* **11**: 43-9.
- Briscoe, J., A. Pierani, T.M. Jessell, and J. Ericson. 2000. A homeodomain protein code specifies progenitor cell identity and neuronal fate in the ventral neural tube. *Cell* **101**: 435-45.
- Brohmann, H., K. Jagla, and C. Birchmeier. 2000. The role of Lbx1 in migration of muscle precursor cells. *Development* **127**: 437-45.
- Carpenter, E.M., J.M. Goddard, O. Chisaka, N.R. Manley, and M.R. Capecchi. 1993. Loss of Hox-A1 (Hox-1.6) function results in the reorganization of the murine hindbrain. *Development* **118**: 1063-75.
- Chen, F., K.C. Liu, and J.A. Epstein. 1999. Lbx2, a novel murine homeobox gene related to the Drosophila ladybird genes is expressed in the developing urogenital system, eye and brain. *Mech Dev* **84**: 181-4.
- Cheng, L., A. Arata, R. Mizuguchi, Y. Qian, A. Karunaratne, P.A. Gray, S. Arata, S. Shirasawa, M. Bouchard, P. Luo, C.L. Chen, M. Busslinger, M. Goulding, H. Onimaru, and Q. Ma. 2004a. Tlx3 and Tlx1 are post-mitotic selector genes determining glutamatergic over GABAergic cell fates. *Nat Neurosci*.
- . 2004b. Tlx3 and Tlx1 are post-mitotic selector genes determining glutamatergic over GABAergic cell fates. *Nat Neurosci* **7**: 510-517.
- Chi, C.L., S. Martinez, W. Wurst, and G.R. Martin. 2003. The isthmus organizer signal FGF8 is required for cell survival in the prospective midbrain and cerebellum. *Development* **130**: 2633-44.
- Chiang, C., Y. Litingtung, E. Lee, K.E. Young, J.L. Corden, H. Westphal, and P.A. Beachy. 1996. Cyclopia and defective axial patterning in mice lacking Sonic hedgehog gene function. *Nature* **383**: 407-13.
- Clark, G. 1981. Staining Procedures. In (ed. J.W. Bartholomew, C.G. E., C. G., C.R. E., D.W. J., K.F. H., M.J. L., P.R. L., S. H., and S. S.S.), pp. 512. Williams & Wilkins, Baltimore, MD 21202, USA.

- Dauger, S., F. Guimiot, S. Renolleau, B. Levacher, B. Boda, C. Mas, V. Nepote, M. Simonneau, C. Gaultier, and J. Gallego. 2001. MASH-1/RET pathway involvement in development of brain stem control of respiratory frequency in newborn mice. *Physiol Genomics* **7**: 149-57.
- Dauger, S., S. Renolleau, G. Vardon, V. Nepote, C. Mas, M. Simonneau, C. Gaultier, and J. Gallego. 1999. Ventilatory responses to hypercapnia and hypoxia in Mash-1 heterozygous newborn and adult mice. *Pediatr Res* **46**: 535-42.
- de Diego, I., K. Kyriakopoulou, D. Karageorgos, and M. Wassef. 2002. Multiple influences on the migration of precerebellar neurons in the caudal medulla. *Development* **129**: 297-306.
- Dolle, P., T. Lufkin, R. Krumlauf, M. Mark, D. Duboule, and P. Chambon. 1993. Local alterations of Krox-20 and Hox gene expression in the hindbrain suggest lack of rhombomeres 4 and 5 in homozygote null Hoxa-1 (Hox-1.6) mutant embryos. *Proc Natl Acad Sci U S A* **90**: 7666-70.
- Dupe, V., M. Davenne, J. Brocard, P. Dolle, M. Mark, A. Dierich, P. Chambon, and F.M. Rijli. 1997. In vivo functional analysis of the Hoxa-1 3' retinoic acid response element (3'RARE). *Development* **124**: 399-410.
- Ericson, J., S. Morton, A. Kawakami, H. Roelink, and T.M. Jessell. 1996. Two critical periods of Sonic Hedgehog signaling required for the specification of motor neuron identity. *Cell* **87**: 661-73.
- Ericson, J., J. Muhr, T.M. Jessell, and T. Edlund. 1995a. Sonic hedgehog: a common signal for ventral patterning along the rostrocaudal axis of the neural tube. *Int J Dev Biol* **39**: 809-16.
- Ericson, J., J. Muhr, M. Placzek, T. Lints, T.M. Jessell, and T. Edlund. 1995b. Sonic hedgehog induces the differentiation of ventral forebrain neurons: a common signal for ventral patterning within the neural tube. *Cell* **81**: 747-56.
- Feinberg, A.P. and B. Vogelstein. 1983. A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Anal Biochem* **132**: 6-13.
- Feldman, J.L., G.S. Mitchell, and E.E. Nattie. 2003. Breathing: rhythmicity, plasticity, chemosensitivity. *Annu Rev Neurosci* **26**: 239-66.
- Fraser, S., R. Keynes, and A. Lumsden. 1990. Segmentation in the chick embryo hindbrain is defined by cell lineage restrictions. *Nature* **344**: 431-5.
- Frohman, M.A., G.R. Martin, S.P. Cordes, L.P. Halamek, and G.S. Barsh. 1993. Altered rhombomere-specific gene expression and hyoid bone differentiation in the mouse segmentation mutant, kreisler (kr). *Development* **117**: 925-36.
- Gaufo, G.O., S. Wu, and M.R. Capecchi. 2004. Contribution of Hox genes to the diversity of the hindbrain sensory system. *Development* **131**: 1259-66.

- Gavalas, A., M. Studer, A. Lumsden, F.M. Rijli, R. Krumlauf, and P. Chambon. 1998. Hoxa1 and Hoxb1 synergize in patterning the hindbrain, cranial nerves and second pharyngeal arch. *Development* **125**: 1123-36.
- Gavrieli, Y., Y. Sherman, and S.A. Ben-Sasson. 1992. Identification of programmed cell death in situ via specific labeling of nuclear DNA fragmentation. *J Cell Biol* **119**: 493-501.
- Ghysen, A. and C. Dambly-Chaudiere. 1990. Early events in the development of Drosophila peripheral nervous system. *J Physiol (Paris)* **84**: 11-20.
- Goddard, J.M., M. Rossel, N.R. Manley, and M.R. Capecchi. 1996. Mice with targeted disruption of Hoxb-1 fail to form the motor nucleus of the VIIth nerve. *Development* **122**: 3217-28.
- Gowan, K., A.W. Helms, T.L. Hunsaker, T. Collisson, P.J. Ebert, R. Odom, and J.E. Johnson. 2001. Crossinhibitory activities of Ngn1 and Math1 allow specification of distinct dorsal interneurons. *Neuron* **31**: 219-32.
- Gross, M.K., M. Dottori, and M. Goulding. 2002. Lbx1 specifies somatosensory association interneurons in the dorsal spinal cord. *Neuron* **34**: 535-49.
- Guillemot, F., L.C. Lo, J.E. Johnson, A. Auerbach, D.J. Anderson, and A.L. Joyner. 1993. Mammalian achaete-scute homolog 1 is required for the early development of olfactory and autonomic neurons. *Cell* **75**: 463-76.
- Hirsch, M.R., M.C. Tiveron, F. Guillemot, J.F. Brunet, and C. Goridis. 1998. Control of noradrenergic differentiation and Phox2a expression by MASH1 in the central and peripheral nervous system. *Development* **125**: 599-608.
- Hogan, B., R. Beddington, F. Constantini, and E. Lacy. 1994. *Manipulating The Mouse Embryo - A Laboratory Manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York 11803-2500, USA.
- Innis, M.A., D.H. Gelfand, and J.J. Sninsky. 1989. PCR Protocols: A Guide To Methods And Applications. In, pp. 482. Academic Press, San Diego, CA 92101-4495, USA.
- Inoue, H., H. Nojima, and H. Okayama. 1990. High efficiency transformation of Escherichia coli with plasmids. *Gene* **96**: 23-8.
- Jagla, K., P. Dolle, M.G. Mattei, T. Jagla, B. Schuhbauer, G. Dretzen, F. Bellard, and M. Bellard. 1995. Mouse Lbx1 and human LBX1 define a novel mammalian homeobox gene family related to the Drosophila lady bird genes. *Mech Dev* **53**: 345-56.
- Jessell, T.M. 2000. Neuronal specification in the spinal cord: inductive signals and transcriptional codes. *Nat Rev Genet* **1**: 20-9.
- Lee, K.J., P. Dietrich, and T.M. Jessell. 2000. Genetic ablation reveals that the roof plate is essential for dorsal interneuron specification. *Nature* **403**: 734-40.
- Lee, K.J. and T.M. Jessell. 1999. The specification of dorsal cell fates in the vertebrate central nervous system. *Annu Rev Neurosci* **22**: 261-94.

- Lee, K.J., M. Mendelsohn, and T.M. Jessell. 1998. Neuronal patterning by BMPs: a requirement for GDF7 in the generation of a discrete class of commissural interneurons in the mouse spinal cord. *Genes Dev* **12**: 3394-407.
- Logan, C., R.J. Wingate, I.J. McKay, and A. Lumsden. 1998. Tlx-1 and Tlx-3 homeobox gene expression in cranial sensory ganglia and hindbrain of the chick embryo: markers of patterned connectivity. *J Neurosci* **18**: 5389-402.
- Lumsden, A. and R. Keynes. 1989. Segmental patterns of neuronal development in the chick hindbrain. *Nature* **337**: 424-8.
- Lumsden, A. and R. Krumlauf. 1996. Patterning the vertebrate neuraxis. *Science* **274**: 1109-15.
- Lumsden, T. 1923. Observations on the respiratory centers in the cat. *The Journal of Physiology* **57**: 153-160.
- Ma, Q., L. Sommer, P. Cserjesi, and D.J. Anderson. 1997. Mash1 and neurogenin1 expression patterns define complementary domains of neuroepithelium in the developing CNS and are correlated with regions expressing notch ligands. *J Neurosci* **17**: 3644-52.
- Maconochie, M.K., S. Nonchev, M. Studer, S.K. Chan, H. Popperl, M.H. Sham, R.S. Mann, and R. Krumlauf. 1997. Cross-regulation in the mouse HoxB complex: the expression of Hoxb2 in rhombomere 4 is regulated by Hoxb1. *Genes Dev* **11**: 1885-95.
- Manzanares, M., S. Bel-Vialar, L. Ariza-McNaughton, E. Ferretti, H. Marshall, M.M. Maconochie, F. Blasi, and R. Krumlauf. 2001. Independent regulation of initiation and maintenance phases of Hoxa3 expression in the vertebrate hindbrain involve auto- and cross-regulatory mechanisms. *Development* **128**: 3595-607.
- Manzanares, M., P.A. Trainor, S. Nonchev, L. Ariza-McNaughton, J. Brodie, A. Gould, H. Marshall, A. Morrison, C.T. Kwan, M.H. Sham, D.G. Wilkinson, and R. Krumlauf. 1999. The role of kreisler in segmentation during hindbrain development. *Dev Biol* **211**: 220-37.
- Marin, F. and L. Puelles. 1995. Morphological fate of rhombomeres in quail/chick chimeras: a segmental analysis of hindbrain nuclei. *Eur J Neurosci* **7**: 1714-38.
- Mark, M., T. Lufkin, P. Dolle, A. Dierich, M. LeMeur, and P. Chambon. 1993. Roles of Hox genes: what we have learnt from gain of function and loss of function mutations in the mouse. *C R Acad Sci III* **316**: 995-1008.
- McDonnell, M.W., M.N. Simon, and F.W. Studier. 1977. Analysis of restriction fragments of T7 DNA and determination of molecular weights by electrophoresis in neutral and alkaline gels. *J Mol Biol* **110**: 119-46.

- McGinnis, W. and R. Krumlauf. 1992. Homeobox genes and axial patterning. *Cell* **68**: 283-302.
- McKay, I.J., I. Muchamore, R. Krumlauf, M. Maden, A. Lumsden, and J. Lewis. 1994. The kreisler mouse: a hindbrain segmentation mutant that lacks two rhombomeres. *Development* **120**: 2199-2111.
- Mizuguchi, R., M. Sugimori, H. Takebayashi, H. Kosako, M. Nagao, S. Yoshida, Y. Nabeshima, K. Shimamura, and M. Nakafuku. 2001. Combinatorial roles of olig2 and neurogenin2 in the coordinated induction of pan-neuronal and subtype-specific properties of motoneurons. *Neuron* **31**: 757-71.
- Morin, X., H. Cremer, M.R. Hirsch, R.P. Kapur, C. Goridis, and J.F. Brunet. 1997. Defects in sensory and autonomic ganglia and absence of locus coeruleus in mice deficient for the homeobox gene Phox2a. *Neuron* **18**: 411-23.
- Muller, T., H. Brohmann, A. Pierani, P.A. Heppenstall, G.R. Lewin, T.M. Jessell, and C. Birchmeier. 2002. The homeodomain factor lbx1 distinguishes two major programs of neuronal differentiation in the dorsal spinal cord. *Neuron* **34**: 551-62.
- Novitch, B.G., A.I. Chen, and T.M. Jessell. 2001. Coordinate Regulation of Motor Neuron Subtype Identity and Pan-Neuronal Properties by the bHLH Repressor Olig2. *Neuron* **31**: 773-89.
- Ogura, T. and R.M. Evans. 1995. A retinoic acid-triggered cascade of HOXB1 gene activation. *Proc Natl Acad Sci U S A* **92**: 387-91.
- Olszewski J, B.D. 1982. Cytoarchitecture of the human Brain Stem. 83-91.
- Pattyn, A., C. Goridis, and J.F. Brunet. 2000a. Specification of the central noradrenergic phenotype by the homeobox gene Phox2b. *Mol Cell Neurosci* **15**: 235-43.
- Pattyn, A., M. Hirsch, C. Goridis, and J.F. Brunet. 2000b. Control of hindbrain motor neuron differentiation by the homeobox gene Phox2b. *Development* **127**: 1349-58.
- Pattyn, A., X. Morin, H. Cremer, C. Goridis, and J.F. Brunet. 1997. Expression and interactions of the two closely related homeobox genes Phox2a and Phox2b during neurogenesis. *Development* **124**: 4065-75.
- . 1999. The homeobox gene Phox2b is essential for the development of autonomic neural crest derivatives. *Nature* **399**: 366-70.
- Pattyn, A., A. Vallstedt, J.M. Dias, O.A. Samad, R. Krumlauf, F.M. Rijli, J.F. Brunet, and J. Ericson. 2003. Coordinated temporal and spatial control of motor neuron and serotonergic neuron generation from a common pool of CNS progenitors. *Genes Dev* **17**: 729-37.

- Qian, Y., B. Fritzsche, S. Shirasawa, C.L. Chen, Y. Choi, and Q. Ma. 2001. Formation of brainstem (nor)adrenergic centers and first-order relay visceral sensory neurons is dependent on homeodomain protein Rnx/Tlx3. *Genes Dev* **15**: 2533-45.
- Roelink, H., J.A. Porter, C. Chiang, Y. Tanabe, D.T. Chang, P.A. Beachy, and T.M. Jessell. 1995. Floor plate and motor neuron induction by different concentrations of the amino-terminal cleavage product of sonic hedgehog autoproteolysis. *Cell* **81**: 445-55.
- Rossel, M. and M.R. Capecchi. 1999. Mice mutant for both Hoxa1 and Hoxb1 show extensive remodeling of the hindbrain and defects in craniofacial development. *Development* **126**: 5027-40.
- Saiki, R.K., S. Scharf, F. Faloona, K.B. Mullis, G.T. Horn, H.A. Erlich, and N. Arnheim. 1985. Enzymatic amplification of beta-globin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. *Science* **230**: 1350-4.
- Sambrook. *Molecular Cloning*.
- Sanger, F., S. Nicklen, and A.R. Coulson. 1977. DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci U S A* **74**: 5463-7.
- Schneider-Maunoury, S., P. Topilko, T. Seitandou, G. Levi, M. Cohen-Tannoudji, S. Pournin, C. Babinet, and P. Charnay. 1993. Disruption of Krox-20 results in alteration of rhombomeres 3 and 5 in the developing hindbrain. *Cell* **75**: 1199-214.
- Sham, M.H., C. Vesque, S. Nonchev, H. Marshall, M. Frain, R.D. Gupta, J. Whiting, D. Wilkinson, P. Charnay, and R. Krumlauf. 1993. The zinc finger gene Krox20 regulates HoxB2 (Hox2.8) during hindbrain segmentation. *Cell* **72**: 183-96.
- Shirasawa, S., A. Arata, H. Onimaru, K.A. Roth, G.A. Brown, S. Horning, S. Arata, K. Okumura, T. Sasazuki, and S.J. Korsmeyer. 2000. Rnx deficiency results in congenital central hypoventilation. *Nat Genet* **24**: 287-90.
- Simeone, A., D. Acampora, V. Nigro, A. Faiella, M. D'Esposito, A. Stornaiuolo, F. Mavilio, and E. Boncinelli. 1991. Differential regulation by retinoic acid of the homeobox genes of the four HOX loci in human embryonal carcinoma cells. *Mech Dev* **33**: 215-27.
- Southern, E.M. 1975. Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J Mol Biol* **98**: 503-17.
- Studer, M., A. Lumsden, L. Ariza-McNaughton, A. Bradley, and R. Krumlauf. 1996. Altered segmental identity and abnormal migration of motor neurons in mice lacking Hoxb-1. *Nature* **384**: 630-4.

- Szucsik, J.C., D.P. Witte, H. Li, S.K. Pixley, K.M. Small, and S.S. Potter. 1997. Altered forebrain and hindbrain development in mice mutant for the Gsh-2 homeobox gene. *Dev Biol* **191**: 230-42.
- Tabor, S. and C.C. Richardson. 1987. DNA sequence analysis with a modified bacteriophage T7 DNA polymerase. *Proc Natl Acad Sci U S A* **84**: 4767-71.
- Thaler, J.P., S.J. Koo, A. Kania, K. Lettieri, S. Andrews, C. Cox, T.M. Jessell, and S.L. Pfaff. 2004. A postmitotic role for Isl-class LIM homeodomain proteins in the assignment of visceral spinal motor neuron identity. *Neuron* **41**: 337-50.
- Timmer, J.R., C. Wang, and L. Niswander. 2002. BMP signaling patterns the dorsal and intermediate neural tube via regulation of homeobox and helix-loop-helix transcription factors. *Development* **129**: 2459-72.
- Vaage, S. 1969. The segmentation of the primitive neural tube in chick embryos (*Gallus domesticus*). A morphological, histochemical and autoradiographical investigation. *Ergeb Anat Entwicklungsgesch* **41**: 3-87.
- Viemari JC, B.M., Burnet H, Coulon P, Pequignot JM, Tiveron MC, Hilaire G. 2004. Phox2a gene, A6 neurons, and noradrenaline are essential for development of normal respiratory rhythm in mice. *J Neurosci* **24(4)**: 928-37.
- Vogelstein, B. and D. Gillespie. 1979. Preparative and analytical purification of DNA from agarose. *Proc Natl Acad Sci U S A* **76**: 615-9.
- von Baer, K.E. 1828. *Entwicklungsgeschichte der Thiere: Beobachtung und Reflexion. Born Träger, Königsberg.*
- Waskiewicz, A.J., H.A. Rikhof, R.E. Hernandez, and C.B. Moens. 2001. Zebrafish Meis functions to stabilize Pbx proteins and regulate hindbrain patterning. *Development* **128**: 4139-51.
- Waskiewicz, A.J., H.A. Rikhof, and C.B. Moens. 2002. Eliminating zebrafish pbx proteins reveals a hindbrain ground state. *Dev Cell* **3**: 723-33.
- Wassarman, K.M., M. Lewandoski, K. Campbell, A.L. Joyner, J.L. Rubenstein, S. Martinez, and G.R. Martin. 1997. Specification of the anterior hindbrain and establishment of a normal mid/hindbrain organizer is dependent on Gbx2 gene function. *Development* **124**: 2923-34.
- Wilkinson, D.G. 1992. *In situ Hybridization: A Practical Approach*. Oxford University Press, Oxford, United Kingdom.
- Wurst, W., A.B. Auerbach, and A.L. Joyner. 1994. Multiple developmental defects in Engrailed-1 mutant mice: an early mid-hindbrain deletion and patterning defects in forelimbs and sternum. *Development* **120**: 2065-75.

Danksagung

M

- Barrow, J.R., H.S. Stadler, and M.R. Capecchi. 2000. Roles of Hoxa1 and Hoxa2 in patterning the early hindbrain of the mouse. In *Development*, pp. 933-44.
- Bell, E., R.J. Wingate, and A. Lumsden. 1999. Homeotic transformation of rhombomere identity after localized Hoxb1 misexpression. In *Science*, pp. 2168-71.
- Birnboim, H.C. and J. Doly. 1979. A rapid alkaline extraction procedure for screening recombinant plasmid DNA. In *Nucleic Acids Res*, pp. 1513-23.
- Briscoe, J. and J. Ericson. 2001. Specification of neuronal fates in the ventral neural tube. In *Curr Opin Neurobiol*, pp. 43-9.
- Briscoe, J., A. Pierani, T.M. Jessell, and J. Ericson. 2000. A homeodomain protein code specifies progenitor cell identity and neuronal fate in the ventral neural tube. In *Cell*, pp. 435-45.
- Brohmann, H., K. Jagla, and C. Birchmeier. 2000. The role of Lbx1 in migration of muscle precursor cells. In *Development*, pp. 437-45.
- Carpenter, E.M., J.M. Goddard, O. Chisaka, N.R. Manley, and M.R. Capecchi. 1993. Loss of Hox-A1 (Hox-1.6) function results in the reorganization of the murine hindbrain. In *Development*, pp. 1063-75.
- Chen, F., K.C. Liu, and J.A. Epstein. 1999. Lbx2, a novel murine homeobox gene related to the Drosophila ladybird genes is expressed in the developing urogenital system, eye and brain. In *Mech Dev*, pp. 181-4.
- Cheng, L., A. Arata, R. Mizuguchi, Y. Qian, A. Karunaratne, P.A. Gray, S. Arata, S. Shirasawa, M. Bouchard, P. Luo, C.L. Chen, M. Busslinger, M. Goulding, H. Onimaru, and Q. Ma. 2004. Tlx3 and Tlx1 are post-mitotic selector genes determining glutamatergic over GABAergic cell fates. In *Nat Neurosci*, pp. 510-517.
- Chi, C.L., S. Martinez, W. Wurst, and G.R. Martin. 2003. The isthmus organizer signal FGF8 is required for cell survival in the prospective midbrain and cerebellum. In *Development*, pp. 2633-44.
- Chiang, C., Y. Litingtung, E. Lee, K.E. Young, J.L. Corden, H. Westphal, and P.A. Beachy. 1996. Cyclopia and defective axial patterning in mice lacking Sonic hedgehog gene function. In *Nature*, pp. 407-13.
- Clark, G. 1981. Staining Procedures. In (ed. S. S.S.), pp. 512. Williams & Wilkins, Baltimore, MD 21202, USA.
- Dauger, S., F. Guimiot, S. Renolleau, B. Levacher, B. Boda, C. Mas, V. Nepote, M. Simonneau, C. Gaultier, and J. Gallego. 2001. MASH-1/RET pathway involvement in development of brain stem control of respiratory frequency in newborn mice. In *Physiol Genomics*, pp. 149-57.
- Dauger, S., S. Renolleau, G. Vardon, V. Nepote, C. Mas, M. Simonneau, C. Gaultier, and J. Gallego. 1999. Ventilatory responses to hypercapnia and hypoxia in Mash-1 heterozygous newborn and adult mice. In *Pediatr Res*, pp. 535-42.
- de Diego, I., K. Kyriakopoulou, D. Karagogeos, and M. Wassef. 2002. Multiple influences on the migration of precerebellar neurons in the caudal medulla. In *Development*, pp. 297-306.
- Dolle, P., T. Lufkin, R. Krumlauf, M. Mark, D. Duboule, and P. Chambon. 1993. Local alterations of Krox-20 and Hox gene expression in the hindbrain suggest

- lack of rhombomeres 4 and 5 in homozygote null *Hoxa-1* (*Hox-1.6*) mutant embryos. In *Proc Natl Acad Sci U S A*, pp. 7666-70.
- Dupe, V., M. Davenne, J. Brocard, P. Dolle, M. Mark, A. Dierich, P. Chambon, and F.M. Rijli. 1997. In vivo functional analysis of the *Hoxa-1* 3' retinoic acid response element (3'RARE). In *Development*, pp. 399-410.
- Ericson, J., S. Morton, A. Kawakami, H. Roelink, and T.M. Jessell. 1996. Two critical periods of Sonic Hedgehog signaling required for the specification of motor neuron identity. In *Cell*, pp. 661-73.
- Ericson, J., J. Muhr, T.M. Jessell, and T. Edlund. 1995a. Sonic hedgehog: a common signal for ventral patterning along the rostrocaudal axis of the neural tube. In *Int J Dev Biol*, pp. 809-16.
- Ericson, J., J. Muhr, M. Placzek, T. Lints, T.M. Jessell, and T. Edlund. 1995b. Sonic hedgehog induces the differentiation of ventral forebrain neurons: a common signal for ventral patterning within the neural tube. In *Cell*, pp. 747-56.
- Feinberg, A.P. and B. Vogelstein. 1983. A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. In *Anal Biochem*, pp. 6-13.
- Feldman, J.L., G.S. Mitchell, and E.E. Nattie. 2003. Breathing: rhythmicity, plasticity, chemosensitivity. In *Annu Rev Neurosci*, pp. 239-66.
- Fraser, S., R. Keynes, and A. Lumsden. 1990. Segmentation in the chick embryo hindbrain is defined by cell lineage restrictions. In *Nature*, pp. 431-5.
- Frohman, M.A., G.R. Martin, S.P. Cordes, L.P. Halamek, and G.S. Barsh. 1993. Altered rhombomere-specific gene expression and hyoid bone differentiation in the mouse segmentation mutant, *kreisler* (*kr*). In *Development*, pp. 925-36.
- Gaufo, G.O., S. Wu, and M.R. Capecchi. 2004. Contribution of Hox genes to the diversity of the hindbrain sensory system. In *Development*, pp. 1259-66.
- Gavalas, A., M. Studer, A. Lumsden, F.M. Rijli, R. Krumlauf, and P. Chambon. 1998. *Hoxa1* and *Hoxb1* synergize in patterning the hindbrain, cranial nerves and second pharyngeal arch. In *Development*, pp. 1123-36.
- Gavrieli, Y., Y. Sherman, and S.A. Ben-Sasson. 1992. Identification of programmed cell death in situ via specific labeling of nuclear DNA fragmentation. In *J Cell Biol*, pp. 493-501.
- Ghysen, A. and C. Dambly-Chaudiere. 1990. Early events in the development of *Drosophila* peripheral nervous system. In *J Physiol (Paris)*, pp. 11-20.
- Goddard, J.M., M. Rossel, N.R. Manley, and M.R. Capecchi. 1996. Mice with targeted disruption of *Hoxb-1* fail to form the motor nucleus of the VIIth nerve. In *Development*, pp. 3217-28.
- Gowan, K., A.W. Helms, T.L. Hunsaker, T. Collisson, P.J. Ebert, R. Odom, and J.E. Johnson. 2001. Crossinhibitory activities of *Ngn1* and *Math1* allow specification of distinct dorsal interneurons. In *Neuron*, pp. 219-32.
- Gross, M.K., M. Dottori, and M. Goulding. 2002. *Lbx1* specifies somatosensory association interneurons in the dorsal spinal cord. In *Neuron*, pp. 535-49.
- Guillemot, F., L.C. Lo, J.E. Johnson, A. Auerbach, D.J. Anderson, and A.L. Joyner. 1993. Mammalian achaete-scute homolog 1 is required for the early development of olfactory and autonomic neurons. In *Cell*, pp. 463-76.
- Hirsch, M.R., M.C. Tiveron, F. Guillemot, J.F. Brunet, and C. Goridis. 1998. Control of noradrenergic differentiation and *Phox2a* expression by *MASH1* in the central and peripheral nervous system. In *Development*, pp. 599-608.

- Hogan, B., R. Beddington, F. Constantini, and E. Lacy. 1994. Manipulating The Mouse Embryo - A Laboratory Manual. In, pp. 497. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York 11803-2500, USA.
- Innis, M.A., D.H. Gelfand, and J.J. Sninsky. 1989. PCR Protocols: A Guide To Methods And Applications. In, pp. 482. Academic Press, San Diego, CA 92101-4495, USA.
- Inoue, H., H. Nojima, and H. Okayama. 1990. High efficiency transformation of Escherichia coli with plasmids. In *Gene*, pp. 23-8.
- Jagla, K., P. Dolle, M.G. Mattei, T. Jagla, B. Schuhbaur, G. Dretzen, F. Bellard, and M. Bellard. 1995. Mouse Lbx1 and human LBX1 define a novel mammalian homeobox gene family related to the Drosophila lady bird genes. In *Mech Dev*, pp. 345-56.
- Jessell, T.M. 2000. Neuronal specification in the spinal cord: inductive signals and transcriptional codes. In *Nat Rev Genet*, pp. 20-9.
- Lee, K.J., P. Dietrich, and T.M. Jessell. 2000. Genetic ablation reveals that the roof plate is essential for dorsal interneuron specification. In *Nature*, pp. 734-40.
- Lee, K.J. and T.M. Jessell. 1999. The specification of dorsal cell fates in the vertebrate central nervous system. In *Annu Rev Neurosci*, pp. 261-94.
- Lee, K.J., M. Mendelsohn, and T.M. Jessell. 1998. Neuronal patterning by BMPs: a requirement for GDF7 in the generation of a discrete class of commissural interneurons in the mouse spinal cord. In *Genes Dev*, pp. 3394-407.
- Logan, C., R.J. Wingate, I.J. McKay, and A. Lumsden. 1998. Tlx-1 and Tlx-3 homeobox gene expression in cranial sensory ganglia and hindbrain of the chick embryo: markers of patterned connectivity. In *J Neurosci*, pp. 5389-402.
- Lumsden, A. and R. Keynes. 1989. Segmental patterns of neuronal development in the chick hindbrain. In *Nature*, pp. 424-8.
- Lumsden, A. and R. Krumlauf. 1996. Patterning the vertebrate neuraxis. In *Science*, pp. 1109-15.
- Lumsden, T. 1923. Observations on the respiratory centers in the cat. In *The Journal of Physiology*, pp. 153-160.
- Ma, Q., L. Sommer, P. Cserjesi, and D.J. Anderson. 1997. Mash1 and neurogenin1 expression patterns define complementary domains of neuroepithelium in the developing CNS and are correlated with regions expressing notch ligands. In *J Neurosci*, pp. 3644-52.
- Maconochie, M.K., S. Nonchev, M. Studer, S.K. Chan, H. Popperl, M.H. Sham, R.S. Mann, and R. Krumlauf. 1997. Cross-regulation in the mouse HoxB complex: the expression of Hoxb2 in rhombomere 4 is regulated by Hoxb1. In *Genes Dev*, pp. 1885-95.
- Manzanares, M., S. Bel-Vialar, L. Ariza-McNaughton, E. Ferretti, H. Marshall, M.M. Maconochie, F. Blasi, and R. Krumlauf. 2001. Independent regulation of initiation and maintenance phases of Hoxa3 expression in the vertebrate hindbrain involve auto- and cross-regulatory mechanisms. In *Development*, pp. 3595-607.
- Manzanares, M., P.A. Trainor, S. Nonchev, L. Ariza-McNaughton, J. Brodie, A. Gould, H. Marshall, A. Morrison, C.T. Kwan, M.H. Sham, D.G. Wilkinson, and R. Krumlauf. 1999. The role of kreisler in segmentation during hindbrain development. In *Dev Biol*, pp. 220-37.
- Marin, F. and L. Puellas. 1995. Morphological fate of rhombomeres in quail/chick chimeras: a segmental analysis of hindbrain nuclei. In *Eur J Neurosci*, pp. 1714-38.

- Mark, M., T. Lufkin, P. Dolle, A. Dierich, M. LeMeur, and P. Chambon. 1993. Roles of Hox genes: what we have learnt from gain of function and loss of function mutations in the mouse. In *C R Acad Sci III*, pp. 995-1008.
- McDonnell, M.W., M.N. Simon, and F.W. Studier. 1977. Analysis of restriction fragments of T7 DNA and determination of molecular weights by electrophoresis in neutral and alkaline gels. In *J Mol Biol*, pp. 119-46.
- McGinnis, W. and R. Krumlauf. 1992. Homeobox genes and axial patterning. In *Cell*, pp. 283-302.
- McKay, I.J., I. Muchamore, R. Krumlauf, M. Maden, A. Lumsden, and J. Lewis. 1994. The kreisler mouse: a hindbrain segmentation mutant that lacks two rhombomeres. In *Development*, pp. 2199-211.
- Mizuguchi, R., M. Sugimori, H. Takebayashi, H. Kosako, M. Nagao, S. Yoshida, Y. Nabeshima, K. Shimamura, and M. Nakafuku. 2001. Combinatorial roles of olig2 and neurogenin2 in the coordinated induction of pan-neuronal and subtype-specific properties of motoneurons. In *Neuron*, pp. 757-71.
- Morin, X., H. Cremer, M.R. Hirsch, R.P. Kapur, C. Goridis, and J.F. Brunet. 1997. Defects in sensory and autonomic ganglia and absence of locus coeruleus in mice deficient for the homeobox gene Phox2a. In *Neuron*, pp. 411-23.
- Muller, T., H. Brohmann, A. Pierani, P.A. Heppenstall, G.R. Lewin, T.M. Jessell, and C. Birchmeier. 2002. The homeodomain factor lbx1 distinguishes two major programs of neuronal differentiation in the dorsal spinal cord. In *Neuron*, pp. 551-62.
- Müller, T., H. Brohmann, A. Pierani, P.A. Heppenstall, G.R. Lewin, T.M. Jessell, and C. Birchmeier. 2002. The homeodomain factor Lbx1 distinguishes two major programs of neuronal differentiation in the dorsal spinal cord. In *Neuron*.
- Novitch, B.G., A.I. Chen, and T.M. Jessell. 2001. Coordinate Regulation of Motor Neuron Subtype Identity and Pan-Neuronal Properties by the bHLH Repressor Olig2. In *Neuron*, pp. 773-89.
- Ogura, T. and R.M. Evans. 1995. A retinoic acid-triggered cascade of HOXB1 gene activation. In *Proc Natl Acad Sci U S A*, pp. 387-91.
- Olszewski J, B.D. 1982. Cytoarchitecture of the human Brain Stem. In, pp. 83-91.
- Pattyn, A., C. Goridis, and J.F. Brunet. 2000a. Specification of the central noradrenergic phenotype by the homeobox gene Phox2b. In *Mol Cell Neurosci*, pp. 235-43.
- Pattyn, A., M. Hirsch, C. Goridis, and J.F. Brunet. 2000b. Control of hindbrain motor neuron differentiation by the homeobox gene Phox2b. In *Development*, pp. 1349-58.
- Pattyn, A., X. Morin, H. Cremer, C. Goridis, and J.F. Brunet. 1997. Expression and interactions of the two closely related homeobox genes Phox2a and Phox2b during neurogenesis. In *Development*, pp. 4065-75.
- . 1999. The homeobox gene Phox2b is essential for the development of autonomic neural crest derivatives. In *Nature*, pp. 366-70.
- Pattyn, A., A. Vallstedt, J.M. Dias, O.A. Samad, R. Krumlauf, F.M. Rijli, J.F. Brunet, and J. Ericson. 2003. Coordinated temporal and spatial control of motor neuron and serotonergic neuron generation from a common pool of CNS progenitors. In *Genes Dev*, pp. 729-37.
- Qian, Y., B. Fritsch, S. Shirasawa, C.L. Chen, Y. Choi, and Q. Ma. 2001. Formation of brainstem (nor)adrenergic centers and first-order relay visceral sensory neurons is dependent on homeodomain protein Rnx/Tlx3. In *Genes Dev*, pp. 2533-45.

- Roelink, H., J.A. Porter, C. Chiang, Y. Tanabe, D.T. Chang, P.A. Beachy, and T.M. Jessell. 1995. Floor plate and motor neuron induction by different concentrations of the amino-terminal cleavage product of sonic hedgehog autoproteolysis. In *Cell*, pp. 445-55.
- Rossel, M. and M.R. Capecchi. 1999. Mice mutant for both Hoxa1 and Hoxb1 show extensive remodeling of the hindbrain and defects in craniofacial development. In *Development*, pp. 5027-40.
- Saiki, R.K., S. Scharf, F. Faloona, K.B. Mullis, G.T. Horn, H.A. Erlich, and N. Arnheim. 1985. Enzymatic amplification of beta-globin genomic sequences and restriction site analysis for diagnosis of sickle cell anemia. In *Science*, pp. 1350-4.
- Sambrook. In *Molecular Cloning*.
- Sanger, F., S. Nicklen, and A.R. Coulson. 1977. DNA sequencing with chain-terminating inhibitors. In *Proc Natl Acad Sci U S A*, pp. 5463-7.
- Schneider-Maunoury, S., P. Topilko, T. Seitandou, G. Levi, M. Cohen-Tannoudji, S. Pournin, C. Babinet, and P. Charnay. 1993. Disruption of Krox-20 results in alteration of rhombomeres 3 and 5 in the developing hindbrain. In *Cell*, pp. 1199-214.
- Sham, M.H., C. Vesque, S. Nonchev, H. Marshall, M. Frain, R.D. Gupta, J. Whiting, D. Wilkinson, P. Charnay, and R. Krumlauf. 1993. The zinc finger gene Krox20 regulates HoxB2 (Hox2.8) during hindbrain segmentation. In *Cell*, pp. 183-96.
- Shirasawa, S., A. Arata, H. Onimaru, K.A. Roth, G.A. Brown, S. Horning, S. Arata, K. Okumura, T. Sasazuki, and S.J. Korsmeyer. 2000. Rnx deficiency results in congenital central hypoventilation. In *Nat Genet*, pp. 287-90.
- Simeone, A., D. Acampora, V. Nigro, A. Faiella, M. D'Esposito, A. Stornaiuolo, F. Mavilio, and E. Boncinelli. 1991. Differential regulation by retinoic acid of the homeobox genes of the four HOX loci in human embryonal carcinoma cells. In *Mech Dev*, pp. 215-27.
- Southern, E.M. 1975. Detection of specific sequences among DNA fragments separated by gel electrophoresis. In *J Mol Biol*, pp. 503-17.
- Studer, M., A. Lumsden, L. Ariza-McNaughton, A. Bradley, and R. Krumlauf. 1996. Altered segmental identity and abnormal migration of motor neurons in mice lacking Hoxb-1. In *Nature*, pp. 630-4.
- Szucsik, J.C., D.P. Witte, H. Li, S.K. Pixley, K.M. Small, and S.S. Potter. 1997. Altered forebrain and hindbrain development in mice mutant for the Gsh-2 homeobox gene. In *Dev Biol*, pp. 230-42.
- Tabor, S. and C.C. Richardson. 1987. DNA sequence analysis with a modified bacteriophage T7 DNA polymerase. In *Proc Natl Acad Sci U S A*, pp. 4767-71.
- Thaler, J.P., S.J. Koo, A. Kania, K. Lettieri, S. Andrews, C. Cox, T.M. Jessell, and S.L. Pfaff. 2004. A postmitotic role for Isl-class LIM homeodomain proteins in the assignment of visceral spinal motor neuron identity. In *Neuron*, pp. 337-50.
- Timmer, J.R., C. Wang, and L. Niswander. 2002. BMP signaling patterns the dorsal and intermediate neural tube via regulation of homeobox and helix-loop-helix transcription factors. In *Development*, pp. 2459-72.
- Vaage, S. 1969. The segmentation of the primitive neural tube in chick embryos (*Gallus domesticus*). A morphological, histochemical and autoradiographical investigation. In *Ergeb Anat Entwicklungsgesch*, pp. 3-87.

- Viemari JC, B.M., Burnet H, Coulon P, Pequignot JM, Tiveron MC, Hilaire G. 2004. Phox2a gene, A6 neurons, and noradrenaline are essential for development of normal respiratory rhythm in mice. In *J Neurosci*, pp. 928-37.
- Vogelstein, B. and D. Gillespie. 1979. Preparative and analytical purification of DNA from agarose. In *Proc Natl Acad Sci U S A*, pp. 615-9.
- von Baer, K.E. 1828. Entwicklungsgeschichte der Thiere: Beobachtung und Reflexion. In *Born Träger, Königsberg*.
- Waskiewicz, A.J., H.A. Rikhof, R.E. Hernandez, and C.B. Moens. 2001. Zebrafish Meis functions to stabilize Pbx proteins and regulate hindbrain patterning. In *Development*, pp. 4139-51.
- Waskiewicz, A.J., H.A. Rikhof, and C.B. Moens. 2002. Eliminating zebrafish pbx proteins reveals a hindbrain ground state. In *Dev Cell*, pp. 723-33.
- Wassarman, K.M., M. Lewandoski, K. Campbell, A.L. Joyner, J.L. Rubenstein, S. Martinez, and G.R. Martin. 1997. Specification of the anterior hindbrain and establishment of a normal mid/hindbrain organizer is dependent on Gbx2 gene function. In *Development*, pp. 2923-34.
- Wilkinson, D.G. 1992. In situ Hybridization: A Practical Approach. In. Oxford University Press, Oxford, United Kingdom.
- Wurst, W., A.B. Auerbach, and A.L. Joyner. 1994. Multiple developmental defects in Engrailed-1 mutant mice: an early mid-hindbrain deletion and patterning defects in forelimbs and sternum. In *Development*, pp. 2065-75.