

References

- Abe, I., Okumoto, K., Tamura, S. and Fujiki, Y. (1998) Clofibrate-inducible, 28-kDa peroxisomal integral membrane protein is encoded by *PEX11*. *FEBS Lett.*, **431**, 468-472.
- Albertini, M., Rehling, P., Erdmann, R., Girzalsky, W., Kiel, J.A., Veenhuis, M. and Kunau, W.H. (1997) Pex14p, a peroxisomal membrane protein binding both receptors of the two PTS-dependent import pathways. *Cell*, **89**, 83-92.
- Anderson SK, G.C., Tanaka A, Kung HJ, Fujita DJ. (1985) Human cellular src gene: nucleotide sequence and derived amino acid sequence of the region coding for the carboxy-terminal two-thirds of pp60c-src. *Mol. Cell Biol*, **5**, 1122-1129.
- Ausubel M. Frederick, B.R., Kingston E. Robert,,Moore D. David, Seidman J.G., Smith A. John, Struhl Kevin. (1987) Current Protocols in Molecular Biology. *Massachusetts General Hospital, Harvard Medical School*.
- Baerends, R.J., Faber, K.N., Kiel, J.A., van der Klei, I.J., Harder, W. and Veenhuis, M. (2000) Sorting and function of peroxisomal membrane proteins. *FEMS Microbiol Rev*, **24**, 291-301.
- Baerends, R.J.S., Rasmussen, S.W., Hilbrands, R.E., van der Heide, M., Faber, K.N., Reuvekamp, P.T.W., Kiel, J.A.K.W., Cregg, J.M., van der Klei, I.J. and Veenhuis, M. (1996) The *Hansenula polymorpha* *PER9* Gene Encodes a Peroxisomal Membrane Protein Essential for Peroxisome Assembly and Integrity. *J. Biol. Chem.*, **271**, 8887-8894.
- Barnett, P., Bottger, G., Klein, A.T., Tabak, H.F. and Distel, B. (2000) The peroxisomal membrane protein Pex13p shows a novel mode of SH3 interaction. *Embo J*, **19**, 6382-6391.
- Barth, P.G., Gootjes, J., Bode, H., Vreken, P., Majoie, C.B. and Wanders, R.J. (2001) Late onset white matter disease in peroxisome biogenesis disorder. *Neurology*, **57**, 1949-1955.
- Baudhuin, C.D.a.P. (1966) Peroxisomes (Microbodies and related particles). *Physio. Rev.*, **46**, 323-357.
- Blobel, F. and Erdmann, R. (1996) Identification of a yeast peroxisomal member of the family of AMP-binding proteins. *Eur. J. Biochem.*, **240**, 468-476.
- Bottger, G., Barnett, P., Klein, A.T., Kragt, A., Tabak, H.F. and Distel, B. (2000) *Saccharomyces cerevisiae* PTS1 receptor Pex5p interacts with the SH3 domain of the peroxisomal membrane protein Pex13p in an unconventional, non-PXXP-related manner. *Mol Biol Cell*, **11**, 3963-3976.
- Braun, A., Kammerer, S., Weissenhorn, W., Weiss, E.H. and Cleve, H. (1994) Sequence of a putative human housekeeping gene (*HK33*) localized on chromosome 1. *Gene*, **146**, 291-295.
- Braverman, N., Dodt, G., Gould, S.J. and Valle, D. (1995) Disorders of peroxisome biogenesis. *Hum. Mol. Genet.*, **4**, 1791-1798.
- Braverman, N., Steel, G., Obie, C., Moser, A., Moser, H., Gould, S.J. and Valle, D. (1997) Human *PEX7* encodes the peroxisomal PTS2 receptor and is responsible for rhizomelic chondrodysplasia punctata. *Nat Genet*, **15**, 369-376.
- Brocard, C., Kragler, F., Simon, M.M., Schuster, T. and Hartig, A. (1994) The tetratricopeptide repeat-domain of the Pas10 protein of *Saccharomyces cerevisiae* is essential for binding the peroxisomal targeting signal-SKL. *Biochem. Biophys. Res. Commun.*, **204**, 1016-1022.

- Brocard, C., Lametschwandtner, G., Koudelka, R. and Hartig, A. (1997) Pex14p is a member of the protein linkage map of Pex5p. *Embo J*, **16**, 5491-5500.
- Brown, T.W., Titorenko, V.I. and Rachubinski, R.A. (2000) Mutants of the *Yarrowia lipolytica* PEX23 gene encoding an integral peroxisomal membrane peroxin mislocalize matrix proteins and accumulate vesicles containing peroxisomal matrix and membrane proteins. *Mol Biol Cell*, **11**, 141-152.
- Chang, C.C., Lee, W.H., Moser, H., Valle, D. and Gould, S.J. (1997) Isolation of the human *PEX12* gene, mutated in group 3 of the peroxisome biogenesis disorders. *Nat. Genet.*, **15**, 385-388.
- Chang, C.C., Warren, D.S., Sacksteder, K.A. and Gould, S.J. (1999) PEX12 interacts with PEX5 and PEX10 and acts downstream of receptor docking in peroxisomal matrix protein import. *J Cell Biol*, **147**, 761-774.
- Crane, D.I., Kalish, J.E. and Gould, S.J. (1994) The *Pichia pastoris* PAS4 gene encodes a ubiquitin-conjugating enzyme required for peroxisome assembly. *J Biol Chem*, **269**, 21835-21844.
- Dammai, V. and Subramani, S. (2001) The human peroxisomal targeting signal receptor, Pex5p, is translocated into the peroxisomal matrix and recycled to the cytosol. *Cell*, **105**, 187-196.
- Das, A.K., Cohen, P.W. and Barford, D. (1998) The structure of the tetratricopeptide repeats of protein phosphatase 5: implications for TPR-mediated protein-protein interactions. *Embo J*, **17**, 1192-1199.
- Distel, B., Erdmann, R., Gould, S.J., Blobel, G., Crane, D.I., Cregg, J.M., Dodt, G., Fujiki, Y., Goodman, J.M., Just, W.W., Kiel, J.A., Kunau, W.H., Lazarow, P.B., Mannaerts, G.P., Moser, H.W., Osumi, T., Rachubinski, R.A., Roscher, A., Subramani, S., Tabak, H.F., Tsukamoto, T., Valle, D., van der Klei, I., van Veldhoven, P.P. and Veenhuis, M. (1996a) A unified nomenclature for peroxisome biogenesis factors. *J Cell Biol*, **135**, 1-3.
- Distel, B., van der Leij, I. and Kos, W. (1996b) Peroxisome isolation. *Methods Mol Biol*, **53**, 133-138.
- Dodt, G., Braverman, N., Wong, C., Moser, A., Moser, H.W., Watkins, P., Valle, D. and Gould, S.J. (1995) Mutations in the PTS1 receptor gene, *PXR1*, define complementation group 2 of the peroxisome biogenesis disorders. *Nat. Genet.*, **9**, 115-125.
- Dodt, G. and Gould, S.J. (1996) Multiple PEX genes are required for proper subcellular distribution and stability of Pex5p, the PTS1 receptor: evidence that PTS1 protein import is mediated by a cycling receptor. *J Cell Biol*, **135**, 1763-1774.
- Douangamath, A., Filipp, F.V., Klein, A.T., Barnett, P., Zou, P., Voorn-Brouwer, T., Vega, M.C., Mayans, O.M., Sattler, M., Distel, B. and Wilmanns, M. (2002) Topography for independent binding of alpha-helical and PPII-helical ligands to a peroxisomal SH3 domain. *Mol Cell*, **10**, 1007-1017.
- Einwachter, H., Sowinski, S., Kunau, W.H. and Schliebs, W. (2001a) *Yarrowia lipolytica* Pex20p, *Saccharomyces cerevisiae* Pex18p/Pex21p and mammalian Pex5pL fulfil a common function in the early steps of the peroxisomal PTS2 import pathway. *EMBO Rep*, **2**, 1035-1039.
- Einwachter, H., Sowinski, S., Kunau, W.H. and Schliebs, W. (2001b) *Yarrowia lipolytica* Pex20p, *Saccharomyces cerevisiae* Pex18p/Pex21p and mammalian Pex5pL fulfil a common function in the early steps of the peroxisomal PTS2 import pathway. *EMBO Rep*, **2**, 1035-1039.
- Eitzen, G.A., Aitchison, J.D., Szilard, R.K., Veenhuis, M., Nuttley, W.M. and Rachubinski, R.A. (1995) The *Yarrowia lipolytica* gene *PAY2* encodes a 42-kDa

- peroxisomal integral membrane protein essential for matrix protein import and peroxisome enlargement but not for peroxisome membrane proliferation. *J Biol Chem*, **270**, 1429-1436.
- Eitzen, G.A., Szilard, R.K. and Rachubinski, R.A. (1997) Enlarged peroxisomes are present in oleic acid-grown *Yarrowia lipolytica* overexpressing the PEX16 gene encoding an intraperoxisomal peripheral membrane peroxin. *J Cell Biol*, **137**, 1265-1278.
- Elgersma, Y., Elgersma-Hooisma, M., Wenzel, T., McCaffery, J.M., Farquhar, M.G. and Subramani, S. (1998) A mobile PTS2 receptor for peroxisomal protein import in *Pichia pastoris*. *J Cell Biol*, **140**, 807-820.
- Elgersma, Y., Kwast, L., Klein, A., Voorn-Brouwer, T., van den Berg, M., Metzlig, B., America, T., Tabak, H.F. and Distel, B. (1996) The SH3 domain of the *Saccharomyces cerevisiae* peroxisomal membrane protein Pex13p functions as a docking site for Pex5p, a mobile receptor for the import of PTS1 containing proteins. *J. Cell Biol.*, **135**, 97-109.
- Elgersma, Y., Kwast, L., van den Berg, M., Snyder, W.B., Distel, B., Subramani, S. and Tabak, H.F. (1997) Overexpression of Pex15p, a phosphorylated peroxisomal integral membrane protein required for peroxisome assembly in *S.cerevisiae*, causes proliferation of the endoplasmic reticulum membrane. *EMBO J.*, **16**, 7326-7341.
- Erdmann, R. and Blobel, G. (1995) Giant peroxisomes in oleic acid-induced *Saccharomyces cerevisiae* lacking the peroxisomal membrane protein Pmp27p. *J. Cell Biol.*, **128**, 509-523.
- Erdmann, R. and Blobel, G. (1996) Identification of Pex13p a peroxisomal membrane receptor for the PTS1 recognition factor. *J Cell Biol*, **135**, 111-121.
- Erdmann, R. and Kunau, W.-H. (1992) A Genetic Approach to the Biogenesis of Peroxisomes in the Yeast *Saccharomyces cerevisiae*. *Cell. Biochem. Funct.*, **10**, 167-174.
- Erdmann, R., Veenhuis, M. and Kunau, W.-H. (1997) Peroxisomes: organelles at the crossroads. *Trends Cell Biol.*, **7**, 400-407.
- Erdmann, R., Veenhuis, M., Mertens, D. and Kunau, W.H. (1989) Isolation of peroxisome-deficient mutants of *Saccharomyces cerevisiae*. *Proc Natl Acad Sci U S A*, **86**, 5419-5423.
- Erdmann, R., Wiebel, F.F., Flessau, A., Rytka, J., Beyer, A., Fr 黷lich, K.U. and Kunau, W.-H. (1991) *PAS1*, a yeast gene required for peroxisome biogenesis, encodes a member of a novel family of putative ATPases. *Cell*, **64**, 499-510.
- Faber, K.N., Heyman, J.A. and Subramani, S. (1998) Two AAA family peroxins, PpPex1p and PpPex6p, interact with each other in an ATP-dependent manner and are associated with different subcellular membranous structures distinct from peroxisomes. *Mol. Cell. Biol.*, **18**, 936-943.
- Feilotter, H.E., Hannon, G.J., Ruddell, C.J. and Beach, D. (1994) Construction of an improved host strain for two hybrid screening. *Nucleic Acids Res*, **22**, 1502-1503.
- Fields, S. and Song, O.K. (1989) A novel genetic system to detect protein-protein interactions. *Nature*, **340**, 245-246.
- Fransen, M., Brees, C., Baumgart, E., Vanhooren, J.C.T., Baes, M., Mannaerts, G.P. and van Veldhoven, P.P. (1995) Identification and characterization of the putative human peroxisomal C-terminal targeting signal import receptor. *J. Biol. Chem.*, **270**, 7731-7736.

- Fransen, M., Terlecky, S.R. and Subramani, S. (1998) Identification of a human PTS1 receptor docking protein directly required for peroxisomal protein import. *Proc. Natl. Acad. Sci. U S A*, **95**, 8087-8092.
- Geisbrecht, B.V. and Gould, S.J. (1999) The human PICD gene encodes a cytoplasmic and peroxisomal NADP(+)-dependent isocitrate dehydrogenase. *J Biol Chem*, **274**, 30527-30533.
- Gietz, R.D. and Sugino, A. (1988) New yeast-E. coli shuttle vectors constructed with *in vitro* mutagenized yeast genes lacking six-base pair restriction sites. *Gene*, **74**, 527-534.
- Girzalsky, W., Rehling, P., Stein, K., Kipper, J., Blank, L., Kunau, W.H. and Erdmann, R. (1999) Involvement of Pex13p in Pex14p localization and peroxisomal targeting signal 2-dependent protein import into peroxisomes. *J Cell Biol*, **144**, 1151-1162.
- Goldfischer, S., Moor, C.L., Johnson, A.B., Spiro, A.J., M.P., V., Wisniewski, H.K., Ritch, R.H., Norton, W.T., Rapin, I. and Gerner, L.M. (1973) Peroxisomal and mitochondrial defects in cerebrohepato-renal syndrome. *Science*, **182**, 62-64.
- Gotte, K., Girzalsky, W., Linkert, M., Baumgart, E., Kammerer, S., Kunau, W.H. and Erdmann, R. (1998) Pex19p, a farnesylated protein essential for peroxisome biogenesis. *Mol Cell Biol*, **18**, 616-628.
- Gould, S.J., Kalish, J.E., Morrell, J.C., Bjorkman, J., Urquhart, A.J. and Crane, D.I. (1996) Pex13p is an SH3 protein of the peroxisome membrane and a docking factor for the predominantly cytoplasmic PTS1 receptor. *J Cell Biol*, **135**, 85-95.
- Hettema, E.H., Distel, B. and Tabak, H.F. (1999) Import of proteins into peroxisomes. *Biochim Biophys Acta*, **1451**, 17-34.
- Hettema, E.H., Girzalsky, W., van Den Berg, M., Erdmann, R. and Distel, B. (2000) *Saccharomyces cerevisiae* pex3p and pex19p are required for proper localization and stability of peroxisomal membrane proteins. *Embo J*, **19**, 223-233.
- Hettema, E.H., Ruigrok, C.C., Koerkamp, M.G., van den Berg, M., Tabak, H.F., Distel, B. and Braakman, I. (1998) The cytosolic DnaJ-like protein djp1p is involved specifically in peroxisomal protein import. *J Cell Biol*, **142**, 421-434.
- Heyman, J.A., Monosov, E. and Subramani, S. (1994) Role of the *PAS1* gene of *Pichia pastoris* in peroxisome biogenesis. *J. Cell. Biol.*, **127**, 1259-1273.
- Hohfeld, J., Veenhuis, M. and Kunau, W.H. (1991) *PAS3*, a *Saccharomyces cerevisiae* gene encoding a peroxisomal integral membrane protein essential for peroxisome biogenesis. *J Cell Biol*, **114**, 1167-1178.
- Holroyd, C. and Erdmann, R. (2001) Protein translocation machineries of peroxisomes. *FEBS Lett*, **501**, 6-10.
- Huhse, B., Rehling, P., Albertini, M., Blank, L., Meller, K. and Kunau, W.H. (1998) Pex17p of *Saccharomyces cerevisiae* is a novel peroxin and component of the peroxisomal protein translocation machinery. *J. Cell Biol.*, **140**, 49-60.
- Kalish, J.E., Keller, G.A., Morrell, J.C., Mihalik, S.J., Smith, B., Cregg, J.M. and Gould, S.J. (1996) Characterization of a novel component of the peroxisomal protein import apparatus using fluorescent peroxisomal proteins. *EMBO J.*, **15**, 3275-3285.
- Kammerer, S., Holzinger, A., Welsch, U. and Roscher, A.A. (1998) Cloning and characterization of the gene encoding the human peroxisomal assembly protein Pex3p. *FEBS Lett.*, **429**, 53-60.
- Kay, L.E. (1995) Pulsed field gradient multi-dimensional NMR methods for the study of protein structure and dynamics in solution. *Prog Biophys Mol Biol*, **63**, 277-299.

- Koller, A., Snyder, W.B., Faber, K.N., Wenzel, T.J., Rangell, L., Keller, G.A. and Subramani, S. (1999) Pex22p of *Pichia pastoris*, essential for peroxisomal matrix protein import, anchors the ubiquitin-conjugating enzyme, Pex4p, on the peroxisomal membrane. *J Cell Biol*, **146**, 99-112.
- Komori, M., Rasmussen, S.W., Kiel, J.A.K.W., Baerends, R.J.S., Cregg, J.M., van der Klei, I.J. and Veenhuis, M. (1997) The *Hansenula polymorpha* PEX14 gene encodes a novel peroxisomal membrane protein essential for peroxisome biogenesis. *EMBO J*, **16**, 44-53.
- Lazarow, P.B. (1995) Peroxisome structure, function, and biogenesis--human patients and yeast mutants show strikingly similar defects in peroxisome biogenesis. *J Neuropathol Exp Neurol*, **54**, 720-725.
- Lazarow, P.B., Black, V., Shio, H., Fujiki, Y., Hajra, A.K., Datta, N.S., Bangaru, B.S. and Dancis, J. (1985) Zellweger syndrome: biochemical and morphological studies on two patients treated with clofibrate. *Pediatr Res*, **19**, 1356-1364.
- Lazarow P.B., F.Y. (1985) Biogenesis of Peroxisomes. *Annu Rev Cell Biol*, **1**, 489-530.
- Lazarow, P.B. and Moser, H.W. (1995) Disorders in peroxisome biogenesis. The metabolic bases of inherited diseases. 7th ed. Scriver C.R., Beaudet, A.L., Sly, W.S. and Valle, D. (eds) McGraw Hill Inc., New York, 7 edition.
- Liu, H., Tan, X., Russell, K.A., Veenhuis, M. and Cregg, J.M. (1995) PER3, a gene required for peroxisome biogenesis in *Pichia pastoris*, encodes a peroxisomal membrane protein involved in protein import. *J Biol Chem*, **270**, 10940-10951.
- Marzioch, M., Erdmann, R., Veenhuis, M. and Kunau, W.H. (1994) PAS7 encodes a novel yeast member of the WD-40 protein family essential for import of 3-oxoacyl-CoA thiolase, a PTS2-containing protein, into peroxisomes. *Embo J*, **13**, 4908-4918.
- Matsumura, T., Otera, H. and Fujiki, Y. (2000) Disruption of the interaction of the longer isoform of Pex5p, Pex5pL, with Pex7p abolishes peroxisome targeting signal type 2 protein import in mammals. Study with a novel Pex5-impaired Chinese hamster ovary cell mutant. *J Biol Chem*, **275**, 21715-21721.
- Mayer, B.J. and Eck, M.J. (1995) SH3-domains. Minding your p's and q's. *Curr. Biol.*, **5**, 364-367.
- McCollum, D., Monosov, E. and Subramani, S. (1993) The pas8 mutant of *Pichia pastoris* exhibits the peroxisomal protein import deficiencies of Zellweger syndrome cells--the PAS8 protein binds to the COOH-terminal tripeptide peroxisomal targeting signal, and is a member of the TPR protein family. *J Cell Biol*, **121**, 761-774.
- McNew, J.A. and Goodman, J.M. (1994) An oligomeric protein is imported into peroxisomes in vivo. *J Cell Biol*, **127**, 1245-1257.
- Morton, C.J. and Campbell, I.D. (1994) SH3 domains. Molecular 'Velcro'. *Curr Biol*, **4**, 615-617.
- Moser, A.B., Rasmussen, M., Naidu, S., Watkins, P.A., McGuinness, M., Hajra, A.K., Chen, G., Raymond, G., Liu, A., Gordon, D., Garnaas, K., Walton, D.S., Skjeldal, O.H., Guggenheim, M.A., Jackson, L.G., Elias, E.R. and Moser, H.W. (1995) Phenotype of patients with peroxisomal disorders subdivided into sixteen complementation groups. *J. Pediatr.*, **127**, 13-22.
- Mosser, J., Lutz, Y., Stoeckel, M.E., Sarde, C.O., Kretz, C., Douar, A.M., Lopez, J., Aubourg, P. and Mandel, J.L. (1994) The gene responsible for adrenoleukodystrophy encodes a peroxisomal membrane protein. *Hum Mol Genet*, **3**, 265-271.
- Motley, A.M., Hetteema, E.H., Hogenhout, E.M., Brites, P., ten Asbroek, A.L.M.A., Wijburg, F.A., Baas, F., Heijmans, H.S., Tabak, H.F., Wanders, R.J.A. and Distel, B. (1997) Rhizomelic chondrodysplasia punctata is a peroxisomal protein targeting disease caused by a non-functional PTS2 receptor. *Nat. Genet.*, **15**, 377-380.

- Musacchio, A., Gibson, T., Lehto, V.P. and Saraste, M. (1992) SH3--an abundant protein domain in search of a function. *FEBS Lett*, **307**, 55-61.
- Neuwald, A.F., Aravind, L., Spouge, J.L. and Koonin, E.V. (1999) AAA+: A class of chaperone-like ATPases associated with the assembly, operation, and disassembly of protein complexes. *Genome Res*, **9**, 27-43.
- Nicolay, K., Veenhuis, M., Douma, A.C. and Harder, W. (1987) A ³¹P NMR study of the internal pH of yeast peroxisomes. *Arch Microbiol*, **147**, 37-41.
- Nuttley, W.M., Brade, A.M., Eitzen, G.A., Veenhuis, M., Aitchison, J.D., Szilard, R.K., Glover, J.R. and Rachubinski, R.A. (1994) *PAY4*, a gene required for peroxisome assembly in the yeast *Yarrowia lipolytica*, encodes a novel member of a family of putative ATPases. *J. Biol. Chem.*, **269**, 556-566.
- Nuttley, W.M., Szilard, R.K., Smith, J.J., Veenhuis, M. and Rachubinski, R.A. (1995) The *PAH2* gene is required for peroxisome assembly in the methylotrophic yeast *Hansenula polymorpha* and encodes a member of the tetratricopeptide repeat family of proteins. *Gene*, **160**, 33-39.
- Osumi, T., Tsukamoto, T., Hata, S., Yokota, S., Miura, S., Fujiki, Y., Hijikata, M., Miyazawa, S. and Hashimoto, T. (1991) Amino-terminal presequence of the precursor of peroxisomal 3-ketoacyl-CoA thiolase is a cleavable signal peptide for peroxisomal targeting. *Biochem. Biophys. Res. Commun.*, **181**, 947-954.
- Otera, H., Harano, T., Honsho, M., Ghaedi, K., Mukai, S., Tanaka, A., Kawai, A., Shimizu, N. and Fujiki, Y. (2000) The mammalian peroxin Pex5pL, the longer isoform of the mobile peroxisome targeting signal (PTS) type 1 transporter, translocates the Pex7p.PTS2 protein complex into peroxisomes via its initial docking site, Pex14p. *J Biol Chem*, **275**, 21703-21714.
- Otera, H., Setoguchi, K., Hamasaki, M., Kumashiro, T., Shimizu, N. and Fujiki, Y. (2002) Peroxisomal targeting signal receptor Pex5p interacts with cargoes and import machinery components in a spatiotemporally differentiated manner: conserved Pex5p WXXXF/Y motifs are critical for matrix protein import. *Mol Cell Biol*, **22**, 1639-1655.
- Parsons, M., Furuya, T., Pal, S. and Kessler, P. (2001) Biogenesis and function of peroxisomes and glycosomes. *Mol Biochem Parasitol*, **115**, 19-28.
- Pawson, T. (1995) Protein modules and signalling networks. *Nature*, **373**, 573-580.
- Portsteffen, H., Beyer, A., Becker, E., Epplen, C., Pawlak, A., Kunau, W.-H. and Dodt, G. (1997) Human PEX1 is mutated in complementation group 1 of the peroxisome biogenesis disorders. *Nat. Genet.*, **17**, 449-452.
- Purdue, P.E. and Lazarow, P.B. (2001) Peroxisome biogenesis. *Annu Rev Cell Dev Biol*, **17**, 701-752.
- Purdue, P.E., Yang, X. and Lazarow, P.B. (1998) Pex18p and Pex21p, a novel pair of related peroxins essential for peroxisomal targeting by the PTS2 pathway. *J Cell Biol*, **143**, 1859-1869.
- Purdue, P.E., Zhang, J.W., Skoneczny, M. and Lazarow, P.B. (1997) Rhizomelic chondrodysplasia punctata is caused by deficiency of human PEX7, a homologue of the yeast PTS2 receptor. *Nat Genet*, **15**, 381-384.
- Rehling, P., Marzioch, M., Niesen, F., Wittke, E., Veenhuis, M. and Kunau, W.H. (1996) The import receptor for the peroxisomal targeting signal 2 (PTS2) in *Saccharomyces cerevisiae* is encoded by the *PAS7* gene. *Embo J*, **15**, 2901-2913.
- Reineke, U., Volkmer-Engert, R. and Schneider-Mergener, J. (2001) Applications of peptide arrays prepared by the SPOT-technology. *Curr Opin Biotechnol*, **12**, 59-64.
- Reuber, B.E., Germain-Lee, E., Collins, C.S., Morrell, J.C., Ameritunga, R., Moser, H.W., Valle, D. and Gould, S.J. (1997) Mutations in *PEX1* are the most common cause of peroxisome biogenesis disorders. *Nat. Genet.*, **17**, 445-448.

- Sacksteder, K.A. and Gould, S.J. (2000) The genetics of peroxisome biogenesis. *Annu Rev Genet*, **34**, 623-652.
- Santos, M.J., Imanaka, T., Shio, H. and Lazarow, P.B. (1988a) Peroxisomal integral membrane proteins in control and Zellweger fibroblasts. *J Biol Chem*, **263**, 10502-10509.
- Santos, M.J., Imanaka, T., Shio, H., Small, G.M. and Lazarow, P.B. (1988b) Peroxisomal membrane ghosts in Zellweger syndrome--aberrant organelle assembly. *Science*, **239**, 1536-1538.
- Schrader, M., Krieglstein, K. and Fahimi, H.D. (1998) Tubular peroxisomes in HepG2 cells: selective induction by growth factors and arachidonic acid. *Eur J Cell Biol*, **75**, 87-96.
- Shimizu, N., Itoh, R., Hirono, Y., Otera, H., Ghaedi, K., Tateishi, K., Tamura, S., Okumoto, K., Harano, T., Mukai, S. and Fujiki, Y. (1999) The peroxin Pex14p. cDNA cloning by functional complementation on a Chinese hamster ovary cell mutant, characterization, and functional analysis. *J Biol Chem*, **274**, 12593-12604.
- Shimozawa, N., Tsukamoto, T., Suzuki, Y., Orii, T., Shirayoshi, Y., Mori, T. and Fujiki, Y. (1992) A human gene responsible for Zellweger syndrome that affects peroxisome assembly. *Science*, **255**, 1132-1134.
- Smith, J.J., Marelli, M., Christmas, R.H., Vizeacoumar, F.J., Dilworth, D.J., Ideker, T., Galitski, T., Dimitrov, K., Rachubinski, R.A. and Aitchison, J.D. (2002) Transcriptome profiling to identify genes involved in peroxisome assembly and function. *J Cell Biol*, **158**, 259-271.
- Smith, J.J. and Rachubinski, R.A. (2001) A role for the peroxin Pex8p in Pex20p-dependent thiolase import into peroxisomes of the yeast *Yarrowia lipolytica*. *J Biol Chem*, **276**, 1618-1625.
- Snyder, W.B., Faber, K.N., Wenzel, T.J., Koller, A., Luers, G.H., Rangell, L., Keller, G.A. and Subramani, S. (1999a) Pex19p interacts with Pex3p and Pex10p and is essential for peroxisome biogenesis in *Pichia pastoris*. *Mol Biol Cell*, **10**, 1745-1761.
- Snyder, W.B., Koller, A., Choy, A.J., Johnson, M.A., Cregg, J.M., Rangell, L., Keller, G.A. and Subramani, S. (1999b) Pex17p is required for import of both peroxisome membrane and luminal proteins and interacts with Pex19p and the peroxisome targeting signal-receptor docking complex in *Pichia pastoris*. *Mol Biol Cell*, **10**, 4005-4019.
- Snyder, W.B., Koller, A., Choy, A.J. and Subramani, S. (2000) The peroxin Pex19p interacts with multiple, integral membrane proteins at the peroxisomal membrane. *J Cell Biol*, **149**, 1171-1178.
- South, S.T. and Gould, S.J. (1999) Peroxisome synthesis in the absence of preexisting peroxisomes. *J Cell Biol*, **144**, 255-266.
- South, S.T., Sacksteder, K.A., Li, X., Liu, Y. and Gould, S.J. (2000) Inhibitors of COPI and COPII do not block PEX3-mediated peroxisome synthesis. *J Cell Biol*, **149**, 1345-1360.
- Spong, A.P. and Subramani, S. (1993) Cloning and characterization of *PAS5*: a gene required for peroxisome biogenesis in the methylotrophic yeast *Pichia pastoris*. *J Cell Biol*, **123**, 535-548.
- Stein, K., Schell-Steven, A., Erdmann, R. and Rottensteiner, H. (2002) Interactions of Pex7p and Pex18p/Pex21p with the peroxisomal docking machinery: implications for the first steps in PTS2 protein import. *Mol Cell Biol*, **22**, 6056-6069.
- Subramani, S. (1998) Components involved in peroxisome import, biogenesis, proliferation, turnover, and movement. *Physiol Rev*, **78**, 171-188.

- Subramani, S., Koller, A. and Snyder, W.B. (2000) Import of peroxisomal matrix and membrane proteins. *Annu Rev Biochem*, **69**, 399-418.
- Swinkels, B.W., Gould, S.J., Bodnar, A.G., Rachubinski, R.A. and Subramani, S. (1991) A novel, cleavable peroxisomal targeting signal at the amino-terminus of the rat 3-ketoacyl-CoA thiolase. *Embo J*, **10**, 3255-3262.
- Szilard, R.K., Titorenko, V.I., Veenhuis, M. and Rachubinski, R.A. (1995) Pay32p of the Yeast *Yarrowia lipolytica* Is an Intraperoxisomal Component of the Matrix Protein Translocation Machinery. *J. Cell Biol.*, **131**, 1453-1469.
- Tam, Y.Y. and Rachubinski, R.A. (2002) *Yarrowia lipolytica* cells mutant for the PEX24 gene encoding a peroxisomal membrane peroxin mislocalize peroxisomal proteins and accumulate membrane structures containing both peroxisomal matrix and membrane proteins. *Mol Biol Cell*, **13**, 2681-2691.
- Tan, X., Waterham, H.R., Veenhuis, M. and Cregg, J.M. (1995) The *Hansenula polymorpha* PER8 gene encodes a novel peroxisomal integral membrane protein involved in proliferation. *J Cell Biol*, **128**, 307-319.
- Tanaka A, G.C., Arthur RR, Anderson SK, Kung HJ, Fujita DJ. (1987) DNA sequence encoding the amino-terminal region of the human c-src protein: implications of sequence divergence among src-type kinase oncogenes. *Mol. Cell Biol*, **7**, 1978-1983.
- Terlecky, S.R., Nuttley, W.M., McCollum, D., Sock, E. and Subramani, S. (1995) The *Pichia pastoris* peroxisomal protein PAS8p is the receptor for the C-terminal tripeptide peroxisomal targeting signal. *Embo J*, **14**, 3627-3634.
- Titorenko, V.I., Ogrzydziak, D.M. and Rachubinski, R.A. (1997) Four distinct secretory pathways serve protein secretion, cell surface growth, and peroxisome biogenesis in the yeast *Yarrowia lipolytica*. *Mol Cell Biol*, **17**, 5210-5226.
- Titorenko, V.I. and Rachubinski, R.A. (1998a) The endoplasmic reticulum plays an essential role in peroxisome biogenesis. *Trends Biochem Sci*, **23**, 231-233.
- Titorenko, V.I. and Rachubinski, R.A. (1998b) Mutants of the yeast *Yarrowia lipolytica* defective in protein exit from the endoplasmic reticulum are also defective in peroxisome biogenesis. *Mol Cell Biol*, **18**, 2789-2803.
- Titorenko, V.I., Smith, J.J., Szilard, R.K. and Rachubinski, R.A. (1998) Pex20p of the yeast *Yarrowia lipolytica* is required for the oligomerization of thiolase in the cytosol and for its targeting to the peroxisome. *J Cell Biol*, **142**, 403-420.
- Titorenko, V.I., Smith, J.J., Szilard, R.K. and Rachubinski, R.A. (2000) Peroxisome biogenesis in the yeast *Yarrowia lipolytica*. *Cell Biochem Biophys*, **32**, 21-26.
- Tong, A.H., Drees, B., Nardelli, G., Bader, G.D., Brannetti, B., Castagnoli, L., Evangelista, M., Ferracuti, S., Nelson, B., Paoluzi, S., Quondam, M., Zucconi, A., Hogue, C.W., Fields, S., Boone, C. and Cesareni, G. (2002) A combined experimental and computational strategy to define protein interaction networks for peptide recognition modules. *Science*, **295**, 321-324.
- Trotter, P.J. (2001) The genetics of fatty acid metabolism in *Saccharomyces cerevisiae*. *Annu Rev Nutr*, **21**, 97-119.
- Tsukamoto, T., Miura, S. and Fujiki, Y. (1991) Restoration by a 35K membrane protein of peroxisome assembly in a peroxisome-deficient mammalian cell mutant. *Nature*, **350**, 77-81.
- Tsukamoto, T., Miura, S., Nakai, T., Yokota, S., Shimozawa, N., Suzuki, Y., Orii, T., Fujiki, Y., Sakai, F., Bogaki, A. and et al. (1995) Peroxisome assembly factor-2, a putative ATPase cloned by functional complementation on a peroxisome-deficient mammalian cell mutant. *Nat Genet*, **11**, 395-401.

- Urquhart, A.J., Kennedy, D., Gould, S.J. and Crane, D.I. (2000) Interaction of Pex5p, the type 1 peroxisome targeting signal receptor, with the peroxisomal membrane proteins Pex14p and Pex13p. *J Biol Chem*, **275**, 4127-4136.
- van der Klei, I.J., Hilbrands, R.E., Kiel, J.A., Rasmussen, S.W., Cregg, J.M. and Veenhuis, M. (1998) The ubiquitin-conjugating enzyme Pex4p of *Hansenula polymorpha* is required for efficient functioning of the PTS1 import machinery. *Embo J*, **17**, 3608-3618.
- van der Klei, I.J., Hilbrands, R.E., Swaving, G.J., Waterham, H.R., Vrieling, E.G., Titorenko, V.I., Cregg, J.M., Harder, W. and Veenhuis, M. (1995) The *Hansenula polymorpha* PER3 gene is essential for the import of PTS1 proteins into the peroxisomal matrix. *J. Biol. Chem.*, **270**, 17229-17236.
- van der Leij, I., Franse, M.M., Elgersma, Y., Distel, B. and Tabak, H.F. (1993) PAS10 is a tetratricopeptide-repeat protein that is essential for the import of most matrix proteins into peroxisomes of *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA*, **90**, 11782-11786.
- van Roermund, C.W., Elgersma, Y., Singh, N., Wanders, R.J. and Tabak, H.F. (1995) The membrane of peroxisomes in *Saccharomyces cerevisiae* is impermeable to NAD(H) and acetyl-CoA under in vivo conditions. *Embo J*, **14**, 3480-3486.
- Veenhuis, M., Mateblowski, M., Kunau, W.-H. and Harder, W. (1987) Proliferation of microbodies in *Saccharomyces cerevisiae*. *Yeast*, **3**, 77-84.
- Voorn-Brouwer, T., van der Leij, I., Hemrika, W., Distel, B. and Tabak, H.F. (1993) Sequence of the PAS8 gene, the product of which is essential for biogenesis of peroxisomes in *Saccharomyces cerevisiae*. *Biochem. Biophys. Acta*, **1216**, 325-328.
- Walton, P.A., Hill, P.E. and Subramani, S. (1995) Import of stably folded proteins into peroxisomes. *Mol Biol Cell*, **6**, 675-683.
- Warren, D.S., Morrell, J.C., Moser, H.W., Valle, D. and Gould, S.J. (1998) Identification of PEX10, the Gene Defective in Complementation Group 7 of the Peroxisome-Biogenesis Disorders. *Am. J. Hum. Genet.*, **63**, 347-359.
- Waterham, H.R., de Vries, Y., Russel, K.A., Xie, W., Veenhuis, M. and Cregg, J.M. (1996) The *Pichia pastoris* PER6 gene product is a peroxisomal integral membrane protein essential for peroxisome biogenesis and has sequence similarity to the Zellweger syndrome protein PAF-1. *Mol Cell Biol*, **16**, 2527-2536.
- Waterham, H.R., Titorenko, V.I., Haima, P., Cregg, J.M., Harder, W. and Veenhuis, M. (1994) The *Hansenula polymorpha* PER1 gene is essential for peroxisome biogenesis and encodes a peroxisomal matrix protein with both carboxy- and amino-terminal targeting signals. *J Cell Biol*, **127**, 737-749.
- Wiebel, F.F. and Kunau, W.H. (1992) The Pas2 protein essential for peroxisome biogenesis is related to ubiquitin-conjugating enzymes. *Nature*, **359**, 73-76.
- Wiemer, E.A., Luers, G.H., Faber, K.N., Wenzel, T., Veenhuis, M. and Subramani, S. (1996) Isolation and characterization of Pas2p, a peroxisomal membrane protein essential for peroxisome biogenesis in the methylotrophic yeast *Pichia pastoris*. *J Biol Chem*, **271**, 18973-18980.
- Wiemer, E.A., Nuttley, W.M., Bertolaet, B.L., Li, X., Francke, U., Wheelock, M.J., Anne, U.K., Johnson, K.R. and Subramani, S. (1995) Human peroxisomal targeting signal-1 receptor restores peroxisomal protein import in cells from patients with fatal peroxisomal disorders. *J Cell Biol*, **130**, 51-65.
- Will, G.K., Soukupova, M., Hong, X., Erdmann, K.S., Kiel, J.A., Dodt, G., Kunau, W.H. and Erdmann, R. (1999) Identification and characterization of the human orthologue of yeast Pex14p. *Mol Cell Biol*, **19**, 2265-2277.

Yahraus, T., Braverman, N., Dodt, G., Kalish, J.E., Morrell, J.C., Moser, H.W., Valle, D. and Gould, S.J. (1996) The peroxisome biogenesis disorder group 4 gene, PXAAA1, encodes a cytoplasmic ATPase required for stability of the PTS1 receptor. *EMBO J.*, **15**, 2914-2923.