

## 7. References

- Abraham CR, McGraw WT, Slot F, Yamin R. Alpha 1-antichymotrypsin inhibits A beta degradation in vitro and in vivo. *Ann N Y Acad Sci.* 2000;920:245-8.
- Aderem A, Ulevitch RJ. Toll-like receptors in the induction of the innate immune response. *Nature.* 2000 Aug 17;406(6797):782-7.
- Akita K, Ohtsuki T, Nukada Y et al. Involvement of caspase-1 and caspase-3 in the production and processing of mature human interleukin 18 in monocytic THP.1 cells. *J Biol Chem* 1997 Oct 17;272(42):26595-603
- Akiyama H, Barger S, Barnum et al. Inflammation and Alzheimer's disease. *Neurobiol Aging* 2000 May-Jun;21(3):383-421.
- Allan SM, Rothwell NJ. Cytokines and acute neurodegeneration. *Nat Rev Neurosci.* 2001 Oct;2(10):734-44.
- Aloisi F, Ria F, Penna G, Adorini L. Microglia are more efficient than astrocytes in antigen processing and in Th1 but not Th2 cell activation. *J Immunol.* 1998 May 15;160(10):4671-80.
- Aloisi F. Immune function of microglia. *Glia.* 2001 Nov;36(2):165-79. Review.
- Arakawa H, Urisaka T, Tsuruta H, Amemiya Y, Kihara H, Ikai A. The kinetics of conformational changes of alpha 2-macroglobulin determined by time resolved X-ray solution scattering. *FEBS Lett.* 1994 Jan 10;337(2):171-4.
- Arendt T. Dysregulation of neuronal differentiation and cell cycle control in Alzheimer's disease. *J Neural Transm Suppl.* 2002;(62):77-85.
- Aschner M, Allen JW, Kimelberg HK, Lopachin RM, Streit WJ. Glial cells in neurotoxicity development. *Annu. Rev. Pharmacol. Toxicol.* 1999; 39:151-73.
- Asensio VC and Campbell IL. Chemokines in the CNS: plurifunctional mediators in diverse states. *Trndz Neurosci.* 1999; 22: 504-512.
- Baldwin AS Jr. The NF-kappa B and I kappa B proteins: new discoveries and insights. *Annu Rev Immunol.* 1996;14:649-83.
- Bales KR, Du Y, Dodel RC, Yan GM, Hamilton-Byrd E, Paul SM. The NF-kappaB/Rel family of proteins mediates Abeta-induced neurotoxicity and glial activation. *Brain Res Mol Brain Res.* 1998 Jun 1;57(1):63-72.
- Banisadr G, Queraud-Lesaux F, Boutterin M et al. Distribution, cellular localization and functional role of CCR2 chemokine receptors in adult rat brain. *J Neurochem.* 2002 Apr;81(2):257-69.
- Barrett AJ. Alpha 2-macroglobulin. *Methods Enzymol.* 1981;80 Pt C:737-54. Review.
- Barger SW, Harmon AD. Microglial activation by Alzheimer amyloid precursor protein and modulation by apolipoprotein E. *Nature* 1997; 388:878-81.
- Bartus RT, Dean RL 3rd, Beer B, Lippa AS. The cholinergic hypothesis of geriatric memory dysfunction. *Science.* 1982 Jul 30;217(4558):408-14. Review.

- Bauer J, Strauss S, Schreiter-Gasser U et al. Interleukin-6 and alpha-2-macroglobulin indicate an acute-phase state in Alzheimer's disease cortices. *FEBS Lett.* 1991 Jul 8;285(1):111-4
- Becher B, Prat A, Antel JP. Brain-immune connection: immuno-regulatory properties of CNS-resident cells. *Glia* 2000 Feb 15;29(4):293-304
- Beckman JS, Koppenol WH. Nitric oxide, superoxide, and peroxynitrite: the good, the bad, and the ugly. *Am J Physiol.* 1996 Nov;271(5 Pt 1):C1424-37. Review.
- Behl C, Davis JB, Lesley R, Schubert D. Hydrogen peroxide mediates amyloid beta protein toxicity. *Cell.* 1994 Jun 17;77(6):817-27.
- Bellinger FP, Madamba SG, Campbell IL, Siggins GR. Reduced long-term potentiation in the dentate gyrus of transgenic mice with cerebral overexpression of interleukin-6. *Neurosci Lett.* 1995 Sep 29;198(2):95-8.
- Beyreuther K, Pollicino P, Multhaup G et al.. Regulation and expression of the Alzheimer's beta/A4 amyloid protein precursor in health, disease, and Down's syndrome. *Ann N Y Acad Sci.* 1993 Sep 24;695:91-102. Review.
- Bhat NR, Zhang P, Lee JC, Hogan EL. Extracellular signal-regulated kinase and p38 subgroups of mitogen-activated protein kinases regulate inducible nitric oxide synthase and tumor necrosis factor-alpha gene expression in endotoxin-stimulated primary glial cultures. *J Neurosci.* 1998 Mar 1;18(5):1633-41.
- Blasko I, Veerhuis R, Stampfer-Kountchev M, et al. Costimulatory effects of interferon-gamma and interleukin-1beta or tumor necrosis factor alpha on the synthesis of Abeta1-40 and Abeta1-42 by human astrocytes. *Neurobiol Dis* 2000 Dec;7(6 Pt B):682-9
- Boyle EA, McGeer PL. Cellular immune response in multiple sclerosis plaques. *Am J Pathol* 1990 Sep;137(3):575-84
- Busciglio J, Gabuzda D, Matsudaira P, Yanker B. Generation of beta-amyloid in the secretory pathway in neuronal and nonneuronal cells. *Proc Natl Acad Sci U S A.* 1993 Mar 1;90(5):2092-6.
- Businaro R, Fabrizi C, Fumagalli L, Lauro GM. Synthesis and secretion of alpha 2-macroglobulin by human glioma established cell lines. *Exp Brain Res.* 1992;88(1):213-8.
- Buxbaum JD, Oishi M, Chen HI, et al. Cholinergic agonists and interleukin 1 regulate processing and the secretion of the Alzheimer beta/A4 amyloid protein precursor. *Proc Natl Acad Sci USA* 1992; 89:10075-8
- Calingasan NY, Park LC, Calo LL, Trifiletti RR, Gandy SE, Gibson GE. Induction of nitric oxide synthase and microglial responses precede selective cell death induced by chronic impairment of oxidative metabolism. *Am J Pathol.* 1998 Aug;153(2):599-610.
- Campbell IL, Abraham CR, Masliah E, et al. Neurologic disease induced in transgenic mice by cerebral overexpression of interleukin 6. *Proc Natl Acad Sci U S A* 1993 Nov 1;90(21):10061-5
- Cao Z, Henzel WJ, Gao X. IRAK: a kinase associated with the interleukin-1 receptor. *Science* 1996a Feb 23;271(5252):1128-31

- Cao Z, Xiong J, Takeuchi M, Kurama T, Goeddel DV. TRAF6 is a signal transducer for interleukin-1. *Nature* 1996b Oct 3;383(6599):443-6
- Chang RC, Chen W, Hudson P et al. Neurons reduce glial responses to lipopolysaccharide (LPS) and prevent injury of microglial cells from over-activation by LPS. *J Neurochem.* 2001 Feb;76(4):1042-9.
- Citron M, Diehl TS, Gordon G, Biere AL, Seubert P, Selkoe DJ. Evidence that the 42- and 40-amino acid forms of amyloid beta protein are generated from the beta-amyloid precursor protein by different protease activities. *Proc Natl Acad Sci U S A.* 1996 Nov 12;93(23):13170-5.
- Colasanti M, Persichini T, Di Pucchio T, Gremo F, Lauro GM. Human ramified microglial cells produce nitric oxide upon Escherichia coli lipopolysaccharide and tumor necrosis factor alpha stimulation. *Neurosci Lett.* 1995 Nov 17;200(2):144-6.
- Colasanti M, Persichini T, Menegazzi M et al. Induction of nitric oxide synthase mRNA expression. Suppression by exogenous nitric oxide. *J Biol Chem.* 1995 Nov 10;270(45):26731-3.
- Collart MA, Baeuerle P, Vassalli P. Regulation of tumor necrosis factor alpha transcription in macrophages: involvement of four kappa B-like motifs and of constitutive and inducible forms of NF-kappa B. *Mol Cell Biol.* 1990 Apr;10(4):1498-506.
- Colasanti M, Persichini T, Di Pucchio T, Gremo F, Lauro GM. Human ramified microglial cells produce nitric oxide upon Escherichia coli lipopolysaccharide and tumor necrosis factor alpha stimulation. *Neurosci Lett.* 1995 Nov 17;200(2):144-6.
- Combs CK, Johnson DE, Cannady SB, Lehman TM, Landreth GE. Identification of microglial signal transduction pathways mediating a neurotoxic response to amyloidogenic fragments of beta-amyloid and prion proteins. *J Neurosci.* 1999 Feb 1;19(3):928-39.
- Combs CK, Karlo JC, Kao SC, Landreth GE. beta-Amyloid stimulation of microglia and monocytes results in TNFalpha-dependent expression of inducible nitric oxide synthase and neuronal apoptosis. *J Neurosci.* 2001 Feb 15;21(4):1179-88.
- Conti B, Park LC, Calingasan NY et al. Cultures of astrocytes and microglia express interleukin 18. *Brain Res Mol Brain Res* 1999 Apr 6;67(1):46-52
- Cotter RL, Burke WJ, Thomas VS et al. Insights into the neurodegenerative process of Alzheimer's disease: a role for mononuclear phagocyte-associated inflammation and neurotoxicity. *J Leukoc Biol.* 1999 Apr;65(4):416-27. Review.
- Coughlan CM, McManus CM, Sharron M et al. Expression of multiple functional chemokine receptors and monocyte chemoattractant protein-1 in human neurons. *Neuroscience.* 2000;97(3):591-600.
- Cuadros MA, Navascues J. The origin and differentiation of microglial cells during development. *Prog Neurobiol.* 1998 Oct;56(2):173-89. Review.

- Daffada AA, Murray EJ, Young SP. Control of activator protein-1 and nuclear factor kappa B activity by interleukin-1, interleukin-6 and metals in HEPG2 cells. *Biochim Biophys Acta*. 1994 Jun 30;1222(2):234-40.
- da Silva Correia J, Soldau K, Christen U, Tobias PS, Ulevitch RJ. Lipopolysaccharide is in close proximity to each of the proteins in its membrane receptor complex; transfer from CD14 to TLR4 and MD-2. *J Biol Chem*. 2001 Jun 15;276 (24):21129-35.
- Dehmer T, Lindenau J, Haid S, Dichgans J, Schulz JB. Deficiency of inducible nitric oxide synthase protects against MPTP toxicity in vivo. *: J Neurochem* 2000 May;74(5):2213-6
- Depboylu C, Lohmuller F, Du Y et al. Alpha2-macroglobulin, lipoprotein receptor-related protein and lipoprotein receptor-associated protein and the genetic risk for developing Alzheimer's disease. *Neurosci Lett*. 2006 Jun 12;400(3):187-90.
- Dickson DW, Lee SC, Mattiace LA, Yen SH, Brosnan C. Microglia and cytokines in neurological disease, with special reference to AIDS and Alzheimer's disease. *Glia* 1993 Jan;7(1):75-83
- Dinarello CA. Role of pro- and anti-inflammatory cytokines during inflammation: experimental and clinical findings. *J Biol Regul Homeost Agents*. 1997 Jul-Sep;11(3):91-103.
- Dinarello CA. Interleukin-1, interleukin-1 receptors and interleukin-1 receptor antagonist. *Int Rev Immunol*. 1998;16(5-6):457-99.
- Dinarello CA, Novick D, Puren AJ et al. Overview of interleukin-18: more than an interferon-gamma inducing factor. *J Leukoc Biol* 1998 Jun;63(6):658-64
- Dovey HF, Suomensaari-Chrysler S, Lieberburg I, Sinha S, Keim PS. Cells with a familial Alzheimer's disease mutation produce authentic beta-peptide. *Neuroreport*. 1993 Aug;4(8):1039-42.
- Draheim HJ, Prinz M, Weber JR, Weiser T, Kettenmann H, Hanisch UK. Induction of potassium channels in mouse brain microglia: cells acquire responsiveness to pneumococcal cell wall components during late development. *Neuroscience* 1999;89(4):1379-90
- Du Y, Ni B, Glinn M et al. alpha2-Macroglobulin as a beta-amyloid peptide-binding plasma protein. *J Neurochem*. 1997 Jul;69(1):299-305.
- Du Y, Bales KR, Dodel RC, Liu X, Glinn MA, Horn JW, Little SP, Paul SM. Alpha2-macroglobulin attenuates beta-amyloid peptide 1-40 fibril formation and associated neurotoxicity of cultured fetal rat cortical neurons. *J Neurochem*. 1998 Mar;70(3):1182-8.
- Du ZY, Li XY. Inhibitory effects of indomethacin on interleukin-1 and nitric oxide production in rat microglia in vitro. *Int J Immunopharmacol*. 1999 Mar;21(3):219-25.
- Dudal S, Morissette C, Lacombe D, Tremblay P, Gervais F. Differences in the amyloid-beta-induced inflammatory response in microglia from C57BL/6 and A/J strains of mice. *J Neuroimmunol*. 2004 Aug;153(1-2):26-35.
- Eder C, Schilling T, Heinemann U, Haas D, Hailer N, Nitsch R. Morphological, immunophenotypical and electrophysiological properties of resting microglia in vitro. *Eur J Neurosci*. 1999 Dec;11(12):4251-61.

- Eikelenboom P, Zhan SS, van Gool WA, Allsop D. Inflammatory mechanisms in Alzheimer's disease. *Trends Pharmacol Sci.* 1994 Dec;15(12):447-50.
- Eikelenboom P, Rozemuller JM, van Muiswinkel FL. Inflammation and Alzheimer's disease: relationships between pathogenic mechanisms and clinical expression. *Exp Neurol* 1998 Nov;154(1):89-98
- El Khoury JB, Moore KJ, Means TK, Leung J, Terada K, Toft M, Freeman MW, Luster AD. CD36 mediates the innate host response to beta-amyloid. *J Exp Med.* 2003 Jun 16;197(12):1657-66.
- Evin G, Sernee MF, Masters CL. Inhibition of gamma-secretase as a therapeutic intervention for Alzheimer's disease: prospects, limitations and strategies. *CNS Drugs.* 2006;20(5):351-72. Review.
- Fabrizi C, Colosati M, Persichini T et al. Interferon gamma up-regulates alpha 2 macroglobulin expression in human astrocytoma cells. *J Neuroimmunol.* 1994 Aug;53(1):31-7.
- Fabrizi C, Businaro R, Lauro GM, Starace G, Fumagalli L. Activated alpha2macroglobulin increases beta-amyloid (25-35)-induced toxicity in LAN5 human neuroblastoma cells. *Exp Neurol.* 1999 Feb;155(2):252-9.
- Farber K, Kettenmann H. Physiology of microglial cells. *Brain Res Brain Res Rev.* 2005 Apr;48(2):133-43. Review.
- Fattori E, Lazzaro D, Musiani P et al. IL-6 expression in neurons of transgenic mice causes reactive astrocytosis and increase in ramified microglial cells but no neuronal damage. *Eur J Neurosci* 1995 Dec 1;7(12):2441-9
- Ferrari D, Chiozzi P, Falzoni S, Hanau S, Di Virgilio F. Purinergic modulation of interleukin-1 beta release from microglial cells stimulated with bacterial endotoxin. *J Exp Med.* 1997 Feb 3;185(3):579-82.
- Fiala M, Zhang L, Gan X, Sherry B, Taub D, Graves MC, Hama S, Way D, Weinand M, Witte M, Lorton D, Kuo YM, Roher AE. Amyloid-beta induces chemokine secretion and monocyte migration across a human blood--brain barrier model. *Mol Med.* 1998 Jul;4(7):480-9.
- Fiebich BL, Hofer TJ, Lieb K et al. The non-steroidal anti-inflammatory drug tepoxalin inhibits interleukin-6 and alpha1-anti-chymotrypsin synthesis in astrocytes by preventing degradation of IkappaB-alpha. *Neuropharmacology.* 1999 Sep;38(9):1325-33.
- Figler NL, Strickland DK, Allietta M, Gonias SL. Immunoelectron microscopy studies with a monoclonal antibody directed against a receptor recognition site epitope in human alpha 2-macroglobulin. *J Struct Biol.* 1991 Jun;106(3):237-42.
- Fillit H, Ding WH, Buee L et al. Elevated circulating tumor necrosis factor levels in Alzheimer's disease. *Neurosci Lett.* 1991 Aug 19;129(2):318-20.
- Gadient RA, Otten UH. Interleukin-6 (IL-6)--a molecule with both beneficial and destructive potentials. *Prog Neurobiol* 1997 Aug;52(5):379-90

- Garter U, Strauss S, Jonas U et al. Alpha 2-macroglobulin synthesis in interleukin-6-stimulated human neuronal (SH-SY5Y neuroblastoma) cells. Potential significance for the processing of Alzheimer beta-amyloid precursor protein. *FEBS Lett.* 1991 Apr 22;282(1):127-31.
- Ganter U, Strauss S, Jonas U, Weidemann A, Beyreuther K, Volk B, Berger M, Bauer J. Alpha 2-macroglobulin synthesis in interleukin-6-stimulated human neuronal (SH-SY5Y neuroblastoma) cells. Potential significance for the processing of Alzheimer beta-amyloid precursor protein. *FEBS Lett.* 1991 Apr 22;282(1):127-31.
- Gao F, Bales KR, Dodel RC et al. NF-kappaB mediates IL-1beta-induced synthesis/release of alpha2-macroglobulin in a human glial cell line. *Brain Res Mol Brain Res.* 2002 Sep 30;105(1-2):108-14.
- Garrington TP, Johnson GL. Organization and regulation of mitogen-activated protein kinase signaling pathways. *Curr Opin Cell Biol.* 1999 Apr;11(2):211-8.
- Garton MJ, Keir G, Lakshmi MV, Thompson EJ. Age-related changes in cerebrospinal fluid protein concentrations. *J Neurol Sci.* 1991 Jul;104(1):74-80.
- Gasic-Milenkovic J, Dukic-Stefanovic S, Deuther-Conrad W, Gartner U, Munch G. Beta-amyloid peptide potentiates inflammatory responses induced by lipopolysaccharide, interferon -gamma and 'advanced glycation endproducts' in a murine microglia cell line. *Eur J Neurosci.* 2003 Feb;17(4):813-21.
- Gebicke-Haerter PJ, Spleiss O, Ren LQ et al. Microglial chemokines and chemokine receptors. *Prog Brain Res.* 2001;132:525-32. Review.
- Gehrmann J, Banati RB, Wiessner C, Hossmann KA, Kreutzberg GW. Reactive microglia in cerebral ischaemia: an early mediator of tissue damage? *Neuropathol Appl Neurobiol.* 1995 Aug;21(4):277-89. Review.
- Gehrmann J, Matsumoto Y, Kreutzberg GW. Microglia: intrinsic immune effector cell of the brain. *Brain Res Brain Res Rev.* 1995 Mar;20(3):269-87. Review.
- Gelman BB. Diffuse microgliosis associated with cerebral atrophy in the acquired immunodeficiency syndrome. *Ann Neurol* 1993 Jul;34(1):65-70
- Gitter BD, Cox LM, Rydel RE, May PC. Amyloid beta peptide potentiates cytokine secretion by interleukin-1 beta-activated human astrocytoma cells. *Proc Natl Acad Sci U S A.* 1995 Nov 7;92(23):10738-41.
- Giulian D, Haverkamp LJ, Yu JH et al. Specific domains of beta-amyloid from Alzheimer plaque elicit neuron killing in human microglia. *J Neurosci.* 1996 Oct 1;16(19):6021-37.
- Giulian D, Haverkamp LJ, Yu J, Karshin W et al. The HHQK domain of beta-amyloid provides a structural basis for the immunopathology of Alzheimer's disease. *J Biol Chem.* 1998 Nov 6;273(45):29719-26.
- Giulian D. Microglia and the immune pathology of Alzheimer disease. *Am J Hum Genet* 1999 Jul;65(1):13-8

- Goldgaber D, Harris HW, Hla T, et al. Interleukin 1 regulates synthesis of amyloid beta-protein precursor mRNA in human endothelial cells. *Proc Natl Acad Sci USA* 1989; 86:7606-10.
- Gollin PA, Kalaria RN, Eikelenboom P, Rozemuller A, Perry G. Alpha 1-antitrypsin and alpha 1-antichymotrypsin are in the lesions of Alzheimer's disease. *Neuroreport*. 1992 Feb;3(2):201-3.
- Good PF, Werner P, Hsu A, Olanow CW, Perl DP. Evidence of neuronal oxidative damage in Alzheimer's disease. *Am J Pathol*. 1996 Jul;149(1):21-8.
- Goodwin JL, Uemura E, Cunnick JE. Microglial release of nitric oxide by the synergistic action of beta-amyloid and IFN-gamma. *Brain Res*. 1995 Sep 18;692(1-2):207-14.
- Grammas P, Ovase R. Inflammatory factors are elevated in brain microvessels in Alzheimer's disease. *Neurobiol Aging*. 2001 Nov-Dec;22(6):837-42.
- Gravina SA, Ho L, Eckman CB et al. Amyloid beta protein (A beta) in Alzheimer's disease brain. Biochemical and immunocytochemical analysis with antibodies specific for forms ending at A beta 40 or A beta 42(43). *J Biol Chem*. 1995 Mar 31;270(13):7013-6.
- Griffin WS, Stanley LC, Ling C, et al. brain interleukin 1 and S-100 immunoreactivity are elevated in Down syndrome and Alzheimer disease. *Proc Natl Acad Sci USA* 1989; 86:7611-5.
- Griffin WS, Sheng JG, Roberts GW, Mrak RE. IL-1 expression in different plaque types in Alzheimer's disease: significance in plaque evolution. *J Neuropathol Exp Neurol* 1995; 54:276-81
- Griffin WS, Sheng JG, Royston MC, et al. Glial-neuronal interactions in Alzheimer's disease: the potential role of a "cytokine cycle" in disease progression. *Brain Pathol* 1998; 8:65-72.
- Haass C, Schlossmacher MG, Hung AY et al. Amyloid beta-peptide is produced by cultured cells during normal metabolism. *Nature*. 1992 Sep 24;359(6393):322-5.
- Hailer NP, Jarhult JD, Nitsch R. Resting microglial cells in vitro: analysis of morphology and adhesion molecule expression in organotypic hippocampal slice cultures. *Glia*. 1996 Dec;18(4):319-31.
- Hailer NP, Wirjatijasa F, Roser N et al. Astrocytic factors protect neuronal integrity and reduce microglial activation in an in vitro model of N-methyl-D-aspartate-induced excitotoxic injury in organotypic hippocampal slice cultures. *Eur J Neurosci*. 2001 Jul;14(2):315-26.
- Hanisch UK, Prinz M, Angstwurm K, et al. The protein tyrosine kinase inhibitor AG126 prevents the massive microglial cytokine induction by pneumococcal cell walls. *Eur J Immunol*. 2001 Jul;31(7):2104-15.
- Hanisch UK. Effects of interleukin-2 and interferons on the nervous system. In: *Psychoneuroimmunology*, 3<sup>rd</sup> edition (Ader R, Felten DL & Cohen N, eds), Academic Press, San Diego, 2001; 585-631.
- Hanisch UK. Microglia as a source and target of cytokines. *Glia* 2002; 40:140-155.
- Hardy JA, Higgins GA. Alzheimer's disease: the amyloid cascade hypothesis. *Science*. 1992 Apr 10;256(5054):184-5. Review.
- Hardy J, Selkoe DJ. The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science*. 2002 Jul 19;297(5580):353-6. Review

- Harper JD, Lansbury PT Jr. Models of amyloid seeding in Alzheimer's disease and scrapie: mechanistic truths and physiological consequences of the time-dependent solubility of amyloid proteins. *Annu Rev Biochem.* 1997;66:385-407. Review.
- Harrison JK, Jiang Y, Chen S, Xia Y et al. Role for neuronally derived fractalkine in mediating interactions between neurons and CX3CR1-expressing microglia. *Proc Natl Acad Sci U S A.* 1998 Sep 1;95(18):10896-901.
- Hausler KG, Prinz M, Nolte C et al. Interferon-gamma differentially modulates the release of cytokines and chemokines in lipopolysaccharide- and pneumococcal cell wall-stimulated mouse microglia and macrophages. *Eur J Neurosci.* 2002 Dec;16(11):2113-22.
- Hesslgrenner J, Horuk R. Chemokine and chemokine receptor \_expression in the central nervous system. *J. Neurovirol.* 1999; 5: 13-26.
- Heyser CJ, Maslia E, Samimi A, Campbell IL, Gold LH. Progressive decline in avoidance learning paralleled by inflammatory neurodegeneration in transgenic mice expressing interleukin 6 in the brain. *Proc Natl Acad Sci U S A.* 1997 Feb 18;94(4):1500-5.
- Higuchi M, Ito T, Imai Y et al. Expression of the alpha 2-macroglobulin-encoding gene in rat brain and cultured astrocytes. *Gene.* 1994 Apr 20;141(2):155-62.
- Hirota H, Kiyama H, Kishimoto T, Taga T. Accelerated Nerve Regeneration in Mice by upregulated expression of interleukin (IL) 6 and IL-6 receptor after trauma. *J Exp Med* 1996 Jun 1;183(6):2627-34
- Hiscott J, Marois J, Garoufalis J et al. Characterization of a functional NF-kappa B site in the human interleukin 1 beta promoter: evidence for a positive autoregulatory loop. *Mol Cell Biol.* 1993 Oct;13(10):6231-40.
- Hoek RM, Ruuls SR, Murphy CA et al. Down-regulation of the macrophage lineage through interaction with OX2 (CD200). *Science.* 2000 Dec 1;290(5497):1768-71.
- Hoffmann A, Kann O, Ohlemeyer C, Hanisch UK, Kettenmann H. Elevation of basal intracellular calcium as a central element in the activation of brain macrophages (microglia): suppression of receptor-evoked calcium signaling and control of release function. *J Neurosci.* 2003 Jun 1;23(11):4410-9.
- Hollenbach E, Ackermann S, Hyman BT, Rebeck GW. Confirmation of an association between a polymorphism in exon 3 of the low-density lipoprotein receptor-related protein gene and Alzheimer's disease. *Neurology.* 1998 Jun;50(6):1905-7.
- Hopkins SJ, Rothwell NJ. Cytokines and the nervous system. I: \_Expression and recognition. *Trends Neurosci* 1995 Feb;18(2):83-8
- Horuk R, Martin AW, Wang Z et al. Expression of chemokine receptors by subsets of neurons in the central nervous system. *J Immunol.* 1997 Mar 15;158(6):2882-90.
- Howlett DR, Jennings KH, Lee DC et al. Aggregation state and neurotoxic properties of Alzheimer beta-amyloid peptide. *Neurodegeneration.* 1995 Mar;4(1):23-32.

- Hughes SR, Khorkova O, Goyal S et al. Alpha2-macroglobulin associates with beta-amyloid peptide and prevents fibril formation. *Proc Natl Acad Sci U S A.* 1998 Mar 17;95(6):3275-80.
- Hull M, Berger M, Volk B, Bauer J. Occurrence of interleukin-6 in cortical plaques of Alzheimer's disease patients may precede transformation of diffuse into neuritic plaques. *Ann N Y Acad Sci.* 1996 Jan 17;777:205-12.
- Ishizuka K, Kimura T, Igata-yi R, Katsuragi S, Takamatsu J, Miyakawa T. Identification of monocyte chemoattractant protein-1 in senile plaques and reactive microglia of Alzheimer's disease. *Psychiatry Clin Neurosci.* 1997 Jun;51(3):135-8.
- Iwatsubo T, Odaka A, Suzuki N, Mizusawa H, Nukina N, Ihara Y. Visualization of A beta 42(43) and A beta 40 in senile plaques with end-specific A beta monoclonals: evidence that an initially deposited species is A beta 42(43). *Neuron.* 1994 Jul;13(1):45-53.
- Iwatsubo T, Mann DM, Odaka A, Suzuki N, Ihara Y. Amyloid beta protein (A beta) deposition: A beta 42(43) precedes A beta 40 in Down syndrome. *Ann Neurol.* 1995 Mar;37(3):294-9.
- Iwatsubo T, Saido TC, Mann DM, Lee VM, Trojanowski JQ. Full-length amyloid-beta (1-42(43)) and amino-terminally modified and truncated amyloid-beta 42(43) deposit in diffuse plaques. *Am J Pathol.* 1996 Dec;149(6):1823-30.
- James K, van den Haan J, Lens S, Farmer K. Preliminary studies on the interaction of TNF alpha and IFN gamma with alpha 2-macroglobulin. *Immunol Lett.* 1992 Mar;32(1):49-57.
- James K, Milne I, Cunningham A, Elliott SF. The effect of alpha 2 macroglobulin in commercial cytokine assays. *J Immunol Methods.* 1994 Jan 12;168(1):33-7.
- Janabi N, Hau I, Tardieu M. Negative feedback between prostaglandin and alpha- and beta-chemokine synthesis in human microglial cells and astrocytes. *J Immunol.* 1999 Feb 1;162(3):1701-6.
- Jantzen PT, Connor KE, DiCarlo G et al. Microglial activation and beta -amyloid deposit reduction caused by a nitric oxide-releasing nonsteroidal anti-inflammatory drug in amyloid precursor protein plus presenilin-1 transgenic mice. *J Neurosci* 2002 Mar 15;22(6):2246-54
- Jarrett JT, Berger EP, Lansbury PT Jr. The carboxy terminus of the beta amyloid protein is critical for the seeding of amyloid formation: implications for the pathogenesis of Alzheimer's disease. *Biochemistry.* 1993 May 11;32(18):4693-7.
- Johnstone M, Gearing AJ, Miller KM. A central role for astrocytes in the inflammatory response to beta-amyloid; chemokines, cytokines and reactive oxygen species are produced. *J Neuroimmunol.* 1999 Jan 1;93(1-2):182-93.
- Kang DE, Saitoh T, Chen X et al. Genetic association of the low-density lipoprotein receptor-related protein gene (LRP), an apolipoprotein E receptor, with late-onset Alzheimer's disease. *Neurology.* 1997 Jul;49(1):56-61.
- Kawai T, Adachi O, Ogawa T, Takeda K, Akira S. Unresponsiveness of MyD88-deficient mice to endotoxin. *Immunity.* 1999 Jul;11(1):115-22.
- Kettenmann H, Banati R, Walz W. Electrophysiological behavior of microglia. *Glia.* 1993 Jan;7(1):93-101.

- Kirschner DA, Inouye H, Duffy LK et al. Synthetic peptide homologous to beta protein from Alzheimer disease forms amyloid-like fibrils in vitro. *Proc Natl Acad Sci U S A.* 1987 Oct;84(19):6953-7.
- Klegeris A, McGeer PL. Cyclooxygenase and 5-lipoxygenase inhibitors protect against mononuclear phagocyte neurotoxicity. *Neurobiol Aging* 2002 Sep-Oct;23(5):787-94
- Klein WL, Krafft GA, Finch CE. Targeting small Abeta oligomers: the solution to an Alzheimer's disease conundrum? *Trends Neurosci* 2001 Apr;24(4):219-24
- Kojima H, Takeuchi M, Ohta T et al. Interleukin-18 activates the IRAK-TRAF6 pathway in mouse EL-4 cells. *Biochem Biophys Res Commun* 1998 Mar 6;244(1):183-6
- Kopf M, Baumann H, Freer G, et al. Impaired immune and acute-phase responses in interleukin-6-deficient mice. *Nature* 1994 Mar 24;368(6469):339-42
- Korotzer AR, Watt J, Cribbs D et al. Cultured rat microglia express C1q and receptor for C1q: implications for amyloid effects on microglia. *Exp Neurol* 1995 Aug;134(2):214-21
- Koshinaga M, Katayama Y, Fukushima M et al. Rapid and widespread microglial activation induced by traumatic brain injury in rat brain slices. *J Neurotrauma* 2000 Mar;17(3):185-92
- Kreutzberg GW. Microglia: a sensor for pathological events in the CNS. *Trends Neurosci* 1996 Aug;19(8):312-8
- LaMarre J, Wollenberg GK, Gonias SL, Hayes MA. Cytokine binding and clearance properties of proteinase-activated alpha 2-macroglobulins. *Lab Invest.* 1991 Jul;65(1):3-14. Review.
- Lambert MP, Barlow AK, Chromy BA et al. Diffusible, nonfibrillar ligands derived from Abeta1-42 are potent central nervous system neurotoxins. *Proc Natl Acad Sci U S A.* 1998 May 26;95(11):6448-53.
- Lanzrein AS, Johnson CM, Perry VH et al. Longitudinal study of inflammatory factors in serum, cerebrospinal fluid, and brain tissue in Alzheimer disease: interleukin-1beta, interleukin-6, interleukin-1 receptor antagonist, tumor necrosis factor-alpha, the soluble tumor necrosis factor receptors I and II, and alpha1-antichymotrypsin. *Alzheimer Dis Assoc Disord.* 1998 Sep;12(3):215-27.
- Lawrence AD, Sahakian BJ. The cognitive psychopharmacology of Alzheimer's disease: focus on cholinergic systems. *Neurochem Res.* 1998 May;23(5):787-94. Review.
- Lawson LJ, Perry VH, Dri P, Gordon S. Heterogeneity in the distribution and morphology of microglia in the normal adult mouse brain. *Neuroscience*. 1990;39(1):151-70.
- Lawson LJ, Perry VH, Gordon S. Turnover of resident microglia in the normal adult mouse brain. *Neuroscience*. 1992;48(2):405-15.
- Lee JK, Kim SH, Lewis EC et al. Differences in signaling pathways by IL-1beta and IL-18. *Proc Natl Acad Sci U S A.* 2004 Jun 8;101(23):8815-20.
- Legres LG, Pochon F, Baray M, Gay F, Chouaib S, Delain E. Evidence for the binding of a biologically active interleukin-2 to human alpha 2-macroglobulin. *J Biol Chem.* 1995 Apr 14;270(15):8381-4.

- Lemere CA, Blusztajn JK, Yamaguchi H et al. Sequence of deposition of heterogeneous amyloid beta-peptides and APO E in Down syndrome: implications for initial events in amyloid plaque formation. *Neurobiol Dis.* 1996 Feb;3(1):16-32.
- Li S. et al., 2002. IRAK-4: a novel member of the IRAK family with the properties of an IRAK-kinase. *Proc Natl Acad Sci U S A.* 99(8):5567-72.
- Li Y, Liu L, Kang J, et al. Neuronal-glial interactions mediated by IL-1 enhance neuronal acetylcholinesterase activity and mRNA \_expression. *J Neurosci.* 2000;20:149-55.
- Li Y, Liu L, Barger SW, Griffin WS. Interleukin-1 mediates pathological effects of microglia on tau phosphorylation and on synaptophysin synthesis in cortical neurons through a p38-MAPK pathway. *J Neurosci* 2003 Mar 1;23(5):1605-11
- Libermann TA, Baltimore D. Activation of interleukin-6 gene expression through the NF-kappa B transcription factor. *Mol Cell Biol.* 1990 May;10(5):2327-34.
- Lichtenthaler SF, Ida N, Multhaup G, Masters CL, Beyreuther K. Mutations in the transmembrane domain of APP altering gamma-secretase specificity. *Biochemistry.* 1997 Dec 9;36(49):15396-403.
- Lieb K, Engels S, Fiebich BL. Inhibition of LPS-induced iNOS and NO synthesis in primary rat microglial cells. *Neurochem Int.* 2003 Jan;42(2):131-7.
- Lim GP, Yang F, Chu T et al. Ibuprofen suppresses plaque pathology and inflammation in a mouse model for Alzheimer's disease. *J Neurosci* 2000 Aug 1;20(15):5709-14
- Lim GP, Chu T, Yang F, Beech W, Frautschy SA, Cole GM. The curry spice curcumin reduces oxidative damage and amyloid pathology in an Alzheimer transgenic mouse. *J Neurosci.* 2001 Nov 1;21(21):8370-7.
- Liu B, Wang K, Gao HM, Mandavilli B, Wang JY, Hong JS. Molecular consequences of activated microglia in the brain: overactivation induces apoptosis. *J Neurochem.* 2001 Apr;77(1):182-9.
- Loddick SA, Turnbull AV, Rothwell NJ. Cerebral interleukin-6 is neuroprotective during permanent focal cerebral ischemia in the rat. *J Cereb Blood Flow Metab* 1998 Feb;18(2):176-9
- Loddick SA, Liu C, Takao T, Hashimoto K, De Souza EB. Interleukin-1 receptors: cloning studies and role in central nervous system disorders. *Brain Res Brain Res Rev.* 1998 May;26(2-3):306-19.
- Lomaga MA, Yeh WC, Sarosi I et al. TRAF6 deficiency results in osteopetrosis and defective interleukin-1, CD40, and LPS signaling. *Genes Dev.* 1999 Apr 15;13(8):1015-24.
- Lomakin A, Teplow DB, Kirschner DA, Benedek GB. Kinetic theory of fibrillogenesis of amyloid beta-protein. *Proc Natl Acad Sci U S A.* 1997 Jul 22;94(15):7942-7.
- Lorenzo A, Yankner BA. Beta-amyloid neurotoxicity requires fibril formation and is inhibited by congo red. *Proc Natl Acad Sci U S A.* 1994 Dec 6;91(25):12243-7.
- Lorton D, Kocsis JM, King L, Madden K, Brunden KR. beta-Amyloid induces increased release of interleukin-1 beta from lipopolysaccharide-activated human monocytes. *J Neuroimmunol.* 1996 Jun;67(1):21-9.

- Lue LF, Rydel R, Brigham EF et al. Inflammatory repertoire of Alzheimer's disease and nondemented elderly microglia in vitro. *Glia*. 2001 Jul;35(1):72-9.
- Lue LF, Walker DG, Rogers J. Modeling microglial activation in Alzheimer's disease with human postmortem microglial cultures. *Neurobiol Aging*. 2001 Nov-Dec;22(6):945-56.
- Lysiak JJ, Hussaini IM, Webb DJ, Glass WF 2nd, Allietta M, Gonias SL. Alpha 2-macroglobulin functions as a cytokine carrier to induce nitric oxide synthesis and cause nitric oxide-dependent cytotoxicity in the RAW 264.7 macrophage cell line. *J Biol Chem*. 1995 Sep 15;270(37):21919-27.
- Mackenzie IR, Munoz DG. Nonsteroidal anti-inflammatory drug use and Alzheimer-type pathology in aging. *Neurology*. 1998 Apr;50(4):986-90.
- Malaguarnera L, Motta M, Di Rosa M, Anzaldi M, Malaguarnera M. Interleukin-18 and transforming growth factor-beta 1 plasma levels in Alzheimer's disease and vascular dementia. *Neuropathology*. 2006 Aug;26(4):307-12.
- Marz P, Herget T, Lang E, Otten U, Rose-John S. Activation of gp130 by IL-6/soluble IL-6 receptor induces neuronal differentiation. *Eur J Neurosci* 1997 Dec;9(12):2765-73
- Marzolo MP, von Bernhardi R, Bu G, Inestrosa NC. Expression of alpha(2)-macroglobulin receptor/low density lipoprotein receptor-related protein (LRP) in rat microglial cells. *J Neurosci Res*. 2000 May 1;60(3):401-11.
- McDonald DR, Bamberger ME, Combs CK, Landreth GE. beta-Amyloid fibrils activate parallel mitogen-activated protein kinase pathways in microglia and THP1 monocytes. *J Neurosci*. 1998 Jun 15;18(12):4451-60.
- McGeer PL, Kawamata T, Walker DG, Akiyama H, Tooyama I, McGeer EG. Microglia in degenerative neurological disease. *Glia* 1993 Jan;7(1):84-92
- McGeer EG, McGeer PL. Brain inflammation in Alzheimer disease and the therapeutic implications. *Curr Pharm Des* 1999 Oct;5(10):821-36
- McGeer PL, Rogers J, McGeer EG. Inflammation, anti-inflammatory agents and Alzheimer disease: the last 12 years. *J Alzheimers Dis*. 2006;9(3 Suppl):271-6. Review.
- McManus CM, Brosnan CF, Berman JW. Cytokine induction of MIP-1 alpha and MIP-1 beta in human fetal microglia. *J Immunol* 1998 Feb 1;160(3):1449-55
- McMillian M, Kong LY, Sawin SM, et al. Selective killing of cholinergic neurons by microglial activation in basal forebrain mixed neuronal/glial cultures. *Biochem Biophys Res Commun*. 1995 Oct 13;215(2):572-7.
- Meda L, Cassatella MA, Szendrei GI et al. Activation of microglial cells by beta-amyloid protein and interferon-gamma. *Nature*. 1995 Apr 13; 374(6523):647-50.
- Mertsch K, Hanisch UK, Kettenmann H, Schnitzer J. Characterization of microglial cells and their response to stimulation in an organotypic retinal culture system. *J Comp Neurol*. 2001 Mar 5;431(2):217-27.

- Miranda S, Opazo C, Larrondo LF et al. The role of oxidative stress in the toxicity induced by amyloid beta-peptide in Alzheimer's disease. *Prog Neurobiol.* 2000 Dec;62(6):633-48. Review.
- Misra UK, Chu CT, Rubenstein DS, Gawdi G, Pizzo SV. Receptor-recognized alpha 2-macroglobulin-methylamine elevates intracellular calcium, inositol phosphates and cyclic AMP in murine peritoneal macrophages. *Biochem J.* 1993 Mar 15;290 ( Pt 3):885-91.
- Misra UK, Chu CT, Gawdi G, Pizzo SV. The relationship between low density lipoprotein-related protein/alpha 2-macroglobulin (alpha 2M) receptors and the newly described alpha 2M signaling receptor. *J Biol Chem.* 1994 Jul 15;269(28):18303-6.
- Misra UK, Pizzo SV. Induction of cyclooxygenase-2 synthesis by ligation of the macrophage alpha(2)-macroglobulin signalling receptor. *Cell Signal.* 2001 Nov;13(11):801-8.
- Misra UK, Pizzo SV. Regulation of cytosolic phospholipase A2 activity in macrophages stimulated with receptor-recognized forms of alpha 2-macroglobulin: role in mitogenesis and cell proliferation. *J Biol Chem.* 2002 Feb 8;277(6):4069-78.
- Morihara T, Teter B, Yang F et al. Ibuprofen suppresses interleukin-1beta induction of pro-amyloidogenic alpha1-antichymotrypsin to ameliorate beta-amyloid (Abeta) pathology in Alzheimer's models. *Neuropsychopharmacology.* 2005 Jun;30(6):1111-20.
- Molina-Holgado E, Ortiz S, Molina-Holgado F, Guaza C. Induction of COX-2 and PGE(2) biosynthesis by IL-1beta is mediated by PKC and mitogen-activated protein kinases in murine astrocytes. *Br J Pharmacol* 2000 Sep;131(1):152-9
- Mrak RE, Sheng JG, Griffin WS. Glial cytokines in Alzheimer's disease: review and pathogenic implications. *Hum Pathol.* 1995 Aug;26(8):816-23. Review.
- Muehlhauser F, Liebl U, Kuehl S et al. Aggregation-Dependent interaction of the Alzheimer's beta-amyloid and microglia. *Clin Chem Lab Med.* 2001 Apr;39(4):313-6.
- Muller JM, Ziegler-Heitbrock HW, Baeuerle PA. Nuclear factor kappa B, a mediator of lipopolysaccharide effects. *Immunobiology.* 1993 Apr;187(3-5):233-56.
- Murphy GM Jr, Yang L, Cordell B. Macrophage colony-stimulating factor augments beta-amyloid-induced interleukin-1, interleukin-6, and nitric oxide production by microglial cells. *J Biol Chem.* 1998 Aug 14;273(33):20967-71.
- Nagai A, Nakagawa E, Hatori K et al. Generation and characterization of immortalized human microglial cell lines: expression of cytokines and chemokines. *Neurobiol Dis.* 2001 Dec;8(6):1057-68.
- Nakamura Y, Si QS, Kataoka K. Lipopolysaccharide-induced microglial activation in culture: temporal profiles of morphological change and release of cytokines and nitric oxide. *Neurosci Res* 1999 Nov;35(2):95-100
- Netland EE, Newton JL, Majocha RE, Tate BA. Indomethacin reverses the microglial response to amyloid beta-protein. *Neurobiol Aging* 1998 May-Jun;19(3):201-4

- Neumann H, Misgeld T, Matsumuro K, Wekerle H. Neurotrophins inhibit major histocompatibility class II inducibility of microglia: involvement of the p75 neurotrophin receptor. *Proc Natl Acad Sci U S A.* 1998 May 12;95(10):5779-84.
- Nimmerjahn A, Kirchhoff F, Helmchen F. Resting microglial cells are highly dynamic surveillants of brain parenchyma in vivo. *Science.* 2005 May 27;308(5726):1314-8.
- Norenberg W, Gebicke-Haerter PJ, Illes P. Inflammatory stimuli induce a new K<sup>+</sup> outward current in cultured rat microglia. *Neurosci Lett.* 1992 Dec 7;147(2):171-4.
- O'Barr S, Cooper NR. The C5a complement activation peptide increases IL-1 $\beta$  and IL-6 release from amyloid-beta primed human monocytes: implications for Alzheimer's disease. *J Neuroimmunol.* 2000 Sep 22;109(2):87-94.
- Ohno M, Higashi Y, Suzuki K. Microglial cell response to neuronal degeneration in the brain of brindled mouse. *Brain Res Dev Brain Res* 1992 May 22;67(1):37-45
- O'Neill LA, Greene C. Signal transduction pathways activated by the IL-1 receptor family: ancient signaling machinery in mammals, insects, and plants. *J Leukoc Biol* 1998 Jun;63(6):650-7
- Otero GC, Merrill JE. Cytokine receptors on glial cells. *Glia.* 1994 Jun;11(2):117-28. Review.
- Pike CJ, Walencewicz-Wasserman AJ, Kosmoski J et al. Structure-activity analyses of beta-amyloid peptides: contributions of the beta 25-35 region to aggregation and neurotoxicity. *J Neurochem.* 1995 Jan;64(1):253-65.
- Pocock JM, Liddle AC. Microglial signalling cascades in neurodegenerative disease. *Prog Brain Res.* 2001;132:555-65. Review.
- Prinz M, Kann O, Draheim HJ, Schumann RR et al. Microglial activation by components of gram-positive and -negative bacteria: distinct and common routes to the induction of ion channels and cytokines. *J Neuropathol Exp Neurol.* 1999 Oct;58(10):1078-89.
- Prinz M, Hanisch UK. Murine microglial cells produce and respond to interleukin-1 $\beta$ . *J Neurochem* 1999 May;72(5):2215-8
- Prinz M, Hausler KG, Kettenmann H, Hanisch U. beta-adrenergic receptor stimulation selectively inhibits IL-12p40 release in microglia. *Brain Res* 2001 Apr 27;899(1-2):264-70
- Pyo H, Jou I, Jung S, Hong S, Joe EH. Mitogen-activated protein kinases activated by lipopolysaccharide and beta-amyloid in cultured rat microglia. *Neuroreport.* 1998 Mar 30;9(5):871-4.
- Qiu WQ, Borth W, Ye Z, Haass C, Teplow DB, Selkoe DJ. Degradation of amyloid beta-protein by a serine protease-alpha2-macroglobulin complex. *J Biol Chem.* 1996 Apr 5;271(14):8443-51.
- Qiu Z, Strickland DK, Hyman BT, Rebeck GW. Alpha2-macroglobulin enhances the clearance of endogenous soluble beta-amyloid peptide via low-density lipoprotein receptor-related protein in cortical neurons. *J Neurochem.* 1999 Oct;73(4):1393-8.

- Raivich G, Bohatschek M, Kloss CU, Werner A, Jones LL, Kreutzberg GW. Neuroglial activation repertoire in the injured brain: graded response, molecular mechanisms and cues to physiological function. *Brain Res Brain Res Rev* 1999 Jul;30(1):77-105
- Ravaglia G, Paola F, Maioli F et al. Interleukin-1beta and interleukin-6 gene polymorphisms as risk factors for AD: a prospective study. *Exp Gerontol.* 2006 Jan;41(1):85-92. Review.
- Ren L, Lubrich B, Biber K, Gebicke-Haerter PJ. Differential expression of inflammatory mediators in rat microglia cultured from different brain regions. *Brain Res Mol Brain Res.* 1999 Mar 5;65(2):198-205.
- Rich JB, Rasmussen DX, Folstein MF et al. Nonsteroidal anti-inflammatory drugs in Alzheimer's disease. *Neurology.* 1995 Jan;45(1):51-5. *Neurology.* 1997 Mar;48(3):626-32.
- Ringheim GE, Szczepanik AM, Petko W et al. Enhancement of beta-amyloid precursor protein transcription and expression by the soluble interleukin-6 receptor/interleukin-6 complex. *Brain Res Mol Brain Res.* 1998 Mar 30;55(1):35-44.
- Rogers J, Kirby LC, Hempelman Sr et al. Clinical trial of indomethacin in Alzheimer's disease. *Neurology.* 1993 Aug;43(8):1609-11.
- Rogers J, Webster S, Lue LF et al. Inflammation and Alzheimer's disease pathogenesis. *eurobiol Aging.* 1996 Sep-Oct;17(5):681-6. Review
- Roher AE, Chaney MO, Kuo YM et al. Morphology and toxicity of Abeta-(1-42) dimer derived from neuritic and vascular amyloid deposits of Alzheimer's disease. *J Biol Chem.* 1996 Aug 23;271(34):20631-5.
- Roher AE, Baudry J, Chaney MO, Kuo YM, Stine WB, Emmerling MR. Oligomerizaiton and fibril asssembly of the mayloid-beta protein. *Biochim Biophys Acta* 2000 Jul 26;1502(1):31-43
- Rothwell NJ, Hopkins SJ. Cytokines and the nervous system II: Actions and mechanisms of action. *Trends Neurosci* 1995 Mar;18(3):130-6
- Rothwell NJ. Cytokines and acute neurodegeneration. *Mol Psychiatry.* 1997 Mar ;2(2):120-1.
- Rothwell NJ, Luheshi GN. Interleukin 1 in the brain: biology, pathology and therapeutic target. *Trends Neurosci* 2000 Dec;23(12):618-25
- Schmeichel DE, Goldgaber D, Burkhardt DS, Gilbert JR, Gajdusek DC, Roses AD. Cellular localization of messenger RNA encoding amyloid-beta-protein in normal tissue and in Alzheimer disease. *Alzheimer Dis Assoc Disord.* 1988;2(2):96-111.
- Schuringa JJ, Jonk LJ, Dokter WH, Vellenga E, Kruijer W. Interleukin-6-induced STAT3 transactivation and Ser727 phosphorylation involves Vav, Rac-1 and the kinase SEK-1/MKK-4 as signal transduction components. *Biochem J.* 2000 Apr 1;347 Pt 1:89-96.
- Seilheimer B, Bohrmann B, Bondolfi L et al. The toxicity of the Alzheimer's beta-amyloid peptide correlates with a distinct fiber morphology. *J Struct Biol.* 1997 Jun;119(1):59-71.
- Selkoe DJ. Normal and abnormal biology of the beta-amyloid precursor protein. *Annu Rev Neurosci* 1994;17:489-517

- Selkoe DJ. Translating cell biology into therapeutic advances in Alzheimer's disease. *Nature*. 1999 Jun 24; 399 (6738 Suppl):A23-31.
- Selkoe DJ. Presenilin, Notch, and the genesis and treatment of Alzheimer's disease. *Proc Natl Acad Sci U S A* 2001 Sep 25;98(20):11039-41
- Seubert P, Vigo-Pelfrey C, Esch F et al. Isolation and quantification of soluble Alzheimer's beta-peptide from biological fluids. *Nature*. 1992 Sep 24;359(6393):325-7.
- Shakhov AN, Collart MA, Vassalli P, Nedospasov SA, Jongeneel CV. Kappa B-type enhancers are involved in lipopolysaccharide-mediated transcriptional activation of the tumor necrosis factor alpha gene in primary macrophages. *J Exp Med*. 1990 Jan 1;171(1):35-47.
- Sheng JG, Mrak RE, Rovnaghi CR, Kozlowska E, Van Eldik LJ, Griffin WS. Human brain S100 beta and S100 beta mRNA \_expression increases with age: pathogenic implications for Alzheimer's disease. *Neurobiology Aging* 1996;17:359-63.
- Sheng JG, Jones RA, Zhou XQ, McGinness JM, Van Eldik LJ, Mrak RE, Griffin WS. Interleukin-1 promotion of MAPK-p38 overexpression in experimental animals and in Alzheimer's disease: potential significance for tau protein phosphorylation. *Neurochem Int*. 2001 Nov-Dec;39(5-6):341-8.
- Shimizu H, Mitomo K, Watanabe T, Okamoto S, Yamamoto K. Involvement of a NF-kappa B-like transcription factor in the activation of the interleukin-6 gene by inflammatory lymphokines. *Mol Cell Biol*. 1990 Feb;10(2):561-8.
- Shoji M, Golde TE, Ghiso J et al. Production of the Alzheimer amyloid beta protein by normal proteolytic processing. *Science*. 1992 Oct 2;258(5079):126-9.
- Singhrao SK, Neal JW, Morgan BP, Gasque P. Increased complement biosynthesis by microglia and complement activation on neurons in Huntington's disease. *Exp Neurol* 1999 Oct;159(2):362-76
- Sivaprakasam K. Towards a unifying hypothesis of Alzheimer's disease: cholinergic system linked to plaques, tangles and neuroinflammation. *Curr Med Chem*. 2006;13(18):2179-88. Review.
- Smith, P.K., Krohn, R.I., Hermanson, G.T et al. Measurement of protein using bicinchoninic acid. *Anal. Biochem*. 1985.150, 76-85.
- Smits HA, Rijsmus A, van Loon JH, Wat JW, Verhoef J, Boven LA, Nottet HS. Amyloid-beta-induced chemokine production in primary human macrophages and astrocytes. *J Neuroimmunol*. 2002 Jun;127(1-2):160-8.
- Song X, Shapiro S, Goldman DL, Casadevall A, Scharff M, Lee SC. Fcgamma receptor I- and III-mediated macrophage inflammatory protein 1alpha induction in primary human and murine microglia. *Infect Immun*. 2002 Sep;70(9):5177-84.
- Sparacio SM, Zhang Y, Vilcek J, Benveniste EN. Cytokine regulation of interleukin-6 gene expression in astrocytes involves activation of an NF-kappa B-like nuclear protein. *J Neuroimmunol* 1992 Aug;39(3):231-42

- Stohwasser R, Giesebricht J, Kraft R et al. Biochemical analysis of proteasomes from mouse microglia: induction of immunoproteasomes by interferon-gamma and lipopolysaccharide. *Glia*. 2000 Feb;15(4):355-65.
- Stoll G, Jander S, Schroeter M. Cytokines in CNS disorders: neurotoxicity versus neuroprotection. *J Neural Transm Suppl* 2000;59:81-9.
- Strauss S, Bauer J, Ganter U, Jonas U, Berger M, Volk B. Detection of interleukin-6 and alpha 2-macroglobulin immunoreactivity in cortex and hippocampus of Alzheimer's disease patients. *Lab Invest*. 1992 Feb;66(2):223-30.
- Streit WJ, Graeber MB, Kreutzberg GW. Functional plasticity of microglia: a review. *Glia*. 1988;1(5):301-7. Review.
- Streit W. J. Microglial cells. In *Neuroglia* (Kettenmann H, Ransom BR eds.) Oxford: Oxford University Press. 1995; pp:85-96.
- Streit WJ, Conde JR, Harrison JK. Chemokines and Alzheimer's disease. *Neurobiol Aging*. 2001 Nov-Dec;22(6):909-13. Review.
- Streit WJ, Walter SA, Pennell NA. Reactive microgliosis. *Prog Neurobiol* 1999 Apr;57(6):563-81
- Sun YX, Minthon L, Wallmark A et al. Inflammatory markers in matched plasma and cerebrospinal fluid from patients with Alzheimer's disease. *Dement Geriatr Cogn Disord*. 2003;16(3):136-44.
- Suzuki N, Cheung TT, Cai XD et al. An increased percentage of long amyloid beta protein secreted by familial amyloid beta protein precursor (beta APP717) mutants. *Science*. 1994 May 27;264(5163):1336-40.
- Szczepanik AM, Rampe D, Ringheim GE. Amyloid-beta peptide fragments p3 and p4 induce pro-inflammatory cytokine and chemokine production in vitro and in vivo. *J Neurochem*. 2001 Apr;77(1):304-17.
- Takata K, Kitamura Y, Umeki M et al. Possible involvement of small oligomers of amyloid-beta peptides in 15-deoxy-delta 12,14 prostaglandin J2-sensitive microglial activation. *J Pharmacol Sci*. 2003 Apr;91(4):330-3.
- Takatsuna H., et al. Identification of TIFA as an adapter protein that links TRAF6 to IRAK-1 in IL-1 receptor signaling. *J Biol Chem*. 2003.
- Tan J, Town T, Mori T et al. CD45 opposes beta-amyloid peptide-induced microglial activation via inhibition of p44/42 mitogen-activated protein kinase. *J Neurosci*. 2000 Oct 15;20(20):7587-94.
- Tarkowski E, Blennow K, Wallin A, Tarkowski A. Intracerebral production of tumor necrosis factor-alpha, a local neuroprotective agent, in Alzheimer disease and vascular dementia. *J Clin Immunol*. 1999 Jul;19(4):223-30.
- Tehranian R, Hasanvan H, Iverfeldt K, Post C, Schultzberg M. Early induction of interleukin-6 mRNA in the hippocampus and cortex of APPsw transgenic mice Tg2576. *Neurosci Lett* 2001 Mar 23;301(1):54-8

- Terry RD, Wisniewski HM. Structural and chemical changes of the aged human brain. *Psychopharmacol Bull.* 1975 Oct;11(4):46.
- Thal DR, Schober R, Birkenmeier G. The subunits of alpha2-macroglobulin receptor/low density lipoprotein receptor-related protein, native and transformed alpha2-macroglobulin and interleukin 6 in Alzheimer's disease. *Brain Res* 1997 Nov 28;777(1-2):223-7
- Thomson AW. *The Cytokine Handbook*. Academic Press Limited. 1994.
- Town T, Nikolic V, Tan J. The microglial "activation" continuum: from innate to adaptive responses. *J Neuroinflammation*. 2005 Oct 31;2:24.
- Uihara T, Akiyama H, Kondo H, Ikeda K. Activated microglia cells are colocalized with perivascular deposits of amyloid-beta protein in Alzheimer's disease brain. *Stroke*. 1997 Oct; 28 (10): 1948-50.
- Vandenabeele P, Fiers W. Is amyloidogenesis during Alzheimer's disease due to an IL-1/IL-6-mediated 'acute phase response' in the brain? *Immunol Today* 1991 Jul;12(7):217-9
- Van Gool D, De Strooper B, Van Leuven F, Triau E, Dom R. alpha 2-Macroglobulin expression in neuritic-type plaques in patients with Alzheimer's disease. *Neurobiol Aging*. 1993 May-Jun;14(3):233-7.
- Van Muiswinkel FL, Veerhuis R, Eikelenboom P. Amyloid beta protein primes cultured rat microglial cells for an enhanced phorbol 12-myristate 13-acetate-induced respiratory burst activity. *J Neurochem* 1996 Jun;66(6):2468-76
- Van Rossum D and Hanisch UK. Microglia. *Metab Brain Dis*. 2004 Dec;19(3-4):393-411.
- Veerhuis R, Van Breemen MJ, Hoozemans JM, et al. Amyloid beta plaque-associated proteins C1q and SAP enhance the Abeta1-42 peptide-induced cytokine secretion by adult human microglia in vitro. *Acta Neuropathol (Berl)*. 2003 Feb;105(2):135-44.
- in t' Veld BA, Ruitenberg A, Hofman A et al. Reduced incidence of AD with NSAID but not H2 receptor antagonists: the Cache County Study. *Neurology*. 2002 Sep 24;59(6):880-6.
- Vigo-Pelfrey C, Lee D, Keim P, Lieberburg I, Schenk DB. Characterization of beta-amyloid peptide from human cerebrospinal fluid. *J Neurochem*. 1993 Nov;61(5):1965-8
- Webb DJ, Wen J, Lysiak JJ, Umans L, Van Leuven F, Gonias SL. Murine alpha-macroglobulins demonstrate divergent activities as neutralizers of transforming growth factor-beta and as inducers of nitric oxide synthesis. A possible mechanism for the endotoxin insensitivity of the alpha2-macroglobulin gene knock-out mouse. *J Biol Chem*. 1996 Oct 4;271(40):24982-8.
- Weldon DT, Rogers SD, Ghilardi JR et al. Fibrillar beta-amyloid induces microglial phagocytosis, expression of inducible nitric oxide synthase, and loss of a select population of neurons in the rat CNS in vivo. *J Neurosci*. 1998 Mar 15;18(6):2161-73.
- Wesche H, Korherr C, Kracht M et al. The interleukin-1 receptor accessory protein (IL-1RAcP) is essential for IL-1-induced activation of interleukin-1 receptor-associated kinase (IRAK) and stress-activated protein kinases (SAP kinases). *J Biol Chem* 1997 Mar 21;272(12):7727-31

- Wollenberg GK, LaMarre J, Rosendal S, Gonias SL, Hayes MA. Binding of tumor necrosis factor alpha to activated forms of human plasma alpha 2 macroglobulin. *Am J Pathol*. 1991 Feb;138(2):265-72.
- Wood JA, Wood PL, Ryan R et al. Cytokine indices in Alzheimer's temporal cortex: no changes in mature IL-1 beta or IL-1RA but increases in the associated acute phase proteins IL-6, alpha 2-macroglobulin and C-reactive protein. *Brain Res*. 1993 Dec 3;629(2):245-52.
- Xie QW, Whisnant R, Nathan C. Promoter of the mouse gene encoding calcium-independent nitric oxide synthase confers inducibility by interferon gamma and bacterial lipopolysaccharide. *J Exp Med*. 1993 Jun 1;177(6):1779-84.
- Xia MQ, Qin SX, Wu LJ, Mackay CR, Hyman BT. Immunohistochemical study of the beta-chemokine receptors CCR3 and CCR5 and their ligands in normal and Alzheimer's disease brains. *Am J Pathol*. 1998 Jul;153(1):31-7.
- Xia M, Hyman BT. GROalpha/KC, a chemokine receptor CXCR2 ligand, can be a potent trigger for neuronal ERK1/2 and PI-3 kinase pathways and for tau hyperphosphorylation-a role in Alzheimer's disease? *J Neuroimmunol*. 2002 Jan;122(1-2):55-64.
- Yamada T, Fujieda S, Yanagi S et al. IL-1 induced chemokine production through the association of Syk with TNF receptor-associated factor-6 in nasal fibroblast lines. *J Immunol*. 2001 Jul 1;167(1):283-8.
- Yamaguchi H, Nakazato Y, Hirai S, Shoji M, Harigaya Y. Electron micrograph of diffuse plaques. Initial stage of senile plaque formation in the Alzheimer brain. *Am J Pathol*. 1989 Oct;135(4):593-7.
- Yates SL, Burgess LH, Kocsis-Angle J et al. Amyloid beta and amylin fibrils induce increases in proinflammatory cytokine and chemokine production by THP-1 cells and murine microglia. *J Neurochem*. 2000 Mar;74(3):1017-25.
- Zassler B, Weis C, Humpel C. Tumor necrosis factor-alpha triggers cell death of sensitized potassium chloride-stimulated cholinergic neurons. *Brain Res Mol Brain Res*. 2003 May 12;113(1-2):78-85.
- Zhang P, Hogan EL, Bhat NR. Activation of JNK/SAPK in primary glial cultures: II. Differential activation of kinase isoforms corresponds to their differential expression. *Neurochem Res* 1998 Feb;23(2):219-25
- Zhao M, Cribbs DH, Anderson AJ, et al. The induction of the TNFalpha death domain signaling pathway in Alzheimer's disease brain. *Neurochem Res*. 2003 Feb;28(2):307-18.
- Zielasek J, Archelos JJ, Toyka KV, Hartung HP. Expression of intercellular adhesion molecule-1 on rat microglial cells. *Neurosci Lett*. 1993 Apr 30;153(2):136-9.