

7 Literaturverzeichnis

- [1] Dobkin, B. H. (2005): Clinical practice. Rehabilitation after stroke, *N.Engl.J.Med.* 352 [16], Seite 1677-1684. URL: PM:15843670
- [2] Brott, T. und Bogousslavsky, J. (2000): Treatment of acute ischemic stroke, *N.Engl.J.Med.* 343 [10], Seite 710-722. URL: PM:10974136
- [3] Statistisches Bundesamt Deutschland (2001): Deutsche Todesursachenstatistik 2001
- [4] Kolominsky-Rabas, P. L.; Sarti, C.; Heuschmann, P. U.; Graf, C.; Siemonsen, S.; Neundoerfer, B.; Katalinic, A.; Lang, E.; Gassmann, K. G. und von Stockert, T. R. (1998): A prospective community-based study of stroke in Germany--the Erlangen Stroke Project (ESPro): incidence and case fatality at 1, 3, and 12 months, *Stroke* 29 [12], Seite 2501-2506. URL: PM:9836758
- [5] Eisenblätter, D.; Heinemann, L. und Classen, E. (1995): Community-based stroke incidence trends from the 1970s through the 1980s in East Germany, *Stroke* 26 [6], Seite 919-923. URL: PM:7762038
- [6] Berger, K.; Schulte, H.; Stogbauer, F. und Assmann, G. (1998): Incidence and risk factors for stroke in an occupational cohort: the PROCAM Study. Prospective Cardiovascular Muenster Study, *Stroke* 29 [8], Seite 1562-1566. URL: PM:9707193
- [7] Taylor, T. N.; Davis, P. H.; Torner, J. C.; Holmes, J.; Meyer, J. W. und Jacobson, M. F. (1996): Lifetime cost of stroke in the United States, *Stroke* 27 [9], Seite 1459-1466. URL: PM:8784113
- [8] Weih, M.; Müller-Nordhorn, J.; Amberger, N.; Masuhr, F.; Lurtzing, F.; Dreier, J. P. und Hetzel, A. (2004): [Risk factors in ischemic stroke. Review of evidence in primary prevention], *Nervenarzt* 75 [4], Seite 324-335. URL: PM:15088088
- [9] Collins, R.; Peto, R.; MacMahon, S.; Hebert, P.; Fiebich, N. H.; Eberlein, K. A.; Godwin, J.; Qizilbash, N.; Taylor, J. O. und Hennekens, C. H. (1990): Blood pressure, stroke, and coronary heart disease. Part 2, Short-term reductions in blood pressure: overview of randomised drug trials in their epidemiological context, *Lancet* 335 [8693], Seite 827-838. URL: PM:1969567
- [10] Reimer, K. A.; Murry, C. E.; Yamasawa, I.; Hill, M. L. und Jennings, R. B. (1986): Four brief periods of myocardial ischemia cause no cumulative ATP loss or necrosis, *Am.J.Physiol* 251 [6 Pt 2], Seite H1306-H1315. URL: PM:3789183
- [11] Murry, C. E.; Jennings, R. B. und Reimer, K. A. (1986): Preconditioning with ischemia: a delay of lethal cell injury in ischemic myocardium, *Circulation* 74 [5], Seite 1124-1136. URL: PM:3769170
- [12] Chen, J.; Graham, S. H.; Zhu, R. L. und Simon, R. P. (1996): Stress proteins and tolerance to focal cerebral ischemia, *J.Cereb.Blood Flow Metab* 16 [4], Seite 566-577. URL: PM:8964795
- [13] Yellon, D. M.; Alkhulaifi, A. M. und Pugsley, W. B. (1993): Preconditioning the human myocardium, *Lancet* 342 [8866], Seite 276-277. URL: PM:8101304
- [14] Jenkins, D. P.; Pugsley, W. B.; Alkhulaifi, A. M.; Kemp, M.; Hooper, J. und Yellon, D. M. (1997): Ischaemic preconditioning reduces troponin T release in patients undergoing coronary artery bypass surgery, *Heart* 77 [4], Seite 314-318. URL: PM:9155608

- [15] Airaksinen, K. E. und Huikuri, H. V. (1997): Antiarrhythmic effect of repeated coronary occlusion during balloon angioplasty, *J.Am.Coll.Cardiol.* 29 [5], Seite 1035-1038. URL: PM:9120156
- [16] Eltchaninoff, H.; Cribier, A.; Tron, C.; Derumeaux, G.; Koning, R.; Hecketsweiler, B. und Letac, B. (1997): Adaptation to myocardial ischemia during coronary angioplasty demonstrated by clinical, electrocardiographic, echocardiographic, and metabolic parameters, *Am.Heart J.* 133 [4], Seite 490-496. URL: PM:9124180
- [17] Ottani, F.; Galvani, M.; Ferrini, D.; Sorbello, F.; Limonetti, P.; Pantoli, D. und Rusticali, F. (1995): Prodromal angina limits infarct size. A role for ischemic preconditioning, *Circulation* 91 [2], Seite 291-297. URL: PM:7805230
- [18] Kloner, R. A.; Shook, T.; Przyklenk, K.; Davis, V. G.; Junio, L.; Matthews, R. V.; Burstein, S.; Gibson, M.; Poole, W. K.; Cannon, C. P. und . (1995): Previous angina alters in-hospital outcome in TIMI 4. A clinical correlate to preconditioning?, *Circulation* 91 [1], Seite 37-45. URL: PM:7805217
- [19] Andreotti, F.; Pasceri, V.; Hackett, D. R.; Davies, G. J.; Haider, A. W. und Maseri, A. (1996): Preinfarction angina as a predictor of more rapid coronary thrombolysis in patients with acute myocardial infarction, *N.Engl.J.Med.* 334 [1], Seite 7-12. URL: PM:7494588
- [20] Dahl, N. A. und Balfour, W. M. (1964): Prolonged anoxic survival due to anoxia pre-exposure: Brain ATP, lactate, and pyruvate, *Am.J.Physiol* 207, Seite 452-456. URL: PM:14205366
- [21] Du, Z. Y.; Hicks, M.; Winlaw, D.; Spratt, P. und Macdonald, P. (1996): Ischemic preconditioning enhances donor lung preservation in the rat, *J.Heart Lung Transplant.* 15 [12], Seite 1258-1267. URL: PM:8981211
- [22] Kume, M.; Yamamoto, Y.; Saad, S.; Gomi, T.; Kimoto, S.; Shimabukuro, T.; Yagi, T.; Nakagami, M.; Takada, Y.; Morimoto, T. und Yamaoka, Y. (1996): Ischemic preconditioning of the liver in rats: implications of heat shock protein induction to increase tolerance of ischemia-reperfusion injury, *J.Lab Clin.Med.* 128 [3], Seite 251-258. URL: PM:8783632
- [23] Peralta, C.; Closa, D.; Hotter, G.; Gelpi, E.; Prats, N. und Rosello-Catafau, J. (1996): Liver ischemic preconditioning is mediated by the inhibitory action of nitric oxide on endothelin, *Biochem.Biophys.Res.Commun.* 229 [1], Seite 264-270. URL: PM:8954116
- [24] Hotter, G.; Closa, D.; Prados, M.; Fernandez-Cruz, L.; Prats, N.; Gelpi, E. und Rosello-Catafau, J. (1996): Intestinal preconditioning is mediated by a transient increase in nitric oxide, *Biochem.Biophys.Res.Commun.* 222 [1], Seite 27-32. URL: PM:8630069
- [25] Islam, C. F.; Mathie, R. T.; Dinneen, M. D.; Kiely, E. A.; Peters, A. M. und Grace, P. A. (1997): Ischaemia-reperfusion injury in the rat kidney: the effect of preconditioning, *Br.J.Urol.* 79 [6], Seite 842-847. URL: PM:9202547
- [26] Pudupakkam, S.; Harris, K. A.; Jamieson, W. G.; DeRose, G.; Scott, J. A.; Carson, M. W.; Schlag, M. G.; Kvietys, P. R. und Potter, R. F. (1998): Ischemic tolerance in skeletal muscle: role of nitric oxide, *Am.J.Physiol* 275 [1 Pt 2], Seite H94-H99. URL: PM:9688900
- [27] Perez-Pinzon, M. A.; Xu, G. P.; Dietrich, W. D.; Rosenthal, M. und Sick, T. J. (1997): Rapid preconditioning protects rats against ischemic neuronal damage after 3 but not 7 days of reperfusion following global cerebral ischemia, *J.Cereb.Blood Flow*

Metab 17 [2], Seite 175-182. URL: PM:9040497

[28] Kitagawa, K.; Matsumoto, M.; Tagaya, M.; Hata, R.; Ueda, H.; Niinobe, M.; Handa, N.; Fukunaga, R.; Kimura, K.; Mikoshiba, K. und . (1990): 'Ischemic tolerance' phenomenon found in the brain, *Brain Res.* 528 [1], Seite 21-24. URL: PM:2245337

[29] Kasischke, K.; Huber, R.; Li, H.; Timmler, M. und Riepe, M. W. (1999): Primary hypoxic tolerance and chemical preconditioning during estrus cycle in mice, *Stroke* 30 [6], Seite 1256-1262. URL: PM:10356109

[30] Cohen, M. V.; Liu, G. S. und Downey, J. M. (1991): Preconditioning causes improved wall motion as well as smaller infarcts after transient coronary occlusion in rabbits, *Circulation* 84 [1], Seite 341-349. URL: PM:2060104

[31] Schott, R. J.; Rohmann, S.; Braun, E. R. und Schaper, W. (1990): Ischemic preconditioning reduces infarct size in swine myocardium, *Circ.Res.* 66 [4], Seite 1133-1142. URL: PM:2317890

[32] Kato, H.; Liu, Y.; Araki, T. und Kogure, K. (1992): MK-801, but not anisomycin, inhibits the induction of tolerance to ischemia in the gerbil hippocampus, *Neurosci.Lett.* 139 [1], Seite 118-121. URL: PM:1407677

[33] Kirino, T.; Tsujita, Y. und Tamura, A. (1991): Induced tolerance to ischemia in gerbil hippocampal neurons, *J.Cereb.Blood Flow Metab* 11 [2], Seite 299-307. URL: PM:1997501

[34] Liu, Y.; Kato, H.; Nakata, N. und Kogure, K. (1992): Protection of rat hippocampus against ischemic neuronal damage by pretreatment with sublethal ischemia, *Brain Res.* 586 [1], Seite 121-124. URL: PM:1380876

[35] Gidday, J. M.; Fitzgibbons, J. C.; Shah, A. R. und Park, T. S. (1994): Neuroprotection from ischemic brain injury by hypoxic preconditioning in the neonatal rat, *Neurosci.Lett.* 168 [1-2], Seite 221-224. URL: PM:8028780

[36] Kawai, K.; Nakagomi, T.; Kirino, T.; Tamura, A. und Kawai, N. (1998): Preconditioning in vivo ischemia inhibits anoxic long-term potentiation and functionally protects CA1 neurons in the gerbil, *J.Cereb.Blood Flow Metab* 18 [3], Seite 288-296. URL: PM:9498845

[37] Barone, F. C.; White, R. F.; Spera, P. A.; Ellison, J.; Currie, R. W.; Wang, X. und Feuerstein, G. Z. (1998): Ischemic preconditioning and brain tolerance: temporal histological and functional outcomes, protein synthesis requirement, and interleukin-1 receptor antagonist and early gene expression, *Stroke* 29 [9], Seite 1937-1950. URL: PM:9731622

[38] Matsushima, K.; Hogan, M. J. und Hakim, A. M. (1996): Cortical spreading depression protects against subsequent focal cerebral ischemia in rats, *J.Cereb.Blood Flow Metab* 16 [2], Seite 221-226. URL: PM:8594053

[39] Schurr, A.; Reid, K. H.; Tseng, M. T.; West, C. und Rigor, B. M. (1986): Adaptation of adult brain tissue to anoxia and hypoxia in vitro, *Brain Res.* 374 [2], Seite 244-248. URL: PM:3719335

[40] Weih, M.; Kallenberg, K.; Bergk, A.; Dirnagl, U.; Harms, L.; Wernecke, K. D. und Einhaupl, K. M. (1999): Attenuated stroke severity after prodromal TIA: a role for ischemic tolerance in the brain?, *Stroke* 30 [9], Seite 1851-1854. URL: PM:10471435

[41] Moncayo, J.; de Freitas, G. R.; Bogouslavsky, J.; Altieri, M. und van, Melle G. (2000): Do transient ischemic attacks have a neuroprotective effect?, *Neurology* 54 [11], Seite 2089-2094. URL: PM:10851368

- [42] Wegener, S.; Gottschalk, B.; Jovanovic, V.; Knab, R.; Fiebach, J. B.; Schellinger, P. D.; Kucinski, T.; Jungehulsing, G. J.; Brunecker, P.; Muller, B.; Banasik, A.; Amberger, N.; Wernecke, K. D.; Siebler, M.; Rother, J.; Villringer, A. und Weih, M. (2004): Transient ischemic attacks before ischemic stroke: preconditioning the human brain? A multicenter magnetic resonance imaging study, *Stroke* 35 [3], Seite 616-621. URL: PM:14963288
- [43] Bruer, U.; Weih, M. K.; Isaev, N. K.; Meisel, A.; Ruscher, K.; Bergk, A.; Trendelenburg, G.; Wiegand, F.; Victorov, I. V. und Dirnagl, U. (1997): Induction of tolerance in rat cortical neurons: hypoxic preconditioning, *FEBS Lett.* 414 [1], Seite 117-121. URL: PM:9305743
- [44] Grabb, M. C. und Choi, D. W. (1999): Ischemic tolerance in murine cortical cell culture: critical role for NMDA receptors, *J.Neurosci.* 19 [5], Seite 1657-1662. URL: PM:10024352
- [45] Riepe, M. W.; Esclaire, F.; Kasischke, K.; Schreiber, S.; Nakase, H.; Kempfski, O.; Ludolph, A. C.; Dirnagl, U. und Hugon, J. (1997): Increased hypoxic tolerance by chemical inhibition of oxidative phosphorylation: "chemical preconditioning", *J.Cereb.Blood Flow Metab* 17 [3], Seite 257-264. URL: PM:9119898
- [46] Sugino, T.; Nozaki, K.; Takagi, Y. und Hashimoto, N. (1999): 3-Nitropropionic acid induces ischemic tolerance in gerbil hippocampus in vivo, *Neurosci.Lett.* 259 [1], Seite 9-12. URL: PM:10027543
- [47] Wiegand, F.; Liao, W.; Busch, C.; Castell, S.; Knapp, F.; Lindauer, U.; Megow, D.; Meisel, A.; Redetzky, A.; Ruscher, K.; Trendelenburg, G.; Victorov, I.; Riepe, M.; Diener, H. C. und Dirnagl, U. (1999): Respiratory chain inhibition induces tolerance to focal cerebral ischemia, *J.Cereb.Blood Flow Metab* 19 [11], Seite 1229-1237. URL: PM:10566969
- [48] Aketa, S.; Nakase, H.; Kamada, Y.; Hiramatsu, K. und Sakaki, T. (2000): Chemical preconditioning with 3-nitropropionic acid in gerbil hippocampal slices: therapeutic window and the participation of adenosine receptor, *Exp.Neurol.* 166 [2], Seite 385-391. URL: PM:11085903
- [49] Shake, J. G.; Peck, E. A.; Marban, E.; Gott, V. L.; Johnston, M. V.; Troncoso, J. C.; Redmond, J. M. und Baumgartner, W. A. (2001): Pharmacologically induced preconditioning with diazoxide: a novel approach to brain protection, *Ann.Thorac.Surg.* 72 [6], Seite 1849-1854. URL: PM:11789759
- [50] Nishio, S.; Yunoki, M.; Chen, Z. F.; Anzivino, M. J. und Lee, K. S. (2000): Ischemic tolerance in the rat neocortex following hypothermic preconditioning, *J.Neurosurg.* 93 [5], Seite 845-851. URL: PM:11059667
- [51] Zimmermann, C.; Ginis, I.; Furuya, K.; Klimanis, D.; Ruetzler, C.; Spatz, M. und Hallenbeck, J. M. (2001): Lipopolysaccharide-induced ischemic tolerance is associated with increased levels of ceramide in brain and in plasma, *Brain Res.* 895 [1-2], Seite 59-65. URL: PM:11259760
- [52] Lobner, D. und Choi, D. W. (1996): Preincubation with protein synthesis inhibitors protects cortical neurons against oxygen-glucose deprivation-induced death, *Neuroscience* 72 [2], Seite 335-341. URL: PM:8737404
- [53] Meloni, B. P.; Majda, B. T. und Knuckey, N. W. (2002): Evaluation of preconditioning treatments to protect near-pure cortical neuronal cultures from in vitro ischemia induced acute and delayed neuronal death, *Brain Res.* 928 [1-2], Seite 69-75. URL: PM:11844473

- [54] Kawahara, N.; Ruetzler, C. A. und Klatzo, I. (1995): Protective effect of spreading depression against neuronal damage following cardiac arrest cerebral ischaemia, *Neurol.Res.* 17 [1], Seite 9-16. URL: PM:7746347
- [55] Kobayashi, S.; Harris, V. A. und Welsh, F. A. (1995): Spreading depression induces tolerance of cortical neurons to ischemia in rat brain, *J.Cereb.Blood Flow Metab* 15 [5], Seite 721-727. URL: PM:7673367
- [56] Taga, K.; Patel, P. M.; Drummond, J. C.; Cole, D. J. und Kelly, P. J. (1997): Transient neuronal depolarization induces tolerance to subsequent forebrain ischemia in rats, *Anesthesiology* 87 [4], Seite 918-925. URL: PM:9357895
- [57] Prass, K.; Ruscher, K.; Karsch, M.; Isaev, N.; Megow, D.; Priller, J.; Scharff, A.; Dirnagl, U. und Meisel, A. (2002): Desferrioxamine induces delayed tolerance against cerebral ischemia in vivo and in vitro, *J.Cereb.Blood Flow Metab* 22 [5], Seite 520-525. URL: PM:11973424
- [58] Ohtsuki, T.; Matsumoto, M.; Kuwabara, K.; Kitagawa, K.; Suzuki, K.; Taniguchi, N. und Kamada, T. (1992): Influence of oxidative stress on induced tolerance to ischemia in gerbil hippocampal neurons, *Brain Res.* 599 [2], Seite 246-252. URL: PM:1337859
- [59] Kirino, T. (2002): Ischemic tolerance, *J.Cereb.Blood Flow Metab* 22 [11], Seite 1283-1296. URL: PM:12439285
- [60] Atochin, D. N.; Clark, J.; Demchenko, I. T.; Moskowitz, M. A. und Huang, P. L. (2003): Rapid cerebral ischemic preconditioning in mice deficient in endothelial and neuronal nitric oxide synthases, *Stroke* 34 [5], Seite 1299-1303. URL: PM:12677017
- [61] Kitagawa, K.; Matsumoto, M.; Kuwabara, K.; Tagaya, M.; Ohtsuki, T.; Hata, R.; Ueda, H.; Handa, N.; Kimura, K. und Kamada, T. (1991): 'Ischemic tolerance' phenomenon detected in various brain regions, *Brain Res.* 561 [2], Seite 203-211. URL: PM:1802339
- [62] Ohno, M. und Watanabe, S. (1996): Ischemic tolerance to memory impairment associated with hippocampal neuronal damage after transient cerebral ischemia in rats, *Brain Res.Bull.* 40 [3], Seite 229-236. URL: PM:8736585
- [63] Shamloo, M. und Wieloch, T. (1999): Changes in protein tyrosine phosphorylation in the rat brain after cerebral ischemia in a model of ischemic tolerance, *J.Cereb.Blood Flow Metab* 19 [2], Seite 173-183. URL: PM:10027773
- [64] Choi, D. W.; Maulucci-Gedde, M. und Kriegstein, A. R. (1987): Glutamate neurotoxicity in cortical cell culture, *J.Neurosci.* 7 [2], Seite 357-368. URL: PM:2880937
- [65] Frandsen, A. und Schousboe, A. (1990): Development of excitatory amino acid induced cytotoxicity in cultured neurons, *Int.J.Dev.Neurosci.* 8 [2], Seite 209-216. URL: PM:1970220
- [66] Barbe, M. F.; Tytell, M.; Gower, D. J. und Welch, W. J. (1988): Hyperthermia protects against light damage in the rat retina, *Science* 241 [4874], Seite 1817-1820. URL: PM:3175623
- [67] Chopp, M.; Chen, H.; Ho, K. L.; Dereski, M. O.; Brown, E.; Hetzel, F. W. und Welch, K. M. (1989): Transient hyperthermia protects against subsequent forebrain ischemic cell damage in the rat, *Neurology* 39 [10], Seite 1396-1398. URL: PM:2797463
- [68] Kitagawa, K.; Matsumoto, M.; Tagaya, M.; Kuwabara, K.; Hata, R.; Handa, N.;

- Fukunaga, R.; Kimura, K. und Kamada, T. (1991): Hyperthermia-induced neuronal protection against ischemic injury in gerbils, *J.Cereb.Blood Flow Metab* 11 [3], Seite 449-452. URL: PM:2016352
- [69] Nishio, S.; Chen, Z. F.; Yunoki, M.; Toyoda, T.; Anzivino, M. und Lee, K. S. (1999): Hypothermia-induced ischemic tolerance, *Ann.N.Y.Acad.Sci.* 890, Seite 26-41. URL: PM:10668411
- [70] Matsushima, K.; Schmidt-Kastner, R.; Hogan, M. J. und Hakim, A. M. (1998): Cortical spreading depression activates trophic factor expression in neurons and astrocytes and protects against subsequent focal brain ischemia, *Brain Res.* 807 [1-2], Seite 47-60. URL: PM:9756993
- [71] Lowenstein, D. H.; Simon, R. P. und Sharp, F. R. (1990): The pattern of 72-kDa heat shock protein-like immunoreactivity in the rat brain following flurothyl-induced status epilepticus, *Brain Res.* 531 [1-2], Seite 173-182. URL: PM:2289119
- [72] Ying, H. S.; Weishaupt, J. H.; Grabb, M.; Canzoniero, L. M.; Sensi, S. L.; Sheline, C. T.; Monyer, H. und Choi, D. W. (1997): Sublethal oxygen-glucose deprivation alters hippocampal neuronal AMPA receptor expression and vulnerability to kainate-induced death, *J.Neurosci.* 17 [24], Seite 9536-9544. URL: PM:9391008
- [73] Nawashiro, H.; Tasaki, K.; Ruetzler, C. A. und Hallenbeck, J. M. (1997): TNF-alpha pretreatment induces protective effects against focal cerebral ischemia in mice, *J.Cereb.Blood Flow Metab* 17 [5], Seite 483-490. URL: PM:9183285
- [74] Weih, M.; Bergk, A.; Isaev, N. K.; Ruscher, K.; Megow, D.; Riepe, M.; Meisel, A.; Victorov, I. V. und Dirnagl, U. (1999): Induction of ischemic tolerance in rat cortical neurons by 3-nitropropionic acid: chemical preconditioning, *Neurosci.Lett.* 272 [3], Seite 207-210. URL: PM:10505617
- [75] Tremblay, R.; Chakravarthy, B.; Hewitt, K.; Tauskela, J.; Morley, P.; Atkinson, T. und Durkin, J. P. (2000): Transient NMDA receptor inactivation provides long-term protection to cultured cortical neurons from a variety of death signals, *J.Neurosci.* 20 [19], Seite 7183-7192. URL: PM:11007874
- [76] Liu, J.; Ginis, I.; Spatz, M. und Hallenbeck, J. M. (2000): Hypoxic preconditioning protects cultured neurons against hypoxic stress via TNF-alpha and ceramide, *Am.J.Physiol Cell Physiol* 278 [1], Seite C144-C153. URL: PM:10644522
- [77] Coles, C. J.; Edmondson, D. E. und Singer, T. P. (1979): Inactivation of succinate dehydrogenase by 3-nitropropionate, *J.Biol.Chem.* 254 [12], Seite 5161-5167. URL: PM:447637
- [78] Fink, S. L.; Ho, D. Y. und Sapolsky, R. M. (1996): Energy and glutamate dependency of 3-Nitropropionic acid neurotoxicity in culture, *Exp.Neurol.* 138 [2], Seite 298-304. URL: PM:8620928
- [79] Pang, Z. und Geddes, J. W. (1997): Mechanisms of cell death induced by the mitochondrial toxin 3-nitropropionic acid: acute excitotoxic necrosis and delayed apoptosis, *J.Neurosci.* 17 [9], Seite 3064-3073. URL: PM:9096141
- [80] Morimoto, T.; Globus, M. Y.; Busto, R.; Martinez, E. und Ginsberg, M. D. (1996): Simultaneous measurement of salicylate hydroxylation and glutamate release in the penumbral cortex following transient middle cerebral artery occlusion in rats, *J.Cereb.Blood Flow Metab* 16 [1], Seite 92-99. URL: PM:8530561
- [81] Wardman, P. und Candeias, L. P. (1996): Fenton chemistry: an introduction, *Radiat.Res.* 145 [5], Seite 523-531. URL: PM:8619017

- [82] Miller, D. M. und Aust, S. D. (1989): Studies of ascorbate-dependent, iron-catalyzed lipid peroxidation, *Arch.Biochem.Biophys.* 271 [1], Seite 113-119. URL: PM:2712569
- [83] Blanchard, B.; Servy, C. und Ducrocq, C. (2001): Chemical evaluation of compounds as nitric oxide or peroxyxynitrite donors using the reactions with serotonin, *Free Radic.Res.* 34 [2], Seite 189-191. URL: PM:11264894
- [84] Stone, J. R. und Marletta, M. A. (1996): Spectral and kinetic studies on the activation of soluble guanylate cyclase by nitric oxide, *Biochemistry* 35 [4], Seite 1093-1099. URL: PM:8573563
- [85] Laurent, M.; Lepoivre, M. und Tenu, J. P. (1996): Kinetic modelling of the nitric oxide gradient generated in vitro by adherent cells expressing inducible nitric oxide synthase, *Biochem.J.* 314 (Pt 1), Seite 109-113. URL: PM:8660270
- [86] Henry, Y.; Lepoivre, M.; Drapier, J. C.; Ducrocq, C.; Boucher, J. L. und Guissani, A. (1993): EPR characterization of molecular targets for NO in mammalian cells and organelles, *FASEB J.* 7 [12], Seite 1124-1134. URL: PM:8397130
- [87] Dawson, V. L.; Dawson, T. M.; London, E. D.; Bredt, D. S. und Snyder, S. H. (1991): Nitric oxide mediates glutamate neurotoxicity in primary cortical cultures, *Proc.Natl.Acad.Sci.U.S.A* 88 [14], Seite 6368-6371. URL: PM:1648740
- [88] Nickel, B. (1987): The antinociceptive activity of flupirtine: a structurally new analgesic, *Postgrad.Med.J.* 63 Suppl 3, Seite 19-28. URL: PM:2833734
- [89] Block, F.; Pergande, G. und Schwarz, M. (1994): Flupirtine protects against ischaemic retinal dysfunction in rats, *Neuroreport* 5 [18], Seite 2630-2632. URL: PM:7696619
- [90] Osborne, N. N.; Schwarz, M. und Pergande, G. (1996): Protection of rabbit retina from ischemic injury by flupirtine, *Invest Ophthalmol.Vis.Sci.* 37 [2], Seite 274-280. URL: PM:8603831
- [91] Rupalla, K.; Cao, W. und Krieglstein, J. (1995): Flupirtine protects neurons against excitotoxic or ischemic damage and inhibits the increase in cytosolic Ca²⁺ concentration, *Eur.J.Pharmacol.* 294 [2-3], Seite 469-473. URL: PM:8750707
- [92] Muller, W. E.; Pergande, G.; Ushijima, H.; Schleger, C.; Kelve, M. und Perovic, S. (1996): Neurotoxicity in rat cortical cells caused by N-methyl-D-aspartate (NMDA) and gp120 of HIV-1: induction and pharmacological intervention, *Prog.Mol.Subcell.Biol.* 16, Seite 44-57. URL: PM:8822791
- [93] Perovic, S.; Pergande, G.; Ushijima, H.; Kelve, M.; Forrest, J. und Muller, W. E. (1995): Flupirtine partially prevents neuronal injury induced by prion protein fragment and lead acetate, *Neurodegeneration.* 4 [4], Seite 369-374. URL: PM:8846229
- [94] Osborne, N. N.; Cazevieuille, C.; Wood, J. P.; Nash, M. S.; Pergande, G.; Block, F.; Kosinski, C. und Schwarz, M. (1998): Flupirtine, a nonopioid centrally acting analgesic, acts as an NMDA antagonist, *Gen.Pharmacol.* 30 [3], Seite 255-263. URL: PM:9510072
- [95] Kornhuber, J.; Bleich, S.; Wiltfang, J.; Maler, M. und Parsons, C. G. (1999): Flupirtine shows functional NMDA receptor antagonism by enhancing Mg²⁺ block via activation of voltage independent potassium channels. Rapid communication, *J.Neural Transm.* 106 [9-10], Seite 857-867. URL: PM:10599868
- [96] Akaike, A.; Tamura, Y.; Sato, Y. und Yokota, T. (1993): Protective effects of a vitamin B12 analog, methylcobalamin, against glutamate cytotoxicity in cultured

cortical neurons, *Eur.J.Pharmacol.* 241 [1], Seite 1-6. URL: PM:7901032

[97] Koh, J. Y. und Choi, D. W. (1987): Quantitative determination of glutamate mediated cortical neuronal injury in cell culture by lactate dehydrogenase efflux assay, *J.Neurosci.Methods* 20 [1], Seite 83-90. URL: PM:2884353

[98] Mosmann, T. (1983): Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays, *J.Immunol.Methods* 65 [1-2], Seite 55-63. URL: PM:6606682

[99] Alley, M. C.; Scudiero, D. A.; Monks, A.; Hursey, M. L.; Czerwinski, M. J.; Fine, D. L.; Abbott, B. J.; Mayo, J. G.; Shoemaker, R. H. und Boyd, M. R. (1988): Feasibility of drug screening with panels of human tumor cell lines using a microculture tetrazolium assay, *Cancer Res.* 48 [3], Seite 589-601. URL: PM:3335022

[100] Scudiero, D. A.; Shoemaker, R. H.; Paull, K. D.; Monks, A.; Tierney, S.; Nofziger, T. H.; Currens, M. J.; Seniff, D. und Boyd, M. R. (1988): Evaluation of a soluble tetrazolium/formazan assay for cell growth and drug sensitivity in culture using human and other tumor cell lines, *Cancer Res.* 48 [17], Seite 4827-4833. URL: PM:3409223

[101] Morley, D. und Keefer, L. K. (1993): Nitric oxide/nucleophile complexes: a unique class of nitric oxide-based vasodilators, *J.Cardiovasc.Pharmacol.* 22 Suppl 7, Seite S3-S9. URL: PM:7504766

[102] Maragos, C. M.; Morley, D.; Wink, D. A.; Dunams, T. M.; Saavedra, J. E.; Hoffman, A.; Bove, A. A.; Isaac, L.; Hrabie, J. A. und Keefer, L. K. (1991): Complexes of .NO with nucleophiles as agents for the controlled biological release of nitric oxide. Vasorelaxant effects, *J.Med.Chem.* 34 [11], Seite 3242-3247. URL: PM:1956043

[103] Ferrero, R.; Rodriguez-Pascual, F.; Miras-Portugal, M. T. und Torres, M. (1999): Comparative effects of several nitric oxide donors on intracellular cyclic GMP levels in bovine chromaffin cells: correlation with nitric oxide production, *Br.J.Pharmacol.* 127 [3], Seite 779-787. URL: PM:10401570

[104] Kuroiwa, T.; Yamada, I.; Endo, S.; Hakamata, Y. und Ito, U. (2000): 3-Nitropropionic acid preconditioning ameliorates delayed neurological deterioration and infarction after transient focal cerebral ischemia in gerbils, *Neurosci.Lett.* 283 [2], Seite 145-148. URL: PM:10739896

[105] Maruoka, N.; Murata, T.; Omata, N.; Fujibayashi, Y.; Yonekura, Y. und Wada, Y. (2002): Hypoxic tolerance induction in rat brain slices following 3-nitropropionic acid pretreatment as revealed by dynamic changes in glucose metabolism, *Neurosci.Lett.* 319 [2], Seite 83-86. URL: PM:11825676

[106] Porter, D. J. und Bright, H. J. (1980): 3-Carbanionic substrate analogues bind very tightly to fumarase and aspartase, *J.Biol.Chem.* 255 [10], Seite 4772-4780. URL: PM:7372610

[107] Schloss, J. V. und Cleland, W. W. (1982): Inhibition of isocitrate lyase by 3-nitropropionate, a reaction-intermediate analogue, *Biochemistry* 21 [18], Seite 4420-4427. URL: PM:7126549

[108] Osman, M. Y. (1982): Effect of B-nitropropionic acid on rat brain acetylcholinesterase, *Biochem.Pharmacol.* 31 [24], Seite 4067-4068. URL: PM:7159482

[109] Zeevalk, G. D.; rr-Yellin, E. und Nicklas, W. J. (1995): NMDA receptor involvement in toxicity to dopamine neurons in vitro caused by the succinate

- dehydrogenase inhibitor 3-nitropropionic acid, *J.Neurochem.* 64 [1], Seite 455-458. URL: PM:7798946
- [110] Nakagawa, I.; Nakase, H.; Aketa, S.; Kamada, Y.; Yamashita, M. und Sakaki, T. (2002): ATP-dependent potassium channel mediates neuroprotection by chemical preconditioning with 3-nitropropionic acid in gerbil hippocampus, *Neurosci.Lett.* 320 [1-2], Seite 33-36. URL: PM:11849757
- [111] von Arnim, C. A.; Timmler, M.; Ludolph, A. C. und Riepe, M. W. (2000): Adenosine receptor up-regulation: initiated upon preconditioning but not upheld, *Neuroreport* 11 [6], Seite 1223-1226. URL: PM:10817596
- [112] Riepe, M. W.; Kasischke, K.; Gericke, C. A.; Lowe, A. und Hellweg, R. (1996): Increase of hypoxic tolerance in rat hippocampal slices following 3-nitropropionic acid is not mediated by endogenous nerve growth factor, *Neurosci.Lett.* 211 [1], Seite 9-12. URL: PM:8809835
- [113] Brambrink, A. M.; Noga, H.; Astheimer, A.; Heimann, A. und Kempfski, O. (2004): Pharmacological preconditioning in global cerebral ischemia, *Acta Neurochir.Suppl* 89, Seite 63-66. URL: PM:15335102
- [114] Brambrink, A. M.; Schneider, A.; Noga, H.; Astheimer, A.; Gotz, B.; Korner, I.; Heimann, A.; Welschof, M. und Kempfski, O. (2000): Tolerance-Inducing dose of 3-nitropropionic acid modulates bcl-2 and bax balance in the rat brain: a potential mechanism of chemical preconditioning, *J.Cereb.Blood Flow Metab* 20 [10], Seite 1425-1436. URL: PM:11043905
- [115] Furuichi, T.; Liu, W.; Shi, H.; Miyake, M. und Liu, K. J. (2005): Generation of hydrogen peroxide during brief oxygen-glucose deprivation induces preconditioning neuronal protection in primary cultured neurons, *J.Neurosci.Res.* URL: PM:15668910
- [116] Andoh, T.; Chock, P. B. und Chiueh, C. C. (2002): Preconditioning-mediated neuroprotection: role of nitric oxide, cGMP, and new protein expression, *Ann.N.Y.Acad.Sci.* 962, Seite 1-7. URL: PM:12076958
- [117] Gidday, J. M.; Shah, A. R.; Maceren, R. G.; Wang, Q.; Pelligrino, D. A.; Holtzman, D. M. und Park, T. S. (1999): Nitric oxide mediates cerebral ischemic tolerance in a neonatal rat model of hypoxic preconditioning, *J.Cereb.Blood Flow Metab* 19 [3], Seite 331-340. URL: PM:10078885
- [118] Atochin, D. N.; Clark, J.; Demchenko, I. T.; Moskowitz, M. A. und Huang, P. L. (2003): Rapid cerebral ischemic preconditioning in mice deficient in endothelial and neuronal nitric oxide synthases, *Stroke* 34 [5], Seite 1299-1303. URL: PM:12677017
- [119] Hashiguchi, A.; Yano, S.; Morioka, M.; Hamada, J.; Ushio, Y.; Takeuchi, Y. und Fukunaga, K. (2004): Up-regulation of endothelial nitric oxide synthase via phosphatidylinositol 3-kinase pathway contributes to ischemic tolerance in the CA1 subfield of gerbil hippocampus, *J.Cereb.Blood Flow Metab* 24 [3], Seite 271-279. URL: PM:15091107
- [120] Laude, K.; Favre, J.; Thuillez, C. und Richard, V. (2003): NO produced by endothelial NO synthase is a mediator of delayed preconditioning-induced endothelial protection, *Am.J.Physiol Heart Circ.Physiol* 284 [6], Seite H2053-H2060. URL: PM:12521944
- [121] von Arnim, C. A.; Timmler, M.; Ludolph, A. C. und Riepe, M. W. (2001): Chemical preconditioning in mice is not mediated by upregulation of nitric oxide synthase isoforms, *Neurosci.Lett.* 299 [1-2], Seite 130-134. URL: PM:11166955
- [122] Centeno, J. M.; Orti, M.; Salom, J. B.; Sick, T. J. und Perez-Pinzon, M. A.

- (1999): Nitric oxide is involved in anoxic preconditioning neuroprotection in rat hippocampal slices, *Brain Res.* 836 [1-2], Seite 62-69. URL: PM:10415405
- [123] Kawahara, K.; Yanoma, J.; Tanaka, M.; Nakajima, T. und Kosugi, T. (2004): Nitric oxide produced during ischemia is toxic but crucial to preconditioning-induced ischemic tolerance of neurons in culture, *Neurochem.Res.* 29 [4], Seite 797-804. URL: PM:15098944
- [124] Block, F.; Pergande, G. und Schwarz, M. (1997): Flupirtine reduces functional deficits and neuronal damage after global ischemia in rats, *Brain Res.* 754 [1-2], Seite 279-284. URL: PM:9134985
- [125] Muller, W. E.; Romero, F. J.; Perovic, S.; Pergande, G. und Pialoglou, P. (1997): Protection of flupirtine on beta-amyloid-induced apoptosis in neuronal cells in vitro: prevention of amyloid-induced glutathione depletion, *J.Neurochem.* 68 [6], Seite 2371-2377. URL: PM:9166730
- [126] Perovic, S.; Schroder, H. C.; Pergande, G.; Ushijima, H. und Muller, W. E. (1997): Effect of flupirtine on Bcl-2 and glutathione level in neuronal cells treated in vitro with the prion protein fragment (PrP106-126), *Exp.Neurol.* 147 [2], Seite 518-524. URL: PM:9344576
- [127] Perovic, S.; Pialoglou, P.; Schroder, H. C.; Pergande, G. und Muller, W. E. (1996): Flupirtine increases the levels of glutathione and Bc1-2 in hNT (human Ntera/D1) neurons: mode of action of the drug-mediated anti-apoptotic effect, *Eur.J.Pharmacol.* 317 [1], Seite 157-164. URL: PM:8982732
- [128] Gassen, M.; Pergande, G. und Youdim, M. B. (1998): Antioxidant properties of the triaminopyridine, flupirtine, *Biochem.Pharmacol.* 56 [10], Seite 1323-1329. URL: PM:9825731
- [129] Osborne, N. N.; Nash, M. S. und Wood, J. P. (1998): Melatonin counteracts ischemia-induced apoptosis in human retinal pigment epithelial cells, *Invest Ophthalmol.Vis.Sci.* 39 [12], Seite 2374-2383. URL: PM:9804146
- [130] Jakob, R. und Krieglstein, J. (1997): Influence of flupirtine on a G-protein coupled inwardly rectifying potassium current in hippocampal neurones, *Br.J.Pharmacol.* 122 [7], Seite 1333-1338. URL: PM:9421279
- [131] Perovic, S.; Bohm, M.; Meesters, E.; Meinhardt, A.; Pergande, G. und Muller, W. E. (1998): Pharmacological intervention in age-associated brain disorders by Flupirtine: Alzheimer's and prion diseases, *Mech.Ageing Dev.* 101 [1-2], Seite 1-19. URL: PM:9593309
- [132] Perovic, S.; Schleger, C.; Pergande, G.; Iskric, S.; Ushijima, H.; Rytik, P. und Muller, W. E. (1994): The triaminopyridine flupirtine prevents cell death in rat cortical cells induced by N-methyl-D-aspartate and gp120 of HIV-1, *Eur.J.Pharmacol.* 288 [1], Seite 27-33. URL: PM:7705465
- [133] Gozlan, H. und Ben-Ari, Y. (1995): NMDA receptor redox sites: are they targets for selective neuronal protection?, *Trends Pharmacol.Sci.* 16 [11], Seite 368-374. URL: PM:8578605
- [134] Nash, M. S.; Wood, J. P.; Melena, J. und Osborne, N. N. (2000): Flupirtine ameliorates ischaemic-like death of rat retinal ganglion cells by preventing calcium influx, *Brain Res.* 856 [1-2], Seite 236-239. URL: PM:10677631
- [135] Zimmer, G.; Balakirev, M.; Zwicker, K.; Hofmann, M.; Woodcock, B. G. und Pergande, G. (1998): Effect of the triaminopyridine flupirtine on calcium uptake, membrane potential and ATP synthesis in rat heart mitochondria, *Br.J.Pharmacol.*

123 [6], Seite 1154-1158. URL: PM:9559899

[136] McBain, C. J. und Mayer, M. L. (1994): N-methyl-D-aspartic acid receptor structure and function, *Physiol Rev.* 74 [3], Seite 723-760. URL: PM:8036251

[137] Novelli, A.; Reilly, J. A.; Lysko, P. G. und Henneberry, R. C. (1988): Glutamate becomes neurotoxic via the N-methyl-D-aspartate receptor when intracellular energy levels are reduced, *Brain Res.* 451 [1-2], Seite 205-212. URL: PM:2472189

[138] Wieloch, T. (1985): Hypoglycemia-induced neuronal damage prevented by an N-methyl-D-aspartate antagonist, *Science* 230 [4726], Seite 681-683. URL: PM:2996146

[139] Mayer, M. L.; Westbrook, G. L. und Guthrie, P. B. (1984): Voltage-dependent block by Mg²⁺ of NMDA responses in spinal cord neurones, *Nature* 309 [5965], Seite 261-263. URL: PM:6325946

[140] Johnson, J. W. und Ascher, P. (1987): Glycine potentiates the NMDA response in cultured mouse brain neurons, *Nature* 325 [6104], Seite 529-531. URL: PM:2433595

[141] Mori, H. und Mishina, M. (1995): Structure and function of the NMDA receptor channel, *Neuropharmacology* 34 [10], Seite 1219-1237. URL: PM:8570021

[142] Stenzel-Poore, M. P.; Stevens, S. L. und Simon, R. P. (2004): Genomics of preconditioning, *Stroke* 35 [11 Suppl 1], Seite 2683-2686. URL: PM:15459430

[143] Stenzel-Poore, M. P.; Stevens, S. L.; Xiong, Z.; Lessov, N. S.; Harrington, C. A.; Mori, M.; Meller, R.; Rosenzweig, H. L.; Tobar, E.; Shaw, T. E.; Chu, X. und Simon, R. P. (2003): Effect of ischaemic preconditioning on genomic response to cerebral ischaemia: similarity to neuroprotective strategies in hibernation and hypoxia-tolerant states, *Lancet* 362 [9389], Seite 1028-1037. URL: PM:14522533

[144] Trendelenburg, G.; Prass, K.; Priller, J.; Kapinya, K.; Polley, A.; Muselmann, C.; Ruscher, K.; Kannbley, U.; Schmitt, A. O.; Castell, S.; Wiegand, F.; Meisel, A.; Rosenthal, A. und Dirnagl, U. (2002): Serial analysis of gene expression identifies metallothionein-II as major neuroprotective gene in mouse focal cerebral ischemia, *J.Neurosci.* 22 [14], Seite 5879-5888. URL: PM:12122050