

# Literaturverzeichnis

- Adler, V.; Zeiler, B.; Kryukov, V.; Kascsak, R.; Rubenstein, R. und Grossman, A. 2003. Small, highly structured RNAs participate in the conversion of human recombinant PrP<sup>Sen</sup> to PrP<sup>Res</sup> in vitro. *J. Mol. Biol.*, **332**: 47–57.
- Alvarez-Martinez, M. T.; Torrent, J.; Lange, R.; Verdier, J.-M.; Balny, C. und Li-autard, J.-P. 2003. Optimized overproduction, purification, characterization and high-pressure sensitivity of the prion protein in the native (PrP<sup>C</sup>-like) or amyloid (PrP<sup>Sc</sup>-like) conformation. *Biochim. Biophys. Acta*, **1645**: 228–240.
- Anfinsen, C. B. 1973. Principles that govern the folding of protein chains. *Science*, **181**: 223–230.
- Armen, R. S.; Alonso, D. O. und Daggett, V. 2004a. Anatomy of an amyloidogenic intermediate: conversion of  $\beta$ -sheet to  $\alpha$ -sheet structure in transthyretin at acidic pH. *Structure (Camb.)*, **12**: 1847–1863.
- Armen, R. S.; DeMarco, M. L.; Alonso, D. O. und Daggett, V. 2004b. Pauling and Corey's  $\alpha$ -pleated sheet structure may define the prefibrillar amyloidogenic intermediate in amyloid disease. *Proc. Natl. Acad. Sci. USA*, **101**: 11622–11627.
- Arrondo, J. L.; Muga, A.; Castresana, J. und Goni, F. M. 1993. Quantitative studies of the structure of proteins in solution by Fourier-transform infrared spectroscopy. *Prog. Biophys. Mol. Biol.*, **59**: 23–56.
- Atkinson, P. H. 2004. Glycosylation of prion strains in transmissible spongiform encephalopathies. *Aust. Vet. J.*, **82**: 292–299.
- Bandekar, J. 1992. Amide modes and protein conformation. *Biochim. Biophys. Acta*, **1120**: 123–143.
- Baron, G. S.; Wehrly, K.; Dorward, D. W.; Chesebro, B. und Caughey, B. 2002. Conversion of raft associated prion protein to the protease-resistant state requires insertion of PrP-res (PrP<sup>Sc</sup>) into contiguous membranes. *EMBO J.*, **21**: 1031–1040.
- Barth, A. 2000. The infrared absorption of amino acid side chains. *Prog. Biophys. Mol. Biol.*, **74**: 141–173.

- Barth, A. und Zscherp, C. 2002. What vibrations tell us about proteins. *Q. Rev. Biophys.*, **35**: 369–430.
- Barton II, F. E.; Himmelsbach, D. S.; Duckworth, J. H. und Smith, M. J. 1992. Two-dimensional vibration spectroscopy: correlation of mid- and near-infrared regions. *Appl. Spectrosc.*, **46**: 420–429.
- Baskakov, I. V.; Aagaard, C.; Mehlhorn, I.; Wille, H.; Groth, D.; Baldwin, M. A.; Prusiner, S. B. und Cohen, F. E. 2000. Self-assembly of recombinant prion protein of 106 residues. *Biochemistry*, **39**: 2792–2804.
- Baskakov, I. V. und Bocharova, O. V. 2005. In vitro conversion of mammalian prion protein into amyloid fibrils displays unusual features. *Biochemistry*, **44**: 2339–2348.
- Baskakov, I. V.; Legname, G.; Baldwin, M. A.; Prusiner, S. B. und Cohen, F. E. 2002. Pathway complexity of prion protein assembly into amyloid. *J. Biol. Chem.*, **277**: 21140–21148.
- Baskakov, I. V.; Legname, G.; Gryczynski, Z. und Prusiner, S. B. 2004. The peculiar nature of unfolding of the human prion protein. *Protein Sci.*, **13**: 586–595.
- Baskakov, I. V.; Legname, G.; Prusiner, S. B. und Cohen, F. E. 2001. Folding of prion protein to its native  $\alpha$ -helical conformation is under kinetic control. *J. Biol. Chem.*, **276**: 19687–19690.
- Basler, K.; Oesch, B.; Scott, M.; Westaway, D.; Walchli, M.; Groth, D. F.; McKinley, M. P.; Prusiner, S. B. und Weissmann, C. 1986. Scrapie and cellular PrP isoforms are encoded by the same chromosomal gene. *Cell*, **46**: 417–428.
- Bocharova, O. V.; Breydo, L.; Parfenov, A. S.; Salnikov, V. V. und Baskakov, I. V. 2005. In vitro conversion of full-length mammalian prion protein produces amyloid form with physical properties of PrP<sup>Sc</sup>. *J. Mol. Biol.*, **346**: 645–659.
- Bruce, M. E. 2003. TSE strain variation. *Br. Med. Bull.*, **66**: 99–108.
- Bucciantini, M.; Giannoni, E.; Chiti, F.; Baroni, F.; Formigli, L.; Zurdo, J.; Taddei, N.; Ramponi, G.; Dobson, C. M. und Stefani, M. 2002. Inherent toxicity of aggregates implies a common mechanism for protein misfolding diseases. *Nature*, **416**: 507–511.
- Callaham, M. A.; Xiong, L.-W. und Caughey, B. 2001. Reversibility of scrapie-associated prion protein aggregation. *J. Biol. Chem.*, **276**: 28022–28028.
- Caughey, B. und Kocisko, D. A. 2003. Prion diseases: a nucleic-acid accomplice? *Nature*, **425**: 673–674.

- Caughey, B. und Lansbury, J.; P. T. 2003. Protofibrils, pores, fibrils, and neurodegeneration: separating the responsible protein aggregates from the innocent bystanders. *Annu. Rev. Neurosci.*
- Caughey, B.; Raymond, G. J. und Bessen, R. A. 1998. Strain-dependent differences in  $\beta$ -sheet conformations of abnormal prion protein. *J. Biol. Chem.*, **273**: 32230–32235.
- Chan, H. S. und Dill, K. A. 1998. Protein folding in the landscape perspective: chevron plots and non-Arrhenius kinetics. *Proteins*, **30**: 2–33.
- Chiesa, R. und Harris, D. A. 2001. Prion diseases: What is the neurotoxic molecule? *Neurobiol. Dis.*, **8**: 743–763.
- Cleary, J. P.; Walsh, D. M.; Hofmeister, J. J.; Shankar, G. M.; Kuskowski, M. A.; Selkoe, D. J. und Ashe, K. H. 2005. Natural oligomers of the amyloid- $\beta$  protein specifically disrupt cognitive function. *Nat. Neurosci.*, **8**: 79–84.
- Cordeiro, Y.; Machado, F.; Juliano, L.; Juliano, M. A.; Brentani, R. R.; Foguel, D. und Silva, J. L. 2001. DNA converts cellular prion protein into the  $\beta$ -sheet conformation and inhibits prion peptide aggregation. *J. Biol. Chem.*, **276**: 49400–49409.
- Couzin, J. 2004. Biomedicine. An end to the prion debate? Don't count on it. *Science*, **305**: 589.
- Critchley, P.; Kazlauskaitė, J.; Eason, R. und Pinheiro, T. J. 2004. Binding of prion proteins to lipid membranes. *Biochem. Biophys. Res. Commun.*, **313**: 559–567.
- Damaschun, G.; Damaschun, H.; Fabian, H.; Gast, K.; Krober, R.; Wieske, M. und Zirwer, D. 2000. Conversion of yeast phosphoglycerate kinase into amyloid-like structure. *Proteins*, **39**: 204–211.
- Deleault, N. R.; Lucassen, R. W. und Supattapone, S. 2003. RNA molecules stimulate prion protein conversion. *Nature*, **425**: 717–720.
- Demuro, A.; Mina, E.; Kaye, R.; Milton, S. C.; Parker, I. und Glabe, C. G. 2005. Calcium dysregulation and membrane disruption as a ubiquitous neurotoxic mechanism of soluble amyloid oligomers. *J. Biol. Chem.*
- Dobson, C. M. 2001. The structural basis of protein folding and its links with human disease. *Philos. Trans. R. Soc. Lond. B*, **356**: 133–145.
- Donne, D. G.; Viles, J. H.; Groth, D.; Mehlhorn, I.; James, T. L.; Cohen, F. E.; Prusiner, S. B.; Wright, P. E. und Dyson, H. J. 1997. Structure of the recombinant full-length hamster prion protein PrP(29–231): The N terminus is highly flexible. *Proc. Natl. Acad. Sci. USA*, **94**: 13452–13457.

- Eberl, H.; Tittmann, P. und Glockshuber, R. 2004. Characterization of recombinant, membrane-attached full-length prion protein. *J. Biol. Chem.*, **279**: 25058–25065.
- Edskes, H. K. und Wickner, R. B. 2004. Transmissible spongiform encephalopathies: prion proof in progress. *Nature*, **430**: 977–979.
- Fabian, H. und Schultz, C. P. Fourier transform infrared spectroscopy in peptide and protein analysis. In *Encyclopedia of Analytical Chemistry*, 5779–5803. John Wiley & Sons Ltd., Chichester, 2000.
- Fernández, A. und Berry, R. S. 2003. Proteins with H-bond packing defects are highly interactive with lipid bilayers: implications for amyloidogenesis. *Proc. Natl. Acad. Sci. USA*, **100**: 2391–2396.
- Gajdusek, D. C. 1977. Unconventional viruses and the origin and disappearance of kuru. *Science*, **197**: 943–960.
- Glockshuber, R. 2001. Folding dynamics and energetics of recombinant prion proteins. *Adv. Protein Chem.*, **57**: 83–105.
- Govaerts, C.; Wille, H.; Prusiner, S. B. und Cohen, F. E. 2004. Evidence for assembly of prions with left-handed  $\beta$ -helices into trimers. *Proc. Natl. Acad. Sci. USA*, **101**: 8342–8347.
- Hagge, S.-O. *Charakterisierung der Eigenschaften rekonstituierter planarer Membranen und deren Änderung durch porenbildende Proteine mit Hilfe von elektrischen und optischen Methoden*. Doktorarbeit, Christian-Albrechts-Universität zu Kiel, 2003.
- Hornemann, S. und Glockshuber, R. 1998. A scrapie-like unfolding intermediate of the prion protein domain PrP(121–231) induced by acidic pH. *Proc. Natl. Acad. Sci. USA*, **95**: 6010–6014.
- Hornemann, S.; Schorn, C. und Wüthrich, K. 2004. NMR structure of the bovine prion protein isolated from healthy calf brains. *EMBO Rep.*, **5**: 1159–1164.
- Hrsg.: Hörnlimann, B.; Riesner, D. und Kretzschmar, H. *Prionen und Prionkrankheiten*. Walter de Gruyter, Berlin und New York, 2001.
- Jackson, G. S.; Hill, A. F.; Joseph, C.; Hosszu, L.; Power, A.; Waltho, J. P.; Clarke, A. R. und Collinge, J. 1999a. Multiple folding pathways for heterologously expressed human prion protein. *Biochim. Biophys. Acta*, **1431**: 1–13.
- Jackson, G. S.; Hosszu, L. L. P.; Power, A.; Hill, A. F.; Kennedy, J.; Saibil, H.; Craven, C. J.; Waltho, J. P.; Clarke, A. R. und Collinge, J. 1999b. Reversible conversion of monomeric human prion protein between native and fibrillogenic conformations. *Science*, **283**: 1935–1937.

- James, T. L.; Liu, H.; Ulyanov, N. B.; Farr-Jones, S.; Zhang, H.; Donne, D. G.; Kaneko, K.; Groth, D.; Mehlhorn, I.; Prusiner, S. B. und Cohen, F. E. 1997. Solution structure of a 142-residue recombinant prion protein corresponding to the infectious fragment of the scrapie isoform. *Proc. Natl. Acad. Sci. USA*, **94**: 10086–10091.
- Jeffrey, M.; Goddbrand, I. A. und Goodsir, C. M. 1995. Pathology of the transmissible spongiform encephalopathies with special emphasis on ultrastructure. *Micron*, **26**: 277–298.
- Jones, C. E.; Abdelraheim, S. R.; Brown, D. R. und Viles, J. H. 2004. Preferential  $\text{Cu}^{2+}$  coordination by His96 and His111 induces  $\beta$ -sheet formation in the unstructured amyloidogenic region of the prion protein. *J. Biol. Chem.*, **279**: 32018–32027.
- Kasczak, R. J.; Rubenstein, R.; Merz, P. A.; Tonna-DeMasi, M.; Fersko, R.; Carp, R. I.; Wisniewski, H. M. und Diringer, H. 1987. Mouse polyclonal and monoclonal antibody to scrapie-associated fibril proteins. *J. Virol.*, **61**: 3688–3693.
- Kayed, R.; Head, E.; Thompson, J. L.; McIntire, T. M.; Milton, S. C.; Cotman, C. W. und Glabe, C. G. 2003. Common structure of soluble amyloid oligomers implies common mechanism of pathogenesis. *Science*, **300**: 486–489.
- Kazlauskaite, J.; Young, A.; Gardner, C. E.; Macpherson, J. V.; Venien-Bryan, C. und Pinheiro, T. J. 2005. An unusual soluble  $\beta$ -turn-rich conformation of prion is involved in fibril formation and toxic to neuronal cells. *Biochem. Biophys. Res. Commun.*, **328**: 292–305.
- Khurana, R. und Fink, A. L. 2000. Do parallel  $\beta$ -helix proteins have a unique Fourier transform infrared spectrum? *Biophys. J.*, **78**: 994–1000.
- Kimberlin, R. H. und Walker, C. A. 1977. Characteristics of a short incubation model of scrapie in the golden hamster. *J. Gen. Virol.*, **34**: 295–304.
- Knaus, K. J.; Morillas, M.; Swietnicki, W.; Malone, M.; Surewicz, W. K. und Yee, V. C. 2001. Crystal structure of the human prion protein reveals a mechanism for oligomerization. *Nat. Struct. Biol.*, **8**: 770–774.
- Kneipp, J.; Miller, L. M.; Spassov, S.; Sokolowski, F.; Lasch, P.; Beekes, M. und Naumann, D. 2004. Scrapie-infected cells, isolated prions, and recombinant prion protein: a comparative study. *Biopolymers*, **74**: 163–167.
- Koppel, D. E. 1972. Analysis of macromolecular polydispersity in intensity correlation spectroscopy: the method of cumulants. *J. Chem. Phys.*, **57**: 4814–4820.
- Koradi, R.; Billeter, M. und Wüthrich, K. 1996. MOLMOL: a program for display and analysis of macromolecular structures. *J. Mol. Graph.*, **14**: 51–55, 29–32.

- Kozin, S. A.; Bertho, G.; Mazur, A. K.; Rabesona, H.; Girault, J.-P.; Haertle, T.; Takahashi, M.; Debey, P. und Hoa, G. H. B. 2001. Sheep prion protein synthetic peptide spanning helix 1 and  $\beta$ -strand 2 (residues 142–166) shows  $\beta$ -hairpin structure in solution. *J. Biol. Chem.*, **276**: 46364–46370.
- Kretschmar, H. A.; Stowring, L. E.; Westaway, D.; Stubblebine, W. H.; Prusiner, S. B. und DeArmond, S. J. 1986. Molecular cloning of a human prion protein cDNA. *DNA*, **5**: 315–324.
- Krimm, S. und Bandekar, J. 1986. Vibrational spectroscopy and conformation of peptides, polypeptides, and proteins. *Adv. Protein Chem.*, **38**: 181–364.
- Kubelka, J. und Keiderling, T. A. 2001. Differentiation of  $\beta$ -sheet-forming structures: ab initio-based simulations of IR absorption and vibrational CD for model peptide and protein  $\beta$ -sheets. *J. Am. Chem. Soc.*, **123**: 12048–12058.
- Lansbury, J.; P. T. 1999. Evolution of amyloid: What normal protein folding may tell us about fibrillogenesis and disease. *Proc. Natl. Acad. Sci. USA*, **96**: 3342–3344.
- Lashuel, H. A.; Hartley, D.; Petre, B. M.; Walz, T. und Lansbury, J.; P. T. 2002. Neurodegenerative disease: amyloid pores from pathogenic mutations. *Nature*, **418**: 291.
- Leclerc, E.; Peretz, D.; Ball, H.; Sakurai, H.; Legname, G.; Serban, A.; Prusiner, S. B.; Burton, D. R. und Williamson, R. A. 2001. Immobilized prion protein undergoes spontaneous rearrangement to a conformation having features in common with the infectious form. *EMBO J.*, **20**: 1547–1554.
- Lee, S. und Eisenberg, D. 2003. Seeded conversion of recombinant prion protein to a disulfide-bonded oligomer by a reduction-oxidation process. *Nat. Struct. Biol.*, **10**: 725–730.
- Legname, G.; Baskakov, I. V.; Nguyen, H. O.; Riesner, D.; Cohen, F. E.; DeArmond, S. J. und Prusiner, S. B. 2004. Synthetic mammalian prions. *Science*, **305**: 673–676.
- Levinthal, C. 1968. Are there pathways for protein folding? *J. Chim. Phys.*, **65**: 44–45.
- Liu, H.; Farr-Jones, S.; Ulyanov, N. B.; Llinas, M.; Marqusee, S.; Groth, D.; Cohen, F. E.; Prusiner, S. B. und James, T. L. 1999. Solution structure of Syrian hamster prion protein rPrP(90–231). *Biochemistry*, **38**: 5362–5377.
- Lu, B.-Y.; Atanasov, I.; Zhou, Z. H. und Chang, J. Y. 2003. Reversible aggregation of mouse prion protein derivatives with PrP<sup>Sc</sup>-like structural properties. *J. Protein Chem.*, **22**: 115–126.

- Lu, B.-Y. und Chang, J.-Y. 2001. Isolation of isoforms of mouse prion protein with PrP<sup>Sc</sup>-like structural properties. *Biochemistry*, **40**: 13390–13396.
- Lu, B.-Y. und Chang, J.-Y. 2002. Isolation and characterization of a polymerized prion protein. *Biochem. J.*, **364**: 81–87.
- Ma, J.; Wollmann, R. und Lindquist, S. 2002. Neurotoxicity and neurodegeneration when PrP accumulates in the cytosol. *Science*, **298**: 1781–1785.
- Maiti, N. R. und Surewicz, W. K. 2001. The role of disulfide bridge in the folding and stability of the recombinant human prion protein. *J. Biol. Chem.*, **276**: 2427–2431.
- Makhatadze, G. I.; Clore, G. M. und Gronenborn, A. M. 1995. Solvent isotope effect and protein stability. *Nat. Struct. Biol.*, **2**: 852–855.
- Mallucci, G.; Dickinson, A.; Linehan, J.; Klohn, P. C.; Brandner, S. und Collinge, J. 2003. Depleting neuronal PrP in prion infection prevents disease and reverses spongiosis. *Science*, **302**: 871–874.
- Masuch, R. und Moss, D. A. 2003. Stopped flow apparatus for time-resolved Fourier transform infrared difference spectroscopy of biological macromolecules in <sup>1</sup>H<sub>2</sub>O. *Appl. Spectrosc.*, **57**: 1407–1418.
- Mausser, H. *Formale Kinetik*. Bertelsmann Universitätsverlag, Düsseldorf, 1974.
- McKinley, M. P.; Meyer, R. K.; Kenaga, L.; Rahbar, F.; Cotter, R.; Serban, A. und Prusiner, S. B. 1991. Scrapie prion rod formation in vitro requires both detergent extraction and limited proteolysis. *J. Virol.*, **65**: 1340–1351.
- Modler, A. J. *Studien zur Kinetik der Fehlfaltung und Aggregation von Proteinen*. Doktorarbeit, Humboldt-Universität zu Berlin, 2003.
- Modler, A. J.; Fabian, H.; Sokolowski, F.; Lutsch, G.; Gast, K. und Damaschun, G. 2004. Polymerization of proteins into amyloid protofibrils shares common critical oligomeric states but differs in the mechanisms of their formation. *Amyloid*, **11**: 215–231.
- Modler, A. J.; Gast, K.; Lutsch, G. und Damaschun, G. 2003. Assembly of amyloid protofibrils via critical oligomers—a novel pathway of amyloid formation. *J. Mol. Biol.*, **325**: 135–148.
- Montal, M. und Mueller, P. 1972. Formation of bimolecular membranes from lipid monolayers and a study of their electrical properties. *Proc. Natl. Acad. Sci. USA*, **69**: 3561–3566.
- Morillas, M.; Vanik, D. L. und Surewicz, W. K. 2001. On the mechanism of  $\alpha$ -helix to  $\beta$ -sheet transition in the recombinant prion protein. *Biochemistry*, **40**: 6982–6987.

- Moritz, R.; Reinstädler, D.; Fabian, H. und Naumann, D. 2002. Time-resolved FTIR difference spectroscopy as tool for investigating refolding reactions of ribonuclease T1 synchronized with trans  $\rightarrow$  cis prolyl isomerization. *Biopolymers*, **67**: 145–155.
- Nandi, P. K.; Leclerc, E.; Nicole, J. C. und Takahashi, M. 2002. DNA-induced partial unfolding of prion protein leads to its polymerisation to amyloid. *J. Mol. Biol.*, **322**: 153–161.
- Nishina, K.; Deleault, N. R.; Lucassen, R. W. und Supattapone, S. 2004. In vitro prion protein conversion in detergent-solubilized membranes. *Biochemistry*, **43**: 2613–2621.
- Noda, I. 1993. Generalized two-dimensional correlation method applicable to infrared, Raman, and other types of spectroscopy. *Appl. Spectrosc.*, **47**: 1329–1336.
- Oberg, K. A.; Ruysschaert, J. M. und Goormaghtigh, E. 2004. The optimization of protein secondary structure determination with infrared and circular dichroism spectra. *Eur. J. Biochem.*, **271**: 2937–2948.
- Pan, K.-M.; Baldwin, M. A.; Nguyen, J.; Gasset, M.; Serban, A.; Groth, D.; Mehlhorn, I.; Huang, Z.; Fletterick, R. J.; Cohen, F. E. und Prusiner, S. B. 1993. Conversion of  $\alpha$ -helices into  $\beta$ -sheets features in the formation of the scrapie prion proteins. *Proc. Natl. Acad. Sci. USA*, **90**: 10962–10966.
- Paramithiotis, E.; Pinard, M.; Lawton, T.; LaBoissiere, S.; Leathers, V. L.; Zou, W. Q.; Estey, L. A.; Lamontagne, J.; Lehto, M. T.; Kondejewski, L. H.; Francoeur, G. P.; Papadopoulos, M.; Haghighat, A.; Spatz, S. J.; Head, M.; Will, R.; Ironside, J.; O'Rourke, K.; Tonelli, Q.; Ledebur, H. C.; Chakrabartty, A. und Cashman, N. R. 2003. A prion protein epitope selective for the pathologically misfolded conformation. *Nat. Med.*, **9**: 893–899.
- Provencher, S. W. 1982. CONTIN: a general purpose constrained regularization program for inverting noisy linear algebraic an integral equations. *Comput. Phys. Commun.*, **27**: 229–242.
- Prusiner, S. B. 1982. Novel proteinaceous infectious particles cause scrapie. *Science*, **216**: 136–144.
- Prusiner, S. B. 1998. Prions. *Proc. Natl. Acad. Sci. USA*, **95**: 13363–13383.
- Reinstädler, D.; Fabian, H.; Backmann, J. und Naumann, D. 1996. Refolding of thermally and urea-denatured ribonuclease A monitored by time-resolved FTIR spectroscopy. *Biochemistry*, **35**: 15822–15830.
- Reinstädler, D.; Fabian, H. und Naumann, D. 1999. New structural insights into the refolding of ribonuclease T1 as seen by time-resolved Fourier-transform infrared spectroscopy. *Proteins*, **34**: 303–316.



- Requena, J. R.; Dimitrova, M. N.; Legname, G.; Teijeira, S.; Prusiner, S. B. und Levine, R. L. 2004. Oxidation of methionine residues in the prion protein by hydrogen peroxide. *Arch. Biochem. Biophys.*, **432**: 188–195.
- Rezaei, H.; Eghiaian, F.; Perez, J.; Doublet, B.; Choiset, Y.; Haertle, T. und Grosclaude, J. 2005. Sequential generation of two structurally distinct ovine prion protein soluble oligomers displaying different biochemical reactivities. *J. Mol. Biol.*, **347**: 665–679.
- Robakis, N. K.; Devine-Gage, E. A.; Jenkins, E. C.; Kascsak, R. J.; Brown, W. T.; Krawczun, M. S. und Silverman, W. P. 1986. Localization of a human gene homologous to the PrP gene on the p arm of chromosome 20 and detection of PrP-related antigens in normal human brain. *Biochem. Biophys. Res. Commun.*, **140**: 758–765.
- Rogers, M.; Serban, D.; Gyuris, T.; Scott, M.; Torchia, T. und Prusiner, S. B. 1991. Epitope mapping of the Syrian hamster prion protein utilizing chimeric and mutant genes in a vaccinia virus expression system. *J. Immunol.*, **147**: 3568–3574.
- Roucou, X. und Leblanc, A. C. 2005. Cellular prion protein neuroprotective function: implications in prion diseases. *J. Mol. Med.*, **83**: 3–11.
- Rudd, P. M.; Endo, T.; Colominas, C.; Groth, D.; Wheeler, S. F.; Harvey, D. J.; Wormald, M. R.; Serban, H.; Prusiner, S. B.; Kobata, A. und Dwek, R. A. 1999. Glycosylation differences between the normal and pathogenic prion protein isoforms. *Proc. Natl. Acad. Sci. USA*, **96**: 13044–13049.
- Rudd, P. M.; Wormald, M. R.; Wing, D. R.; Prusiner, S. B. und Dwek, R. A. 2001. Prion glycoprotein: structure, dynamics, and role for the sugars. *Biochemistry*, **40**: 3759–3766.
- Saborio, G. P.; Permanne, B. und Soto, C. 2001. Sensitive detection of pathological prion protein by cyclic amplification of protein misfolding. *Nature*, **411**: 810–813.
- Safar, J. G.; Geschwind, M. D.; Deering, C.; Didorenko, S.; Sattavat, M.; Sanchez, H.; Serban, A.; Vey, M.; Baron, H.; Giles, K.; Miller, B. L.; DeArmond, S. J. und Prusiner, S. B. 2005. Diagnosis of human prion disease. *Proc. Natl. Acad. Sci. USA*, **102**: 3501–3506.
- Sanghera, N. und Pinheiro, T. J. T. 2002. Binding of prion protein to lipid membranes and implications for prion conversion. *J. Mol. Biol.*, **315**: 1241–1256.
- Sarver, R. W. und Krueger, W. C. 1991. Protein secondary structure from Fourier transform infrared spectroscopy: a data base analysis. *Anal. Biochem.*, **194**: 89–100.

- Sattler, M.; Schleucher, J. und Griesinger, C. 1999. Heteronuclear multidimensional NMR experiments for the structure determination of proteins in solution employing pulsed field gradients. *Prog. NMR Spectrosc.*, **34**: 93–158.
- Savitzky, A. und Golay, M. J. E. 1964. Smoothing and differentiation of data by simplified least squares procedures. *Anal. Chem.*, **36**: 1627–1639.
- Schmieder, P.; Leidert, M.; Kelly, M. und Oschkinat, H. 1998. Multiplicity-selective coherence transfer steps for the design of amino acid-selective experiments—a triple-resonance experiment selective for Asn and Gln. *J. Magn. Reson.*, **131**: 199–202.
- Selkoe, D. J. 2002. Alzheimer’s disease is a synaptic failure. *Science*, **298**: 789–791.
- Shaked, G. M.; Meiner, Z.; Avraham, I.; Taraboulos, A. und Gabizon, R. 2001. Reconstitution of prion infectivity from solubilized protease-resistant PrP and nonprotein components of prion rods. *J. Biol. Chem.*, **276**: 14324–14328.
- Sokolowski, F. *Expression, Reinigung und FTIR-spektroskopische Strukturuntersuchung des rekombinanten Prionproteins des Syrischen Hamsters (SHaPrP<sup>90-233</sup>)*. Diplomarbeit, Freie Universität Berlin, 2001.
- Sokolowski, F.; Modler, A. J.; Masuch, R.; Zirwer, D.; Baier, M.; Lutsch, G.; Moss, D. A.; Gast, K. und Naumann, D. 2003. Formation of critical oligomers is a key event during conformational transition of recombinant Syrian hamster prion protein. *J. Biol. Chem.*, **278**: 40481–40492.
- Stöckel, J. und Hartl, F. U. 2001. Chaperonin-mediated de novo generation of prion protein aggregates. *J. Mol. Biol.*, **313**: 861–872.
- Stefani, M. und Dobson, C. M. 2003. Protein aggregation and aggregate toxicity: new insights into protein folding, misfolding diseases, and biological evolution. *J. Mol. Med.*, **81**: 678–699.
- Stryer, L. *Biochemie*. Spektrum Akad. Verl., Heidelberg, Berlin, Oxford, 1996.
- Susi, H. 1972. Infrared spectroscopy – conformation. *Meth. Enzymol.*, **26**: 455–472.
- Swietnicki, W.; Morillas, M.; Chen, S. G.; Gambetti, P. und Surewicz, W. K. 2000. Aggregation and fibrilization of the recombinant human prion protein huPrP<sup>90-231</sup>. *Biochemistry*, **39**: 424–431.
- Swietnicki, W.; Petersen, R.; Gambetti, P. und Surewicz, W. K. 1997. pH-dependent stability and conformation of the recombinant human prion protein PrP(90–231). *J. Biol. Chem.*, **272**: 27517–27520.
- Thomzig, A.; Spassov, S.; Friedrich, M.; Naumann, D. und Beekes, M. 2004. Discriminating scrapie and bovine spongiform encephalopathy isolates by infrared spectroscopy of pathological prion protein. *J. Biol. Chem.*, **279**: 33847–33854.

- Torrent, J.; Alvarez-Martinez, M. T.; Harricane, M. C.; Heitz, F.; Liautard, J. P.; Balny, C. und Lange, R. 2004. High pressure induces scrapie-like prion protein misfolding and amyloid fibril formation. *Biochemistry*, **43**: 7162–7170.
- Torrent, J.; Alvarez-Martinez, M. T.; Heitz, F.; Liautard, J. P.; Balny, C. und Lange, R. 2003. Alternative prion structural changes revealed by high pressure. *Biochemistry*, **42**: 1318–1325.
- Tsui, F. C.; Ojcius, D. M. und Hubbell, W. L. 1986. The intrinsic pK<sub>a</sub> values for phosphatidylserine and phosphatidylethanolamine in phosphatidylcholine host bilayers. *Biophys. J.*, **49**: 459–468.
- Tycko, R. 2004. Progress towards a molecular-level structural understanding of amyloid fibrils. *Curr. Opin. Struct. Biol.*, **14**: 96–103.
- Vanik, D. L. und Surewicz, W. K. 2002. Disease-associated F198S mutation increases the propensity of the recombinant prion protein for conformational conversion to scrapie-like form. *J. Biol. Chem.*, **277**: 49065–49070.
- Venyaminov, S. Y. und Kalnin, N. N. 1990. Quantitative IR spectrophotometry of peptide compounds in water (H<sub>2</sub>O) solutions. II. Amide absorption bands of polypeptides and fibrous proteins in  $\alpha$ -,  $\beta$ -, and random coil conformations. *Biopolymers*, **30**: 1259–1271.
- Volles, M. J. und Lansbury, J.; P. T. 2002. Vesicle permeabilization by protofibrillar  $\alpha$ -synuclein is sensitive to Parkinson's disease-linked mutations and occurs by a pore-like mechanism. *Biochemistry*, **41**: 4595–4602.
- Walsh, D. M. und Selkoe, D. J. 2004. Oligomers on the brain: the emerging role of soluble protein aggregates in neurodegeneration. *Protein Pept. Lett.*, **11**: 213–228.
- Welker, E.; Raymond, L. D.; Scheraga, H. A. und Caughey, B. 2002. Intramolecular versus intermolecular disulfide bonds in prion proteins. *J. Biol. Chem.*, **277**: 33477–33481.
- Wildegger, G.; Liemann, S. und Glockshuber, R. 1999. Extremely rapid folding of the C-terminal domain of the prion protein without kinetic intermediates. *Nat. Struct. Biol.*, **6**: 550–553.
- Wille, H.; Michelitsch, M. D.; Guenebaut, V.; Supattapone, S.; Serban, A.; Cohen, F. E.; Agard, D. A. und Prusiner, S. B. 2002. Structural studies of the scrapie prion protein by electron crystallography. *Proc. Natl. Acad. Sci. USA*, **99**: 3563–3568.
- Winter, R. *Methoden der Biophysikalischen Chemie*. Teubner, Stuttgart, 1998.
- Wüthrich, K. und Riek, R. 2001. Three-dimensional structures of prion proteins. *Adv. Protein Chem.*, **57**: 55–82.

- Xiong, L.-W.; Raymond, L. D.; Hayes, S. F.; Raymond, G. J. und Caughey, B. 2001. Conformational change, aggregation and fibril formation induced by detergent treatments of cellular prion protein. *J. Neurochem.*, **79**: 669–678.
- Yon, J. M. 2002. Protein folding in the post-genomic era. *J. Cell. Mol. Med.*, **6**: 307–327.
- Zahn, R.; Liu, A.; Lührs, T.; Riek, R.; Schroetter, C.von; García, F. L.; Billeter, M.; Calzolari, L.; Wider, G. und Wüthrich, K. 2000. NMR solution structure of the human prion protein. *Proc. Natl. Acad. Sci. USA*, **97**: 145–150.
- Zhang, H.; Stöckel, J.; Mehlhorn, I.; Groth, D.; Baldwin, M. A.; Prusiner, S. B.; James, T. L. und Cohen, F. E. 1997. Physical studies of conformational plasticity in a recombinant prion protein. *Biochemistry*, **36**: 3543–3553.