
8 References

Aarestrup FM. 1998. Association between decreased susceptibility to a new antibiotic for treatment of human diseases, everninomicin (SCH 27899), and resistance to an antibiotic used for growth promotion in animals, avilamycin. *Microb Drug Resist* 4: 137-141.

Aarestrup FM, Kruse H, Tast E, Hammerum AM and Jensen LB. 2000. Associations between the use of antimicrobial agents for growth promotion and the occurrence of resistance among *Enterococcus faecium* from broilers and pigs in Denmark, Finland, and Norway. *Microb Drug Resist* 6: 63-70.

Aarestrup FM, Seyfarth AM, Emborg HD, Pedersen K, Hendriksen RS and Bager F. 2001. Effect of abolishment of the use of antimicrobial agents for growth promotion on occurrence of antimicrobial resistance in fecal enterococci from food animals in Denmark. *Antimicrob Agents Chemother* 45: 2054-2059.

Akkermans ADL, Konstantinov SR, Zhu Wei-Yun, Favier CF and Williams BA. 2003. Postnatal development of the intestinal microbiota of the pig. *Digestive Physiology in Pigs*: 49-56.

Amann R, Binder BJ, Olson RJ, Chisholm SW, Devereux R and Stahl DA. 1990. Combination of 16S rRNA-targeted oligonucleotide probes with flow cytometry for analyzing mixed microbial populations. *Appl Environ Microbiol* 56: 1919-1925.

Asche GL, Lewis AJ and Peo ER Jr. 1989. Protein digestion in weanling pigs: Effect of feeding regimen and endogenous protein secretion. *J Nutr* 119: 1083-1092.

Bach Knudsen KE, Jensen BB, Andersen JO and Hansen I. 1991. Gastrointestinal implications in pigs of wheat and oat fractions 2. Microbial activity in the gastrointestinal tract. *Br J Nutr* 65: 233-248.

Bardocz S. 1993. The role of dietary polyamines. *European Journal of Clinical Nutrition* 47: 683-690.

Bardocz S, White A, Grant G, Walker TJ, Brown DS and Pusztai A. 2001. The role of polyamines in intestinal function and gut maturation. *Gut Environment in Pigs*: 29-42.

Bartelt J, Bergner H, Drochner W, Kijora C and Gotz KP. 1994. The effect of a different protein and fiber intake on the ileal flow of endogenous protein in ¹⁵N-labeled swine. *Arch Anim Nutr* 46: 305-319.

Blachier F. 1997. Intestinal polyamines. *Digestive Physiology in Pigs* 88: 222-233.

Blank R, Mosenthin R and Sauer WC. 1998. Gastrointestinal response of early-weaned pigs to supplementation of wheat-soyabean meal diets with fumaric acid and sodium bicarbonate. *Journal of Animal and Feed Sciences* 7: 185-189.

Bolduan G, Schneider R, Jung H and Klenke B. 1986. The ammonia content in the digesta of pigs. *Arch Anim Nutr* 36: 281-285.

Bover-Cid S and Holzapfel WH. 1999. Improved screening procedure for biogenic amine production by lactic acid bacteria. *Int J Food Microbiol* 53: 33-41.

Bover-Cid S, Izquierdo-Pulido M and Vidal-Carou MC. 2001. Effect of the interaction between a low tyramine-producing *Lactobacillus* and proteolytic staphylococci on biogenic amine production during ripening and storage of dry sausages. *Int J Food Microbiol* 65: 113-123.

Caine WR, Verstegen MWA, Tamminga S, Sauer WC and Schulze H. 1997. Bacterial contributions of N and indispensable amino acids to ileal digesta of newly-weaned piglets fed diets with protease-treated soybean meal. *Digestive Physiology in Pigs* 88: 454-457.

Canibe N, Engberg RM and Jensen BB. 2001. An overview of the effects of organic acids on gut flora and gut health. Workshop on Alternatives to Feed Antibiotics and Anticoccidials in the Pig and Poultry Meat Production.

Canibe N, Steien SH, Overland M and Jensen BB. 2001. Effect of K-diformate in starter diets on acidity, microbiota, and the amount of organic acids in the digestive tract of piglets, and on gastric alterations. *J Anim Sci* 79: 2123-2133.

Chung TK and Baker DH. 1992. Apparent and true amino acid digestibility of a crystalline amino acid mixture and of casein: comparison of values obtained with ileal-cannulated pigs and cecectomized cockerels. *J Anim Sci* 70: 3781-3790.

Collier CT, Smiricky-Tjardes MR, Albin DM, Wubben JE, Gabert VM, Deplancke B, Bane D, Anderson DB and Gaskins HR. 2003. Molecular ecological analysis of porcine ileal microbiota responses to antimicrobial growth promoters. *J Anim Sci* 81: 3035-3045.

Conway PL. 1994. Function and regulation of the gastrointestinal microbiota of the pig. *Digestive physiology in pigs*: 231-240.

Darragh AJ and Hodgkinson SM. 2000. Quantifying the digestibility of dietary protein. *J Nutr* 130: 1850S-1856S.

de Lange CF, Souffrant WB and Sauer WC. 1990. Real ileal protein and amino acid digestibilities in feedstuffs for growing pigs as determined with the ¹⁵N-isotope dilution technique. *J Anim Sci* 68: 409-418.

Decuypere J and Van der HH. 1972. Study of the gastro-intestinal microflora of suckling piglets and early weaned piglets reared using different feeding systems. *Zentralbl Bakteriol [Orig A]* 221: 492-510.

Dersjant-Li Y, Verstegen MW, Schulze H, Zandstra T, Boer H, Schrama JW and Verreth JA. 2001. Performance, digesta characteristics, nutrient flux, plasma composition, and organ weight in pigs as affected by dietary cation anion difference and nonstarch polysaccharide. *J Anim Sci* 79: 1840-1848.

Eckel B, Roth FX, Kirchgessner M and Eidelsburger U. 1992. Zum Einfluss von Ameisensäure auf die Konzentrationen an Ammoniak und biogenen Aminen im Gastrointestinaltrakt. 4. Mitteilung: Untersuchungen zur nutritiven Wirksamkeit von organischen Säuren in der Ferkelaufzucht. *J Anim Physiol Anim Nutr* 67: 198-205.

Favier CF, Vaughan EE, de Vos WM and Akkermans AD. 2002. Molecular Monitoring of succession of bacterial communities in human neonates. *Appl Environ Microbiol* 68: 219-226.

Franklin MA, Mathews AG, Vickers JR and Clift RA. 2002. Characterization of microbial populations and volatile fatty acid concentrations in the jejunum, ileum and cecum of pigs weaned at 17 vs 24 days of age. *J Anim Sci* 80: 2904-2910.

Fuller MF and Reeds PJ. 1998. Nitrogen cycling in the gut. *Annu Rev Nutr* 18:411.

Garrett JE, Goodrich RD and Meiske JC. 1987. Measurement and use of D-alanine as a bacterial marker. *Can J Anim Sci* 67: 735-743.

Gaskins H.R., Collier C.T. and Anderson D.B. 2002. Antibiotics as growth promotants. mode of action. *Animal Biotechnology* 13: 29-42.

Gedek B, Kirchgessner M, Wiehler S, Bott A, Eidelsburger U and Roth FX. 1993. The nutritive effect of *Bacillus cereus* as a probiotic in the raising of piglets. 2. Effect and microbial count, composition and resistance determination of gastrointestinal and fecal microflora. *Arch Anim Nutr* 44: 215-226.

Geissler Ch., Hoffmann M. and Hickel B. 1976. Ein Beitrag zur gaschromatographischen Bestimmung flüchtiger Fettsäuren. *Arch Anim Nutr* 26: 123-129.

Gollnisch K. 1998. Influence of an antimicrobial feed additive (avilamycin) and a non-starch-polysaccharide (NSP)-hydrolyzing enzyme alone or in combination on the intestinal microflora of pigs. *PhD Thesis, Veterinary School Hannover.*

Grala W, Verstegen MW, Jansman AJ, Huisman J and van Leeusen P. 1998. Ileal apparent protein and amino acid digestibilities and endogenous nitrogen losses in pigs fed soybean and rapeseed products. *J Anim Sci* 76: 557-568.

Greife HA, Rooke JA and Armstrong DG. 1985. The digestion by cattle of silage-containing diets fed at two dry matter intakes. 2. Digestion of total amino acids and of D-alanine and D-glutamic acid. *Br J Nutr* 54: 483-492.

Haacker K, Block H-J and Weissbach F. 1983. Zur kolorimetrischen Milchsäurebestimmung in Silagen mit p-Hydroxydiphenyl. *Arch Anim Nutr* 33: 505-512.

Hagemeister H and Erbersdobler H. 1985. Chemical labelling of dietary protein by transformation of lysine to homoarginine: A new technique to follow intestinal digestion and absorption. *Proc Nutr Soc* 44: 133A+.

Harmsen HJM and Welling GW. 2002. Fluorescence *in situ* hybridisation as a tool in intestinal bacteriology. *Probiotics and Prebiotics: Where are we going?:* 40-58.

Heilig HG, Zoetendal EG, Vaughan EE, Marteau P, Akkermans AD and de Vos WM. 2002. Molecular diversity of *Lactobacillus spp.* and other lactic acid bacteria in the human intestine as determined by specific amplification of 16S ribosomal DNA. *Appl Environ Microbiol* 68: 114-123.

Hennig U, Schönhusen U and Souffrant WB. 1999. Nitrogen and amino acids originated from the small intestinal flora of pigs determined by means of D-alanine. *Protein Metabolism and Nutrition:* 42-42.

Hennig U, Noel R., Herrmann U., Wünsche J and Mehnert E. 1986. Ernährungsphysiologische Untersuchungen an Schweinen mit Ileo-Rektal-Anastomosen. *Arch Anim Nutr* 36: 585-596.

Hill IR, Kenworthy R and Porter P. 1969. Dietary lactobacilli and amine production in the alimentary tract of the pig in relation to weaning and postweaning diarrhoea. *J Gen Microbiol* 59: i-ii.

Hill IR, Kenworthy R and Porter P. 1970. The production of amines in the alimentary tract of the pig in relation to dietary lactobacilli, weaning and post-weaning diarrhoea. 2. In vitro studies of the catabolic activity of the small intestinal microflora. *Proc Nutr Soc* 29: Suppl.:14A+.

Højberg O, Canibe N, Knudsen B and Jensen BB. 2003. Potential rates of fermentation in digesta from the gastrointestinal tract of pigs: Effect of feeding fermented liquid feed. *Appl Environ Microbiol* 69: 408-418.

Houdijk JG, Bosch MW, Tamminga S, Verstegen MW, Berenpas EB and Knoop H. 1999. Apparent ileal and total-tract nutrient digestion by pigs as affected by dietary nondigestible oligosaccharides. *J Anim Sci* 77: 148-158.

Huisman J, Heinz T, van der Poel AF, van Leeuwen P, Souffrant WB and Verstegen MW. 1992. True protein digestibility and amounts of endogenous protein measured with the ¹⁵N-dilution technique in piglets fed on peas (*Pisum sativum*) and common beans (*Phaseolus vulgaris*). *Br J Nutr* 68: 101-110.

Ishikawa S. 1966. Reliability of polyethylenglycol as an indicator for digestion studies with swine, Part I. rate of passage of polyethylenglycol through the digestive tract. *Agr Biol Chem* 30: 278-284.

Jansman AJ, Verstegen MW, Huisman J and van den Berg JW. 1995. Effects of hulls of faba beans (*Vicia faba* L.) with a low or high content of condensed tannins on the apparent ileal and fecal digestibility of nutrients and the excretion of endogenous protein in ileal digesta and feces of pigs. *J Anim Sci* 73: 118-127.

Jensen BB. 1998. The impact of feed additives on the microbial ecology of the gut in young pigs. *Journal of Animal and Feed Sciences* 7: 45-64.

Kamphues J, Tabeling R and Stuke O. 2003. Untersuchungen zu möglichen diätetisch interessanten Effekten von Lactulose als Futterzusatz in der Schweinefütterung. *Dtsch Tierärztl Wochenschr* 110: 365-368.

Konstantinov SR, Zhu Wei-Yun, Williams BA, Tamminga S, de Vos WM and Akkermans AD. 2003. Effect of fermentable carbohydrates on piglet faecal bacterial communities as revealed by denaturing gradient gel electrophoresis analysis of 16S ribosomal DNA. *FEMS Microbiology Ecology* 43: 225-235.

Konstantinov SR, Fitzsimon N, Vaughan EE and Akkermans ADL. 2002. From composition to functionality of the intestinal microflora. *Probiotics and Prebiotics: Where are we going?:* 59-84.

Konstantinov SR, Awati A, Smidt H, Williams BA, Akkermans ADL and de Vos WM. 2004a. Specific response of a novel and abundant *Lactobacillus amylovorus*-like phylotype to dietary prebiotics in the gut of weaning piglets. *Appl Environ Microbiol* 70: 3821-3830.

Konstantinov SR, Favier CF, Wei Yun Zhu, Williams BA, Klüß J, Souffrant WB, de Vos WM, Akkermans ADL and Smidt H. 2004b. Microbial diversity studies of the porcine gastrointestinal ecosystem during weaning transition. *Animal Research* 53: 1-8.

Kovacs F, Nagy B and Sinkovics G. 1972. The gut bacterial flora of healthy early weaned piglets, with special regard to factors influencing its composition. *Acta Vet Acad Sci Hung* 22: 327-338.

Krause DO, Easter RA, White BA and Mackie RI. 1995. Effect of weaning diet on the ecology of adherent lactobacilli in the gastrointestinal tract of the pig. *J Anim Sci* 73: 2347-2354.

Kreienbring F and Wünsche J. 1974. Bericht über die Stickstoff- und Aminosäurenbestimmungs-Enquete 1971/72 im Themekreis und über weitere methodische Arbeiten. *Akad Landwirtsch -Wiss DDR* 124: 19-34.

Kreienbring F. 1987. Weitere Ergebnisse zur vergleichenden Bestimmung von Aminosäuren. *Die Nahrung* 31: 855-862.

Kulla S. 2001. Effects of K-diformiate on the intestinal microflora and parameters of microbial metabolism as well as various digestive processes in chyme of weaned piglets. *PhD Thesis, Veterinary School Hannover.*

Laplace JP, Darcy-Vrillon B, Perez JM, Henry Y, Giger S and Sauvant D. 1989. Associative effects between two fibre source on ileal and overall digestibilities of amino acids, energy and cell-wall components in growing pigs. *Br J Nutr* 61: 75-87.

Laplace JP, Souffrant WB, Hennig U, Chabeauti E and Fevrier C. 1994. Measurement of precaecal dietary protein and plant cell wall digestion in pigs; Comparison of four surgical procedures for ileorectal anastomosis. *Livest Prod Sci* 40: 313-328.

- Leser TD, Lindecrona RH, Jensen TK, Jensen BB and Møller K.** 2000. Changes in bacterial community structure in the colon of pigs fed different experimental diets and after infection with *Brachyspira hyodysenteriae*. *Appl Environ Microbiol* 66: 3290-3296.
- Leser TD, Amenuvor JZ, Jensen TK, Lindecrona RH, Boye M and Møller K.** 2002. Culture-independent analysis of gut bacteria: the pig gastrointestinal tract microbiota revisited. *Appl Environ Microbiol* 68: 673-690.
- Li S, Sauer WC, Huang SX and Gabert VM.** 1996. Effect of beta-glucanase supplementation to hullless barley- or wheat-soybean meal diets on the digestibilities of energy, protein, beta-glucans, and amino acids in young pigs. *J Anim Sci* 74: 1649-1656.
- Maribo H, Jensen BB and Hedemann MS.** 2000. Different doses of organic acids for weaned piglets. *The National Committee for Pig Production*.
- Mariotti F, Huneau J-F, Mahe S and Tome D.** 2000. Protein metabolism and the gut. *Curr Opin Clin Nutr Metab Care* 3: 45-50.
- Mariscal-Landin G, Seve B, Colleaux Y and Lebreton Y.** 1995. Endogenous amino nitrogen collected from pigs with end-to-end ileorectal anastomosis is affected by the method of estimation and altered by dietary fiber. *J Nutr* 125: 136-146.
- Mathew AG, Sutton AL, Scheidt AB, Patterson JA, Kelly DT and Meyerholtz KA.** 1993. Effect of galactan on selected microbial populations and pH and volatile fatty acids in the ileum of the weanling pig. *J Anim Sci* 71: 1503-1509.
- Mathew AG, Robbins CM, Chattin SE and Quigley JD 3rd.** 1997. Influence of galactosyl lactose on energy and protein digestibility, enteric microflora, and performance of weanling pigs. *J Anim Sci* 75: 1009-1016.
- Mathew AG, Chattin SE, Robbins CM and Golden DA.** 1998. Effects of a direct-fed yeast culture on enteric microbial populations, fermentation acids, and performance of weanling pigs. *J Anim Sci* 76: 2138-2145.

- Mathew AG, Jones T and Franklin MA.** 1994. Effect of creep feeding on selected microflora and short-chain fatty acids in the ileum of weanling pigs. *J Anim Sci* 72: 3163-3168.
- Mavromichalis I, Parr TM, Gabert VM and Baker DH.** 2001. True ileal digestibility of amino acids in sow's milk for 17-day-old pigs. *J Anim Sci* 79: 707-713.
- McCartney AL.** 2002. Application of molecular biological methods for studying probiotics and the gut flora. *Br J Nutr* 88 (Suppl.): S29-S37.
- McCracken BA, Gaskins HR, Ruwe-Kaiser PJ, Klasing KC and Jewell DE.** 1995. Diet-dependent and diet-independent metabolic responses underlie growth stasis of pigs at weaning. *J Nutr* 125: 2838-2845.
- Méndez-Álvarez S, Pérez-Hernández X and Claverie-Martín F.** 2000. Glycopeptide resistance in enterococci. *Internatl Microbiol* 3: 71-80.
- Mikkelsen LL and Jensen BB.** 1998. Performance and microbial activity in the gastrointestinal tract of piglets fed fermented liquid feed at weaning. *Journal of Animal and Feed Sciences* 7: 211-215.
- Mikkelsen LL, Bendixen C, Jakobsen M and Jensen BB.** 2003. Enumeration of bifidobacteria in gastrointestinal samples from piglets. *Appl Environ Microbiol* 69: 654-658.
- Moeser AJ, Kim IB, van Heugten E and Kempen TA.** 2002. The nutritional value of degermed, dehulled corn for pigs and its impact on the gastrointestinal tract and nutrient excretion. *J Anim Sci* 80: 2629-2638.
- Moughan PJ, Pedraza M, Smith WC, Williams M and Wilson MN.** 1990. An evaluation with piglets of bovine milk, hydrolyzed bovine milk, and isolated soybean proteins included in infant milk formulas. I. Effect on organ development, digestive enzyme activities, and amino acid digestibility. *Journal of Pediatric Gastroenterology and Nutrition* 10: 385-394.

- Moughan PJ, Souffrant WB and Hodgkinson SM.** 1998. Physiological approaches to determining gut endogenous amino acid flows in the mammal. *Arch Anim Nutr* 51: 237-252.
- Muyzer G, De Waal EC and Uitterlinden AG.** 1993. Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA. *Appl Environ Microbiol* 59: 695-700.
- Naito S, Hayashidani H, Kaneko K, Ogawa M and Benno Y.** 1995. Development of intestinal lactobacilli in normal piglets. *J Appl Bacteriol* 79: 230-236.
- Naumann C and Bassler R.** 1993. Die chemische Untersuchung von Futtermitteln. Methodebnuch. III. Ergänzungslieferung. *VDLUFA-Verlag. Darmstadt.*
- Noack J, Kleesen B, Proll J, Dongowski G and Blaut M.** 1998. Dietary guar gum and pectin stimulate intestinal microbial polyamine synthesis in rats. *J Nutr* 128: 1385-1391.
- NRC.** 1998. Nutrient Requirements of Swine:10th revised edition. *National Academy Press. Washington DC.*
- Nyachoti CM, de Lange CF, McBride BW, Leeson S and Gabert VM.** 2000. Endogenous gut nitrogen losses in growing pigs are not caused by increased protein synthesis rates in the small intestine. *J Nutr* 130: 566-572.
- Nyachoti CM, de Lange CF and Schulze H.** 1997. Estimating endogenous amino acid flows at the terminal ileum and true ileal amino acid digestibilities in feedstuffs for growing pigs using the homoarginine method. *J Anim Sci* 75: 3206-3213.
- O'Sullivan DJ.** 2000. Methods for analysis of the intestinal microflora. *Curr Issues Intest Microbiol* 1: 39-50.
- Piva A, Prandini A, Fiorentini L, Morlacchini M, Galvano F and Luchansky JB.** 2002. Tributyrin and lactitol synergistically enhanced the trophic status of the intestinal mucosa and reduced histamine levels in the gut of nursery pigs. *J Anim Sci* 80: 670-680.

- Porter P, Kenworthy R and Hill IR.** 1970. The production of amines in the alimentary tract of the pig in relation to dietary lactobacilli, weaning and post-weaning diarrhoea. 1. In vivo studies of intestinal and urinary amines. *Proc Nutr Soc* 29: Suppl. 13A+.
- Pryde S, Richardson AJ, Stewart CS and Flint HJ.** 1999. Molecular analysis of the microbial diversity present in the colonic wall, colonic lumen, and cecal lumen of a pig. *Appl Environ Microbiol* 65: 5372-5377.
- Quigley JD and Schwab CG.** 1988. Comparison of D-alanine and diaminopimelic acid as bacterial markers in young calves. *J Anim Sci* 66: 758-763.
- Redlich J, Souffrant WB, Laplace JP, Hennig U, Berg R and Mouwen JMVM.** 1997. Morphometry of the small intestine in pigs with ileo-rectal anastomosis. *Can J Vet Res* 61: 21-27.
- Risley CR, Kornegay ET, Lindemann MD, Wood CM and Eigel WN.** 1992. Effect of feeding organic acids on selected intestinal content measurements at varying times postweaning in pigs. *J Anim Sci* 70: 196-206.
- Rutherford SM and Moughan PJ.** 1998. The digestible amino acid composition of several milk proteins: application of a new bioassay. *J Dairy Sci* 81: 909-917.
- Sanguinetti CJ, Dias Neto E and Simpson AJ.** 1994. Rapid silver staining and recovery of PCR-products separated on polyacrylamide gels. *Biotechniques* 17: 915-919.
- Savage DC.** 1977. Microbial ecology of the gastrointestinal tract. *Annu Rev Microbiol* 31: 107-133.
- Schleifer K-H and Kandler O.** 1972. Peptidoglycan types of bacterial cell walls and their taxonomic implications. *Bacteriological Reviews* 36: 407-477.
- Schnabel E, Bolduan G, Klenke B, Borchert E and Thiele I.** 1985. Mikrobielle Metaboliten in der Digesta von Ferkeln nach der Mahlzeit. *Arch Anim Nutr* 35: 623-628.

Schneider R, Kreienbring F, Bolduan G and Beck M. 1989. Biogene Amine in der Digesta von Schweinen. *Arch Anim Nutr* 39: 1021-1029.

Scholten RH, van der Peet-Schwering CM, den Hartog LA, Balk M, Schrama JW and Verstegen MW. 2002. Fermented wheat in liquid diets: effects on gastrointestinal characteristics in weanling piglets. *J Anim Sci* 80: 1179-1186.

Schönhusen U, Voigt J and Kreienbring F. 1995. Evaluation of different markers for the determination of microbial nitrogen flow into the duodenum of dairy cows. *Arch Anim Nutr* 48: 147-158.

Schulze H, van Leeuwen P, Verstegen MW and van den Berg JW . 1995. Dietary level and source of neutral detergent fiber and ileal endogenous nitrogen flow in pigs. *J Anim Sci* 73: 441-448.

Schulze H, van Leeuwen P, Verstegen MW, Huisman J, Souffrant WB and Ahrens F. 1994. Effect of level of dietary neutral detergent fiber on ileal apparent digestibility and ileal nitrogen losses in pigs. *J Anim Sci* 72: 2362-2368.

Schulze H, Savelkoul FH, Verstegen MW, van der Poel AF, Tamminga S and Groot NS. 1997. Nutritional evaluation of biologically treated white kidney beans (*Phaseolus vulgaris* L.) in pigs: ileal and amino acid digestibility. *J Anim Sci* 75: 3187-3194.

Seabra M, Carvalho S, Freire JP, Ferreira RB, Mourato M, Cunha L, Cabral F, Teixeira A and Aumaitre A. 2001. *Lupinus luteus*, *Vicia sativa* and *Lathyrus cicera* as protein sources for piglets: ileal and total tract apparent digestibility of amino acids and antigenic effects. *Animal Feed Science & Technology* 89: 1-16.

Sève B and Lahaye L. 2004. Nitrogen and amino acid cost of digestion: dietary factors affecting ileal endogenous losses and consequences on the net availability of dietary amino acids for maintenance and growth. *Digestive Physiology in Pigs* 1: 263a-263y.

Sghir A, Antonopoulos D and Mackie RI. 1998. Design and evaluation of a *Lactobacillus* group-specific ribosomal RNA-targeted hybridization probe and its application to the study of intestinal microecology in pigs. *Syst Appl Microbiol* 21: 291-296.

Simon O, Bergner H and Partridge IG. 1987. Estimation of the endogenous N proportions in ileal digesta and faeces in ¹⁵N-labelled pigs. *Arch Anim Nutr* 37: 851-859.

Simon O. 2001. The influence of feed composition on protein metabolism in the gut. *Gut Environment in Pigs*: 63-84.

Simpson JM, McCracken VJ, Gaskins HR and Mackie RI. 2000. Denaturing gradient gel electrophoresis analysis of 16S ribosomal DNA amplicons to monitor changes in fecal bacterial populations of weaning pigs after introduction of *Lactobacillus reuteri* strain MM53. *Appl Environ Microbiol* 66: 4705-4714.

Simpson JM, Kocherginskaya SA, Aminov RI, Skerlos LT, Bradley TM, Mackie RI and White BA. 2002. Comparative microbial diversity in the gastrointestinal tracts of food animal species. *Integrative and Comparative Biology* 42: 327-331.

Simpson JM, McCracken VJ, White BA, Gaskins HR and Mackie RI. 1999. Application of denaturant gradient gel electrophoresis for the analysis of the porcine gastrointestinal microbiota. *Journal of Microbiological Methods* 36: 167-179.

Sohn KS, Maxwell CV, Southern LL and Buchanan DS. 1994. Improved soybean protein sources for early-weaned pigs: II. Effects on ileal amino acid digestibility. *J Anim Sci* 72: 631-637.

Souffrant WB, Köhler R, Matkowitz R, Gebhardt G and Schmandke H. 1981. Nutritional and physiological studies of swine for the evaluation of modified proteins. 2. Determination of endogenous N in the contents of the small intestines using the ¹⁵N-tracer method. *Arch Anim Nutr* 31: 675-683.

Souffrant WB, Köhler R and Gebhardt G. 1982. Determination de l'azote endogene dans les contenus digestifs par la technique isotopique (¹⁵N). *Le Colloques de l'INRA* 12.

Souffrant WB, Verstegen MWAed, Huisman Jed and den Hartog LA. 1991. Endogenous nitrogen losses during digestion in pigs. *Digestive Physiology in Pigs*: 147-166.

- Spreeuwenberg MAM, Verdonk JMAJ, Gaskins HR and Verstegen M.W.A.** 2001. Small intestine epithelial barrier function is compromised in pigs with low feed intake at weaning. *J Nutr* 131: 1520-1527.
- Tamminga S, Schulze H, Van Bruchem J and Huisman J.** 1995. The nutritional significance of endogenous N-losses along the gastro-intestinal tract of farm animals. *Arch Anim Nutr* 48: 9-22.
- Tannock GW, Fuller R and Pedersen K.** 1990. Lactobacillus succession in the piglet digestive tract demonstrated by plasmid profiling. *Appl Environ Microbiol* 56: 1310-1316.
- Thorpe J.** 1999. The effects of two in-feed protease enzymes on the digestibility of soy proteins in the early weaned pig. *Thesis. University of Bristol.*
- Voigt J and Steger H.** 1967. Zur quantitativen Bestimmung von Ammoniak, Harnstoff und Ketokörpern in biologischem Material mit Hilfe eines modifizierten Mikrodiffusionsgefäßes. *Arch Anim Nutr* 17: 289-293.
- Wang W and Higuchi CM.** 2000. Dietary soy protein is associated with reduced intestinal mucosal polyamine concentration in male Wistar rats. *J Nutr* 130: 1815-1820.
- Wegener HC, Aarestrup FM, Jensen LB, Hammerum AM and Bager F.** 1998. The association between the use of antimicrobial growth promoters and development of resistance in pathogenic bacteria towards growth promoting and therapeutic antimicrobials. *Journal of Animal and Feed Sciences* 7: 7-14.
- Welling GW, Efferich P, Raangs GC, Wildeboer-Veloo AC, Jansen GJ and Degener JE.** 1997. 16S ribosomal RNA-targeted oligonucleotide probes for monitoring of intestinal tract bacteria. *Scand J Gastroenterol Suppl* 32: 17-19.
- Williams BA, Awati A, Bosch MW, Konstantinov SR and Akkermans AD.** 2003. The effect of dietary fermentable carbohydrates on lactic acid concentration and presence of lactobacilli in small intestinal digesta of piglets at ten days after weaning. *Tropical and Subtropical Agroecosystems* 3: 233-235.

Wu G, Flynn NE, Knabe DA and Jaeger LA. 2000. A cortisol surge mediates the enhanced polyamine synthesis in porcine enterocytes during weaning. *Am J Physiol Regulatory Integrative Comp Physiol* 279: R554-R559.

Wu G, Flynn NE and Knabe DA. 2000. Enhanced intestinal synthesis of polyamines from proline in cortisol-treated piglets. *Am J Physiol Endocrinol Metab* 279: E395-E402.

Wünsche J, Volker T, Souffrant WB and Borgmann E. 1991. Determination of bacterial N portions in feces and differently collected ileum chymus from swine. *Arch Anim Nutr* 41: 703-716.

Zhu Wei-Yun, Williams BA, Konstantinov SR, Tamminga S, De Vos WM and Akkermans AD. 2003. Analysis of 16S rDNA reveals bacterial shift during in vitro fermentation of fermentable carbohydrate using piglet faeces as inoculum. *Anaerobe* 9: 175-180.

Zoetendal EG, Collier CT, Koike S, Mackie RI and Gaskins HR. 2004. Molecular ecological analysis of the gastrointestinal microbiota: a review. *J Nutr* 134: 465-472.