

## 8. Literaturverzeichnis

- [1] Dickenmann M, Gurke L (2005): Kidney transplantation. *Ther Umsch.* 62(7): 443-447.
- [2] Schnuelle P, Lorenz D, Trede M et al. (1998): Impact of renal cadaveric transplantation on survival in end-stage renal failure: evidence for reduced mortality risk compared with hemodialysis during long-term follow-up. *J Am Soc Nephrol.* 9(11): 2135-2141.
- [3] Winkelmayr WC, Weinstein MC, Mittleman MA, Gi et al. (2002): Health economic evaluations: the special case of end-stage renal disease treatment. *Med Decis Making.* 22(5): 417-430.
- [4] Niu SF, Li IC. (2005): Quality of life of patients having renal replacement therapy. *J Adv Nurs.* 51(1): 15-21.
- [5] Bittencourt ZZ, Alves Filho G, Mazzali M et al. (2004): Quality of life in renal transplant patients: impact of a functioning graft. *Rev Saude Publica.* 38(5): 732-734.
- [6] Deutsche Stiftung Organspende: Bericht über die Tätigkeit der Transplantationszentren 2005. URL: [http://www.dso.de/pdf/berichte2005/Gesamt\\_Taetigkeitsbericht2005.pdf](http://www.dso.de/pdf/berichte2005/Gesamt_Taetigkeitsbericht2005.pdf) (abgerufen am 06.07.2007).
- [7] Deutsche Stiftung Organspende: Bundesweiter Jahresbericht 2006. URL: [http://www.dso.de/pdf/dso\\_jb2006\\_d.pdf](http://www.dso.de/pdf/dso_jb2006_d.pdf) (abgerufen am 06.07.2007).
- [8] Diethelm AG, Deierhoi MH, Hudson SL et al. (1995): Progress in renal transplantation. A single center study of 3359 patients over 25 years. *Ann Surg.* 221: 446-458.
- [9] Lee HM, Posner MP, King AL et al. (1993): Status of long-term (25 years) survival of kidney transplant patients. *Transplant Proc.* 25: 1336-1337.
- [10] Hariharan S, Johnson CP, Bresnahan BA et al. (2000): Improved graft survival after renal transplantation in the United States, 1988 to 1996. *N Engl J Med.* 342(9): 605-12.
- [11] Alvarez G, Gonzalez-Molina M, Cabello M et al. (1991): Pulsed and continuous Doppler evaluation of renal dysfunction after kidney transplantation. *Eur J Radiol.* 12: 108-112.

- [12] Arima M, Ogino T, Hosokawa S et al. (1989): Functional image diagnosis of kidney transplants using ultrasonic Doppler flowmetry and magnetic resonance imaging. *Transplant Proc.* 21: 1907-1911.
- [13] De-Gaetano AM, Boldrini G, Nanni G et al. (1989): Noninvasive surveillance of allografted kidneys by ultrasonic duplex scanning. *Angiology.* 40: 705-715.
- [14] Flückner F, Steiner H, Horn S et al. (1990): Farbkodierte Duplexsonographie and Widerstandsindex bei Nierentransplantaten mit Dysfunction. *RöFo.* 153: 692-697.
- [15] Grenier N, Douws C, Morel D et al. (1991): Detection of vascular complications in renal allografts with color Doppler flow imaging. *Radiology.* 178: 217-223.
- [16] Hollenbeck M (1994): New diagnostic techniques in clinical nephrology. Colour coded duplex sonography of renal transplants – tool or toy for the nephrologist? *Nephrol Dial Transplant.* 9: 1822-1828.
- [17] Steinberg HV, Nelson RC, Murphy FB et al. (1987): Renal allograft rejection: evaluation by Doppler US and MR Imaging. *Radiology.* 162: 337-342.
- [18] Radermacher J, Mengel M, Ellis S et al. (2003): The renal arterial resistance index and renal allograft survival. *N Engl J Med.* 349(2): 182-184.
- [19] Restrepo-Schafer IK, Schwerk WB, Muller TF et al. (1999): Intrarenal doppler flow analysis in patients with kidney transplantation and stable transplant function. *Ultraschall Med.* 20(3): 87-92.
- [20] Letourneau JG, Day DL, Feinberg SB (1987): Ultrasound and computed tomographic evaluation of renal transplantation. *Radiol Clin North Am.* 25: 267-279.
- [21] Dodd GD, Tublin ME, Shah A et al. (1991): Imaging of vascular complications associated with renal transplants. *Am J Roentgenol.* 157: 449-459.
- [22] Hollenbeck M, Kutkuhn B, Grabensee B (1994): Colour Doppler ultrasound in the diagnosis of transplant renal artery stenosis. *Bildgebung.* 61: 248-254.
- [23] Chan L, Kam I: Outcomes and Complications of Renal Transplantation, in Schrier RW, Gottschalk CW (Hrsg): *Diseases of the Kidney*, Boston, 1997, S. 2726-2738.
- [24] Tilney NL, Kupiec-Weglinski JW (1991): The biology of acute transplant rejection. *Ann Surg.* 214(2): 98-106.
- [25] Madan AK, Slakey DP, Becker A et al. (2000): Treatment of antibody-mediated accelerated rejection using plasmapheresis. *J Clin Apher.* 15(3): 180-183.
- [26] Opelz G (1994): Prognostische Faktoren im Verlauf der Nierentransplantation. *Urologe A.* 33(5): 377-382.

- [27] Vincenti F, de Andrés A, Becker T et al. (2006): Interleukin-2 receptor antagonist induction in modern immunosuppression regimens for renal transplant recipients. *Transpl Int.* 19(6): 446-457.
- [28] Boom H, Mallat MJ, de Fijter JW et al. (2000): Delayed graft function influences renal function, but not survival. *Kidney Int.* 58(2): 859-866.
- [29] McLaren AJ, Jassem W, Gray DW et al. (1999): Delayed graft function: risk factors and the relative effects of early function and acute rejection on long-term survival in cadaveric renal transplantation. *Clin Transplant.* 13(3): 266-272.
- [30] Bude RO, Rubin JM, Adler ED (1994): Power versus conventional color Doppler sonography: Comparison in the depiction of normal intrarenal vasculature. *Radiology.* 192: 777-780.
- [31] Genkins SM, Sanfilippo FP, Carroll BA (1989): Duplex Doppler sonography of renal transplants: Lack of sensitivity and specificity in establishing pathologic diagnosis. *Am J Roentgenol.* 152: 535-539.
- [32] Hollenbeck M, Hilbert N, Meusel F et al. (1994): Increasing sensitivity and specificity of Doppler sonographic detection of renal transplant rejection with serial investigation technique. *Clin Investig.* 72: 609-615.
- [33] Townsend RR, Tomlanowich SJ, Goldstein RB et al. (1990): Combined Doppler and morphologic sonographic evaluation of renal transplant rejection. *J Ultrasound Med.* 9: 199-206.
- [34] Weskott HP (1995): Power US: Phantom measurement of slow flow. *Radiology.* 197: 337.
- [35] Waiser J, Schreiber M, Budde K et al. (2000): Prognostic value of the Banff classification. *Transpl Int.* 13 (Suppl 1): 106-111.
- [36] Borrego J, Burgos FJ, Galmes I et al. (1994): Surgical complications in 479 renal transplantations. *Actas Urol Esp.* 18(4): 253-257.
- [37] Khauli RB, Stoff JS, Lovewell T et al. (1993): Post transplant lymphoceles: a critical look into the risk factors, pathophysiology and management. *J Urol.* 150: 22-26.
- [38] Reek C, Tenschert W, Fernandez S et al. (1997): Occurrence of lymphoceles after allogenic kidney transplantation and their therapy. *Urologe A.* 36(4): 313-317.
- [39] Pozniak MA, Dodd GD, Kelcz F (1992): Ultrasonographic evaluation of renal transplantation. *Radiol Clin North Am.* 30(5): 1053-1066.

- [40] Fahlenkamp D, Schönberger B, Lindeke A et al. (1993): Laparoskopische Lymphozelendrainage nach Nierentransplantation. *Z Urologie*. 2: 55-56.
- [41] Benoit G, Bensadoun H, Jardin A (1993): Surgical complications of kidney transplantation. *European Urology Update Series*. 2: 178-183.
- [42] Martinoli C, Bertolotto M, Crespi G et al. (1998): Duplex Doppler analysis of interlobular arteries in transplanted kidneys. *Eur Radiol*. 8: 765-769.
- [43] Schwerk WB, Restrepo IK, Prinz H (1993): Semiquantitative analysis of intrarenal arterial Doppler flow spectra in healthy adults. *Ultraschall Med*. 14: 117-122.
- [44] Delbeke D, Sacks GA, Sandler MP (1989): Diagnosis of allograft renal vein thrombosis. *Clin Nucl Med*. 14: 415-420.
- [45] Jordan ML, Cook GT, Cardelle CJ (1982): Ten years of experience with vascular complications in renal transplants. *J Urol*. 128: 689-692.
- [46] Maia CR, Bittar AE, Goldani JC et al. (1992): Doppler ultrasonography for the detection of renal artery stenosis in transplanted kidneys. *Hypertension*. 19: 207-209.
- [47] Snider JF, Hunter DW, Moradian GP et al. (1989): Transplant renal artery stenosis: evaluation with duplex sonography. *Radiology*. 172: 1027-1030.
- [48] Duckett T, Bretan PN, Cochran ST (1991): Noninvasive radiological diagnosis of renal vein thrombosis in renal transplantation. *J Urol*. 46: 403-406.
- [49] Laplante S, Patriquin HB, Robitaille P et al. (1993): Renal vein thrombosis in children: evidence of early flow recovery with Doppler US. *Radiology*. 189: 37-42.
- [50] Tanaka T, Kyo M, Kokado et al. (2004): Correlation between the Banff 97 classification of renal allograft biopsies and clinical outcome. *Transpl Int*. 17(2): 59-64.
- [51] Bates WD, Davies DR, Welsh K et al. (1999): An evaluation of the Banff classification of early renal allograft biopsies and correlation with outcome. *Nephrol Dial Transplant*. 14(10): 2364-2369.
- [52] Pizzolitto S, Boscutti G, Falconieri G et al. (2004): Clinicopathologic correlations in acute renal graft rejection. *G Ital Nefrol*. 21 (Suppl 26): 19-27.
- [53] Racusen LC, Solez K, Colvin RB et al. (1999): The Banff 97 working classification of renal allograft pathology. *Kidney Int*. 55(2): 713-723.
- [54] Lechevallier E, Dussol B, Luccioni et al (1998): Posttransplantation acute tubular necrosis: risk factors and implications for graft survival. *Am J Kidney Dis*. 32(6): 984-991.

- [55] Stevens PE, Gwyther SJ, Hanson ME et al. (1993): Interpretation of duplex Doppler ultrasound in renal transplants in the early postoperative period. *Nephrol Dial Transplant.* 8: 255-258.
- [56] Keogan MT, Kliewer MA, Hertzberg BS et al. (1996): Renal resistive indexes: variability in Doppler US measurement in a healthy population. *Radiology.* 199(1): 165-9.
- [57] Baxter GM, Ireland H, Moss JG et al. (1995): Colour Doppler ultrasound in renal transplant artery stenosis: which Doppler index? *Clin Radiol.* 50: 618-622.
- [58] Hayry P, Yilmaz S (1994): Chronic allograft rejection: An update. *Transplant Proc.* 26: 3159-3160.
- [59] Rubin JM, Bude RO, Carson PL et al. (1994): Power Doppler US: potentially useful alternative to mean frequency-based color Doppler US. *Radiology.* 190: 853-856.
- [60] Mutze S, Overhoff M, Filimonow S et al. (1996): Ultraschalldiagnostik der Transplantatniere. *Radiologe.* 36: 31-37.
- [61] Blane CE, Gagnadoux MF, Brunelle F et al. (1993): Doppler ultrasonography in the early postoperative evaluation of renal transplants in children. *Can Assoc Radiol J.* 44: 176-178.
- [62] Saarinen O (1991): Diagnostic value of resistive index of renal transplants in the early postoperative period. *Acta Radiol.* 32: 166-169.
- [63] Griffin JF, McNicholas MM (1992): Morphological appearance of renal allografts in transplant failure. *J Clin Ultrasound.* 20: 529-537.
- [64] Jansen O, Rob PM, Schmidtke V et al. (1992): Follow-up study of renal transplants by duplex Doppler and grey scale ultrasound. *Eur J Radiol.* 15: 26-31.
- [65] Renowden SA, Blethyn J, Cochlin DL (1992): Renal transplant sonography: correlation of Doppler and biopsy results in cellular rejection. *Clin Radiol.* 46: 265-269.
- [66] Phillips AO, Deane C, O'Donnell P et al. (1994): Evaluation of Doppler ultrasound in primary non-function of renal transplants. *Clin Transplant.* 8: 83-86.
- [67] Rigsby CM, Burns PM, Weltin GG et al. (1987): Doppler signal quantitation in renal allografts: comparison in normal and rejecting transplants, with pathologic correlation. *Radiology.* 162: 39-42.

- [68] Mallek R, Mostbeck G, Kain R et al. (1990): Vaskuläre Nierentransplantatabstoßung – ist eine duplexsonographische Diagnose möglich? RÖFo. 152: 283.
- [69] Hollenbeck M, Stuhmann M, Trapp R et al. (1994): Farbkodierte Dopplersonographie zur Früherkennung von Abstoßungsreaktionen nach allogener Nierentransplantation. Dtsch med Wochenschr. 116: 921-927.
- [70] Pallardó LM, Sánchez J, Puig N et al. (1995): Chronic rejection in 500 kidney transplant patients treated with cyclosporine: incidence and risk factors. Transplant Proc. 27: 2215-2216.
- [71] Basilio R, Blomley MJ, Harvey CJ et al. (2002): Which continuous US scanning mode is optimal for the detection of vascularity in liver lesions when enhanced with a second generation contrast agent? Eur J Radiol. 41(3): 184-191.
- [72] Schlosser T, Pohl C, Veltmann C (2001): Feasibility of the flash-replenishment concept in renal tissue: which parameters affect the assessment of the contrast replenishment? Ultrasound Med Biol. 27: 937-944.
- [73] Wei K, Le E, Bin JP et al. (2001): Quantification of renal blood flow with contrast-enhanced ultrasound. J Am Coll Cardiol. 37(4): 1135-1140.
- [74] Girard MS, Mattrey RF, Baker KG et al. (2000): Comparison of Standard and Second Harmonic B-mode Sonography in the Detection of Segmental Renal Infarction with Sonographic Contrast in a Rabbit Model. J Ultrasound Med. 19: 185-192.
- [75] Raisinghani A, DeMaria AN et al. (2002): Physical principles of microbubble ultrasound contrast agents. Am J Cardiol. 90(10A): 3J-7J.
- [76] Solbiati L, Martegani A, Leen E et al.: Contrast-Enhanced Ultrasound of Liver Diseases, Mailand 2003
- [77] Bokor D, Chambers JB, Rees PJ et al. (2001): Clinical safety of SonoVue, a new contrast agent for ultrasound imaging, in healthy volunteers and in patients with chronic obstructive pulmonary disease. Invest Radiol. 36(2): 104-109.
- [78] Calliada F, Campani R, Bottinelli O et al. (1998): Ultrasound contrast agents: basic principles. Eur J Radiol. 27(Suppl 2): 157-160.
- [79] Correias JM, Claudon M, Tranquart F et al. (2003): Contrast-enhanced ultrasonography: renal applications. J Radiol. 84(12Pt2): 2041-2054.

- [80] Lucidarme O, Franchi-Abella S, Correas JM et al. (2003): Blood flow quantification with contrast-enhanced US: "entrance in the section" phenomenon-phantom and rabbit study. *Radiology*. 228(2): 298-299.
- [81] Schlosser T, Veltmann C, Lohmaier S et al. (2004): Determination of the renal blood flow in macro- and microcirculation by means of pulse inversion imaging. *RöFo*. 176(5): 724-730.
- [82] Nahar T, Li P, Kuersten B et al. (2003): Detection of resting myocardial perfusion defects by SonoVue myocardial contrast echocardiography. *Echocardiography*. 20(6): 511-517.
- [83] Olszewski R, Timperley J, Cezary S et al. (2007): The clinical applications of contrast echocardiography. *Eur J Echocardiogr*. 8(3): 13-23.
- [84] Kang DH, Kang SJ, Song JM et al. (2005): Efficacy of myocardial contrast echocardiography in the diagnosis and risk stratification of acute coronary syndrome. *Am J Cardiol*. 96(11): 1498-1502.
- [85] Oldenburg A, Hohmann J, Foert E et al. (2005): Detection of hepatic metastases with low MI real time contrast enhanced sonography and SonoVue. *Ultraschall Med*. 26(4): 277-284.
- [86] Bryant TH, Blomley MJ, Albrecht T et al. (2004): Improved characterization of liver lesions with liver-phase uptake of liver-specific microbubbles: prospective multicenter study. *Radiology*. 232(3): 799-809.
- [87] Cassano E, Rizzo S, Bozzini A et al. (2006): Contrast enhanced ultrasound of breast cancer. *Cancer Imaging*. 6: 4-6.
- [88] Schroder RJ, Maurer J, Hidajat N et al. (1999): Diagnostic value of signal-enhanced color Doppler ultrasound in space-occupying lesions of the skin and skin appendages. *Rofo*. 171(3): 211-218.
- [89] Schmid-Wendtner MH, Partscht K, Korting HC et al. (2002): Improved differentiation of benign and malignant lymphadenopathy in patients with cutaneous melanoma by contrast-enhanced color Doppler sonography. *Arch Dermatol*. 138(4): 491-497.
- [90] Halpern EJ. (2006): Contrast-enhanced ultrasound imaging of prostate cancer. *Rev Urol*. 8(Suppl 1): 29-37.
- [91] Valentino M, Serra C, Zironi G (2006): Blunt abdominal trauma: emergency contrast-enhanced sonography for detection of solid organ injuries. *Am J Roentgenol*. 186(5): 1361-1367.

- [92] Poletti PA, Platon A, Becker CD (2004): Blunt abdominal trauma: does the use of a second-generation sonographic contrast agent help to detect solid organ injuries? *Am J Roentgenol.* 183(5): 1293-1301.
- [93] Siracusano S, Quaia E, Bertolotto M et al. (2004): The application of ultrasound contrast agents in the characterization of renal tumors. *World J Urol.* 22(5): 316-322.
- [94] Lacourciere Y, Levesque J, Onrot JM et al. (2002): Impact of Levovist ultrasonographic contrast agent on the diagnosis and management of hypertensive patients with suspected renal artery stenosis: a Canadian multicentre pilot study. *Can Assoc Radiol J.* 53(4): 219-227.
- [95] Nilsson A. (2004): Contrast-enhanced ultrasound of the kidneys. *Eur Radiol.* 14(Suppl8): 104-109.
- [96] Cosgrove D (2006): Ultrasound contrast agents: an overview. *Eur J Radiol.* 60(3): 324-330.
- [97] Ponticelli C (2004): Renal transplantation 2004: where do we stand today? *Nephrol Dial Transplant.* 19(12): 2937-47.
- [98] Terasaki PI, Gjertson DW, Cecka JM et al. (1997): Significance of the donor age effect on kidney transplants. *Clin Transplant.* 11(5 Pt 1): 366-72.
- [99] Dijkmans PA, Visser CA, Kamp O (2005): Adverse reactions to ultrasound contrast agents: is the risk worth the benefit? *Eur J Echocardiogr.* 6(5): 363-6.