

## 5. Literaturverzeichnis

- 1) ABBRACCHIO, M.P. & BURNSTOCK, G. (1994). Purinoceptors: are there families of P2X and P2Y purinoceptors? *Pharmacol Ther*, **64**, 445-75.
- 2) ALDERMANN, M.H. (1994). Non-pharmacological treatment of hypertension. *Lancet*, **344**, 307-311
- 3) BEILIN, L.J. & PUDEY, I.B. (1992). Alcohol and hypertension. *Clin Exp Hypertens [A]*, **14**, 119-38.
- 4) BIANCHI, G., FOX, U., DI FRANCESCO, G.F., GIOVANETTI, A.M. & PAGETTI, D. (1974). Blood pressure changes produced by kidney cross-transplantation between spontaneously hypertensive rats and normotensive rats. *Clin Sci Mol Med*, **47**, 435-48.
- 5) BO, X., ZHANG, Y., NASSAR, M., BURNSTOCK, G. & SCHOEPFER, R. (1995). A P2X purinoceptor cDNA conferring a novel pharmacological profile. *FEBS Lett*, **375**, 129-33.
- 6) BO, X., SCHOEPFER, R. & BURNSTOCK, G. (2000). Molecular cloning and characterization of a novel ATP P2X receptor subtype from embryonic chick skeletal muscle. *J Biol Chem*, **275**, 14401-7.
- 7) BOGDANOV, Y., RUBINO, A. & BURNSTOCK, G. (1998). Characterisation of subtypes of the P2X and P2Y families of ATP receptors in the foetal human heart. *Life Sci*, **62**, 697-703.
- 8) BURNSTOCK, G. (1972). Purinergic nerves. *Pharmacol Rev*, **24**, 509-81.
- 9) BURNSTOCK, G. (1989). Vascular control by purines with emphasis on the coronary system. *Eur Heart J*, **10 Suppl F**, 15-21.
- 10) BURNSTOCK, G., COCKS, T. & CROWE, R. (1978). Evidence for purinergic innervation of the anococcygeus muscle. *Br J Pharmacol*, **64**, 13-20.

- 11) CHAN, C.M., UNWIN, R.J., BARDINI, M., OGLESBY, I.B., FORD, A.P., TOWNSEND-NICHOLSON, A. & BURNSTOCK, G. (1998). Localization of P2X<sub>1</sub> purinoceptors by autoradiography and immunohistochemistry in rat kidneys. *Am J Physiol*, **274**, F799-804.
- 12) CINKILIC, O., KING, B.F., VAN DER GIET, M., SCHLÜTER, H., ZIDEK, W. & BURNSTOCK, G. (2001). Selective agonism of group I P2X receptors by dinucleotides dependent on a single adenine moiety. *J Pharmacol Exp Ther*. 2001; **299**:131-6.
- 13) CLINKINGBEARD, C., SESSIONS, C. & SHENKER, Y. (1990). The physiological role of atrial natriuretic hormone in the regulation of aldosterone and salt and water metabolism. *J Clin Endocrinol Metab*, **70**, 582-9.
- 14) COCKCROFT, S. & GOMPERTS, B.D. (1979). ATP induces nucleotide permeability in rat mast cells. *Nature*, **279**, 541-2.
- 15) CURTIS, J.J., LUKE, R.G., DUSTAN, H.P., KASHGARIAN, M., WHELCHL, J.D., JONES, P. & DIETHELM, A.G. (1983). Remission of essential hypertension after renal transplantation. *N Engl J Med*, **309**, 1009-15.
- 16) DAHL, L.K. & HEINE, M. (1975). Primary role of renal homografts in setting chronic blood pressure levels in rats. *Circ Res*, **36**, 692-6.
- 17) DANIELSON, M. & DAMMSTROM, B. (1981). The prevalence of secondary and curable hypertension. *Acta Med Scand*, **209**, 451-5.
- 18) DAVIES, G., MACALLISTER, R.J., BOGLE, R.G. & VALLANCE, P. (1995). Effect of diadenosine phosphates on human umbilical vessels: novel platelet-derived vasoconstrictors. *Br J Clin Pharmacol*, **40**, 170-2.
- 19) Drury, A.N. and Szent-Gyorgyi, A. (1929). The physiological of adenine compounds with especial reference to their action upon mammalian heart. *J. Physiol. (London)* 68, 213-237

- 20) EVANS, R.J., SURPRENANT, A. & NORTH, R.A. (1998): P2X receptors. In: Turner, J.T., Weisman, G.A. & Fedan, J.S. (Hrsg.). The P2 nucleotide receptor. Totowa, New Jersey: Humana Press; 43-61.
- 21) FALKNER, B. (1991). Blood pressure response to mental stress. *Am J Hypertens*, **4**, 621S-623S.
- 22) HARDEN, T.K., NICHOLAS, R.A., SCHACHTER, J.B., LAZAROWSKI, E.R. & BOYER, J.L. (1998): Pharmacological selectivities of molecularly defined subtypes of P2Y receptors. In: Turner, J.T., Weisman, G.A. & Fedan, J.S. (Hrsg.). The P2 nucleotide receptor. Totowa, New Jersey: Humana Press; 109-134.
- 23) HEIDENREICH, S., TEPEL, M., SCHLUTER, H., HARRACH, B. & ZIDEK, W. (1995). Regulation of rat mesangial cell growth by diadenosine phosphates. *J Clin Invest*, **95**, 2862-7.
- 24) HERRMANN, J.M. (1986). Essentielle Hypertonie. In: Psychosomatische Medizin. Uexküll, Th. (Hrsg.). 3. Auflage, München: Urban und Schwarzenberg Verlag; 715-742.
- 25) HUMPHREY, P.P.A., KHAKH, B.S., KENNEDY, C., KING, B.F., & BURNSTOCK, G. (1998). Nucleotide receptors: P2X receptors. In: IUPHAR Compendium of Receptor Characterization and Classification. IUPHAR Media Publications, pp 198-208.
- 26) INSCHO, E.W., COOK, A.K., MUI, V. & MILLER, J. (1998). Direct assessment of renal microvascular responses to P2-purinoceptor agonists. *Am J Physiol*, **274**, F718-27.
- 27) KANNEL, W.B. (1975). Role of blood pressure in cardiovascular disease: the Framingham Study. *Angiology*, **26**, 1-14.
- 28) KATSUYA, T., KOIKE, G., YEE, T.W., SHARPE, N., JACKSON, R., NORTON, R., HORIUCHI, M., PRATT, R.E., DZAU, V.J. & MACMAHON, S. (1995). Association of angiotensinogen gene T235 variant with increased risk of coronary heart disease. *Lancet*, **345**, 1600-3.

- 29) KAWABE, K., WATANABE, T.X., SHIONO, K. & SOKABE, H. (1979). Influence of blood pressure on renal isografts between spontaneously hypertensive and normotensive rats. *Jap Heart Journal*, **20**, 886-894.
- 30) KHAKH, B.S., SURPRENANT, A. & HUMPHREY, P.P. (1995). A study on P2X purinoceptors mediating the electrophysiological and contractile effects of purine nucleotides in rat vas deferens. *Br J Pharmacol*, **115**, 177-85.
- 31) KHAKH, B.S., BURNSTOCK, G., Kennedy, C., KING, B.F., NORTH, R.A., SEGUELA, P., VOIGT, M., HUMPHREY, P.P. (2001). International union of pharmacology. XXIV. Current status of the nomenclature and properties of P2X receptors and their subunits. *Pharmacol Rev.*, **53**, 107-18.
- 32) KING, B.F., LIU, M., PINTOR, J., GUALIX, J., MIRAS-PORTUGAL, M.T. & BURNSTOCK, G. (1999). Diinosine pentaphosphate (IP5I) is a potent antagonist at recombinant rat P2X1 receptors. *Br J Pharmacol*, **128**, 981-8.
- 33) LANDSBERG, L. (1992). Obesity and hypertension: experimental data. *J Hypertens Suppl*, **10**, S195-201.
- 34) LE, K.T., BABINSKI, K. & SEGUELA, P. (1998). Central P2X4 and P2X6 channel subunits coassemble into a novel heteromeric ATP receptor. *J Neurosci*, **18**, 7152-9.
- 35) LEWIS, C., NEIDHART, S., HOLY, C., NORTH, R.A., BUELL, G. & SURPRENANT, A. (1995). Coexpression of P2X2 and P2X3 receptor subunits can account for ATP-gated currents in sensory neurons. *Nature*, **377**, 432-5.
- 36) LEWIS, C.J. & EVANS, R.J. (2000a). Comparison of P2X receptors in rat mesenteric, basilar and septal (coronary) arteries. *J Auton Nerv Syst*, **81**, 69-74.
- 37) LEWIS, C.J., GITTERMAN, D.P., SCHLUTER, H. & EVANS, R.J. (2000b). Effects of diadenosine polyphosphates (Ap(n)As) and adenosine polyphospho guanosines (Ap(n)Gs) on rat mesenteric artery P2X receptor ion channels. *Br J Pharmacol*, **129**, 124-30.

- 38) LIEBERMANN, E. (1990). Hypertension in childhood and adolescence. In: Kaplan, N.M. (Hrsg.). Clinical Hypertension, 5.Auflage, Baltimore: Williams and Wilkins; 407-433.
- 39) LUDWIG, E., CORNELI, P.S., ANDERSON, J.L., MARSHALL, H.W., LALOUEL, J.M. & WARD, R.H. (1995). Angiotensin-converting enzyme gene polymorphism is associated with myocardial infarction but not with development of coronary stenosis [see comments]. *Circulation*, **91**, 2120-4.
- 40) MAGRINI, F. & REGGIANI, P. (1992). Epidemiology of hypertension in Europe. *Eur Heart J*, **13 Suppl H**, 27-34.
- 41) MANGER, W.M., PAGE, H.I. (1986). Zur Pathogenese und Pathophysiologie der essentiellen Hypertonie. In. Rosenthal, J. (Hrsg.). Arterielle Hypertonie, 3. Auflage, Berlin: Springer Verlag; 3-49.
- 42) MATTU, R.K., NEEDHAM, E.W., GALTON, D.J., FRANGOS, E., CLARK, A.J. & CAULFIELD, M. (1995). A DNA variant at the angiotensin-converting enzyme gene locus associates with coronary artery disease in the Caerphilly Heart Study [see comments]. *Circulation*, **91**, 270-4.
- 43) NAKAI, K., SYOH, T. & HIRAMORI, K. (1995). Angiotensin-I converting enzyme. *Nippon Rinsho*, **53**, 1232-6.
- 44) NORI, S., FUMAGALLI, L., BO, X., BOGDANOV, Y. & BURNSTOCK, G. (1998). Coexpression of mRNAs for P2X1, P2X2 and P2X4 receptors in rat vascular smooth muscle: an in situ hybridization and RT-PCR study. *J Vasc Res*, **35**, 179-85.
- 45) RAFF, T., VAN DER GIET, M., ENDEMANN, D., WIEDERHOLT, T. & PAUL, M. (1997). Design and testing of beta-actin primers for RT-PCR that do not co-amplify processed pseudogenes. *Biotechniques*, **23**, 456-60.
- 46) RALEVIC, V. & BURNSTOCK, G. (1996). Discrimination by PPADS between endothelial P2Y- and P2U-purinoceptors in the rat isolated mesenteric arterial bed. *Br J Pharmacol*, **118**, 428-34.

- 47) RALEVIC, V., HOYLE, C.H. & BURNSTOCK, G. (1995). Pivotal role of phosphate chain length in vasoconstrictor versus vasodilator actions of adenine dinucleotides in rat mesenteric arteries. *J Physiol (Lond)*, **483**, 703-13.
- 48) RIEDE, U.N. & SCHÄFER, H.E. (1995). Lokalisierte Kreislaufstörungen. In: Riede/Schäfer (Hrsg.). Allgemeine und spezielle Pathologie, 4. Auflage, Stuttgart: Georg Thieme Verlag; 418-429.
- 49) RUDNICK, K.V., SACKETT, D.L., HIRST, S. & HOLMES, C. (1977). Hypertension in a family practice. *Can Med Assoc J*, **117**, 492-7.
- 50) SCHLÜTER, H., GROBETA, I., BACHMANN, J., KAUFMANN, R., VAN DER GIET, M., TEPEL, M., NOFER, J.R., ASSMANN, G., KARAS, M., JANKOWSKI, J. & ZIDEK, W. (1998). Adenosine(5') oligophospho-(5') guanosines and guanosine(5') oligophospho-(5') guanosines in human platelets. *J Clin Invest*, **101**, 682-8.
- 51) SCHLÜTER, H., OFFERS, E., BRUGGEMANN, G., VAN DER GIET, M., TEPEL, M., NORDHOFF, E., KARAS, M., SPIEKER, C., WITZEL, H. & ZIDEK, W. (1994). Diadenosine phosphates and the physiological control of blood pressure. *Nature*, **367**, 186-8.
- 52) SINCLAIR, A.M., ISLES, C.G., BROWN, I., CAMERON, H., MURRAY, G.D. & ROBERTSON, J.W. (1987). Secondary hypertension in a blood pressure clinic. *Arch Intern Med*, **147**, 1289-93.
- 53) SINGER, D.R., MISSOURIS, C.G. & JEFFERY, S. (1996). Angiotensin-converting enzyme gene polymorphism. What to do about all the confusion. *Circulation*, **94**, 236-9.
- 54) SNEDDON, P., WESTFALL, T.D., TODOROV, L.D., TODOROVA, S.M., WESTFALL, D.P., NICKEL, P. & KENNEDY, C. (2000). The effect of P2 receptor antagonists and ATPase inhibition on sympathetic purinergic neurotransmission in the guinea-pig isolated vas deferens. *Br J Pharmacol*, **129**, 1089-94.

- 55) SORENSEN, S.S., DANIELSEN, H., AMDISEN, A. & PEDERSEN, E.B. (1989). Atrial natriuretic peptide and exaggerated natriuresis during acute hypertonic volume expansion in essential hypertension. *J Hypertens*, **7**, 21-9.
- 56) SOTO, F., LAMBRECHT, G., NICKEL, P., STUHMER, W. & BUSCH, A.E. (1999). Antagonistic properties of the suramin analogue NF023 at heterologously expressed P2X receptors. *Neuropharmacology*, **38**, 141-9.
- 57) STIEBER, J., DÖRING, A., KEIL, U. (1982). Häufigkeit, Bekanntheits- und Behandlungsgrad der Hypertonie in einer Großstadtbevölkerung. *Münchener Medizinische Wochenschrift*. **124**: 747.
- 58) SURPRENANT, A., RASSENDREN, F., KAWASHIMA, E., NORTH, R.A. & BUELL, G. (1996a). The cytolytic P2Z receptor for extracellular ATP identified as a P2X receptor (P2X7). *Science*, **272**, 735-8.
- 59) TORRES, G.E., EGAN, T.M. & VOIGT, M.M. (1999). Hetero-oligomeric assembly of P2X receptor subunits. Specificities exist with regard to possible partners. *J Biol Chem*, **274**, 6653-9.
- 60) TSUDA, M., UENO, S. & INOUE, K. (1999). In vivo pathway of thermal hyperalgesia by intrathecal administration of alpha,beta-methylene ATP in mouse spinal cord: involvement of the glutamate-NMDA receptor system. *Br J Pharmacol*, **127**, 449-56.
- 61) VAN DER GIET, M., CINKILIC, O., JANKOWSKI, J., TEPEL, M., ZIDEK, W. & SCHLUTER, H. (1999). Evidence for two different P2X-receptors mediating vasoconstriction of Ap5A and Ap6A in the isolated perfused rat kidney. *Br J Pharmacol*, **127**, 1463-9.
- 62) VAN DER GIET, M., KHATTAB, M., BORGEL, J., SCHLUTER, H. & ZIDEK, W. (1997). Differential effects of diadenosine phosphates on purinoceptors in the rat isolated perfused kidney. *Br J Pharmacol*, **120**, 1453-60.

- 63) VAN DER GIET, M., WESTHOFF, T., CINKILIC, O., JANKOWSKI, J., SCHLUTER, H., ZIDEK, W. & TEPEL, M. (2001). The critical role of adenosine and guanosine in the affinity of dinucleoside polyphosphates to P(2X)-receptors in the isolated perfused rat kidney. *Br J Pharmacol*, **132**, 467-74.
- 64) VIAL, C., HECHLER, B., LEON, C., CAZENAVE, J.P. & GACHET, C. (1997). Presence of P2X1 purinoceptors in human platelets and megakaryoblastic cell lines. *Thromb Haemost*, **78**, 1500-4.
- 65) VIRGINIO, C., Robertson, G., Surprenant, A., North, R.A., (1998). Trinitrophenyl-substituted nucleotides are potent antagonists selective for P2X1, P2X3, and heteromeric P2X2/3 receptors. *Mol Pharmacol*, **53**, 969-973.
- 66) VULCHANOV, L., RIEDL, M.S., SHUSTER, S.J., BUELL, G., SURPRENANT, A., NORTH, R.A. & ELDE, R. (1997). Immunohistochemical study of the P2X2 and P2X3 receptor subunits in rat and monkey sensory neurons and their central terminals. *Neuropharmacology*, **36**, 1229-42.
- 67) WILDMAN, S.S., BROWN, S.G., KING, B.F. & BURNSTOCK, G. (1999). Selectivity of diadenosine polyphosphates for rat P2X receptor subunits. *Eur J Pharmacol*, **367**, 119-23.
- 68) YAMAMOTO, K., KORENAGA, R., KAMIYA, A., QI, Z., SOKABE, M. & ANDO, J. (2000). P2X(4) receptors mediate ATP-induced calcium influx in human vascular endothelial cells. *Am J Physiol Heart Circ Physiol*, **279**, H285-92.
- 69) ZIDEK, W., HECKMANN, U., LANGE-ASSCHENFELDT, H., LOSSE, H. & VETTER, H. (1985). Humoral factors in the pathogenesis of primary hypertension. *Klin Wochenschr*, **63**, 94-6.
- 70) ZIDEK, W., HECKMANN, U., LOSSE, H. & VETTER, H. (1986). Effects on blood pressure of cross circulation between spontaneously hypertensive and normotensive rats. *Clin Exp Hypertens*, **8**, 347-54.