

6. Sonderdrucke der eigenen Arbeiten:

Hohenberger P, Allenberg JR, Schlag PM, **Reichardt P** [Impact Faktor 3.941]
Results of surgery and multimodal therapy for patients with soft tissue sarcoma
invading to vascular structures
Cancer 85: 396-408, 1999

Reichardt P, Oechsle K, Pink D, Bokemeyer C, Schneller F, Issels R, Kanz L,
Hartmann J T [Impact Faktor 3.82]
An open label, non-comparative phase II study of topotecan as salvage treatment
for patients with soft tissue sarcoma
Invest New Drugs 21: 481-486, 2003

Reichardt P, Tilgner J, Hohenberger P, Dörken B [Impact Faktor 9.868]
Dose-intensive chemotherapy with ifosfamide, epirubicin, and filgrastim for adult
patients with metastatic or locally advanced soft tissue sarcoma: a phase II study
J Clin Oncol 16: 1438-1443, 1998

Reichardt P, Verweij J, Crowther D [Impact Faktor 3.562]
Should high-dose chemotherapy be used in the treatment of soft tissue
sarcoma?
Eur J Cancer 33: 1351-1360, 1997

Reichardt P [Impact Faktor 2.758]
High-dose chemotherapy in adult soft tissue sarcoma
Crit Rev Oncol Hematol 41: 157-167, 2002

Reichardt P, Pink D, Tilgner J, Kretschmar A, Thuss-Patience P C, Dörken B
[Impact Faktor 0.855]
Oral Trofosfamide: An active and well tolerated maintenance therapy for adult
patients with advanced bone and soft tissue sarcomas. Results of a retrospective
analysis
Onkologie 25: 541-546, 2002

Reichardt P, Lindner T, Pink D, Thuss-Patience P C, Kretschmar A, Dörken B
[Impact Faktor 3.562]
Chemotherapy in alveolar soft part sarcoma: What do we know?
Eur J Cancer 39: 1511-1516, 2003

Reichardt P, Schneider U, Stroszcynski C, Pink D, Hohenberger P [Impact
Faktor 2.549]
Histological response pattern of gastrointestinal stromal tumor following
treatment with tyrosine kinase inhibitor imatinib mesylate
J Clin Path 57: 215-217, 2004

Reichardt P, Pink D, Mrozek A, Lindner T, Hohenberger P [Impact Faktor 0.837]
Gastrointestinale Stromatumoren (GIST)
Z Gastroenterol 2004 (in press)