

6. Literaturverzeichnis

- Anderson, R. M. & May, R. M. (1981): The population dynamics of microparasites and their invertebrate hosts. *Philosophical Transactions of the Royal Society of London*: B291, 451–524.
- Anderson, R. M. (1982): Theoretical basis for the use of pathogens as biological control agents of pest species. *Parasitology*: 84, 3–33.
- Anderson, R. M. & May, R. M. (1992): Infectious Diseases of Humans. Dynamics and Control. Oxford University Press, Oxford, New York, Tokyo.
- Andreadis, T. G. & Hall, D. W. (1979): Significance of transovarial infections of *Amblyospora* sp. (Microspora: Thelohaniidae) in relation to parasite maintenance in the *Culex salinarius*. *Journal of Invertebrate Pathology*: 34, 152–157.
- Andreadis, T. G. (1981): Impact of *Nosema pyrausta* on field populations of *Macrocentrus grandii*, an introduced parasite of the European corn borer, *Ostrinia nubilalis*. *Journal of Invertebrate Pathology* 39, 298–302.
- Alger, N. E. & Undeen, A. H. (1970): The control of a microsporidian, *Nosema* sp., in an anopheline colony by an egg rinsing technique. *Journal of Invertebrate Pathology*: 15, 321–327.
- Bathon, H. (1996): Zur Bedeutung der Parasitoide beim Zusammenbruch der Schwammspinner-Gradation in Südhessen. *Mitteilungen aus der Biologischen Bundesanstalt für Land- und Forstwirtschaft*: 322, 74–90.
- Bauer, L. S. & Nordin, O. L. (1989): Effect of *Nosema fumiferanae* (Microsporidia) on fecundity, fertility, and progeny performance of *Choristoneura fumiferana* (Lepidoptera: Tortricidae). *Environmental Entomology*: 18, 261–264.
- Becnel, J. J. & Andreadis, T. G. (1999): Microsporidia in Insects. In: The Microsporidia and Microsporidiosis (Wittner, M. & Weiss, L. M, Eds.). ASM Press, Washington, D.C., 447-501.
- Begon, M.; Harper, J. L. & Townsend, C. R. (1996): Ecology. 3rd ed., Blackwell Science, Oxford.
- Bell, R. A.; Owens, C. D.; Shapiro, M. & Tardiff, J. R. (1981): Mass rearing and virus production. In: The Gypsy Moth: Research toward Integrated Pest Management (C. C: Doane & M. L. McManus, Eds.), USDA Washington. [U.S. Dept. Agric. For. Serv. Tech. Bull. 1584], 599–600.
- Berryman, A. A.; Stenseth, N. Chr. & Isaev, A. S. (1987): Natural regulation of herbivorous forest insect populations. *Oecologia*: 71, 174–184.

- Bordat, D.; Goudegnon, A. E. & Bouix, G. (1993): Influence of *Nosema bordati*, (Microsporidia: Nosematidae) on the biological cycle of *Chilo partellus* (Lep: Pyralidae), a stem borer of cultivated gramineae. *Entomophaga*: 38, 551–564.
- Briggs, C. J. & Godfray, H. C. (1995): Models of intermediate complexity in insect-pathogen interactions: population dynamics of the microsporidian pathogen, *Nosema pyrausta*, of the European corn borer, *Ostrinia nubilalis*. *Parasitology*: 111, 71–89.
- Brooks, W. M. (1988): Entomogenous protozoa. In: CRC Handbook of Natural Pesticides: Volume V, Microbial Insecticides, Part A, Entomogenous Protozoa and Fungi. (C. M. Ignoffo, Eds.), CRC Press, Boca Raton, Florida, 1–149.
- Brown, G. C. & Nordin, G. L. (1982): An epizootic model of an insect-fungal pathogen system. *Bulletin of Mathematical Biology*: 44, 731–739.
- Brown, G. C. (1984): Stability in an epizoological model with age-dependent immunity and seasonal host reproduction. *Bulletin of Mathematical Biology*: 46, 139–153.
- Campbell, R. W. (1967): The analysis of numerical change in gypsy moth populations. *Forest Science Monographs*: 15, 1–33.
- Campbell, R. W. (1975): The gypsy moth and its natural enemies. *United States Department of Agriculture Bulletin*: 381.
- Campbell, R. W. (1981): Population dynamics. In: The Gypsy Moth: Research toward Integrated Pest Management (C. C. Doane & M. L. McManus, Eds.), USDA Washington. [U.S. Dept. Agric. For. Serv. Tech. Bull. 1584], 65–66.
- Canning, E. U. & Hulls, R. H. (1970): A microsporidian infection of *Anopheles gambiae* Giles, from Tanzania: interpretation of its mode of transmission and notes on *Nosema* infection in mosquitoes. *Journal of Protozoology*: 17, 71–81.
- Canning, E. U. (1982): An evaluation of protozoal characteristics in relation to biological control of pests. *Parasitology*: 84, 119–149.
- Dettner, K. & Zwölfer, H. (1999): Biologische, chemische und biotechnische Schädlingsbekämpfung. In: Lehrbuch der Entomologie (Peters, W. & Dettner, K., Hrsg.). Gustav Fischer, Stuttgart, 1–52.
- Diss, A. L.; Kunkel, J. G.; Montgomery & Leonard, D. E. (1996): Effects of maternal nutrition and egg provisioning on parameters of larval hatch, survival and dispersal in the gypsy moth, *Lymantria dispar* L. *Oecologia*: 106, 470–477.
- Dwyer, G. (1991): The roles of density, stage, and patchiness in the transmission of an insect virus. *Ecology*: 72, 559–574.

- Dwyer, G. (1994): Density dependence and spatial structure in the dynamics of insect pathogens. *The American Naturalist*: 143, 533–562.
- Dwyer, G.; Elkington, J. S. & Bounacorsi, J. P. (1997): Host heterogeneity in susceptibility and disease dynamics: tests of a mathematical model. *The American Naturalist*: 150, 685–707.
- Dwyer, G.; Dushoff, J.; Elkington, J. S. & Levin, S. A. (2000): Pathogen-driven outbreaks in forest defoliators revisited: building models from experimental data. *The American Naturalist*: 156, 105–120.
- Erelli, M. C. & Elkington, J. S. (2000): Maternal effects on gypsy moth (Lepidoptera: Lymantriidae) population dynamics: a field experiment. *Environmental Entomology*: 29, 476–488.
- Federici, B. A. (1993): Viral pathobiology in relation to insect control. In: Parasites and Pathogens of Insects 2: Pathogens (Beckage, N. E.; Thompson, S. N. & Federici, B. A., Eds.). Academic Press, San Diego, 8–101.
- Fine, P. E. M. (1975): Vectors and vertical transmission: an epidemiologic perspective. *Annals of the New York Academy of Sciences*: 266, 173–194.
- Forbush, E. H. & Fernald, C. H. (1896): The Gypsy Moth. Wright and Potter, Boston.
- Franz, J. M. & Huger, A. M. (1971): Microsporidia causing the collapse of an outbreak of the green tortrix (*Tortrix viridana* L.) in Germany. Proceedings of the International Colloquium of the Insect Pathology 4th, College Park, Md. 48–55.
- Fuxa, J. R.; Ayyappath, R. & Goyer, R. A. (1998): Pathogens and Microbial Control of North America Forest Insect Pest. Forest Health Enterprise Team FHTET-97-27.
- Goertz, D.; Pilarska, D.; Linde, A. (2002): Comparative Studies on the Impact of two Microsporidian Isolates from Bulgaria on the Gypsy Moth, *Lymantria dispar* L. USDA General Technical Report NE-300, 31.
- Graser, E. (1996): Clusteranalysen mit RAPD-PCR-Mustern: Eine Möglichkeit zur Rassendifferenzierung beim Schwammspinner. *Mitteilungen aus der Biologischen Bundesanstalt für Land- und Forstwirtschaft*: 322, 117–125.
- Hajek, A. E. (1996): *Entomophaga maimaiga*: A fungal pathogen of gypsy moth in the limelight. Proceedings of the Cornell Community Conference on Biological Control, Internet: <http://www.nysaes.cornell.edu/ent/bccconf/Talks/hajek.html>, 2001.
- Heinrich, G. (1957): Fibel der histologischen Technik. 2. Aufl., VEB Gustav Fischer Verlag, Jena.

- Henn, W. M. & Solter, L. F. (2000): Food utilization values of gypsy moth *Lymantria dispar* (Lepidoptera: Lymantriidae) larvae infected with a microsporidium *Vairimorpha* sp. (Microsporidia: Burenellida). *Journal of Invertebrate Pathology*: 76, 263-269.
- Heynen, C. (1983): Untersuchungen zum Einfluss des Chitinsynthesehemmers Dimilin auf zwei Wirt-Parasit-Systeme. Dissertation, Georg-August-Universität Göttingen.
- Hill, R. E. & Gary, W. J. (1979): Effects of the microsporidium, *Nosema pyrausta*, on field populations of European corn borer in Nebraska. *Journal of Environmental Entomology*: 8, 91-95.
- Hoch, G. (1998): Wechselwirkungen zwischen einer entomopathogenen Mikrosporidie und dem Endoparasiten *Glyptapanteles liparidis* in ihrem gemeinsamen Wirt, der *Lymantria dispar* Larve. Dissertation. Universität für Bodenkultur Wien.
- Hoch, G.; Zubrik, M.; Novotny, J. & Schopf, A. (2001): The natural enemy complex of the gypsy moth, *Lymantria dispar* (Lep., Lymantriidae) in different phases of its population dynamics in eastern Austria and Slovakia – a comparative study. *Journal of Applied Entomology*: 125, 217–227.
- Hochberg, M. E. (1989): The potential role of pathogens in biological control. *Nature*: 337, 262–265.
- Iwano, H. & Ishihara, R. (1991): Dimorphism of spores of *Nosema* spp. in cultured cell. *Journal of Invertebrate Pathology*: 57, 211-219.
- Jones, C. G.; Ostfeld, R. S., Richard, M. P.; Schauber, E. M. & Wolff, J. O. (1998): Chain reactions linking acorns to gypsy moth outbreaks and lyme disease risk, *Science*: 279, 1023–1026.
- Jeffords, M. R., Maddox, J. V., O'Hayer, K. W. (1987): Microsporidian spores in gypsy moth larval silk: a possible route of horizontal transmission. *Journal of Invertebrate Pathology*: 49, 332–333.
- Jeffords, M. R.; Maddox, J. V.; McManus, M.L.; Webb, R. E. & Wieber, A. (1988): Egg contamination as a method for the inoculative release of exotic microsporidia of the gypsy moth. *Journal of Invertebrate Pathology*: 51, 190–196.
- Jeffords, M. R.; Maddox, J. V.; McManus, M.L.; Webb, R. E. & Wieber, A. (1989): Evaluation of the overwintering success of two European microsporidia inoculatively released into gypsy moth populations in Maryland. *Journal of Invertebrate Pathology*: 53, 253–240.
- Kellen, W. R. & Lindegreen, J.E. (1971): Modes of transmission of *Nosema plodiae* Kellen and Lindegreen, a pathogen of *Plodia interpunctella* (Hübner). *Journal of Stored Product Research*: 7, 31–34.

- Kellen, W. R. & Lindegreen, J.E. (1973): Transovarian transmission of *Nosema plodiae* in the Indian meal moth *Plodia interpunctella*. *Journal of Invertebrate Pathology*: 21, 248–254.
- Kramer, J. P. (1959): Some relationships between *Perezia pyraustae* Paillot (Sporozoa, Nosematidae) and *Pyrausta nubilalis* (Hübner) (Lepidoptera, Pyralidae). *Journal of Insect Pathology*: 1, 25–33.
- Krieg, A. & Franz, J. M. (1989): Lehrbuch der biologischen Schädlingsbekämpfung. 4. Aufl., Paul Parey, Berlin, Hamburg.
- Leonard, D. E. (1967): Silking behavior of the gypsy moth, *Lymantria dispar*. *Canadian Entomology*: 99, 1145–1149.
- Leonard, D. E. (1968): Effects of density of larvae on the biology of gypsy moth, *Lymantria dispar*. *Entomologia Experimentalis et Applicata*: 11, 291–304.
- Leonard, D. E. (1970): Effects of starvation on behavior, number of larval instars and developmental rate of *Lymantria dispar*. *Journal of Insect Physiology*: 16, 25–31.
- Leonard, D. E. (1981): Bioecology of the gypsy moth. In: The Gypsy Moth: Research toward Integrated Pest Management (C. C. Doane & M. L. McManus, Eds.), USDA Washington. [U.S. Dept. Agric. For. Serv. Tech. Bull. 1584], 9–29.
- Lewis, L. C. & Lynch, R. E. (1978): Foliar application of *Nosema pyrausta* for suppression of populations of European corn borer. *Entomophaga*: 23, 83–88.
- Liebold, A.; Elkington, J.; Williams, D. & Muzika, R.-M. (2000): What causes outbreaks of the gypsy moth in North America? *Population Ecology*: 42, 257–266.
- Linde, A. (1993): Können Mikrosporidien (Protozoa) zur Kontrolle des Schwammspinnens (*Lymantria dispar* L.) beitragen? *Mitteilungen aus der Biologischen Bundesanstalt für Land- und Forstwirtschaft*: 293, 125–129.
- Linde, A.; Genthe, C.; Lacker, J. (1998): The effects of artificial infections with microsporidia on gypsy moth (*Lymantria dispar*) and nun moth (*Lymantria monacha*) - First results. Proceedings: Population Dynamics, Impacts, and Integrated Management of forest Defoliating Insects" (M. L. McManus & A. M. Liebold, Eds.). *USDA Forest Service General Technical Report, NE - 247*, 198–205.
- Linde, A.; Golldack, J. & Goertz, D. (2000): Evaluation of the potential of a microsporidian isolate for the biological control of *Lymantria dispar* L. *IOBC wprs Bulletin*: 23, 285–290.
- Lipa, J. J. & Madziara-Borusiewicz, K. (1976): Microsporidian parasitizing the green tortrix (*Tortrix viridana* L.) in Poland and their role in the collapse of the tortrix outbreak in Puszczza Niepolomicka during 1970 – 1974. *Acta Protozoologica*: 31, 529–536.

- Long, D. W.; Groden, E.; Drummond, F. A. (2000a): Horizontal transmission of *Beauveria bassiana* (Bals.) Vuill. *Agricultural and Forest Entomology*: 2, 11–17.
- Long, D. W.; Drummond, F. A.; Groden, E. & Donahue, D. W. (2000b): Modelling *Beauveria bassiana* horizontal transmission. *Agricultural and Forest Entomology*: 2, 19–32.
- Maddox, J. M.; McManus, M. L. & Solter, L. F. (1998): Microsporidia affecting forest lepidoptera. Proceedings: Population Dynamics, Impacts, and Integrated Management of Forest Defoliating Insects (M. L. McManus & A. M. Liebold, Eds.). *USDA Forest Service General Technical Report, NE - 247*, 198–205.
- Maddox, J. V., Baker, M. D., Jeffords, M. R., Kuras, M., Linde, A., Solter, L. F., McManus, M. L., Vávra, J., Vossbrinck, C. R. (1999): *Nosema Portugal*, n. sp., isolated from gypsy moth (*Lymantria dispar* L.) collected in Portugal. *Journal of Invertebrate Pathology*: 73, 1–14.
- McManus, M. L.; Maddox, J. V.; Jeffords, M. R. & Webb, R. E. (1988): Evaluation and selection of candidate microsporidia for introduction into U.S. gypsy moth populations. Proceedings: Lymantriidae: A comparison of features of New and Old World tussock moths (Wallner, W. E. & McManus, K. A., Eds.). *USDA Forest Service General Technical Report, NE-123*, 455–468.
- Mitchell, M. J. & Cali, A. (1994): *Vairimorpha necatrix* (Microsporidia: Burenellidae) affect growth and development of *Heliothis zea* (Lepidoptera: Noctuidae) raced at various temperatures, *Journal of Invertebrate Pathology*: 87, 933–940.
- Norten, E.; Linde, A.; Happe, M. & Madel, G. (1988): Infektion und Pathologie einer Mikrosporidiose von *Diadegma semiclausum* HELLEN (Hym., Ich.). *Mitteilungen der deutschen Gesellschaft für allgemeine und angewandte Entomologie*: 6, 284–285.
- Norten, E.; Linde, A. & Strick, H. (1989): Zur Diagnostik entomopathogener Mikrosporidien. *Mitteilungen der deutschen Gesellschaft für allgemeine und angewandte Entomologie*: 7, 169–170.
- Ogwang, J. A. & Odindo, M. O. (1993): Vertical transmission of a microsporidium *Nosema marucae* in the cereal stem borer *Chilo partellus*. *Insect Science and its Application*: 14, 489–482.
- Onstad, D. W. & Maddox, J. V. (1988): Dealing the effects of the microsporidium, *Nosema pyrausta*, on the population dynamics of the insect, *Ostrinia nubilalis*. *Journal of Invertebrate Pathology*: 53, 410–421.
- Onstad, D. W. & Carruthers, R. I. (1990): Epizoological models of insect diseases. *Annual Review of Entomology*: 35, 399–419.

- Onstad, D. W. & Maddox, J. V. (1990): Simulation model of *Tribolium confusum* and its pathogen, *Nosema whithei*. *Ecological Modelling*: 51, 143–160.
- Peters, W. (1999): Integument. In: Lehrbuch der Entomologie (Peters, W. & Dettner, K., Hrsg.). Gustav Fischer, Stuttgart, 1–52.
- Pierce, C. M. F.; Solter, L. F.; Weinzierl, R. A. (2001): Interactions between *Nosema pyrausta* (Microsporidia: Nosematidae) and *Bacillus thuringiensis* subsp. *kurstaki* in the European corn borer (Lepidoptera: Pyralidae). *Journal of Economic Entomology*: 94, 1361–1386.
- Reardon, (1981): Alternative Controls. In: The Gypsy Moth: Research toward Integrated Pest Management (C. C: Doane & M. L. McManus, Eds.), USDA Washington. [U.S. Dept. Agric. For. Serv. Tech. Bull. 1584], 299–421.
- Regnieri, J. (1984): Vertical transmission of disease and population dynamics of insects with discrete generations: a model. *Journal of Theoretical Biology*: 107, 287–301.
- Richardson, K. C.; Jarrett, L. & Finke, E. H. (1960): Embedding in epoxy resins for ultrathin sectioning in electron microscopy. *Stain Technology*: 35, 313.
- Rozkhov, A. S. & Vasilyeva, T. G. (1982): Gypsy moth in Middle and Eastern Siberia. In: Neparny Shelkoprayo v Srednai i Vostochnoi Sibiri. Russian Academy of Sciences, Siberian Branch, Novosibirsk, 4–19.
- Ruf, A. & Römbke, J. (1996): Erste Ergebnisse einer Untersuchung zur Wirkung von Dimilin und B.t.k. auf Bodenfauna und Streuabbau. *Mitteilungen aus der Biologischen Bundesanstalt für Land- und Forstwirtschaft*: 322, 175–187.
- Sachs, L. (1997): Angewandte Statistik. 8. Aufl., Springer Verlag, Berlin, Heidelberg.
- Schaaf, C. & Vogt, H. (1996): Nebenwirkungen von Dimilin bei Bekämpfungsmaßnahmen gegen den Schwammspinner, *Lymantria dispar* L. *Mitteilungen aus der Biologischen Bundesanstalt für Land- und Forstwirtschaft*: 322, 173–174.
- Siegel, J. P., Maddox, J. V., Ruesink, W. G. (1986): Lethal and sublethal effects of *Nosema pyrausta* on the European corn borer (*Ostrinia nubilalis*) in Central Illinois. *Journal of Invertebrate Pathology*: 48, 167–173.
- Skatulla, U (1975a): Successful trial for controlling *Lymantria dispar* and *L. monacha* by help of PH 60-40, prohibiting the molting of insect larvae. *Anzeiger für Schädlingskunde, Pflanzenschutz und Umweltschutz*: 48, 17–18.
- Skatulla, U (1975b): Effect of Dimilin, a new selective insecticide on forest insects. *Anzeiger für Schädlingskunde, Pflanzenschutz und Umweltschutz*: 48, 145–147.

- Solter, L. F. & Becnel, J. J. (2000): Entomopathogenic microsporidia. In Field Manuscript of Techniques for the Evaluation of Entomopathogens (Lacey, L. & Kaya, H., Eds.). Kluwer Academic Publishers, Dordrecht.
- Solter, L. F.; Siegel, J. P.; Pilarska, D. K. Higgs, M. C. (2002): The impact of mixed infection of three species of microsporidia isolated from the gypsy moth, *Lymantria dispar* L. (Lepidoptera: Lymantriidae), *Journal of Invertebrate Pathology*: 81, 103–113.
- SPSS für WINDOWS. Release 11.0.1. 15.11.2001. SPSS Inc. 1989–2001.
- Steinhaus, E. A. & Martignoni, M. E. (1970): An Abridged Glossary of Terms used in Invertebrate Pathology. 2nd ed., Pacific Northwest Forest and Range Experiment Station, USDA, Forest Service, Washington DC.
- Tanada, Y. & Kaya, H. K. (1992): Insect Pathology. Academic Press, San Diego.
- Thomson, H. M. (1958): Some aspects of the epidemiology of a microsporidian parasite of the spruce budworm, *Choristoneura fumiferanae* (Clem.). *Canadian Journal of Zoology*: 36, 309–316.
- Thomas, M. B.; Wood, S. N. & Lomer, C. J. (1995): Biological control of locusts and grasshoppers using a fungal pathogen: the importance of secondary cycle. *Proceedings of the Royal Society of London*: B259, S. 265–270.
- Vavra, J. & Larsson, J. I. R. (1999): Structure of the microsporidia. In: The Microsporidia and Microsporidiosis (Wittner, M. & Weiss, L. M, Eds.). ASM Press, Washington, D.C., 7–84.
- Waldbauer, G. P. (1968): The consumption and utilization of food by insects. *Advances in Insect Physiology*: 5, 229–288.
- Wallner, W. E. (1988): An overview of pest lymantriids of North America. In: Proceedings Lymantriidae: A Comparison of Features of New and Old World Tussock Moths USDA Forest Service General Technical Report, NE - 123, 65–80.
- Weiser, J. (1957): Mikrosporidien des Schwammspinners und Goldafters. *Zeitschrift für angewandte Entomologie*: 40, 509–527.
- Weiser, J. (1958): Transovariale Übertragung der *Nosema otiorrhynchi* W. *Vestnik Ceskoslovenske Spolecnosti Zoologicke*: 22, 10–12.
- Weiser, J. (1961): Die Mikrosporidien als Parasiten der Insekten. *Monografien zur angewandten Entomologie. Beihefte zur Zeitschrift für angewandte Entomologie*: 17, 1–149.
- Weiser, J. & Novotny, J. (1993): Transovarial transmission of *Nosema lymantriae* (Protozoa, Microsporidia) in the gypsy moth *Lymantria dispar* L. *Biologia (Bratislava)*: 48, 125–129.
- Weiser (1998): Pathogens of the gypsy moth in Central Europe: host range and interactions. In: Proceedings: Population Dynamics, Impacts, and Integrated Management of forest

- Defoliating Insects (M. L. McManus & A. M. Liebhold, Eds.). *USDA Forest Service General Technical Report, NE - 247*, 198–205.
- Weiser, J. & Linde, A. (1998): Microsporidian early spores as confusing factor in taxonomy: *Nosema muscularis* vs. *N. lymantriae*. *Abstracts of the VIth European Congress of Entomology*, Ceské Budějovice, 571.
- Weseloh, R. M. & Andreadis, T. A. (1992a): Mechanisms of transmission of the gypsy moth (Lepidoptera: Lymantriidae) fungus, *Entomophaga maimaiga* (Entomophthorales: Entomophthoraceae) and effects of site conditions on its prevalence. *Journal of Environmental Entomology*: 21, 901–906.
- Weseloh, R. M. & Andreadis, T. A. (1992b): Epizootiology of the fungus *Entomophaga maimaiga*, and its impact on gypsy moth populations. *Journal of Invertebrate Pathology*: 59, 133–141.
- Wilson, G. G. (1973): Incidence of microsporidia in a field population of spruce budworm. *Bi-monthly Research Notes*: 29, 35–36.
- Wilson, G. G. (1977): Observations of the incidence rates of *Nosema fumiferanae* in a spruce budworm, *Choristoneura fumiferana* population. *Proceedings of the Entomological Society Ont.*: 108, 144–145.
- Wilson, G. G. (1978): Microsporidian infection in spruce budworm (*Choristoneura fumiferana*) 1 and 2 years after application. *Bi-monthly Research Notes*: 34, 1–16.
- Wilson, G. G. (1981): *Nosema fumiferana*, a natural pathogen of a forest pest: potential for pest management. In: *Microbial Control of Pests and Plant Diseases 1970–1980* (H. D. Burges, Eds.), Academic Press Inc., New York, 595–601.
- Wittner, M. (1999): Historic perspective on the microsporidia: expanding horizons. In: *The Microsporidia and Microsporidiosis* (Wittner, M. & Weiss, L. M, Eds.). Washington, D.C., 1–6.
- Wulf, A. (1993): Schwammspinner-Massenvermehrung 1993 – Resümee und Ausblick. *Mitteilungen aus der Biologischen Bundesanstalt für Land- und Forstwirtschaft*: 293, 278–282.
- Zelinskaya, L. M. (1980): The role of microsporidia in the population dynamics of *Porthetria dispar* L. in plantations of Lower Pridneprovije. *Vestnik zoologii*: 1, 57–62.
- Zelinskaya, L. M. (1981): The use of parasitation index of gypsy moth by microsporidia spores for forecasting gypsy moth outbreaks. *Lesnoje Khozyaystvo*: 14, 58–60.
- Zub, P.; Nässig, W. A. & Kristal, P. M. (1996): Lepidopterologische Begleituntersuchung zur Bekämpfung einer Gradation des Schwammspinnners (*Lymantria dispar*) mit *Bacillus*

thuringiensis kurstaki (B.t.k.) und dem Häutungshemmer Dimilin im Jahr 1994 im Staatsforst bei Lampertheim. *Mitteilungen aus der Biologischen Bundesanstalt für Land- und Forstwirtschaft*: 322, 188–195.