

6. Literatur:

- Amerena J, Julius S. The role of the automatic nervous system in hypertension. *Hypertens Res* 1995; 18: 99-110
- Anderson J, Rodier HE, et al. Comparative effects of beta-adrenergic blocking drugs on experimental ventricular fibrillation threshold. *Am J Cardiol* 1983; 51: 1196-1202
- Andresen D, Tietze U, von Leitner ER: Spontanvariabilität tachykarder Rhythmusstörungen. *Z Kardiol* 69 (1980): 214
- Aronow WS, Epstein S, Koenigsberg M. Usefulness of echocardiographic left ventricular hypertrophy and silent ischaemia in predicting new cardiac events in elderly patients with systemic hypertension or coronary artery disease. *Angiology* 1990; 3: 189-193
- Asmar RG, Pannier B, Santoni JP, Laurent S, London GM, et al. Reversion of cardiac hypertrophy and reduced arterial compliance after converting enzyme inhibition in essential hypertension. *Circulation* 1988; 78: 941-50
- Aurigemma GP, Reichek NR, Venogopal R, et al. Automated left ventricular mass, volume, and shape from 3-dimensional magnetic resonance imaging: in vitro validation. *Am J Cardiac Imaging* 1991; 5: 257-263
- Brown J, Brown MJ. CAPPP trial. Captopril prevention project. *Lancet* 1999 May 22;353(9166):1795-6
- Bugaisky L, Zak R. Biological mechanism of hypertrophy. In: Fozzard H.A, Jennings R.B., Haber H., Katz A.M.(Hrsg.): *The heart and cardiovascular system*. Raven Press, New York 1986, S. 1491-1506
- Caputo GR, Tscholakoff D, Sechtem U, Higgins CB. Measurement of canine left ventricular mass by using MR imaging. *Am J Roentgenol* 1987; 148: 33-38
- Carr AA, Prisant LM, Watkins LO. Detection of hypertensive left ventricular hypertrophy. *Hypertension* 1985; 7: 948-954
- Casale PN, Devereux RB, Alonso DR, Campo E, Kligfield P. Improved sex-specific criteria of left ventricular hypertrophy for clinical and computer interpretation of electrocardiograms: Validation with autopsy findings. *Circulation* 1987; 75: 565-72
- Chanutin A, Barksdale E. Experimental renal insufficiency produced by partial nephrectomy. *Arch Intern Med* 52 (1933), 739-751
- Cooper RS, Simmons BE, Castener A, Santhanam V, Ghali J, Mar M. Left ventricular hypertrophy is associated with worse survival independent of ventricular function and number of coronary arteries severely narrowed. *Am J Cardiol* 1990; 65: 441-445
- Cruickshank JM, Lewis J, Moore V, et al. Reversibility of left ventricular hypertrophy by different types of antihypertensive therapy. *J Hum Hypertens* 1992; 6: 85-90
- Dahlöf B, Pennert K, Hanson L. Reversal of left ventricular hypertrophy in hypertensive patients. A metaanalysis of 109 treatment studies. *Am J Hypertens* 1992; 5: 95-110
- Dalton GR, Jones JV, Evans SJ, Levi AJ. Wall stress-induced arrhythmias in the working rat heart as left ventricular hypertrophy regresses during captopril treatment. *Cardiovasc Res* 1997 Mar; 33 (3): 561-72

- Devereux RB, Reichek N. Echocardiographic determination of left ventricular mass in man: Anatomic validation of the method. *Circulation* 1977; 55: 613-618
- Devereux RB, Casale PN, Wallerson DC, et al. Costeffectiveness of echocardiography for detection of left ventricular hypertrophy in patients with systemic hypertension. *Hypertension* 1987; 9 (Suppl II): II-69-II-76
- Dunn FG, Oigman W, Ventura HO, Messerli FH, Kobrin I, et al. Enalapril improves systemic and renal haemodynamics and allows regression of left ventricular mass in essential hypertension. *Am J Cardio* 1984; 53 : 105-8
- Dunn FG, Ventura HO, Messerli FH, Kobrin I, Frohlich ED. Time course of regression of left ventricular hypertrophy in hypertensive patients treated with atenolol. *Circulation* 1989; 76 (2): 254-58
- Eichstädt H. Welche Bedeutung hat die Echokardiographie zur Beurteilung der linksventrikulären Hypertrophie ? *Verh Dtsch Ges Kreislaufforschung* 1977, 573
- Eichstädt H, Krämer R, Gutmann M, et al. Regression einer linksventrikulären Hypertrophie unter chronischer Betablockade mit Penbutolol. *Herz-Kreislauf* (1985a);Bd 17, Nr 7, 356-362
- Eichstädt H, Steiner G, Lange M, Schultze E, Felix R (1985b) Eine einfache Methode zur Abbildung von Herzwandkontrakturstörungen mittels MRT. *Fortschr Röntgenstr* 143:186-189
- Eichstädt H, Felix R, Langer M, et al. Use of nuclear magnetic resonance imaging to show regression of hypertrophy with ramipril treatment. *Am J Cardiol* 1987; 59: 98D-103D
- Eichstädt H, Mertens D, Schröder R, et al. Linksventrikuläre Hypertrophieregression unter Lisinopril-Therapie bei Patienten mit leichter bis mittelschwerer Hypertonie. *Perfusion* 1991; 10/4; 338-342
- Eichstädt H, Gatz G, Schröder R, et al. Linksventrikuläre Hypertrophieregression unter einer Therapie mit Moxonidin. *Journal Pharmacol u Ther* 1/1992a: 20-26
- Eichstädt H, Gatz G, Schröder R, et al. Induktion einer linksventrikulären Hypertrophieregression unter der Therapie mit Moxonidin. *Perfusion* 2/1992b:8-12
- Eichstädt H, Moersler J, Klefisch F, et al. Regression der linksventrikulären Myokardhypertrophie unter Spirapril-Therapie, belegt durch magnetresonanztomographische Wanddickenmessungen. *Perfusion* 9/1996: 338-343
- Eichstädt H, Moersler J, Möckel M, et al. Regression der linksventrikulären Hypertrophie durch Gallopamil retard bei Patienten mit leichter bis mittelschwerer Hypertonie. *Herz/Kreisl* 2/1997; 29: 51-55
- Florentine MS, Grosskreut CL, Chang W, et al. Measurement of left ventricular mass in vivo using gated nuclear magnetic resonance imaging. *J Am Coll Cardiol* 1986; 8:107-112
- Foster E, Johnson D, Francesco B, Backwell G, Orr R, Roney M, Stanley A, Pohost G.: Changes in left ventricular mass and volumes in patients receiving angiotensin-converting enzyme inhibitor therapy for left ventricular dysfunction after Q-Wave myocardial infarction. *Am Heart J* 1998;136:269-275
- Fouad FM, Nakashima Y, Tarazi RC, Salcedo EE. Reversal of left ventricular hypertrophy in hypertensive patients treated with methyldopa: lack of association with blood pressure control. *Am J Cardio* 1982; 49: 795-801

- Fouad FM, Tarazi RL, Liebson PR. Echocardiographic studies of regression of left ventricular hypertrophy in hypertension. *Hypertension* 1987; 9 (Suppl II):II65-II68
- Frohlich ED. Cardiac hypertrophy in hypertension. *New Engl J Med* 1987; 317: 831-33
- Frohlich ED, Pfeffer MA. Heart and hypertension: The magnitude of the problem. In: Safar M.E., Fouad-Tarazi F.M. (Hrsg): *The heart in hypertension*. Kluwer Academic Publishers, Dordrecht-Boston-London 1989, S.169
- Garavaglia GE, Messerli FH, Nunez BD, Schmieder RE, Frohlich ED. Angiotensin converting enzyme inhibitors: disparities in the mechanism of their antihypertensive effect. *Am J Hypertens* 1988; 1: 214-6
- Gatzoulis KA, Vyssoulis GP, Apostolopoulos T, et al. Mild left ventricular hypertrophy in essential hypertension: Is it really arrhythmogenic? *Am J Hypertens* 2000; 13: 340-345
- Georgiou D. Regression of ventricular mass in systemic hypertension. *Clin Cardiol* 1992, Jan 15 (1), 5-16
- Genovesi-Ebert F. Left ventricular filling: relationship with arterial blood pressure, left ventricular mass, age, heart rate and body build. *J Hypertens* 1991, Apr 9(4), 345-353
- Gerstenblith. Influence of left ventricular mass regression on cardiac function in hypertensive elderly individuals. *Eur J Clin Pharmacol* 1990, 39, 25-28
- Ghali JK, Youlian L, Simmons B, Castener A, Guichan C, Cooper RS. The prognostic role of left ventricular hypertrophy with or without coronary artery disease. *Ann Intern Med* 1992; 117: 831-6
- Gottdiener JS. Left ventricular mass, diastolic dysfunction and hypertension. *Adv Intern Med* 1993; 38: 31-56
- Grandi AM, Venco A, Barizza F, Casadei B, Marchesi E, et al. Effect of Enalapril on left ventricular mass and performance in essential hypertension. *Am J Cardiol* 1989; 63: 1093-97
- Grossman W, Jones D, Mc Laurin LP. Wall stress and patterns of hypertrophy in the human left ventricle. *J Clin Invest* 1975; 56: 56-64
- Hammond IW, Devereux RB, Alderman MH, Lutas EM, Spitzer MC, Crowley JS, Laragh JH. The prevalence and correlates of echocardiographic left ventricular hypertrophy among employed patients with uncomplicated hypertension. *J Am Coll Cardiol* 7 (1986), 639-650
- Hammond IW, Devereux RB, Alterman MH, et al. The prevalence and correlates of echocardiographic left ventricular hypertrophy among employed patients with uncomplicated hypertension. *J Am Coll Cardiol* 1987; 7: 639-650
- Harrison DG, Marcus ML, Dellperger KC, et al. Pathophysiology of myocardial perfusion in hypertension. *Circulation* 1991; 83 (Suppl III): III-14-III-15
- Hoffmann U, Globits S, Steffenelli T, Loewe C, Kostner K, Frank H. The effects of ACE Inhibitor Therapy on left ventricular myocardial mass and diastolic filling in previously untreated hypertensive patients: A cine mri study. *J Magn Reson Imaging* 2001; 14: 16-22
- Houghton JL, Carr AA, Prisant LM, et al. Morphologic, haemodynamic and coronary perfusion characteristics in severe left ventricular hypertrophy secondary to systemic hypertension and evidence for nonatherosclerotic myocardial ischemia. *Am J Cardiol* 1992; 69: 219-224

- James MA, Jones JV. Systolic wall stress and ventricular arrhythmia: the role of acute change in blood pressure in the isolated working rat heart. *Clin Sci* 1990; 79: 499-504
- Julien J, Dufloux MA, Prasquier R, Chatellier G, Menard D, et al. Effects of captopril and minoxidil on left ventricular hypertrophy in resistant hypertensive patients: a 6 month double blind comparsion. *Journal of the American Collage of Cardiology* 1990; 16: 137-42
- Just H, Holubarsch C, Friedburg H. Estimation of left ventricular volume and mass by magnetic resonance imaging: Comparsion with quantitative biplane angiocardiology. *Cardiovasc Intervent Radiol* 10 (1987): 1-4
- Kalusche D. Plötzlicher Herztod, Prävention und Therapie. In: Roskamm H, Reindell H (Hrsg.): Herzkrankheiten: Pathophysiologie, Diagnostik und Therapie, 718-725, Springer Verlag, Wien/ New York 1989
- Katz J, Milleken MC, Stray-Gunderson J, Buja LM, Parkey RW, Mitchell IH, Peshock RM. Estimation of human cardial mass with MRI. *Radiology* 169 (1988a) 495-498
- Keller AU, Peshock RM, Malloy CR, Buja LM, Nunally R, Parkey RW, Willerson JT. In vivo measurment of myocardial mass using nuclear magnetic resonance imaging. *JACC* 8 (1986) 113-117
- Kennedy HL, Chandra V, Sayther KL, Caralis DG: Effectiveness of increasing hours of continious ambulatory electrocardiography in detecting maximal ventricular ectopy. *Am J Cardiol* 42 (1978) 925-930
- Komajda M, Klimeczak K, Boutin B, Brackman F, Guez D, et al. Effects of Indapamide on left ventricular mass and function in systemic hypertension with left ventricular hypertrophy. *Am J Cardiol* 1990; 65: 37H- 42H
- Koren MJ, Devereux RB, Casale P, Savage DD, Laragh JH. Relation of left ventricular mass and geometry to morbidity in uncomplicated essential hypertension. *Ann Inter Med* 1991; 114: 345-52
- Lavie CJ, Ventura HO, Messerli FH. Regression of increased left ventriculat mass by anti-hypertensives. *Drugs* 1991; 42 (6): 945-61
- Le Heuzey J, Guize L, et al: Cardiac prognosis in hypertensive patients. Incidence of sudden death and ventricular arrhythmias. *Am J Med* 1988; Suppl 1B: 65-68
- Leitner ER, Schröder RR. Das Langzeit-EKG beim Herzgesunden. *DMW* 1983; 108: 523-526
- Leren P, Foss PO, Helgeland A, Hjermann I, Holme I, et al. Effect of propanolol and prazosin on blood lipids: the Oslo Study. *Lancet* 1980; 2:4-6
- Levy D, Anderson KM, Savage DD, Balkus SA, Kannel WB, Castelli WP. Risk of Ventricular arrhythmias in left ventricular hypertrophy: The Framingham heart study. *Am J Cardiol* 1987; 60: 560-5
- Levy D, Garrison RJ, Savage DD, Kannel WB, Castelli W. Left ventricular mass and incidence of coronary heart disease in an elderly cohort. *Ann Intern Med* 1989;110: 101-7
- Levy D, Garrison RJ, Savage DD, Kannel WB, Castell WP. Prognostic Implications of echocardiographically determined left ventricular mass in the Framingham heart study. *N Engl J Med* 1990; 322: 1561-6
- Levy D, Salomon M, Agostino D, et al. Prognostic implications of baseline electrocardiographic features and their serial changes in subjects with left ventricular hypertrophy. *Circulation* 1994; 90: 1786-1793

- Liebson PR. Clinical studies of drug reversal of hypertensive left ventricular hypertrophy. *Am J Hypertens* 1990; 3: 512-517
- Lievré M, Gueret P, Gayet C, Roudaut R, Haugh M, Delair S, Boissel JP. Ramipril-induced regression of left ventricular hypertrophy in treated hypertensive individuals. *Hypertension* 1995; 25: 92-97
- Lucarini AR, Picano E, Salvetti A. Coronary microvascular disease in hypertensives. *Clin Exp Hypertens* 1992; A 14: 55-66
- Lumme AJ, Jounela AJ. Left ventricular mass, serum electrolyte levels and cardiac arrhythmias in patients with mild hypertension treated with cilazapril or hydrochlorthiazide. *Int Jour Cardio* 1993; 42: 71-8
- Mace PJE, Littler WA, Glover DR, Rowlands DB, Stallard TJ. Regression of left ventricular hypertrophy in hypertension: comparative effects of three different drugs. *Journal of cardiovascular pharmacology* 1985; 7: 52-55
- Malerba M, Muisan M, Zulli R, Rizzoni D, et al. Ventricular arrhythmias in hypertensive patients with and without hypertrophy, effect of antihypertensive treatment. *J Am Coll Cardiol* 1991; 17: 177A
- Manolis AJ, Kolovou G, Handanis S, Athanasopoulos G, Antonopoulos S, Cokkinos S, Cokkinos DV. Regression of left ventricular hypertrophy with Israpidine in previously untreated hypertensive Patients. *Am J Hypertens* 1993; 6: 86-88
- Manolis A, Beldekos D, Handanis S, et al. Comparsion of Spirapril, Isradipine, or combination in hypertensive patients with left ventricular hypertrophy. *American Journal of Hypertens* 1998; 11: 640-648
- Marcus ML, Harrison DG, Chilian WM, Koyanagi S, Inon T, et al. *Circulation* 1987; 75 (Suppl I): 19-25
- Maron BJ, Savage DD, Wolfson JK, Epstein SE. Prognostic significance of 24-hour ambulatory electrocardiographic monitoring in patients with hypertrophic cardiomyopathy: a prospectiv study. *Amer J Cardiol* 1981; 48: 252- 57
- Mathew J, Sleight P, Lonn E, Johnstone D, Pogue J, Yi Q, Bosch J, Sussex B, Probstfield J, Yusuf S:Reduction of cardiovascular risk by regression of electrocardiographic markers of left ventricular hypertrophy by the angiotensin-converting enzyme inhibitor ramipril. *Circulation* 2001; 104: 1615-1621
- Mc Lenahan JM, Henderson E, Morris KI, Dargie HJ. Ventricular arrhythmias in patients with hypertensive left ventricular hypertrophy. *N Engl J Med* 1987; 317: 787-92
- Mc Veigh GE. Effects of Perindopril on cardiovascular remodeling. *Am J Cardiol* 2001; 88 (Suppl): 28i-35i
- Messerli FH, Ventura HO, Elizardie DJ, Dunn FG, Frohlich ED. Hypertension and sudden death: increased ventricular ectopic activity in left ventricular hypertrophy. *Am J Med* 1984; 77: 18-22
- Messerli FH, Nunez BD, Ventura HO, Snyder DW. Oberweight and sudden death: Increased ventricular ectopy in cardiomyopathy of obesity. *Arch Intern Med* 1988a; 147: 1725-1728
- Messerli FH, Oren S, Grossman E. Left ventricular hypertrophy and antihypertensive therapy. *Drug* 1988b; 35 (Suppl.5): 27-33

- Messerli FH, Boris DN, Nunez BD, Mireya MN, Garavaglia GE, Schmieder RE, Ventura HO. Hypertension and sudden death: Disparate Effects of calcium entry blocker and diuretic therapy on cardiac dysrhythmias. *Arch Intern Med* 1989;149: 1263-7
- Messerli FH. Antihypertensive therapy: going to the heart of the matter. *Circulation* 1990; 81: 1128-35
- Nielsen F, Sato A, Tarnow L, Smidt U, Kastrup J, Parving H. Beneficial impact of ramipril on left ventricular hypertrophy in normotensive nonalbuminuric NIDDM patients. *Diabetes care*, Vol 21, nr 5, 5/1998
- Novo S, Abrignani M, Novo G, Nardi E, Dominguez L, Strano A, Babagallo M. Effect of drug therapy on cardiac arrhythmias and ischemia in hypertensives with LVH. *Am J Hypertens* 2001; 14: 637-643
- Pickering G. Systemic arterial hypertension. In: Fishman AP, Richards DW (Hrsg): *Circulation of blood: Men and ideas*. American Physiol. Society 1982, 487-541
- Pitt B, Reichek N, Willenbrock R, Zannad F, et al. Effects of Eplerenone, Enalapril and Eplerenone / Enalapril in patients with essential hypertension and left ventricular hypertrophy. *Circulation* 2003; 108: 1831-1838
- Raasch W, Bartels T, Schwartz C, Häuser W, Rütten H, Dominiak P. Regression of ventricular and vascular hypertrophy: are there differences between structurally different angiotensin-converting enzyme inhibitors? *Journal of hypertension* 2002; 20: 2495-2504
- Reichek N, Devereux RB. Left ventricular hypertrophy: Relationship of anatomic, echocardiographic and electrocardiographic findings. *Circulation* 1981; 63: 1391-98
- Rials SJ, Wu Y, Ford N, Paulette FJ, Abramson SV, Rubin AM, Marinchak RA, Kowey PR. Effect of left ventricular hypertrophy and its regression on ventricular electrophysiology and vulnerability to inducible arrhythmia in the feline heart. *Circulation* 1995; 91: 426-30
- Rials SJ, Xiaoping X, Ying W, Marinchak R, Kowey P. Regression of LVH with captopril normalises membrane currents in rabbits. *Am Phys Soc*; 1998: H1216
- Rodstein M, Wolloch L, Grubner RS. Mortality study of the significance of extrasystoles in an insured population. *Circulation* 1971; 44: 617-625
- Rorive GL, Carlier PG, Foidart JM. The structural responses of the vascular wall in experimental hypertension. Zanchetti A, Tarazi RC (Hrsg): *Handbook of hypertension*, vol 7: *Pathophysiologie of hypertension* 1986, 428-453
- Rberman W, Weinblatt E, Goldberg JD, Frank CW, Shapiro S. Ventricular premature beats and mortality after myocardial infarction. *New Engl J Med* 1977; 297: 750-758
- Saragoca MA, Cassiolatto JL, Vanetta AM, Canziani ME, Abreu P, Ventura RT, Ramos OL. Reversal of left ventricular hypertrophy with Isradipine induces diminution of cardiac dysrhythmias. *Am J Hypertens* 1993; 6: 89-91
- Savage DD, Drayer BJ, Henry WL, Mathews EC Jr, Ware JH, Gardin JM, et al. Echocardiographic assessment of cardiac anatomy and function in hypertensive subjects. *Circulation* 1979; 59: 623-32
- Schmieder RE, Rüddel H, Grube E, Schulte W. Depressed myocardial contractility in early left ventricular hypertrophy. *Circulation* 1988 (Suppl II): 75
- Schmieder RE, Nunez BD, Messerli FH. Determinants of left ventricular ectopy in hypertensive cardiac disease. *Amer Heart J* 1992; 123: 89-95

- Schmieder RE, Schlaich M, Klingbeil A, et al. Update on reversal of left ventricular hypertrophy in essential hypertension (a meta-analysis of all randomized double-blind studies until December 1996). *Nephrol Dial Transplant* 1998; 13: 564-9
- Schreiber SS, Evans CD, Oratz M, Rothschild MA. Protein synthesis and degradation in cardiac stress. *Circ Res* 48 (1981), 601-611
- Schulte KL, Meyer-Sabellek W, Liederwald K, van Gemmeren D, Lenz T, Gotzen R. Relation of regression of left ventricular hypertrophy to changes in ambulatory blood pressure after long term therapy with perindopril versus nifedipine. *Am J Cardiol* 1992; 70: 468-473
- Schuster H. *Langzeit-Elektrokardiographie*. Gustav Fischer Verlag 1990: 146-147
- Sen S. Alteration in myocardial phenotypes in SHR (abstract). *J Mol Cell Cardiol* 14 (1982), 60
- Shimada Y, Gunasegaram S, Yokohama H, Avkiran M. Inhibition of angiotensin-converting enzyme reduces susceptibility of hypertrophied rat myokardium to ventricular fibrillation. *Circulation* 2002; 11: 1045-53
- Sideris DA, Kontoyannis DA, Michali L, Adractas A, Moulopoulos SD. Acute changes in blood pressure as a cause of cardiac arrhythmias. *Europ Heart J* 1987; 8: 45-52
- Sideris DA, Toumanidis ST, Kostis EB, et al. Arrhythmogenic effect of high blood pressure: some observations on its mechanism. *Cardiovasc Res* 1989; 23: 983-992
- Simone G, Verdecchia P, Pede S, Gorini M, Maggioni AP. Prognosis of inappropriate left ventricular mass in hypertension: the MAVI Study. *Hypertension* 2002 Oct; 40 (4): 470-6
- Stamler R, Stamler J, Gosch FC, Berkson DM, Dyer A, et al. Initial antihypertensive drug therapy. Alpha-blocker or diuretic: interim report of a randomized controlled trial. *Am J Med* 1990; 80 (Suppl 2A): 90-93
- Strauer BE. Ventricular functions and coronary haemodynamics in hypertensive heart disease. *Amer J Cardiol* 1979; 44: 999-1006
- Tanaka M, Fujiwara H, Onodera T, et al. Quantitative analysis of myocardial fibrosis in normals, hypertensive hearts and hypertrophic cardiomyopathy. *Brit Heart J* 1986; 55: 575-581
- Tarazi RC, Sen S. Catecholamines and cardiac hypertrophy. *Catecholamines and the heart, No 8 (Royal Society of Medicine International Congress and Symposium Series)*. Academic Press and Royal Society of Medicine, London 1979
- Tarazi RC, Fouad FM. Reversal of cardiac hypertrophy in humans. *Hypertension* 1984; 6 (Suppl III): 140-46
- Toyoshima H, Park YD, Ishikawa W, et al. Effects of ventricular hypertrophy on conduction velocity of activation front in the ventricular myocardium. *Am J Cardiol* 1982; 49: 1938-1945
- Tubau JF, Szlachcic J, Braun S, Massie BM. Impaired left ventricular functional reserve in hypertensive patients with left ventricular hypertrophy. *Hypertension* 1989; 14: 1-8
- Umemura K, Zierhut W, Rudin M, Novosel D, Robertson E, Pedersen B, Hof RP. Effect of Spirapril on left ventricular hypertrophy due to volume overload in rats. *Journal of cardiovascular pharmacology* 1992; 19: 375-3

- Valdez AS, Motta JA, London E, et al. Evaluation of the echocardiogram as an epidemiologic tool in an asymptomatic population. *Circulation* 1979; 60: 921-929
- Ventura HO, Frohlich ED, Messerli FH, Kobrin I, Kardon BM. Cardiovascular effects and regional blood flow distribution associated with angiotensin converting enzyme inhibitor (captopril) in essential hypertension. *Am J Cardiol* 1985; 55: 1023-1026
- Verdecchia P, Angeli R, Gattobigio R, Guerrieri G, Porcellati C. Does the reduction in systolic blood pressure alone explain the regression of left ventricular hypertrophy? *J of hum hypertens* 2004/ 18: 23-28
- Wallerson DC, Devereux RB. Reproducibility of echocardiographic left ventricular measurements. *Hypertension* 1987; 9 (Suppl II): II-6-II-18
- Weber KT, Janicki JS, Pick R, et al. Collagen in the hypertrophied, pressure overloaded myocardium. *Circulation* 75 (1987), 140-147
- White CW, Mirro MJ, Lund DD, Skorton DJ, Pandian NG, Kerber RE. Alterations in ventricular excitability in conscious dogs during the development of chronic heart failure. *Amer J Physiol* 1986; 250: H 1022-1029
- Wong M, Shah PM, Taylor RD. Reproducibility of left ventricular internal dimensions with M-mode echocardiography: Effects of heart size, body position and transducer angulation. *Am J Cardiol* 1981; 47: 1068-1074
- Yokoshiki H, Tetsuro K, Fumishi T, Noritsugu T, Haruaki N, Kanno M, Kitabatake A. Restoration of action potential duration and transient outward current by regression of left ventricular hypertrophy. *J Mol Cell Cardiol* 1997; 29, 1331-1339