

## 8 Literaturverzeichnis<sup>2</sup>

- Affolter, V.K. und Moore, P.F. (2000). Canine cutaneous and systemic Histiocytosis: reactive Histiocytosis of dermal dendritic Cells. Am J Dermatopathol. 22 (1), 40-48.
- Ando, R., Ueki, H., Oyamada, T., Kurotaki, T., Arai, S. und Yoshikawa, H. (2003). Histopathological Examination of canine cutaneous Histiocytoma. J. Jpn. Vet. Med. Assoc. 56, 23-27.
- Baines, S.J., Bujdoso, R., Blacklaws, B.A., Mc Innes, E., Moore, P.F. und McConnel, I.M. (2000a). Maturation States of dendritic Cells in canine cutaneous Histiocytoma. Vet Dermatol. 11 (Suppl. 1), 9-10.
- Baines, S.J., Bujdoso, R., Blacklaws, B.A., Mc Innes, E., Moore, P.F. und Mcconnell, I.M. (2000b). Maturation States of dendritic Cells in canine cutaneous Histiocytoma. Vet. Dermatol. 11 (Suppl. 1), 9-10.
- Balkwill, F. (2002). Tumor necrosis Factor or Tumor promoting Factor? Cytokine Growth Factor Rev. 13, 135-141.
- Banchereau, J. und Steinman, R.M. (1998). Dendritic Cells and the Control of Immunity. Nature. 392, 245-252.
- Baumal, R., Kahn, H.J., Bailey, D., Phillips, M.J. und Hanna, W. (1984). The Value of Immunohistochemistry in increasing diagnostic Precision of undifferentiated Tumours by the surgical Pathologist. Histochemical Journal. 16, 1061-1078.
- Berek, C. und Ziegner, M. (1993). The Maturation of the immune Response. Immunol Today. 14, 400-404.
- Bingle, L., Brown, N.J. und Lewis, C.E. (2002). The Role of Macrophages in Tumor Progression: Implications for new Anticancer Therapies. J Pathol. 196, 254-265.
- Birbeck, M.S.C., Breathnach, A.S. und Everall, J.D. (1961). An Electron microscopic Study of basal Melanocytes and high Level clear Cells (Langerhans Cells) in Vitiligo. J Invest Dermatol. 37, 51-63.
- Boenisch, T. (1989). Handbuch II immunchemischer Färbemethoden. DAKO corporation: Hamburg. S. 11, 13, 22-27
- Bourne, J.A. (1983). Handbuch I der Immunperoxidase Färbrmethoden. Dako Diagnostika: Hamburg. S. 30-31
- Boyd, W. (1966). The spontaneous Regression of Cancer. Adv Immunol. 18, 67-137.

---

<sup>2</sup> Zeitschriften-Abkürzungen gemäß:

1. List of journals indexed in Index Medicus. Bethesda, Md.: Nat. Library of Medicine, 1989  
 2. Serial Sources for the BIOSIS Data Base. Philadelphia, Pa.: BioScience Information Services, 1988

- Broll, R., Mahlke, C., Best, R., Schimmelpenning, H., Strik, M.W., Schiedeck, T., Bruch, H.-P. und Duchrow, M. (1998). Assessment of the Proliferation Index in gastric Carcinomas with the monoclonal Antibody MIB1. *J Cancer Res Clin Oncol.* 124, 49-54.
- Brown, E.J. (1995). Phagocytosis. *Bioessays.* 17, 109-117.
- Bruno, S. und Darzynkiewicz, Z. (1992). Cell Cycle dependent Expression and Stability of the nuclear Protein detected by Ki-67 Antibody in HL-60 Cells. *Cell Prolif.* 25, 31-40.
- Bühl, A. und Zöfel, P. (1996). Professionelle Datenanalyse mit SPSS für Windows. Addison-Wesley: Bonn. S. 311
- Cangul, I.T., Teske, E. und Ingh, T., S, G. A. M. van den. (1998). Immunhistochemical staining Results of frozen and paraffin embedded Sections of 14 canine malignant Lymphoma Cases: a comparative Study. *Europ J of Vet Pathol.* 4 (1), 17-22.
- Cattoretti, G., Becker, M.H.G., Key, G., Duchrow, M., Schlüter, C., Galle, J. und Gerdes, J. (1992). Monoclonal Antibodies against recombinant Parts of the Ki-67 Antigen (MIB1 and MIB3) detect proliferating Cells in Microwave-processed Formalin-fixed Paraffin Sections. *J Pathol.* 168, 357-363.
- Cattoretti, G., Pileri, S.A., Parravicini, C., Becker, M.H.G., Poggi, S., Bifulco, C., Key, G., D'Amato, L., Sabattini, E., Feudale, E., Reynolds, F., Gerdes, J. und Rilke, F. (1993). Antigen unmasking on Formalin-fixed, Paraffin-embedded Tissue Sections. *J Pathol.* 171, 83-98.
- Chaiet, L. und Wolf, F.J. (1964). The Properties of Streptavidin, a Biotin-binding Protein by Streptomyces. *Arch Biochem Biophys.* 106, 1-5.
- Cobbold, S. und Metcalfe, S. (1994). Monoklonal Antibodies that define Canine Homologous of human CD Antigens: Summary of the First international Canine Leucocyte Antigen Workshop (CLAW). *Tissue Antigens.* 43, 137-154.
- Darbes, J., Majzoub, M. und Hermanns, W. (1997). Evaluation of the Cross-reactivity between Human and Feline or Canine Leucocyte Antigens using commercially available Antibodies. *J Vet Diagn Invest.* 9, 94-97.
- Dierendonck, J.H.v., Wijsman, J.H., Keijzer, R., Velde, C.J.H.v.d. und Cornelisse, C.J. (1991). Cell-cycle-related Staining Patterns of Anti-proliferating Cell Nuclear Antigen monoclonal Antibodies. *Am J Pathol.* 138, 1165-1172.
- Dobson, J.M., Samuel, S., Milstein, H., Rogers, K. und Wood, J.L.N. (2002). Canine Neoplasia in the UK: Estimates of incidence Rates from a Population of insured Dogs. *Journal of Small Animal Practice.* 43, 240-246.
- Dongen, J.J.M.v., Krissansen, G.W., Wolfers-Tettero, J.L.M., Comans-Bitter, W.M., Adri-aansen, H.J., Hooijkaas, H., Wering, E.R.v. und Terhorst, C. (1988). Cytoplasmic Expression of the CD3 Antigen as a diagnostic Marker for immature T cell Malignancies. *Blood.* 71, 603-612.

- Dongen, J.J.M.v., Quertermous, T., Bartram, C.R., Gold, D.P., Wolfers-Tettero, J.L.M., Comans-Bitter, W.M., Hooijkaas, H., Adriaansen, H.J., de Klein, A., Raghavachar, A., Ganser, A., Duby, A.D., Seidman, J.G., Van Den Elsen, P. und Terhorst, C. (1987). T Cell Receptor-CD3 Complex during early T Cell Differentiation: Analysis of immature T Cell acute lymphoblastic Leukemias (T-all) at DNA, RNA and Cell Membrane Level. *J Immunol.* 138, 1260-1265.
- Drommer, W. und Schulz, L.-C. (1969). Vergleichende licht- und elektronenmikroskopische Untersuchungen am übertragbaren venerischen Sarkom und Histiozytom des Hundes. *Path vet.* 6, 273-286.
- Duncan, J.R. und Prasse, K.W. (1979). Cytology of Canine cutaneous round Cell Tumours. Mast Cell Tumour, Histiocytoma, Lymphosarcoma and transmissible venereal Tumor. *Vet Pathol.* 16 (6), 673-9.
- Er, J. und Sutton, R. (1989). A Survey of Skin Neoplasms in dogs from the Brisbane Region. *Australian Veterinary Journal.* 66, 225-227.
- Falini, B., Flenghi, L., Fagioli, M., Stein, H., Schwarting, R., Riccardi, C., Manocchio, I., Pileri, S., Pelicci, P.-G. und Lanfrancone, L. (1989). Evolutionary Conservation in various Mammalian Species of the Human Proliferation-associated Epitope recognized by the Ki-67 monoclonal Antibody. *J Histochem Cytochem.* 37, 1471-1478.
- Falini, B. und Taylor, C.R. (1983). New Developments in Immunoperoxidase Techniques and their Applications. *Arch Pathol Lab Med.* 107, 105-117.
- Ferrer, L., Fondevila, D., Rabanal, R. und Ramis, A. (1992). Detection of T-Lymphocytes in Canine Tissue embedded in Paraffin Wax by means of Antibody to CD3 Antigen. *J Comp Path.* 106, 311-314.
- Finkelman, F.D., Lees, A. und Morris, S.C. (1992). Antigen Presentation by B Lymphocytes to CD4+ T Lymphocytes in vivo: Importance for B Lymphocyte und T Lymphocyte Activation. *Semin Immunol.* 4, 247-255.
- Flavell, D.J., Jones, D.B. und Wright, D.H. (1987). Identification of Tissue Histiocytes on Paraffin Sections by a New Monoclonal Antibody. *J Histochem Cytochem.* 35 (11), 1217-1226.
- Foss, F.M. (2002). Immunologic Mechanisms of Antitumor Activity. *Semin Oncol.* 29 (7), 5-11.
- Frese, K., Eskens, U. und Frank, H. (1982). Hauttumore des Hundes. In *28. Jahrestagung der Fachgruppe Kleintierkrankheiten.* Deutsche Veterinärmedizinische Gesellschaft e. v.: Frankfurt/M, S. 188-208
- Gerdes, J., Becker, M.H.G., Key, G. und Cattoretti, G. (1992). Letters to the Editor: Immunohistological Detection of Tumour Growth Fraction (Ki-67 antigen) in formalin-fixed and routinely processed Tissues. *J Pathol.* 168, 85-86.
- Gerdes, J., Lemke, H., Baisch, H., Wacker, H.H., Schwab, U. und Stein, H. (1984). Cell Cycle Analysis of a Cell Proliferation-associated human nuclear Antigen defined by the Monoclonal Antibody Ki-67. *J Immunol.* 133, 1710-1715.

- Gerdes, J., Li, L., Schlueter, C., Duchrow, M., Wohlenberg, C., Gerlach, C., Stahmer, I., Kloth, S., Brandt, E. und Flad, H.D. (1991). Immunobiochemical and Molecular Biologic Characterization of the Cell Proliferation-associated Nuclear Antigen that is defined by Monoclonal Antibody Ki-67. *American Journal of Pathology*. 138 (4), 867-873.
- Glick, A.D., Holscher, M. und Campbell, G.R. (1976). Canine cutaneous Histiocytoma: ultrastructural and cytochemical Observations. *Vet Pathol.* 13 (5), 374-80.
- Gonzales, C.M., Griffey, S.M., Naydan, D.K., Flores, E., Cepeda, R., Cattaneo, G. und Madewell, B.R. (2000). Canine Transmissible Venereal Tumour: a morphological and immunohistochemical Study of 11 Tumours in Growth Phase and during Regression after Chemotherapy. *J Comp Path.* 122, 241-248.
- Greene, J.F.J., Townsend, J.S.I. und Amoss, M.S.J. (1994). Histopathology of Regression in Sinclair Swine Model of Melanoma. *Lab Invest.* 71, 17-24.
- Griffey, S.M., Kraegel, S.A. und Madewell, B.R. (1999). Proliferation Indices in Spontaneous Canine Lung Cancer: Proliferating Cell Nuclear Antigen (PCNA), Ki-67 (MIB1) and Mitotic Counts. *J Comp Path.* 120, 321-332.
- Guvenc, T., Haligur, M., Orman, M.N. und Haziroglu, R. (2002). Mitosis and Apoptosis in Canine cutaneous Histiozytoma and Transmissible venereal Tumour. *Acta Veterinaria Hungarica.* 50 (3), 315-321.
- Hald, J., Rasmussen, N. und Claesson, M.H. (1995). Tumor-infiltrating Lymphocytes mediate Lysis of autologous squamous-cell Carcinomas of the Head and Neck. *Cancer Immunol Immunother.* 41, 243-250.
- Hall, P.A. und Woods, A.L. (1990). Invited Review: immunohistochemical Markers of cellular Proliferation: Achievements, Problems and Prospects. *Cell Tissue Kinet.* 23, 505-522.
- Hamada, M., Takechi, M. und Itakura, C. (1992). Langerhans' Cells in Eqine cutaneous Papillomas and normal Skin. *Vet Pathol.* 29, 152-160.
- Hart, D.N.J. (1997). Dendritic Cells: unique Leukocyte Populations which control the Primary Immune Response. *Blood.* 90, 3245-3287.
- Hombach, J., Lottspeich, F. und Reth, M. (1990). Identification of the Genes encoding the IgM-a Components of the IgM Antigen Receptor Complex by amino terminal Sequencing. *Eur J Immunol.* 20, 2795-2799.
- Howard, E.B. und Nielsen, S.W. (1969). Cutaneous histiocytomas of Dogs. *National Cancer Institute Monograph.* 32, 321-327.
- Hsiao, Y.W., Liao, K.W., Hung, S.W. und Chu, R.M. (2002). Effect of Tumor infiltrating Lymphocytes on the Expression of MHC Molecules in canine transmissible venereal Tumor Cells. *Vet Immunol Immunopathol.* 87, 19-27.
- Hsu, S.M., Raine, L. und Fanger, H. (1981). Use of Avidin-biotin-Peroxidase Complex (ABC) in immunoperoxidase Techniques: A Comparison between ABC and unlabeled Antibody (PAP) Procedures. *J Histochem Cytochem.* 29 (4), 577-580.

- Hung, K., Hayashi, R., Lafond-Walker, A., Lowenstein, C., Pardoll, D. und Levitsky, H. (1998). The central Role of CD4+ T Cells in the Antitumor Immune Response. *J Exp Med.* 188 (12), 2357-2368.
- Jenkins, D.C., Charles, I.G., Thomsen, L.L., Moss, D.W., Holmes, L.S., Baylis, S.A., Rhodes, P., Westmore, K., Emson, P.C. und Moncada, S. (1995). The Role of nitric Oxide in Tumor Growth. *Proc Nat Acad Sci USA* 92, 4392-4396.
- Johnston, R.B. (1988). Current Concepts: Immunology. Monocytes and Macrophages. *N Engl J Med.* 318, 747-752.
- Jones, M., Cordell, J.L., Beyers, A.D., Tse, A.G.D. und Mason, D.Y. (1993). Detection of T- and B-cells in many Animal Species using cross-reactive anti-peptide Antibodies. *J Immunol.* 150, 5429-5435.
- Jöstens, M. (1995): Differenzierung zwischen Geschwulst-Riesenzellen und osteoklastenähnlichen Riesenzellen in Neoplasien von Katzen und Hunden am immun- und enzymhistochemisch gefärbten Paraffinschnitt. Dissertation, Fachbereich Veterinärmedizin der Freien Universität Berlin.
- Jöstens, M., Rudolph, R., Stein, H. und Walter, J.H. (1993). Immun- and enzymhistochemische Differenzierung zwischen Geschwulst-Riesenzellen und osteoklastenähnlichen Riesenzellen in Neoplasien von Katze und Hund am Paraffinschnitt. Proceeding, 42. Jahrestagung der Europ. Ges. für Vet.-Pathol. (ESVP), Würzburg 01.06. - 02.06.1993.
- Kägi, D., Vignaux, F., Ledermann, B., Bürki, K., Depraetere, V., Nagata, S., Hengartner, H. und Golstein, P. (1994). Fas and Perforin Pathways as major Mechanisms of T cell-mediated Cytotoxicity. *Science.* 265, 528-530.
- Kaim, U. (2003): Die Bedeutung von Entzündungszellen und Zytokinen für die Regression des kaninen kutanen Histiozytoms. Dissertation, Justus-Liebig-Universität Gießen.
- Katz, S.I., Cooper, K.D., Masafumi, I. und Tetsuo, T. (1985). The Role of Langerhans Cells in Antigen Presentation. *J Invest Dermatol.* 85 (1), 96-98.
- Kelly, D.F. (1970). Canine cutaneous Histiocytoma. A Light and Electron microscopic Study. *Vet Pathol.* 7 (1), 12-27.
- Kelm, G. (1982): Licht- und elektronenmikroskopische Untersuchungen über die Morphologie und die formale Pathogenese des kutanen Histiozytoms des Hundes. Dissertation, Freie Universität Berlin.
- Key, G., Becker, M.H.G., Baron, B., Duchrow, M., Schlüter, C. und Flad, H.D. (1993). New Ki-67-equivalent Murine monoclonal Antibodies (MIB 1-3) generated against bacterially expressed Parts of the Ki-67 cDNA containing three 62 Base Pair repetitive Elements encoding for the Ki-67 Epitope. *Lab Invest.* 68, 629-636.
- Khalil, M.M., Nitiuthai, S. und Allen, J.R. (1982). Alkaline phosphatase positive Langerhans Cells in the Epidermis of Cattle. *J Invest Dermatol.* 79, 47-51.
- Kipar, A. (1994): Klassifizierung und Dignität der Rundzellentumoren der Haut und kutanen Schleimhaut des Hundes. Dissertation, Justus-Liebig-Universität Gießen.

- Kipar, A., Baumgartner, W., Kremmer, E., Frese, K. und Weiss, E. (1998). Expression of major Histocompatibility Complex class II Antigen in neoplastic Cells of Canine cutaneous Histiocytoma. *Vet Immunol Immunopathol.* 62 (1), 1-13.
- Kramer, L., Campanini, N., Ferrari, C. und Cabassi, E. (1996). Ultrastructural Identification of Langerhans Cells in normal Feline Epidermis. *J Anat.* 189, 683-684.
- Lacroix, J.V. und Riser, W.H. (1947). Transmissible Lymphosarcoma of the Dog. *North Am Vet.* 28, 451-453.
- Leong, A.S.-Y. und Gilham, P.N. (1989). The Effects of progressiv Formaldehyd Fixation on the Preservation of Tissue Antigens. *Pathology.* 21, 266-268.
- Levitsky, H.I. (1996). Tumors derived from Antigen presenting Cells. *Semin Immunol.* 8, 281-287.
- Marchal, T., Chabanne, L., Kaplanski, C., Rigal, D. und Magnol, J.P. (1997). Immunophenotype of the canine transmissible venereal Tumor. *Vet Immunol Immunopathol.* 57, 1-11.
- Marchal, T., Dezutter-Dambuyant, C., Fournel, C., Magnol, J.P. und Schmitt, D. (1995a). Electronmicroscopic Study of Canine cutaneous Histiocytoma: a benign Langerhans Cell Tumor. *Adv Exp Med Biol.* 378, 549-51.
- Marchal, T., Dezutter-Dambuyant, C., Fournel, C., Magnol, J.P. und Schmitt, D. (1995b). Immunophenotypic and ultrastructural Evidence of the Langerhans Cell Origin of the Canine cutaneous Histiocytoma. *Acta Anat.* 153 (3), 189-202.
- Martin de las Mulas, J., Millan, Y., Ruiz-Villamor, E. und Bautista, M.J. (1999). Apoptosis and Mitosis in Tumours of the Skin and subcutaneous Tissues of the Dog. *Res Vet sci.* 66, 139-146.
- Mason, D.Y., Cordell, J., Brown, M., Pallesen, G., Ralfkiaer, E., Rothbard, J., Crumpton, M. und Gatter, K.C. (1989). Detection of T Cells in Paraffin Wax embedded Tissue using Antibodies against a Peptide Sequence from CD3 Antigen. *J clin Pathol.* 42, 194-200.
- Mason, D.Y., Cordell, J.L., Brown, M.H., Borst, J., Jones, M., Pulford, K., Jaffe, E., Ralfkiaer, E., Dallenbach, F. und Stein, H. (1995). CD79a: a novel Marker for B-Cell-Neoplasms in routinely processed Tissue Samples. *Blood.* 86, 1453-1459.
- Mason, D.Y., Cordell, J.L., Tse, A.G.D., Dongen, J.J.M.v., Noesel, C.J.M.v., Micklem, K., Pulford, K.A.F., Valensi, F., Comans-Bitter, W.M., Borst, J. und Gatter, K.C. (1991). The IgM-associated Protein mb-1 as a Marker of normal and neoplastic B-Cells. *J Immunol.* 147, 2474-2482.
- Mason, D.Y. und Sammons, R. (1978). Alkaline Phophatase and Peroxidase for double immunoenzymatic Labelling of cellular Constituents. *J clin Pathol.* 31, 454-460.
- Mason, J.T. und O'Leary, T.J. (1991). Effects of Formaldehyd Fixation on Protein secondary Structure: a Calorimetric and infrared Spectroscopic Investigation. *J Histochem Cytochem.* 39, 225-229.

- Matsumoto, Y., Iwasaka, T., Yamasaki, F. und Sugimori, H. (1999). Apoptosis and Ki-67 Expression in adenomyotic Lesions and in the corresponding eutopic Endometrium. *Obstet Gynecol.* 94, 71-77.
- McClelland, R.B. (1940). Neoplasms in Dogs. *Cornell Vet.* 30, 161-166.
- Metlay, J.P., Pure, E. und Steinman, R.M. (1989). Control of the immune Response at the Level of Antigen presenting Cells: a Comparison of the Function of dendritic Cells and B Lymphocytes. *Adv Immunol.* 47, 45-116.
- Milner, R.J., Pearson, J., Nesbit, J.W. und Close, P. (1996). Immunphenotypic Classification of Canine malignant Lymphoma on Formalin-fixed Paraffin-embedded Tissue by means of CD3 and CD79a Cell Markers. *Onderstepoort J Vet Res.* 63, 309-313.
- Mizuno, S., Fujinaga, T. und Hagio, M. (1994). Role of Lymphocytes in spontaneous Regression of experimentally transplanted canine transmissible venereal Sarcoma. *J Vet Med Sci.* 56, 15-20.
- Moore, P.F. (1986). Utilization of cytoplasmic Lysozyme Immunoreactivity as a histiocytic Marker in Canine histiocytic Disorders. *Vet Pathol.* 23 (6), 757-62.
- Moore, P.F. und Mariassy, A.T. (1986). Dendritic (Langerhans) Cells in Canine Epidermis: Ultrastructure and Distribution. *Anat Histol Embryol.* 15 (2), 178-179.
- Moore, P.F. und Schrenzel, M.D. (1991). Canine cutaneous Histiocytoma represents a Langerhans Cell proliferative Disorder based on an immunophenotypic Analysis. *Vet Pathol.* 28, 119.
- Moore, P.F., Schrenzel, M.D., Affolter, V.K., Olivry, T. und Naydan, D. (1996). Canine cutaneous Histiocytoma is an epidermotropic Langerhans Cell Histiocytosis that expresses CD1 and specific beta 2-integrin Molecules. *Am J Pathol.* 148 (5), 1699-708.
- Morgan, J.M., Navabi, H., Schmid, K.W. und Jasani, B. (1994). Possible Role of Tissue-bound Calcium Ions in Citrate-mediated high Temperature Antigen Retrieval. *J Pathol.* 174, 301-307.
- Moulton, J.E. (1961). Tumors of the Skin and Subcutis. In *Tumors of domestic animals*. University of California Press: Berkeley and Los Angeles, S. 32-33
- Mozos, E., Méndez, A., Gómez-Villamandos, J.C., Martín de las Mulas, J. und Pérez, J. (1996). Immunhistochemical Characterization of canine transmissible venereal Tumor. *Vet Pathol.* 33, 257-263.
- Mulligan, R.M. (1948). Diseases of Dogs. II. Mast Cell Sarcoma, Lymphosarcoma, Histiocytoma. *Arch Pathol.* 46, 477-492.
- Murphy, G.F., Radu, A., Kaminer, M. und Berd, D. (1993). Autologous Melanoma Vaccine induces Inflammatory Responses in Melanoma Metastases: Relevance to immunologic Regression and Immunotherapy. *J Invest Dermatol.* 100, 335-341.
- Myers, C.D. (1991). Role of B Cell Antigen Processing and Presentation in the humoral immune Response. *FASEB J.* 5, 2547-2553.

- Neumüller, O.A. (1973). Römps Chemie-Lexikon. Franckh'sche Verlagshandlung: Stuttgart.
- Nicholls, P.K., Moore, P.F., Anderson, D.M., Moore, R.A., Parry, N.R., Gough, G.W. und Stanley, M.S. (2001). Regression of canine oral Papillomas is associated with Infiltration of CD4+ and CD8+ Lymphocytes. *Virology*. 283, 31-39.
- Noesel, C.J.M.v., Lier, R.A.W.v., Cordell, J., Tse, A.G.D., Schijndel, G.M.W.v., Vries, E.F.R.d., Mason, D.Y. und Borst, J. (1991). The Membrane IgM-associated Heterodimer on human B Cells is a newly defined B Cell Antigen that contains the Protein Product of the mb-1 Gene. *J Immunol.* 146, 3881-3888.
- Norton, A.J., Jordan, S. und Yeomans, P. (1994). Brief, high-temperature Heat Denaturation (Pressure Cooking): a simple and effective Method of Antigen Retrieval for routinely processed Tissues. *J Pathol.* 173, 371-379.
- Ottosen, H.E. (1949). Studies on Sarcomata of the Skin in Dogs. *Nord Veterinarmed.* 1, 7-30.
- Parnes, J.R. (1990). Molecular Biology an Function of CD4 und CD8. *Adv Immunol.* 44, 265-311.
- Pérez, J., Day, M.J., Martín, M.P., González, S. und Mozos, E. (1999). Immunohistochemical Study of the Inflammatory Infiltrate associated with feline cutaneous squamous Cell Carcinomas and precancerous Lesions (actinic Keratosis). *Vet Immunol Immunopathol.* 69, 33-46.
- Pérez, J., Day, M.J. und Mozos, E. (1998). Immunhistochemical Study of the local Inflammatory Infiltrate in spontaneous canine transmissible venereal Tumour at different Stages of Growth. *Vet Immunol Immunopathol.* 64, 133-147.
- Pérez-Torres, A. und Ustarroz-Cano, M. (2001). Demanstration of Birbeck (Langerhans cells) Granules in the normal Chicken Epidermis. *J Anat.* 199, 493-497.
- Pierrepont, C.G. (1985). Possible Benefits to Veterinary Medicine of considering the Dog as a Model for human Cancer. *J small Anim Pract.* 26, 43-47.
- Pileri, S.A., Roncador, G., Ceccarelli, C., Piccioli, M., Briskomatis, A., Sabattini, E., Ascani, S., Santini, D., Piccaluga, P.P., Leone, O., Damiani, S., Ercolelli, C., Sandri, F., Pieri, F., Leoncini, L. und Falini, B. (1997). Antigen retrieval Techniques in Immunhistochemistry: Comparison of different Methods. *J Pathol.* 183, 116-123.
- Pleiman, C.M., D'Ambrosio, D. und Cambier, J.C. (1994). The B-Cell Antigen Receptor Complex: Structure and Signal Transduction. *Immunol Today.* 15, 393-399.
- Rabanal, R.M., Lluis, F. und Else, R.W. (1995). Immunhistochemical Detection of Canine Leucocyte Antigens by specific monoclonal Antibodies in Canine normal Tissues. *Vet Immunol Immunopathol.* 47, 13-23.
- Roels, S., Tilmant, K. und Ducatelle, R. (1999). PCNA and Ki-67 Proliferation Markers as Criteria for Prediction of Clinical Behaviour of Melanocytic Tumours in Cats and Dogs. *J Comp Path.* 121, 13-24.
- Roitt, I. (1993). Leitfaden der Immunologie. Monographie, Blackwell-Wissenschafts-Verlag.

- Romano, J. und Balaguer, L. (1991). Ultrastructural Identification of Langerhans Cells in normal Swine Epidermis. *J Anat.* 179, 43-46.
- Romeis, B. und Böck, P. (1989a). Fixierung histologischer Präparate. In Romeis, Benno: Mikroskopische Technik, Böck, P. Urban und Schwarzenberg: München (17. Auflage), S. 69-80.
- Romeis, B. und Böck, P. (1989b). Färbemethoden. In Romeis, Benno: Mikroskopische Technik, Böck, P. Urban und Schwarzenberg: München (17. Auflage), Kapitel 8.2, S. 208-247.
- Rosenberg, S.A. (2001). Progress in human Tumour Immunology and Immunotherapy. *Nature.* 411, 380-384.
- Saalmüller, A. und Bryant, J. (1994). Characteristics of Porcine T-Lymphocytes and T-Cell Lines. *Vet Immunol Immunopathol.* 43, 45-52.
- Sadanaga, N., Kuwano, H., Watanabe, M., Maekawa, S., Mori, M. und Sugimachi, K. (1994). Local immune Response to Tumor Invasion in esophageal squamous Cell Carcinoma. The Expression of human Leukocyte Antigen-DR and Lymphocyte Infiltration. *Cancer.* 74, 586-591.
- Sakaguchi, N., Kashiwamura, S., Kimoto, M., Thalmann, P. und Melchers, F. (1988). B-lymphocyte lineage-restricted Expression of mb-1, a Gene with CD3-like structural Properties. *EMBO J.* 7, 3457-3464.
- Sandusky, G.E., Carlton, W.W. und Wightman, K.A. (1987). Diagnostic Immunohistochemistry of Canine round Cell Tumors. *Vet Pathol.* 24 (6), 495-9.
- Sarli, G., Benazzi, C., Preziosi, R. und Marcato, P.S. (1994). Proliferative Activity assessed by Anti-PCNA and Ki-67 monoclonal Antibodies in canine testicular Tumours. *J Comp Path.* 110, 357-368.
- Sawhney, N. und Hall, P.A. (1992). Ki-67-Structure, Function and new Antibodies. *J Pathol.* 168, 161-162.
- Scholzen, T. und Gerdés, J. (2000). The Ki-67 Protein: from the Known and the Unknown. *J Cell Physiol.* 182, 311-322.
- Schulden, U., Walter, J.H., Gutberlet, K. und Rudolph, R. (1998). Immunhistochemical Identification of B Lymphocytes in Canine Lymphnodes embedded in Paraffin Wax by using an Antibody to CD79a Antigen. *Europ J of Vet Pathol.* 4 (2), 1-3.
- Seiter, S. und Marincola, F.M. (2000). The multiple Ways to Tumor Tolerance. *Mod Asp Immunobiol.* 1, 121-124.
- Shi, S.-R., Imam, S.A., Young, L., Cote, R.J. und Taylor, C.R. (1995). Antigen Retrieval Immunohistochemistry under the Influence of pH using monoclonal Antibodies. *J Histochem Cytochem.* 43, 193-201.

- Shi, S.-R., Key, M.E. und Kalra, K.L. (1991). Antigen Retrieval in Formalin-fixed, Paraffin-embedded Tissues: an Enhancement Method for immunohistochemical Staining based on Microwave Oven Heating of Tissue Sections. *J Histochem Cytochem.* 39 (6), 741-748.
- Smith, H.A. und Jones, T.C. (1966). Neoplasia. In *Veterinary Pathology*. Lea and Febiger: Philadelphia, S. 220-222.
- Soini, Y., Pääkkö, P. und V.-P., L. (1998). Histopathological Evaluation of Apoptosis in Cancer. *Am J Pathol.* 153, 1041-1053.
- Stannard, A.A. und Pulley, L.T. (1978). Tumors of the Skin and soft Tissues. In *Tumors in domestic animals*, Moulton, E. University of California press: Berkeley, S. 34-36.
- Steinman, R.M. (1991). The dendritic Cell System and its Role in Immunogenicity. *Annu Rev Immunol.* 9, 271-296.
- Sternberger, L.A., Hardy, P.H.J., J., C.J. und Meyer, H.G. (1970). The unlabeled Antibody Enzyme Method of Immunohistochemistry. Preparation and Properties of soluble Antigen-Antibody Complex (horseradish peroxidase-antihorseradish Peroxidase) and its Use in Identification of Spirochetes. *J Histochem Cytochem.* 18, 315-334.
- Stingl, G., Tamki, K. und Katz, S.I. (1980). Origin and Function of epidermal Langerhans Cells. *Immunol Rev.* 53, 149-174.
- Straus, W. (1971). Inhibition of Peroxidase by Methanol and by Methanol-Nitroferricyanide for use in Immunoperoxidase Procedures. *J Histochem Cytochem.* 19, 682-688.
- Sutton, R.H. und McLennan, M.W. (1987). Cutaneous Histiocytoma in a Heifer. *J Comp Pathol.* 97 (4), 463-8.
- Taylor, D.O., Dorn, C.R. und Luis, O.H. (1969). Morphologic and biologic Characteristics of the Canine cutaneous Histiocytoma. *Cancer Res.* 29 (1), 83-92.
- Thome, M., Hirt, W. und Pfaff, E. (1994). Porcine T-Cell Receptor: molecular and biochemical Charakterisation. *Vet Immunol Immunopathol.* 43, 13-18.
- Tizard, I.R. und Schubot, R.M. (2000a). MHC: The Antigen-Presenting Receptors. In *Veterinary Immunology*. W. B. Saunders Company: Philadelphia, S. 47-57.
- Tizard, I.R. und Schubot, R.M. (2000b). T Cell Function and the Destruction of Cell-Associated Antigen. In *Veterinary Immunology*. W. B. Saunders Company: Philadelphia, S. 161-169.
- Wagner, S.N., Schultewolter, T., Wagner, C., Briedigkeit, L., Becker, J.C., Kwasnicka, H.M. und Goos, M. (1998). Immune Response against human primary malignant Melanoma: a ditinct cytokine mRNA Profile associated with spontaneous Regression. *Lab Invest.* 78, 541-550.
- Walter, J.H., Gutberlet, K., Schwegler, K. und Rudolph, R. (1997). Vorkommen und Häufigkeiten caniner Neoplasien im Sektionsgut. *Kleintierpraxis.* 42 (4), 269-356.

- Watts, C. (1997). Capture and Processing of exogenous Antigens for Presentation on MHC Molecules. *Annu Rev Immunol.* 15, 821-850.
- Weber, T. (1985): Enzymhistochemische Untersuchung am kutanen Histiozytom des Hundes. Dissertation, Fachbereich Veterinärmedizin der Freien Universität Berlin.
- Weinstein, M.J., Carpenter, J.L. und Schunk, C.J. (1989). Nonangiogenic and nonlymphomatous Sarcomas of the Canine Spleen: 57 Cases (1975-1987). *J Am Vet Med Assoc.* 195 (6), 784-8.
- Weiβ, E. (1974). Tumors of the soft (mesenchymal) Tissues. *Bull Wld Hlth Org.* 50, 101-110.
- Williamson, L.H., Steward, M., Milton, I., Parr, A., Krajewski, A.S., Angus, B. und Horne, C.H.W. (1998). Technical Advance: New Antibodies to the T Cell Antigens CD4 and CD8. Production and Charakterization in Formalin-Fixed Paraffin-Embedded Tissue. *Am J Pathol.* 152 (6), 1421-1426.
- Woods, G.S. und Warnke, R. (1981). Suppression of endogenous avidinbinding Activity in Tissues and its Relevance to biotin-avidin Detection System. *J Histochem Cytochem.* 29 (10), 1196-1204.