

CONTENTS

| | |
|--|----------|
| ABBREVIATIONS..... | v |
| 1 INTRODUCTION | 1 |
| 2 OVERALL OBJECTIVE | 6 |
| 2.1 Specific objectives (cross-sectional study)..... | 6 |
| 2.2 Specific objectives (longitudinal study) | 6 |
| 3 LITERATURE REVIEW | 7 |
| 3.1 Trypanosomosis..... | 7 |
| 3.1.1 Introduction | 7 |
| 3.1.2 Aetiology and transmission of AAT..... | 8 |
| 3.1.3 Epidemiology..... | 8 |
| 3.1.4 Pathogenesis | 9 |
| 3.1.5 Economic importance..... | 10 |
| 3.1.6 Diagnosis of trypanosomosis..... | 11 |
| 3.1.7 Control..... | 12 |
| 3.1.7.1 Chemotherapy and chemoprophylaxis of AAT | 12 |
| 3.1.7.2 Diminazene aceturate | 12 |
| 3.1.7.3 Isometamidium chloride | 13 |
| 3.1.7.4 Homidium | 13 |
| 3.1.7.5 Trypanocidal resistance | 14 |
| 3.1.7.6 Tsetse control..... | 15 |
| 3.2 Tick-borne diseases | 16 |
| 3.2.1 Anaplasmosis..... | 16 |
| 3.2.2 Babesiosis | 17 |
| 3.2.3 Theileriosis-East Coast Fever | 19 |
| 3.2.3.1 Aetiology | 19 |
| 3.2.3.2 Transmission..... | 20 |
| 3.2.3.3 Clinical syndrome..... | 20 |
| 3.2.3.4 Diagnosis | 20 |
| 3.2.3.5 Chemotherapy..... | 21 |
| 3.2.4 Epidemiology of tick-borne diseases..... | 22 |
| 3.2.5 Economic importance of TBDs | 22 |
| 3.2.6 Control of ticks and TBDs..... | 23 |
| 3.2.6.1 Acaricides | 24 |
| 3.2.6.2 Methods of application | 26 |

| | | |
|----------|---|-----------|
| 3.2.6.3 | Other methods of control..... | 27 |
| 3.2.6.4 | Integrated control of ticks and TBDs | 27 |
| 3.3 | Helminthosis..... | 28 |
| 3.3.1 | Nematodes | 28 |
| 3.3.1.1 | Life cycle of nematodes | 29 |
| 3.3.2 | Cestodes..... | 30 |
| 3.3.2.1 | Life cycle of cestodes | 31 |
| 3.3.3 | Trematodes | 32 |
| 3.3.3.1 | Life cycle of trematodes | 32 |
| 3.3.4 | Epidemiology of helminthosis..... | 33 |
| 3.3.5 | Interaction of helminth- and trypanosome-infection | 34 |
| 3.3.6 | Economic importance | 35 |
| 3.3.7 | Diagnosis | 36 |
| 3.3.8 | Control of helminthosis | 36 |
| 3.3.8.1 | Anthelmintics..... | 36 |
| 3.3.8.2 | Control of nematodes..... | 38 |
| 3.3.8.3 | Control of cestodes and metacestodes | 39 |
| 3.3.8.4 | Control of fasciolosis..... | 39 |
| 4 | MATERIALS AND METHODS..... | 40 |
| 4.1 | Description of the study area..... | 40 |
| 4.2 | Cross-sectional survey..... | 42 |
| 4.2.1 | Study animals | 42 |
| 4.2.2 | Animal sampling and examination | 42 |
| 4.2.3 | Data collection..... | 43 |
| 4.2.3.1 | Questionnaire survey | 43 |
| 4.2.3.2 | Determination of Packed Cell Volume | 43 |
| 4.2.3.3 | Detection and identification of trypanosomes | 43 |
| 4.2.3.4 | Detection of Theileria, Anaplasma and Babesia parasites | 44 |
| 4.2.4 | Entomological survey..... | 44 |
| 4.2.4.1 | Sampling..... | 44 |
| 4.2.4.2 | Dissection of flies to determine infection rate..... | 44 |
| 4.2.5 | Faecal sampling and determination of FEC | 45 |
| 4.3 | Longitudinal study..... | 46 |
| 4.3.1 | Study area | 46 |
| 4.3.2 | Study design | 47 |
| 4.3.2.1 | Study population and sampling | 47 |
| 4.3.3 | Determination of trypanosome incidence density | 49 |
| 4.3.4 | Faecal sample examination..... | 49 |
| 4.3.5 | Examination for trematode eggs..... | 49 |
| 4.3.6 | Weighing of calves | 50 |
| 4.3.7 | Production parameters | 50 |
| 4.3.8 | Entomological survey..... | 51 |

| | | |
|----------|---|-----------|
| 4.3.8.1 | Collection and identification of bloodmeal samples | 51 |
| 4.4 | Data management and analysis..... | 51 |
| 4.4.1 | Cross-sectional study | 51 |
| 4.4.2 | Longitudinal study | 52 |
| 5 | RESULTS..... | 55 |
| 5.1 | Cross-sectional study | 55 |
| 5.1.1 | Questionnaire survey | 55 |
| 5.1.1.1 | Household data | 55 |
| 5.1.1.2 | Animal husbandry and management | 55 |
| 5.1.1.3 | Management of animals trypanosomosis | 56 |
| 5.1.1.4 | Control of helminths..... | 58 |
| 5.1.1.5 | Tick control..... | 58 |
| 5.1.2 | Results of cross-sectional disease survey | 59 |
| 5.1.3 | PCV profiles | 66 |
| 5.1.4 | Entomological survey | 67 |
| 5.1.5 | Tick infestation | 68 |
| 5.1.6 | Helminthosis in calves..... | 69 |
| 5.2 | Longitudinal study | 70 |
| 5.2.1 | Trypanosome incidence..... | 70 |
| 5.2.2 | Survival analysis for trypanosomosis: multiple failure-time data analysis | 72 |
| 5.2.3 | Entomological survey | 76 |
| 5.2.3.1 | Apparent tsetse density | 76 |
| 5.2.3.2 | Trypanosome species and infection rates in tsetse flies | 78 |
| 5.2.3.3 | Results of bloodmeal identification | 80 |
| 5.2.3.4 | Feeding patterns..... | 81 |
| 5.2.3.5 | Tsetse challenge..... | 82 |
| 5.2.4 | Tick-borne diseases | 83 |
| 5.2.5 | Faecal egg counts | 86 |
| 5.2.5.1 | Descriptive statistics..... | 86 |
| 5.2.5.2 | Faecal egg count reduction | 95 |
| 5.2.5.3 | Results of negative binomial regression analysis | 96 |
| 5.2.5.4 | Shedding of trematode eggs | 98 |
| 5.2.6 | Packed cell volume profiles..... | 101 |
| 5.2.7 | Weight changes in calves | 106 |
| 5.2.8 | Conception rate..... | 110 |
| 5.2.9 | Calving rate..... | 111 |
| 5.2.10 | Abortions and stillbirths | 111 |
| 5.2.11 | Mortality | 112 |
| 5.2.12 | Milk production | 114 |

| | | |
|-----------|--|------------|
| 6 | DISCUSSION..... | 116 |
| 6.1 | Cross-sectional study..... | 116 |
| 6.1.1 | Questionnaire survey..... | 116 |
| 6.1.2 | Trypanosome infection..... | 118 |
| 6.1.3 | Tick-borne diseases..... | 124 |
| 6.1.4 | Helminthosis in calves..... | 125 |
| 6.2 | Longitudinal study..... | 126 |
| 6.2.1 | Trypanosome incidence..... | 126 |
| 6.2.2 | Multivariable analysis for failure-time in trypanosomosis..... | 129 |
| 6.2.3 | Entomological survey and bloodmeal analysis..... | 133 |
| 6.2.4 | Tick-borne diseases..... | 136 |
| 6.2.5 | Helminthosis..... | 138 |
| 6.2.6 | Shedding of trematode eggs..... | 142 |
| 6.2.7 | Packed cell volume profiles..... | 146 |
| 6.2.8 | Weight changes in calves..... | 147 |
| 6.2.9 | Conception, calving and abortion rates..... | 148 |
| 6.2.10 | Milk production..... | 150 |
| 7 | CONCLUSIONS..... | 152 |
| 8 | SUMMARY..... | 153 |
| 9 | ZUSAMMENFASSUNG..... | 155 |
| 10 | REFERENCES..... | 158 |
| 11 | ANNEXES..... | 183 |
| | ACKNOWLEDGEMENT..... | 188 |
| | CURRICULUM VITAE..... | 190 |
| | SELBSTÄNDIGKEITSERKLÄRUNG..... | 191 |