

Literaturverzeichnis

- Agnello, V., Abel, G., Elfahal, M., Knight, G. B., und Zhang, Q. X. (1999). Hepatitis c virus and other flaviviridae viruses enter cells via low density lipoprotein receptor. *Proc Natl Acad Sci U S A*, 96(22):12766–71.
- Aizaki, H., Saito, S., Ogino, T., Miyajima, N., Harada, T., Matsuura, Y., Miyamura, T., und Kohase, M. (2000). Suppression of interferon-induced antiviral activity in cells expressing hepatitis c virus proteins. *J Interferon Cytokine Res*, 20(12):1111–20.
- Alcami, A. und Koszinowski, U. H. (2000). Viral mechanisms of immune evasion. *Immunol Today*, 21(9):447–55.
- Alcami, A. und Smith, G. L. (1995). Vaccinia, cowpox, and camelpox viruses encode soluble gamma interferon receptors with novel broad species specificity. *J Virol*, 69(8):4633–9.
- Ali, S., Pellerin, C., Lamarre, D., und Kukolj, G. (2004). Hepatitis c virus subgenomic replicons in the human embryonic kidney 293 cell line. *J Virol*, 78(1):491–501.
- Antonelli, G., Simeoni, E., Turriziani, O., Tesoro, R., Redaelli, A., Roffi, L., Antonelli, L., Pistello, M., und Dianzani, F. (1999). Correlation of interferon-induced expression of mxa mrna in peripheral blood mononuclear cells with the response of patients with chronic active hepatitis c to ifn-alpha therapy. *J Interferon Cytokine Res*, 19(3):243–51.
- Asabe, S. I., Tanji, Y., Satoh, S., Kaneko, T., Kimura, K., und Shimotohno, K. (1997). The n-terminal region of hepatitis c virus-encoded ns5a is important for ns4a-dependent phosphorylation. *J Virol*, 71(1):790–6.
- Aus dem Siepen, M., Lohmann, V., Wiese, M., Ross, S., Roggendorf, M., und Viazov, S. (2005). Nonstructural protein 5a does not contribute to the resistance of hepatitis c virus replication to interferon alpha in cell culture. *Virology*, 336(2):131–6.
- Babine, R. E., Chen, S.-H., Lamar, J. E., Snyder, N. J., Sun, X. D., Tebbe, M. J., Victor, F., Wang, Q. M., Yip, Y. Y. M., und Collado, I. e. (2002). International patent wo 0218369.
- Banner, B. F., Allan, C., Savas, L., Baker, S., Barnard, G., und Bonkovsky, H. L. (1997). Inflammatory markers in chronic hepatitis c. *Virchows Arch*, 431(3):181–7.
- Barnard, P. und McMillan, N. A. (1999). The human papillomavirus e7 oncoprotein abrogates signaling mediated by interferon-alpha. *Virology*, 259(2):305–13.
- Barret, T. und Inglis, S. (1985). Gowth and purification of influenza viruses. In Mahy, J., editor, *Virology: A Practical Approach*, pages 119–150. IRL Press Oxford.

- Bartenschlager, R., Kaul, A., und Sparacio, S. (2003). Replication of the hepatitis c virus in cell culture. *Antiviral Res*, 60(2):91–102.
- Bartenschlager, R. und Lohmann, V. (2000). Replication of the hepatitis c virus. *Baillieres Best Pract Res Clin Gastroenterol*, 14(2):241–54.
- Bartenschlager, R., Lohmann, V., Wilkinson, T., und Koch, J. O. (1995). Complex formation between the ns3 serine-type proteinase of the hepatitis c virus and ns4a and its importance for polyprotein maturation. *J Virol*, 69(12):7519–28.
- Bartosch, B., Dubuisson, J., und Cosset, F. L. (2003). Infectious hepatitis c virus pseudo-particles containing functional e1-e2 envelope protein complexes. *J Exp Med*, 197(5):633–42.
- Basler, C. F., Mikulasova, A., Martinez-Sobrido, L., Paragas, J., Muhlberger, E., Bray, M., Klenk, H. D., Palese, P., und Garcia-Sastre, A. (2003). The ebola virus vp35 protein inhibits activation of interferon regulatory factor 3. *J Virol*, 77(14):7945–56.
- Basler, C. F., Wang, X., Muhlberger, E., Volchkov, V., Paragas, J., Klenk, H. D., Garcia-Sastre, A., und Palese, P. (2000). The ebola virus vp35 protein functions as a type i ifn antagonist. *Proc Natl Acad Sci U S A*, 97(22):12289–94.
- Basu, A., Meyer, K., Ray, R. B., und Ray, R. (2001). Hepatitis c virus core protein modulates the interferon-induced transacting factors of jak/stat signaling pathway but does not affect the activation of downstream irf-1 or 561 gene. *Virology*, 288(2):379–90.
- Beattie, E., Denzler, K. L., Tartaglia, J., Perkus, M. E., Paoletti, E., und Jacobs, B. L. (1995). Reversal of the interferon-sensitive phenotype of a vaccinia virus lacking e31 by expression of the reovirus s4 gene. *J Virol*, 69(1):499–505.
- Beitz, E. (2000). Texshade: shading and labeling of multiple sequence alignments using latex2 epsilon. *Bioinformatics*, 16(2):135–9.
- Bejarano, M. T. und Masucci, M. G. (1998). Interleukin-10 abrogates the inhibition of epstein-barr virus-induced b-cell transformation by memory t-cell responses. *Blood*, 92(11):4256–62.
- Bellows, C. F., Garry, R. F., und Jaffe, B. M. (2003). Vaccinia virus-induced inhibition of nitric oxide production. *J Surg Res*, 111(1):127–35.
- Benezra, M., Chevallier, N., Morrison, D. J., MacLachlan, T. K., El-Deiry, W. S., und Licht, J. D. (2003). Brca1 augments transcription by the nf-kappab transcription factor by binding to the rel domain of the p65/rela subunit. *J Biol Chem*, 278(29):26333–41.
- Berg, T. und Hopf, U. (2001). Therapie der hepatitis-c-virus (hcv)-infektion. In Häussinger, D. und Niederau, C., editors, *Hepatitis C*, volume 2. Auflage of *Ex Libris (Roche)*, pages 233–284. Blackwell Wissenschafts-Verlag, Berlin, Wien, bd. 8 edition.
- Berg, T., Mas Marques, A., Hohne, M., Wiedenmann, B., Hopf, U., und Schreier, E. (2000). Mutations in the e2-pephd and ns5a region of hepatitis c virus type 1 and the dynamics of hepatitis c viremia decline during interferon alfa treatment. *Hepatology*, 32(6):1386–95.
- Biernz, K., Egger, D., Pfister, T., und Troxler, M. (1992). Structural and functional characterization of the poliovirus replication complex. *J Virol*, 66(5):2740–7.

- Black, T. L., Barber, G. N., und Katze, M. G. (1993). Degradation of the interferon-induced 68,000-m(r) protein kinase by poliovirus requires rna. *J Virol*, 67(2):791–800.
- Blight, K. J., Kolykhalov, A. A., und Rice, C. M. (2000). Efficient initiation of hcv rna replication in cell culture. *Science*, 290(5498):1972–4.
- Blindenbacher, A., Duong, F. H., Hunziker, L., Stutvoet, S. T., Wang, X., Terracciano, L., Moradpour, D., Blum, H. E., Alonzi, T., Tripodi, M., La Monica, N., und Heim, M. H. (2003). Expression of hepatitis c virus proteins inhibits interferon alpha signaling in the liver of transgenic mice. *Gastroenterology*, 124(5):1465–75.
- Bode, J. G., Ludwig, S., Ehrhardt, C., Albrecht, U., Erhardt, A., Schaper, F., Heinrich, P. C., und Haussinger, D. (2003). Ifn-alpha antagonistic activity of hcv core protein involves induction of suppressor of cytokine signaling-3. *Faseb J*, 17(3):488–90.
- Bolten, R., Egger, D., Gosert, R., Schaub, G., Landmann, L., und Bienz, K. (1998). Intracellular localization of poliovirus plus- and minus-strand rna visualized by strand-specific fluorescent in situ hybridization. *J Virol*, 72(11):8578–85.
- Bossert, B., Marozin, S., und Conzelmann, K. K. (2003). Nonstructural proteins ns1 and ns2 of bovine respiratory syncytial virus block activation of interferon regulatory factor 3. *J Virol*, 77(16):8661–8.
- Bowie, A. G. und Haga, I. R. (2005). The role of toll-like receptors in the host response to viruses. *Mol Immunol*, 42(8):859–67.
- Bradley, D., McCaustland, K., Krawczynski, K., Spelbring, J., Humphrey, C., und Cook, E. H. (1991). Hepatitis c virus: buoyant density of the factor viii-derived isolate in sucrose. *J Med Virol*, 34(3):206–8.
- Brass, V., Bieck, E., Montserret, R., Wolk, B., Hellings, J. A., Blum, H. E., Penin, F., und Moradpour, D. (2002). An amino-terminal amphipathic alpha-helix mediates membrane association of the hepatitis c virus nonstructural protein 5a. *J Biol Chem*, 277(10):8130–9.
- Breiman, A., Grandvaux, N., Lin, R., Ottone, C., Akira, S., Yoneyama, M., Fujita, T., Hiscott, J., und Meurs, E. F. (2005). Inhibition of rig-i-dependent signaling to the interferon pathway during hepatitis c virus expression and restoration of signaling by ikkepsilon. *J Virol*, 79(7):3969–78.
- Bukh, J., Pietschmann, T., Lohmann, V., Krieger, N., Faulk, K., Engle, R. E., Govindarajan, S., Shapiro, M., St Claire, M., und Bartenschlager, R. (2002). Mutations that permit efficient replication of hepatitis c virus rna in huh-7 cells prevent productive replication in chimpanzees. *Proc Natl Acad Sci U S A*, 99(22):14416–21.
- Bukh, J., Purcell, R. H., und Miller, R. H. (1992). Sequence analysis of the 5' noncoding region of hepatitis c virus. *Proc Natl Acad Sci U S A*, 89(11):4942–6.
- Butkiewicz, N. J., Wendel, M., Zhang, R., Jubin, R., Pichardo, J., Smith, E. B., Hart, A. M., Ingram, R., Durkin, J., Mui, P. W., Murray, M. G., Ramanathan, L., und Dasmahapatra, B. (1996). Enhancement of hepatitis c virus ns3 proteinase activity by association with ns4a-specific synthetic peptides: identification of sequence and critical residues of ns4a for the cofactor activity. *Virology*, 225(2):328–38.

- Carlsson, T., Lindahl, K., Schvarcz, R., Wejstral, R., Uhnoo, I., Shev, S., und Reichard, O. (2000). Hcv rna levels during therapy with amantadine in addition to interferon and ribavirin in chronic hepatitis c patients with previous nonresponse or response/relapse to interferon and ribavirin. *J Viral Hepat*, 7(6):409–13.
- Carrick, R. J., Schlauder, G. G., Peterson, D. A., und Mushahwar, I. K. (1992). Examination of the buoyant density of hepatitis c virus by the polymerase chain reaction. *J Virol Methods*, 39(3):279–89.
- Cassady, K. A., Gross, M., und Roizman, B. (1998). The second-site mutation in the herpes simplex virus recombinants lacking the gamma134.5 genes precludes shutoff of protein synthesis by blocking the phosphorylation of eif-2alpha. *J Virol*, 72(9):7005–11.
- Castet, V. und Moradpour, D. (2003). A model for the study of hepatitis c virus entry. *Hepatology*, 38(3):771–4.
- Cayley, P. J., Davies, J. A., McCullagh, K. G., und Kerr, I. M. (1984). Activation of the ppp(a²'p)na system in interferon-treated, herpes simplex virus-infected cells and evidence for novel inhibitors of the ppp(a²'p)na-dependent rnase. *Eur J Biochem*, 143(1):165–74.
- CDC, C. f. D. C. (1998). Recommendations for prevention and control of hepatitis c virus (hcv) infection and hcv-related chronic disease. *Mortality and Morbidity Weekly Report*, 47:1–39.
- Chambers, T. J., Fan, X., Droll, D. A., Hembrador, E., Slater, T., Nickells, M. W., Dustin, L. B., und Dabisceglie, A. M. (2005). Quasispecies heterogeneity within the e1/e2 region as a pretreatment variable during pegylated interferon therapy of chronic hepatitis c virus infection. *J Virol*, 79(5):3071–83.
- Chambers, T. J., Weir, R. C., Grakoui, A., McCourt, D. W., Bazan, J. F., Fletterick, R. J., und Rice, C. M. (1990). Evidence that the n-terminal domain of nonstructural protein ns3 from yellow fever virus is a serine protease responsible for site-specific cleavages in the viral polyprotein. *Proc Natl Acad Sci U S A*, 87(22):8898–902.
- Chan, L., Das, S. K., Reddy, T. J., Poisson, C., Proulx, M., Pereira, O., Courchesne, M., Roy, C., Wang, W., und Siddiqui, A. e. (2004). Discovery of thiophene-2-carboxylic acids as potent inhibitors of hcv ns5b polymerase and hcv subgenomic rna replication. part 1: Sulfonamides. *Bioorg Med Chem Lett*, 14(3):793–6.
- Chang, H. W., Watson, J. C., und Jacobs, B. L. (1992). The e3l gene of vaccinia virus encodes an inhibitor of the interferon-induced, double-stranded rna-dependent protein kinase. *Proc Natl Acad Sci U S A*, 89(11):4825–9.
- Chang, J., Yang, S. H., Cho, Y. G., Hwang, S. B., Hahn, Y. S., und Sung, Y. C. (1998). Hepatitis c virus core from two different genotypes has an oncogenic potential but is not sufficient for transforming primary rat embryo fibroblasts in cooperation with the h-ras oncogene. *J Virol*, 72(4):3060–5.
- Chien, D. Y., Choo, Q. L., Tabrizi, A., Kuo, C., McFarland, J., Berger, K., Lee, C., Shuster, J. R., Nguyen, T., Moyer, D. L., und et al. (1992). Diagnosis of hepatitis c virus (hcv) infection using an immunodominant chimeric polyprotein to capture

- circulating antibodies: reevaluation of the role of hcv in liver disease. *Proc Natl Acad Sci U S A*, 89(21):10011–5.
- Choo, Q. L., Kuo, G., Weiner, A. J., Overby, L. R., Bradley, D. W., und Houghton, M. (1989). Isolation of a cdna clone derived from a blood-borne non-a, non-b viral hepatitis genome. *Science*, 244(4902):359–62.
- Choo, Q. L., Richman, K. H., Han, J. H., Berger, K., Lee, C., Dong, C., Gallegos, C., Coit, D., Medina-Selby, R., Barr, P. J., und et al. (1991). Genetic organization and diversity of the hepatitis c virus. *Proc Natl Acad Sci U S A*, 88(6):2451–5.
- Chung, K. M., Lee, J., Kim, J. E., Song, O. K., Cho, S., Lim, J., Seedorf, M., Hahm, B., und Jang, S. K. (2000). Nonstructural protein 5a of hepatitis c virus inhibits the function of karyopherin beta3. *J Virol*, 74(11):5233–41.
- Chung, Y. M., Park, K. J., Choi, S. Y., Hwang, S. B., und Lee, S. Y. (2001). Hepatitis c virus core protein potentiates tnf-alpha-induced nf-kappab activation through traf2-ikkbeta-dependent pathway. *Biochem Biophys Res Commun*, 284(1):15–9.
- Dale, T. C., Rosen, J. M., Guille, M. J., Lewin, A. R., Porter, A. G., Kerr, I. M., und Stark, G. R. (1989). Overlapping sites for constitutive and induced dna binding factors involved in interferon-stimulated transcription. *Embo J*, 8(3):831–9.
- Dansako, H., Naganuma, A., Nakamura, T., Ikeda, F., Nozaki, A., und Kato, N. (2003). Differential activation of interferon-inducible genes by hepatitis c virus core protein mediated by the interferon stimulated response element. *Virus Res*, 97(1):17–30.
- Dauber, B., Heins, G., und Wolff, T. (2004). The influenza b virus nonstructural ns1 protein is essential for efficient viral growth and antagonizes beta interferon induction. *J Virol*, 78(4):1865–72.
- Davies, M. V., Elroy-Stein, O., Jagus, R., Moss, B., und Kaufman, R. J. (1992). The vaccinia virus k3l gene product potentiates translation by inhibiting double-stranded-rna-activated protein kinase and phosphorylation of the alpha subunit of eukaryotic initiation factor 2. *J Virol*, 66(4):1943–50.
- Davis, G. L. (1999). Hepatitis c virus genotypos and quasispecies. *Am J Med*, 107(6B):21S–26S.
- de Lucas, S., Bartolome, J., und Carreno, V. (2005). Hepatitis c virus core protein down-regulates transcription of interferon-induced antiviral genes. *J Infect Dis*, 191(1):93–9.
- Del Porto, P., Punzoriero, G., Scotta, C., Nicosia, A., und Piccolella, E. (2000). High prevalence of hypervariable region 1-specific and -cross-reactive cd4(+) t cells in hcv-infected individuals responsive to ifn-alpha treatment. *Virology*, 269(2):313–24.
- Didcock, L., Young, D. F., Goodbourn, S., und Randall, R. E. (1999). The v protein of simian virus 5 inhibits interferon signalling by targeting stat1 for proteasome-mediated degradation. *J Virol*, 73(12):9928–33.
- Dimitrova, M., Imbert, I., Kieny, M. P., und Schuster, C. (2003). Protein-protein interactions between hepatitis c virus nonstructural proteins. *J Virol*, 77(9):5401–14.
- Dittmann, S., Roggendorf, M., Durkop, J., Wiese, M., Lorbeer, B., und Deinhardt, F. (1991). Long-term persistence of hepatitis c virus antibodies in a single source outbreak. *J Hepatol*, 13(3):323–7.

- Donelan, N. R., Dauber, B., Wang, X., Basler, C. F., Wolff, T., und Garcia-Sastre, A. (2004). The n- and c-terminal domains of the ns1 protein of influenza b virus can independently inhibit irf-3 and beta interferon promoter activation. *J Virol*, 78(21):11574–82.
- Dubois, F., Desenclos, J. C., Mariotte, N., und Goudeau, A. (1997). Hepatitis c in a french population-based survey, 1994: seroprevalence, frequency of viremia, genotype distribution, and risk factors. the collaborative study group. *Hepatology*, 25(6):1490–6.
- DuBridge, R. B., Tang, P., Hsia, H. C., Leong, P. M., Miller, J. H., und Calos, M. P. (1987). Analysis of mutation in human cells by using an epstein-barr virus shuttle system. *Mol Cell Biol*, 7(1):379–87.
- Dubuisson, J. und Rice, C. M. (1996). Hepatitis c virus glycoprotein folding: disulfide bond formation and association with calnexin. *J Virol*, 70(2):778–86.
- Duong, F. H., Filipowicz, M., Tripodi, M., La Monica, N., und Heim, M. H. (2004). Hepatitis c virus inhibits interferon signaling through up-regulation of protein phosphatase 2a. *Gastroenterology*, 126(1):263–77.
- Duverlie, G., Khorsi, H., Castelain, S., Jaillon, O., Izopet, J., Lunel, F., Eb, F., Penin, F., und Wychowski, C. (1998). Sequence analysis of the ns5a protein of european hepatitis c virus 1b isolates and relation to interferon sensitivity. *J Gen Virol*, 79 (Pt 6):1373–81.
- Egger, D., Wolk, B., Gosert, R., Bianchi, L., Blum, H. E., Moradpour, D., und Bienz, K. (2002). Expression of hepatitis c virus proteins induces distinct membrane alterations including a candidate viral replication complex. *J Virol*, 76(12):5974–84.
- Engler, S., Flechtenmacher, C., Wiedemann, K. H., Gugler, R., Stremmel, W., und Kallinowski, B. (2004). Interferon alfa2a induction therapy in combination with ribavirin and amantadine for the treatment of naive patients with chronic hcv infection. *J Viral Hepat*, 11(1):60–8.
- Enomoto, N., Kurosaki, M., Tanaka, Y., Marumo, F., und Sato, C. (1994). Fluctuation of hepatitis c virus quasispecies in persistent infection and interferon treatment revealed by single-strand conformation polymorphism analysis. *J Gen Virol*, 75 (Pt 6):1361–9.
- Enomoto, N., Sakuma, I., Asahina, Y., Kurosaki, M., Murakami, T., Yamamoto, C., Izumi, N., Marumo, F., und Sato, C. (1995). Comparison of full-length sequences of interferon-sensitive and resistant hepatitis c virus 1b. sensitivity to interferon is conferred by amino acid substitutions in the ns5a region. *J Clin Invest*, 96(1):224–30.
- Enomoto, N., Sakuma, I., Asahina, Y., Kurosaki, M., Murakami, T., Yamamoto, C., Ogura, Y., Izumi, N., Marumo, F., und Sato, C. (1996). Mutations in the nonstructural protein 5a gene and response to interferon in patients with chronic hepatitis c virus 1b infection. *N Engl J Med*, 334(2):77–81.
- Evans, M. J., Rice, C. M., und Goff, S. P. (2004). Phosphorylation of hepatitis c virus nonstructural protein 5a modulates its protein interactions and viral rna replication. *Proc Natl Acad Sci U S A*, 101(35):13038–43.

- Ezelle, H. J., Balachandran, S., Sicheri, F., Polyak, S. J., und Barber, G. N. (2001). Analyzing the mechanisms of interferon-induced apoptosis using crma and hepatitis c virus ns5a. *Virology*, 281(1):124–37.
- Failla, C., Tomei, L., und De Francesco, R. (1994). Both ns3 and ns4a are required for proteolytic processing of hepatitis c virus nonstructural proteins. *J Virol*, 68(6):3753–60.
- Falcon, V., Garcia, C., de la Rosa, M. C., Menendez, I., Seoane, J., und Grillo, J. M. (1999). Ultrastructural and immunocytochemical evidences of core-particle formation in the methylotrophic pichia pastoris yeast when expressing hcv structural proteins (core-e1). *Tissue Cell*, 31(2):117–25.
- Farci, P., Shimoda, A., Wong, D., Cabezon, T., De Gioannis, D., Strazzera, A., Shimizu, Y., Shapiro, M., Alter, H. J., und Purcell, R. H. (1996). Prevention of hepatitis c virus infection in chimpanzees by hyperimmune serum against the hypervariable region 1 of the envelope 2 protein. *Proc Natl Acad Sci U S A*, 93(26):15394–9.
- Fernandez-Gutierrez, B., Zamorano, J., Batlle, E., Alfonso, F., Conde, A., Sanchez-Harguindeguy, L., und Jover, J. A. (1999). Coronary dissection associated with hepatitis c virus-related cryoglobulinaemia. *Rheumatology (Oxford)*, 38(12):1299–301.
- Flint, M., Maidens, C., Loomis-Price, L. D., Shotton, C., Dubuisson, J., Monk, P., Higginbottom, A., Levy, S., und McKeating, J. A. (1999). Characterization of hepatitis c virus e2 glycoprotein interaction with a putative cellular receptor, cd81. *J Virol*, 73(8):6235–44.
- Foy, E., Li, K., Sumpter, R., J., Loo, Y. M., Johnson, C. L., Wang, C., Fish, P. M., Yoneyama, M., Fujita, T., Lemon, S. M., und Gale, M., J. (2005). Control of antiviral defenses through hepatitis c virus disruption of retinoic acid-inducible gene-i signaling. *Proc Natl Acad Sci U S A*, 102(8):2986–91.
- Foy, E., Li, K., Wang, C., Sumpter, R., J., Ikeda, M., Lemon, S. M., und Gale, M., J. (2003). Regulation of interferon regulatory factor-3 by the hepatitis c virus serine protease. *Science*, 300(5622):1145–8.
- Francki, R. I. B., Fauquet, C. M., Knudson, D. L., und Brown, F. (1991). *Fifth Report of the International Committee on Taxonomy of Viruses*. Classification and Nomenclature of Viruses. Springer Verlag; (September 1991), Wien New York.
- Francois, C., Duverlie, G., Rebouillat, D., Khorsi, H., Castelain, S., Blum, H. E., Gagnon, A., Wychowski, C., Moradpour, D., und Meurs, E. F. (2000). Expression of hepatitis c virus proteins interferes with the antiviral action of interferon independently of pkr-mediated control of protein synthesis. *J Virol*, 74(12):5587–96.
- Frank, C., Mohamed, M. K., Strickland, G. T., Lavanchy, D., Arthur, R. R., Magder, L. S., El Khoby, T., Abdel-Wahab, Y., Aly Ohn, E. S., Anwar, W., und Sallam, I. (2000). The role of parenteral antischistosomal therapy in the spread of hepatitis c virus in egypt. *Lancet*, 355(9207):887–91.
- Frese, M., Pietschmann, T., Moradpour, D., Haller, O., und Bartenschlager, R. (2001). Interferon-alpha inhibits hepatitis c virus subgenomic rna replication by an mxa-independent pathway. *J Gen Virol*, 82(Pt 4):723–33.

- Fukuma, T., Enomoto, N., Marumo, F., und Sato, C. (1998). Mutations in the interferon-sensitivity determining region of hepatitis c virus and transcriptional activity of the nonstructural region 5a protein. *Hepatology*, 28(4):1147–53.
- Gale, M., Blakely, C. M., Kwieciszewski, B., Tan, S. L., Dossett, M., Tang, N. M., Korth, M. J., Polyak, S. J., Gretch, D. R., und Katze, M. G. (1998). Control of pkr protein kinase by hepatitis c virus nonstructural 5a protein: molecular mechanisms of kinase regulation. *Mol Cell Biol*, 18(9):5208–18.
- Gale, M. J., J., Korth, M. J., Tang, N. M., Tan, S. L., Hopkins, D. A., Dever, T. E., Polyak, S. J., Gretch, D. R., und Katze, M. G. (1997). Evidence that hepatitis c virus resistance to interferon is mediated through repression of the pkr protein kinase by the nonstructural 5a protein. *Virology*, 230(2):217–27.
- Garcia-Sastre, A., Egorov, A., Matassov, D., Brandt, S., Levy, D. E., Durbin, J. E., Palese, P., und Muster, T. (1998). Influenza a virus lacking the ns1 gene replicates in interferon-deficient systems. *Virology*, 252(2):324–30.
- Geiss, G. K., Carter, V. S., He, Y., Kwieciszewski, B. K., Holzman, T., Korth, M. J., Lazaro, C. A., Fausto, N., Bumgarner, R. E., und Katze, M. G. (2003). Gene expression profiling of the cellular transcriptional network regulated by alpha/beta interferon and its partial attenuation by the hepatitis c virus nonstructural 5a protein. *J Virol*, 77(11):6367–75.
- Gerlach, J. T., Diepolder, H. M., Zachoval, R., Gruener, N. H., Jung, M. C., Ulsenheimer, A., Schraut, W. W., Schirren, C. A., Waechtler, M., Backmund, M., und Pape, G. R. (2003). Acute hepatitis c: high rate of both spontaneous and treatment-induced viral clearance. *Gastroenterology*, 125(1):80–8.
- Ghosh, A. K., Majumder, M., Steele, R., Meyer, K., Ray, R., und Ray, R. B. (2000a). Hepatitis c virus ns5a protein protects against tnf-alpha mediated apoptotic cell death. *Virus Res*, 67(2):173–8.
- Ghosh, A. K., Majumder, M., Steele, R., Yaciuk, P., Chrivia, J., Ray, R., und Ray, R. B. (2000b). Hepatitis c virus ns5a protein modulates transcription through a novel cellular transcription factor srcap. *J Biol Chem*, 275(10):7184–8.
- Ghosh, A. K., Steele, R., Meyer, K., Ray, R., und Ray, R. B. (1999). Hepatitis c virus ns5a protein modulates cell cycle regulatory genes and promotes cell growth. *J Gen Virol*, 80 (Pt 5):1179–83.
- Girard, S., Shalhoub, P., Lescure, P., Sabile, A., Misek, D. E., Hanash, S., Brechot, C., und Beretta, L. (2002). An altered cellular response to interferon and up-regulation of interleukin-8 induced by the hepatitis c viral protein ns5a uncovered by microarray analysis. *Virology*, 295(2):272–83.
- Gong, G., Waris, G., Tanveer, R., und Siddiqui, A. (2001). Human hepatitis c virus ns5a protein alters intracellular calcium levels, induces oxidative stress, and activates stat-3 and nf-kappa b. *Proc Natl Acad Sci U S A*, 98(17):9599–604.
- Gonzalez-Peralta, R. P., Qian, K., She, J. Y., Davis, G. L., Ohno, T., Mizokami, M., und Lau, J. Y. (1996). Clinical implications of viral quasispecies heterogeneity in chronic hepatitis c. *J Med Virol*, 49(3):242–7.

- Goodbourn, S., Didcock, L., und Randall, R. E. (2000). Interferons: cell signalling, immune modulation, antiviral response and virus countermeasures. *J Gen Virol*, 81(Pt 10):2341–64.
- Grakoui, A., McCourt, D. W., Wychowski, C., Feinstone, S. M., und Rice, C. M. (1993). A second hepatitis c virus-encoded proteinase. *Proc Natl Acad Sci U S A*, 90(22):10583–7.
- Griffin, S. D., Beales, L. P., Clarke, D. S., Worsfold, O., Evans, S. D., Jaeger, J., Harris, M. P., und Rowlands, D. J. (2003). The p7 protein of hepatitis c virus forms an ion channel that is blocked by the antiviral drug, amantadine. *FEBS Lett*, 535(1-3):34–8.
- Gunnery, S., Rice, A. P., Robertson, H. D., und Mathews, M. B. (1990). Tat-responsive region rna of human immunodeficiency virus 1 can prevent activation of the double-stranded-rna-activated protein kinase. *Proc Natl Acad Sci U S A*, 87(22):8687–91.
- Guo, J. T., Bichko, V. V., und Seeger, C. (2001). Effect of alpha interferon on the hepatitis c virus replicon. *J Virol*, 75(18):8516–23.
- Haller, O., Frese, M., und Kochs, G. (1998). Mx proteins: mediators of innate resistance to rna viruses. *Rev Sci Tech*, 17(1):220–30.
- Hanahan, D., Jesse, J., und Bloom, F. R. (1991). Plasmid transformation of escherichia coli and other bacteria. *Methods Enzymol*, 204:63–113.
- Haushofer, A. C., Kopty, C., Hauer, R., Brunner, H., und Halbmayer, W. M. (2001). Hcv genotypt and age distribution in patients of vienna and surrounding areas. *J Clin Virol*, 20(1-2):41–7.
- He, Y. und Katze, M. G. (2002). To interfere and to anti-interfere: the interplay between hepatitis c virus and interferon. *Viral Immunol*, 15(1):95–119.
- He, Y., Tan, S. L., Tareen, S. U., Vijaysri, S., Langland, J. O., Jacobs, B. L., und Katze, M. G. (2001). Regulation of mrna translation and cellular signaling by hepatitis c virus nonstructural protein ns5a. *J Virol*, 75(11):5090–8.
- Heim, M. H., Moradpour, D., und Blum, H. E. (1999). Expression of hepatitis c virus proteins inhibits signal transduction through the jak-stat pathway. *J Virol*, 73(10):8469–75.
- Heintges, T. und Wands, J. R. (1997). Hepatitis c virus: epidemiology and transmission. *Hepatology*, 26(3):521–6.
- Herion, D. und Hoofnagle, J. H. (1997). The interferon sensitivity determining region: all hepatitis c virus isolates are not the same. *Hepatology*, 25(3):769–71.
- Hertzog, P. J., Hwang, S. Y., und Kola, I. (1994). Role of interferons in the regulation of cell proliferation, differentiation, and development. *Mol Reprod Dev*, 39(2):226–32.
- Hofmann, W. P., Zeuzem, S., und Sarrazin, C. (2005). Hepatitis c virus-related resistance mechanisms to interferon alpha-based antiviral therapy. *J Clin Virol*, 32(2):86–91.
- Hohne, M., Schreier, E., und Roggendorf, M. (1994). Sequence variability in the env-coding region of hepatitis c virus isolated from patients infected during a single source outbreak. *Arch Virol*, 137(1-2):25–34.

- Hoofnagle, J. H., Mullen, K. D., Jones, D. B., Rustgi, V., Di Bisceglie, A., Peters, M., Waggoner, J. G., Park, Y., und Jones, E. A. (1986). Treatment of chronic non-a.non-b hepatitis with recombinant human alpha interferon. a preliminary report. *N Engl J Med*, 315(25):1575–8.
- Hornemann, S., Harlin, O., Staib, C., Kisling, S., Erfle, V., Kaspers, B., Hacker, G., und Sutter, G. (2003). Replication of modified vaccinia virus ankara in primary chicken embryo fibroblasts requires expression of the interferon resistance gene e3l. *J Virol*, 77(15):8394–407.
- Hosui, A., Ohkawa, K., Ishida, H., Sato, A., Nakanishi, F., Ueda, K., Takehara, T., Kasahara, A., Sasaki, Y., Hori, M., und Hayashi, N. (2003). Hepatitis c virus core protein differently regulates the jak-stat signaling pathway under interleukin-6 and interferon-gamma stimuli. *J Biol Chem*, 278(31):28562–71.
- Houldsworth, A., Metzner, M., Rossol, S., Shaw, S., Kaminski, E., Demaine, A. G., und Cramp, M. E. (2005). Polymorphisms in the il-12b gene and outcome of hcv infection. *J Interferon Cytokine Res*, 25(5):271–6.
- Hsu, M., Zhang, J., Flint, M., Logvinoff, C., Cheng-Mayer, C., Rice, C. M., und McKeating, J. A. (2003). Hepatitis c virus glycoproteins mediate ph-dependent cell entry of pseudotyped retroviral particles. *Proc Natl Acad Sci U S A*, 100(12):7271–6.
- Hu, K. Q., Vierling, J. M., und Redeker, A. G. (2001). Viral, host and interferon-related factors modulating the effect of interferon therapy for hepatitis c virus infection. *J Viral Hepat*, 8(1):1–18.
- Hughes, M. G., J., Rudy, C. K., Chong, T. W., Smith, R. L., Evans, H. L., Iezzoni, J. C., Sawyer, R. G., und Pruett, T. L. (2004). E2 quasispecies specificity of hepatitis c virus association with allografts immediately after liver transplantation. *Liver Transpl*, 10(2):208–16.
- Hugle, T., Fehrman, F., Bieck, E., Kohara, M., Krausslich, H. G., Rice, C. M., Blum, H. E., und Moradpour, D. (2001). The hepatitis c virus nonstructural protein 4b is an integral endoplasmic reticulum membrane protein. *Virology*, 284(1):70–81.
- Ide, Y., Zhang, L., Chen, M., Inchauspe, G., Bahl, C., Sasaguri, Y., und Padmanabhan, R. (1996). Characterization of the nuclear localization signal and subcellular distribution of hepatitis c virus nonstructural protein ns5a. *Gene*, 182(1-2):203–11.
- Ihle, J. N. (1996). Stats and mapks: obligate or opportunistic partners in signaling. *Bioessays*, 18(2):95–8.
- Inchauspe, G., Zebedee, S., Lee, D. H., Sugitani, M., Nasoff, M., und Prince, A. M. (1991). Genomic structure of the human prototype strain h of hepatitis c virus: comparison with american and japanese isolates. *Proc Natl Acad Sci U S A*, 88(22):10292–6.
- Isaacs, A. und Lindenmann, J. (1957). Virus interference. i. the interferon. *Proc R Soc Lond B Biol Sci*, 147(927):258–67.
- Isaacs, A. und Lindenmann, J. (1987). Virus interference. i. the interferon. by a. isaacs and j. lindenmann, 1957. *J Interferon Res*, 7(5):429–38.
- Ishihara, K. und Hirano, T. (2002). Molecular basis of the cell specificity of cytokine action. *Biochim Biophys Acta*, 1592(3):281–96.

- Jacobs, B. L. und Langland, J. O. (1998). Reovirus sigma 3 protein: dsrna binding and inhibition of rna-activated protein kinase. *Curr Top Microbiol Immunol*, 233(Pt 1):185–96.
- Jaeckel, E., Cornberg, M., Wedemeyer, H., Santantonio, T., Mayer, J., Zankel, M., Pastore, G., Dietrich, M., Trautwein, C., und Manns, M. P. (2001). Treatment of acute hepatitis c with interferon alfa-2b. *N Engl J Med*, 345(20):1452–7.
- Joo, M., Hahn, Y. S., Kwon, M., Sadikot, R. T., Blackwell, T. S., und Christman, J. W. (2005). Hepatitis c virus core protein suppresses nf-kappab activation and cyclooxygenase-2 expression by direct interaction with ikappab kinase beta. *J Virol*, 79(12):7648–57.
- Jung, M. C., Diepolder, H. M., und Pape, G. R. (1994). T cell recognition of hepatitis b and c viral antigens. *Eur J Clin Invest*, 24(10):641–50.
- Kaito, M., Watanabe, S., Tsukiyama-Kohara, K., Yamaguchi, K., Kobayashi, Y., Konishi, M., Yokoi, M., Ishida, S., Suzuki, S., und Kohara, M. (1994). Hepatitis c virus particle detected by immunoelectron microscopic study. *J Gen Virol*, 75 (Pt 7):1755–60.
- Kanazawa, Y., Hayashi, N., Mita, E., Li, T., Hagiwara, H., Kasahara, A., Fusamoto, H., und Kamada, T. (1994). Influence of viral quasispecies on effectiveness of interferon therapy in chronic hepatitis c patients. *Hepatology*, 20(5):1121–30.
- Kaplan, M., Gawrieh, S., Cotler, S. J., und Jensen, D. M. (2003). Neutralizing antibodies in hepatitis c virus infection: a review of immunological and clinical characteristics. *Gastroenterology*, 125(2):597–604.
- Kato, N., Hijikata, M., Ootsuyama, Y., Nakagawa, M., Ohkoshi, S., Sugimura, T., und Shimotohno, K. (1990). Molecular cloning of the human hepatitis c virus genome from japanese patients with non-a, non-b hepatitis. *Proc Natl Acad Sci U S A*, 87(24):9524–8.
- Kato, N., Lan, K. H., Ono-Nita, S. K., Shiratori, Y., und Omata, M. (1997). Hepatitis c virus nonstructural region 5a protein is a potent transcriptional activator. *J Virol*, 71(11):8856–9.
- Kato, N., Ootsuyama, Y., Ohkoshi, S., Nakazawa, T., Sekiya, H., Hijikata, M., und Shimotohno, K. (1992). Characterization of hypervariable regions in the putative envelope protein of hepatitis c virus. *Biochem Biophys Res Commun*, 189(1):119–27.
- Kato, N., Yoshida, H., Kioko Ono-Nita, S., Kato, J., Goto, T., Otsuka, M., Lan, K., Matsushima, K., Shiratori, Y., und Omata, M. (2000). Activation of intracellular signaling by hepatitis b and c viruses: C-viral core is the most potent signal inducer. *Hepatology*, 32(2):405–12.
- Kato, T., Date, T., Miyamoto, M., Furusaka, A., Tokushige, K., Mizokami, M., und Wakita, T. (2003). Efficient replication of the genotype 2a hepatitis c virus subgenomic replicon. *Gastroenterology*, 125(6):1808–17.
- Kato, T., Date, T., Miyamoto, M., Zhao, Z., Mizokami, M., und Wakita, T. (2005). Nonhepatic cell lines hela and 293 support efficient replication of the hepatitis c virus genotype 2a subgenomic replicon. *J Virol*, 79(1):592–6.

- Katze, M. G., He, Y., und Gale, M., J. (2002). Viruses and interferon: a fight for supremacy. *Nat Rev Immunol*, 2(9):675–87.
- Khabar, K. S., Al-Zoghaibi, F., Al-Ahdal, M. N., Murayama, T., Dhalla, M., Mukaida, N., Taha, M., Al-Sedairy, S. T., Siddiqui, Y., Kessie, G., und Matsushima, K. (1997). The alpha chemokine, interleukin 8, inhibits the antiviral action of interferon alpha. *J Exp Med*, 186(7):1077–85.
- Khan, S., Zimmermann, A., Basler, M., Groettrup, M., und Hengel, H. (2004). A cytomegalovirus inhibitor of gamma interferon signaling controls immunoproteasome induction. *J Virol*, 78(4):1831–42.
- Kim, J. L., Morgenstern, K. A., Griffith, J. P., Dwyer, M. D., Thomson, J. A., Murcko, M. A., Lin, C., und Caron, P. R. (1998). Hepatitis c virus ns3 rna helicase domain with a bound oligonucleotide: the crystal structure provides insights into the mode of unwinding. *Structure*, 6(1):89–100.
- Kim, W. H., Hong, F., Jaruga, B., Hu, Z., Fan, S., Liang, T. J., und Gao, B. (2001). Additive activation of hepatic nf-kappab by ethanol and hepatitis b protein x (hbx) or hcv core protein: involvement of tnf-alpha receptor 1-independent and -dependent mechanisms. *Faseb J*, 15(13):2551–3.
- Knapp, S., Yee, L. J., Frodsham, A. J., Hennig, B. J., Hellier, S., Zhang, L., Wright, M., Chiaramonte, M., Graves, M., Thomas, H. C., Hill, A. V., und Thursz, M. R. (2003). Polymorphisms in interferon-induced genes and the outcome of hepatitis c virus infection: roles of mxa, oas-1 and pkr. *Genes Immun*, 4(6):411–9.
- Kobayashi, Y., Watanabe, S., Konishi, M., Yokoi, M., Kakehashi, R., Kaito, M., Kondo, M., Hayashi, Y., Jomori, T., und Suzuki, S. (1993). Quantitation and typing of serum hepatitis c virus rna in patients with chronic hepatitis c treated with interferon-beta. *Hepatology*, 18(6):1319–25.
- Koch, J. O. und Bartenschlager, R. (1999). Modulation of hepatitis c virus ns5a hyperphosphorylation by nonstructural proteins ns3, ns4a, and ns4b. *J Virol*, 73(9):7138–46.
- Koizumi, K., Enomoto, N., Kurosaki, M., Murakami, T., Izumi, N., Marumo, F., und Sato, C. (1995). Diversity of quasispecies in various disease stages of chronic hepatitis c virus infection and its significance in interferon treatment. *Hepatology*, 22(1):30–5.
- Kuiken, C., Yusim, K., Boykin, L., und Richardson, R. (2005). The los alamos hepatitis c sequence database. *Bioinformatics*, 21(3):379–84.
- Kutay, U., Hartmann, E., und Rapoport, T. A. (1993). A class of membrane proteins with a c-terminal anchor. *Trends Cell Biol*, 3(3):72–5.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage t4. *Nature*, 227(5259):680–5.
- Lamarre, D., Anderson, P. C., Bailey, M., Beaulieu, P., Bolger, G., Bonneau, P., Bos, M., Cameron, D. R., Cartier, M., und Cordingley, M. G. e. (2003). An ns3 protease inhibitor with antiviral effects in humans infected with hepatitis c virus. *Nature*, 426(6963):186–9.

- Lan, K. H., Sheu, M. L., Hwang, S. J., Yen, S. H., Chen, S. Y., Wu, J. C., Wang, Y. J., Kato, N., Omata, M., Chang, F. Y., und Lee, S. D. (2002). Hcv ns5a interacts with p53 and inhibits p53-mediated apoptosis. *Oncogene*, 21(31):4801–11.
- Lanford, R. E., Guerra, B., Lee, H., Averett, D. R., Pfeiffer, B., Chavez, D., Notvall, L., und Bigger, C. (2003). Antiviral effect and virus-host interactions in response to alpha interferon, gamma interferon, poly(i)-poly(c), tumor necrosis factor alpha, and ribavirin in hepatitis c virus subgenomic replicons. *J Virol*, 77(2):1092–104.
- Langland, J. O., Pettiford, S., Jiang, B., und Jacobs, B. L. (1994). Products of the porcine group c rotavirus nsp3 gene bind specifically to double-stranded rna and inhibit activation of the interferon-induced protein kinase pkr. *J Virol*, 68(6):3821–9.
- Lauer, G. M. und Walker, B. D. (2001). Hepatitis c virus infection. *N Engl J Med*, 345(1):41–52.
- Lavillette, D., Tarr, A. W., Voisset, C., Donot, P., Bartosch, B., Bain, C., Patel, A. H., Dubuisson, J., Ball, J. K., und Cosset, F. L. (2005). Characterization of host-range and cell entry properties of the major genotypes and subtypes of hepatitis c virus. *Hepatology*, 41(2):265–274.
- Leaman, D. W., Leung, S., Li, X., und Stark, G. R. (1996). Regulation of stat-dependent pathways by growth factors and cytokines. *Faseb J*, 10(14):1578–88.
- Lei, M., Liu, Y., und Samuel, C. E. (1998). Adenovirus vai rna antagonizes the rna-editing activity of the adar adenosine deaminase. *Virology*, 245(2):188–96.
- Lennette, E. H. (1951). Interference between animal viruses. *Annu Rev Microbiol*, 5:277–94.
- Leonard, G. T. und Sen, G. C. (1996). Effects of adenovirus e1a protein on interferon-signaling. *Virology*, 224(1):25–33.
- Lerat, H., Berby, F., Trabaud, M. A., Vidalin, O., Major, M., Trepo, C., und Inchauspe, G. (1996). Specific detection of hepatitis c virus minus strand rna in hematopoietic cells. *J Clin Invest*, 97(3):845–51.
- Leung, S., Qureshi, S. A., Kerr, I. M., Darnell, J. E., J., und Stark, G. R. (1995). Role of stat2 in the alpha interferon signaling pathway. *Mol Cell Biol*, 15(3):1312–7.
- Lin, C., Lin, K., Luong, Y. P., Rao, B. G., Wei, Y. Y., Brennan, D. L., Fulghum, J. R., Hsiao, H. M., Ma, S., Maxwell, J. P., Cottrell, K. M., Perni, R. B., Gates, C. A., und Kwong, A. D. (2004). In vitro resistance studies of hepatitis c virus serine protease inhibitors, vx-950 and biln 2061: structural analysis indicates different resistance mechanisms. *J Biol Chem*, 279(17):17508–14.
- Lin, C., Lindenbach, B. D., Pragai, B. M., McCourt, D. W., und Rice, C. M. (1994). Processing in the hepatitis c virus e2-ns2 region: identification of p7 and two distinct e2-specific products with different c termini. *J Virol*, 68(8):5063–73.
- Lin, W., Choe, W. H., Hiasa, Y., Kamegaya, Y., Blackard, J. T., Schmidt, E. V., und Chung, R. T. (2005). Hepatitis c virus expression suppresses interferon signaling by degrading stat1. *Gastroenterology*, 128(4):1034–41.

- Liu, Y., Wolff, K. C., Jacobs, B. L., und Samuel, C. E. (2001). Vaccinia virus e3l interferon resistance protein inhibits the interferon-induced adenosine deaminase a-to-i editing activity. *Virology*, 289(2):378–87.
- Lohmann, V., Korner, F., Koch, J., Herian, U., Theilmann, L., und Bartenschlager, R. (1999). Replication of subgenomic hepatitis c virus rnas in a hepatoma cell line. *Science*, 285(5424):110–3.
- Lu, L., Pilot-Matias, T. J., Stewart, K. D., Randolph, J. T., Pithawalla, R., He, W., Huang, P. P., Klein, L. L., Mo, H., und Molla, A. (2004). Mutations conferring resistance to a potent hepatitis c virus serine protease inhibitor in vitro. *Antimicrob Agents Chemother*, 48(6):2260–6.
- Lu, Y., Wambach, M., Katze, M. G., und Krug, R. M. (1995). Binding of the influenza virus ns1 protein to double-stranded rna inhibits the activation of the protein kinase that phosphorylates the elf-2 translation initiation factor. *Virology*, 214(1):222–8.
- Löwer, J. (1998). Über die ergebnisse des stufenplanverfahrens zur verminderung des risikos von heptitis b-, heptitis c- und hiv-infektionen bei empfängern von erythrozytenkonzentraten. Technical Report BAnz. Nr. 63, Paul-Ehrlich-Institut.
- Macdonald, A. und Harris, M. (2004). Hepatitis c virus ns5a: tales of a promiscuous protein. *J Gen Virol*, 85(Pt 9):2485–502.
- Major, E. M., Rehermann, B., und Feinstone, S. M. (2001). Hepatitis c viruses. In Knipe, D. M., Howley, P. M., und Griffin, D. E., editors, *Fields Virology*, pages 1127–1161. Lippincott Williams & Wilkins, fourth edition edition.
- Majumder, M., Ghosh, A. K., Steele, R., Ray, R., und Ray, R. B. (2001). Hepatitis c virus ns5a physically associates with p53 and regulates p21/waf1 gene expression in a p53-dependent manner. *J Virol*, 75(3):1401–7.
- Majumder, M., Ghosh, A. K., Steele, R., Zhou, X. Y., Phillips, N. J., Ray, R., und Ray, R. B. (2002). Hepatitis c virus ns5a protein impairs tnf-mediated hepatic apoptosis, but not by an anti-fas antibody, in transgenic mice. *Virology*, 294(1):94–105.
- Mangia, A., Santoro, R., Minerva, N., Ricci, G. L., Carretta, V., Persico, M., Vinelli, F., Scotto, G., Bacca, D., Annese, M., Romano, M., Zechini, F., Sogari, F., Spirito, F., und Andriulli, A. (2005). Peginterferon alfa-2b and ribavirin for 12 vs. 24 weeks in hcv genotype 2 or 3. *N Engl J Med*, 352(25):2609–17.
- Martell, M., Esteban, J. I., Quer, J., Genesca, J., Weiner, A., Esteban, R., Guardia, J., und Gomez, J. (1992). Hepatitis c virus (hcv) circulates as a population of different but closely related genomes: quasispecies nature of hcv genome distribution. *J Virol*, 66(5):3225–9.
- Martinand, C., Montavon, C., Salehzada, T., Silhol, M., Lebleu, B., und Bisbal, C. (1999). Rnase 1 inhibitor is induced during human immunodeficiency virus type 1 infection and down regulates the 2-5a/rnase 1 pathway in human t cells. *J Virol*, 73(1):290–6.
- Matsumoto, M., Hwang, S. B., Jeng, K. S., Zhu, N., und Lai, M. M. (1996). Homotypic interaction and multimerization of hepatitis c virus core protein. *Virology*, 218(1):43–51.

- McHutchison, J. G., Gordon, S. C., Schiff, E. R., Shiffman, M. L., Lee, W. M., Rustgi, V. K., Goodman, Z. D., Ling, M. H., Cort, S., und Albrecht, J. K. (1998). Interferon alfa-2b alone or in combination with ribavirin as initial treatment for chronic hepatitis c. hepatitis interventional therapy group. *N Engl J Med*, 339(21):1485–92.
- McLauchlan, J. (2000). Properties of the hepatitis c virus core protein: a structural protein that modulates cellular processes. *J Viral Hepat*, 7(1):2–14.
- Medzhitov, R. (2001). Toll-like receptors and innate immunity. *Nat Rev Immunol*, 1(2):135–45.
- Medzhitov, R. und Janeway, C., J. (2000). Innate immunity. *N Engl J Med*, 343(5):338–44.
- Meisel, H., Reip, A., Faltus, B., Lu, M., Porst, H., Wiese, M., Roggendorf, M., und Kruger, D. H. (1995). Transmission of hepatitis c virus to children and husbands by women infected with contaminated anti-d immunoglobulin. *Lancet*, 345(8959):1209–11.
- Mercer, D. F., Schiller, D. E., Elliott, J. F., Douglas, D. N., Hao, C., Rinfret, A., Addison, W. R., Fischer, K. P., Churchill, T. A., Lakey, J. R., Tyrrell, D. L., und Kneteman, N. M. (2001). Hepatitis c virus replication in mice with chimeric human livers. *Nat Med*, 7(8):927–33.
- Mihm, S., Fayyazi, A., Hartmann, H., und Ramadori, G. (1997). Analysis of histopathological manifestations of chronic hepatitis c virus infection with respect to virus genotype. *Hepatology*, 25(3):735–9.
- Miller, D. M., Rahill, B. M., Boss, J. M., Lairmore, M. D., Durbin, J. E., Waldman, J. W., und Sedmak, D. D. (1998). Human cytomegalovirus inhibits major histocompatibility complex class ii expression by disruption of the jak/stat pathway. *J Exp Med*, 187(5):675–83.
- Miller, K., McArdle, S., Gale, M. J., J., Geller, D. A., Tenover, B., Hiscott, J., Gretch, D. R., und Polyak, S. J. (2004). Effects of the hepatitis c virus core protein on innate cellular defense pathways. *J Interferon Cytokine Res*, 24(7):391–402.
- Miyamoto, H., Okamoto, H., Sato, K., Tanaka, T., und Mishiro, S. (1992). Extraordinarily low density of hepatitis c virus estimated by sucrose density gradient centrifugation and the polymerase chain reaction. *J Gen Virol*, 73 (Pt 3):715–8.
- Mizushima, H., Hijikata, M., Asabe, S., Hirota, M., Kimura, K., und Shimotohno, K. (1994). Two hepatitis c virus glycoprotein e2 products with different c termini. *J Virol*, 68(10):6215–22.
- Monazahian, M., Bohme, I., Bonk, S., Koch, A., Scholz, C., Grethe, S., und Thomssen, R. (1999). Low density lipoprotein receptor as a candidate receptor for hepatitis c virus. *J Med Virol*, 57(3):223–9.
- Montalbano, M. und Neff, G. W. (2005). An update in liver transplantation in patients with hepatitis b and hepatitis c. *Minerva Gastroenterol Dietol*, 51(1):109–26.
- Moradpour, D., Englert, C., Wakita, T., und Wands, J. R. (1996). Characterization of cell lines allowing tightly regulated expression of hepatitis c virus core protein. *Virology*, 222(1):51–63.

- Moradpour, D., Gosert, R., Egger, D., Penin, F., Blum, H. E., und Bienz, K. (2003). Membrane association of hepatitis c virus nonstructural proteins and identification of the membrane alteration that harbors the viral replication complex. *Antiviral Res*, 60(2):103–9.
- Mosley, J. W., Operkalski, E. A., Tobler, L. H., Andrews, W. W., Phelps, B., Dockter, J., Giachetti, C., und Busch, M. P. (2005). Viral and host factors in early hepatitis c virus infection. *Hepatology*, 42(1):86–92.
- Mueller, T., Mas-Marques, A., Sarrazin, C., Wiese, M., Halangk, J., Witt, H., Ahlenstiel, G., Spengler, U., Goebel, U., Wiedenmann, B., Schreier, E., und Berg, T. (2004). Influence of interleukin 12b (il12b) polymorphisms on spontaneous and treatment-induced recovery from hepatitis c virus infection. *J Hepatol*, 41(4):652–8.
- Murashima, S., Kumashiro, R., Ide, T., Miyajima, I., Hino, T., Koga, Y., Ishii, K., Ueno, T., Sakisaka, S., und Sata, M. (2000). Effect of interferon treatment on serum 2',5'-oligoadenylate synthetase levels in hepatitis c-infected patients. *J Med Virol*, 62(2):185–90.
- Murashima, S., Kumashiro, R., Ide, T., und Sata, M. (1999). Hcv ns5a 2209-2248 amino acid substitutions affect serum 2,5as activities. *J Hepatol*, 31(2):382.
- Nakamoto, Y., Kaneko, S., Ohno, H., Honda, M., Unoura, M., Murakami, S., und Kobayashi, K. (1996). B-cell epitopes in hypervariable region 1 of hepatitis c virus obtained from patients with chronic persistent hepatitis. *J Med Virol*, 50(1):35–41.
- Namba, K., Naka, K., Dansako, H., Nozaki, A., Ikeda, M., Shiratori, Y., Shimotohno, K., und Kato, N. (2004). Establishment of hepatitis c virus replicon cell lines possessing interferon-resistant phenotype. *Biochem Biophys Res Commun*, 323(1):299–309.
- Neddermann, P., Quintavalle, M., Di Pietro, C., Clementi, A., Cerretani, M., Altamura, S., Bartholomew, L., und De Francesco, R. (2004). Reduction of hepatitis c virus ns5a hyperphosphorylation by selective inhibition of cellular kinases activates viral rna replication in cell culture. *J Virol*, 78(23):13306–14.
- Nelson, N., Marks, M. S., Driggers, P. H., und Ozato, K. (1993). Interferon consensus sequence-binding protein, a member of the interferon regulatory factor family, suppresses interferon-induced gene transcription. *Mol Cell Biol*, 13(1):588–99.
- Neumann, A. U., Lam, N. P., Dahari, H., Gretch, D. R., Wiley, T. E., Layden, T. J., und Perelson, A. S. (1998). Hepatitis c viral dynamics in vivo and the antiviral efficacy of interferon-alpha therapy. *Science*, 282(5386):103–7.
- Niwa, H., Yamamura, K., und Miyazaki, J. (1991). Efficient selection for high-expression transfectants with a novel eukaryotic vector. *Gene*, 108(2):193–9.
- Nolandt, O., Kern, V., Muller, H., Pfaff, E., Theilmann, L., Welker, R., und Krausslich, H. G. (1997). Analysis of hepatitis c virus core protein interaction domains. *J Gen Virol*, 78 (Pt 6):1331–40.
- Nouri-Aria, K. T., Sallie, R., Mizokami, M., Portmann, B. C., und Williams, R. (1995). Intrahepatic expression of hepatitis c virus antigens in chronic liver disease. *J Pathol*, 175(1):77–83.

- Nousbaum, J., Polyak, S. J., Ray, S. C., Sullivan, D. G., Larson, A. M., Carithers, R. L., J., und Gretch, D. R. (2000). Prospective characterization of full-length hepatitis c virus ns5a quasispecies during induction and combination antiviral therapy. *J Virol*, 74(19):9028–38.
- Offergeld, R., Stark, K., und Hamouda, O. (2003). Infektionen bei blutspendern. *Ges.forsch. Ges.schutz*, 46:775–9.
- Okada, S., Akahane, Y., Suzuki, H., Okamoto, H., und Mishiro, S. (1992). The degree of variability in the amino terminal region of the e2/ns1 protein of hepatitis c virus correlates with responsiveness to interferon therapy in viremic patients. *Hepatology*, 16(3):619–24.
- Okuda, M., Hino, K., Korenaga, M., Yamaguchi, Y., Katoh, Y., und Okita, K. (1999). Differences in hypervariable region 1 quasispecies of hepatitis c virus in human serum, peripheral blood mononuclear cells, and liver. *Hepatology*, 29(1):217–22.
- Oleksyk, T. K., Thio, C. L., Truelove, A. L., Goedert, J. J., Donfield, S. M., Kirk, G. D., Thomas, D. L., O'Brien, S. J., und Smith, M. W. (2005). Single nucleotide polymorphisms and haplotypes in the il10 region associated with hcv clearance. *Genes Immun*, 6(4):347–57.
- Palitzsch, K. D., Hottentrager, B., Schlottmann, K., Frick, E., Holstege, A., Scholmerich, J., und Jilg, W. (1999). Prevalence of antibodies against hepatitis c virus in the adult german population. *Eur J Gastroenterol Hepatol*, 11(11):1215–20.
- Pascu, M., Martus, P., Hohne, M., Wiedenmann, B., Hopf, U., Schreier, E., und Berg, T. (2004). Sustained virological response in hepatitis c virus type 1b infected patients is predicted by the number of mutations within the ns5a-isdr: a meta-analysis focused on geographical differences. *Gut*, 53(9):1345–51.
- Paterson, M., Laxton, C. D., Thomas, H. C., Ackrill, A. M., und Foster, G. R. (1999). Hepatitis c virus ns5a protein inhibits interferon antiviral activity, but the effects do not correlate with clinical response. *Gastroenterology*, 117(5):1187–97.
- Pawlotsky, J. M. (2003). Mechanisms of antiviral treatment efficacy and failure in chronic hepatitis c. *Antiviral Res*, 59(1):1–11.
- Pawlotsky, J. M., Germanidis, G., Neumann, A. U., Pellerin, M., Frainais, P. O., und Dhumeaux, D. (1998a). Interferon resistance of hepatitis c virus genotype 1b: relationship to nonstructural 5a gene quasispecies mutations. *J Virol*, 72(4):2795–805.
- Pawlotsky, J. M., Pellerin, M., Bouvier, M., Roudot-Thoraval, F., Germanidis, G., Bastie, A., Darthuy, F., Remire, J., Soussy, C. J., und Dhumeaux, D. (1998b). Genetic complexity of the hypervariable region 1 (hvr1) of hepatitis c virus (hcv): influence on the characteristics of the infection and responses to interferon alfa therapy in patients with chronic hepatitis c. *J Med Virol*, 54(4):256–64.
- Pflugheber, J., Frederickson, B., Sumpter, R., J., Wang, C., Ware, F., Sodora, D. L., und Gale, M., J. (2002). Regulation of pkr and irf-1 during hepatitis c virus rna replication. *Proc Natl Acad Sci U S A*, 99(7):4650–5.
- Piai, G., Rocco, P., Tartaglione, M. T., Ciarleglio, A., Focareta, R., Grimaldi, E., Ievoli, F., Iuliano, D., Pacelli, M., und Forte, G. (2003). Triple (interferon, ribavirin,

- amantadine) versus double (interferon, ribavirin) re-therapy for interferon relapser genotype 1b hcv chronic active hepatitis patients. *Hepatol Res*, 25(4):355–363.
- Pileri, P., Uematsu, Y., Campagnoli, S., Galli, G., Falugi, F., Petracca, R., Weiner, A. J., Houghton, M., Rosa, D., Grandi, G., und Abrignani, S. (1998). Binding of hepatitis c virus to cd81. *Science*, 282(5390):938–41.
- Podevin, P., Sabile, A., Gajardo, R., Delhem, N., Abadie, A., Lozach, P. Y., Beretta, L., und Brechot, C. (2001). Expression of hepatitis c virus ns5a natural mutants in a hepatocytic cell line inhibits the antiviral effect of interferon in a pkr-independent manner. *Hepatology*, 33(6):1503–11.
- Polyak, S. J., Khabar, K. S., Paschal, D. M., Ezelle, H. J., Duverlie, G., Barber, G. N., Levy, D. E., Mukaida, N., und Gretch, D. R. (2001a). Hepatitis c virus nonstructural 5a protein induces interleukin-8, leading to partial inhibition of the interferon-induced antiviral response. *J Virol*, 75(13):6095–106.
- Polyak, S. J., Khabar, K. S., Rezeiq, M., und Gretch, D. R. (2001b). Elevated levels of interleukin-8 in serum are associated with hepatitis c virus infection and resistance to interferon therapy. *J Virol*, 75(13):6209–11.
- Polyak, S. J., Paschal, D. M., McArdle, S., Gale, M. J., J., Moradpour, D., und Gretch, D. R. (1999). Characterization of the effects of hepatitis c virus nonstructural 5a protein expression in human cell lines and on interferon-sensitive virus replication. *Hepatology*, 29(4):1262–71.
- Polywka, S., Feucht, H., Zollner, B., und Laufs, R. (1997). Hepatitis c virus infection in pregnancy and the risk of mother-to-child transmission. *Eur J Clin Microbiol Infect Dis*, 16(2):121–4.
- Powell, P. P., Dixon, L. K., und Parkhouse, R. M. (1996). An ikappab homolog encoded by african swine fever virus provides a novel mechanism for downregulation of proinflammatory cytokine responses in host macrophages. *J Virol*, 70(12):8527–33.
- Power, J. P., Lawlor, E., Davidson, F., Holmes, E. C., Yap, P. L., und Simmonds, P. (1995). Molecular epidemiology of an outbreak of infection with hepatitis c virus in recipients of anti-d immunoglobulin. *Lancet*, 345(8959):1211–3.
- Prince, A. M., Brotman, B., Huima, T., Pascual, D., Jaffery, M., und Inchauspe, G. (1992). Immunity in hepatitis c infection. *J Infect Dis*, 165(3):438–43.
- Prince, A. M., Huima-Byron, T., Parker, T. S., und Levine, D. M. (1996). Visualization of hepatitis c virions and putative defective interfering particles isolated from low-density lipoproteins. *J Viral Hepat*, 3(1):11–7.
- Proud, C. G. (1995). Pkr: a new name and new roles. *Trends Biochem Sci*, 20(6):241–6.
- Radun, D. und Hamouda, O. (2004). Epidemiologie der hepatitis b und c in deutschland. *Med Welt*, 55(7-8):206–10.
- Ralston, R., Thudium, K., Berger, K., Kuo, C., Gervase, B., Hall, J., Selby, M., Kuo, G., Houghton, M., und Choo, Q. L. (1993). Characterization of hepatitis c virus envelope glycoprotein complexes expressed by recombinant vaccinia viruses. *J Virol*, 67(11):6753–61.

- Ray, R. B., Lagging, L. M., Meyer, K., und Ray, R. (1996). Hepatitis c virus core protein cooperates with ras and transforms primary rat embryo fibroblasts to tumorigenic phenotype. *J Virol*, 70(7):4438–43.
- Reed, K. E. und Rice, C. M. (1999). Identification of the major phosphorylation site of the hepatitis c virus h strain ns5a protein as serine 2321. *J Biol Chem*, 274(39):28011–8.
- Rice, C. M. (2003). Virology: fresh assault on hepatitis c. *Nature*, 426(6963):129–31.
- Rios, M., Munoz, M., Torrence, P. F., und Spencer, E. (1995). Effect of interferon and 2',5'-oligoadenylates on rotavirus rna synthesis. *Antiviral Res*, 26(2):133–43.
- Roberts, R. M., Liu, L., Guo, Q., Leaman, D., und Bixby, J. (1998). The evolution of the type i interferons. *J Interferon Cytokine Res*, 18(10):805–16.
- Robertson, B., Myers, G., Howard, C., Brettin, T., Bukh, J., Gaschen, B., Gojobori, T., Maertens, G., Mizokami, M., und Nainan, O. e. (1998). Classification, nomenclature, and database development for hepatitis c virus (hcv) and related viruses: proposals for standardization. international committee on virus taxonomy. *Arch Virol*, 143(12):2493–503.
- Robertson, H. D. und Mathews, M. B. (1996). The regulation of the protein kinase pkr by rna. *Biochimie*, 78(11-12):909–14.
- Roccasecca, R., Ansuini, H., Vitelli, A., Meola, A., Scarselli, E., Acali, S., Pezzanera, M., Ercole, B. B., McKeating, J., Yagnik, A., Lahm, A., Tramontano, A., Cortese, R., und Nicosia, A. (2003). Binding of the hepatitis c virus e2 glycoprotein to cd81 is strain specific and is modulated by a complex interplay between hypervariable regions 1 and 2. *J Virol*, 77(3):1856–67.
- Ronco, L. V., Karpova, A. Y., Vidal, M., und Howley, P. M. (1998). Human papillomavirus 16 e6 oncoprotein binds to interferon regulatory factor-3 and inhibits its transcriptional activity. *Genes Dev*, 12(13):2061–72.
- Rosa, D., Campagnoli, S., Moretto, C., Guenzi, E., Cousens, L., Chin, M., Dong, C., Weiner, A. J., Lau, J. Y., Choo, Q. L., Chien, D., Pileri, P., Houghton, M., und Abrignani, S. (1996). A quantitative test to estimate neutralizing antibodies to the hepatitis c virus: cytofluorimetric assessment of envelope glycoprotein 2 binding to target cells. *Proc Natl Acad Sci U S A*, 93(5):1759–63.
- Rosmorduc, O., Sirma, H., Soussan, P., Gordien, E., Lebon, P., Horisberger, M., Brechot, C., und Kremsdorff, D. (1999). Inhibition of interferon-inducible mxa protein expression by hepatitis b virus capsid protein. *J Gen Virol*, 80 (Pt 5):1253–62.
- Rost, B., Casadio, R., Fariselli, P., und Sander, C. (1995). Transmembrane helices predicted at 95% accuracy. *Protein Sci*, 4(3):521–33.
- Roy, S., Katze, M. G., Parkin, N. T., Edery, I., Hovanessian, A. G., und Sonenberg, N. (1990). Control of the interferon-induced 68-kilodalton protein kinase by the hiv-1 tat gene product. *Science*, 247(4947):1216–9.
- Sakai, A., Claire, M. S., Faulk, K., Govindarajan, S., Emerson, S. U., Purcell, R. H., und Bukh, J. (2003). The p7 polypeptide of hepatitis c virus is critical for infectivity and contains functionally important genotype-specific sequences. *Proc Natl Acad Sci U S A*, 100(20):11646–51.

- Sakamoto, M., Akahane, Y., Tsuda, F., Tanaka, T., Woodfield, D. G., und Okamoto, H. (1994). Entire nucleotide sequence and characterization of a hepatitis c virus of genotype v/3a. *J Gen Virol*, 75 (Pt 7):1761–8.
- Sambrook, J. und Russell, D. (2001). *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press.
- Samuel, C. E. (2001). Antiviral actions of interferons. *Clin Microbiol Rev*, 14(4):778–809, table of contents.
- Sanger, F., Nicklen, S., und Coulson, A. R. (1977). Dna sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci U S A*, 74(12):5463–7.
- Santolini, E., Migliaccio, G., und La Monica, N. (1994). Biosynthesis and biochemical properties of the hepatitis c virus core protein. *J Virol*, 68(6):3631–41.
- Santolini, E., Pacini, L., Fipaldini, C., Migliaccio, G., und Monica, N. (1995). The ns2 protein of hepatitis c virus is a transmembrane polypeptide. *J Virol*, 69(12):7461–71.
- Sato, M., Hata, N., Asagiri, M., Nakaya, T., Taniguchi, T., und Tanaka, N. (1998). Positive feedback regulation of type i ifn genes by the ifn-inducible transcription factor irf-7. *FEBS Lett*, 441(1):106–10.
- Scarselli, E., Ansuini, H., Cerino, R., Roccasecca, R. M., Acali, S., Filocamo, G., Traboni, C., Nicosia, A., Cortese, R., und Vitelli, A. (2002). The human scavenger receptor class b type i is a novel candidate receptor for the hepatitis c virus. *Embo J*, 21(19):5017–25.
- Schmidt-Mende, J., Bieck, E., Hugle, T., Penin, F., Rice, C. M., Blum, H. E., und Moradpour, D. (2001). Determinants for membrane association of the hepatitis c virus rna-dependent rna polymerase. *J Biol Chem*, 276(47):44052–63.
- Schreier, E., Radun, D., Neuhauser, H., und Stark, K. (2003). Hepatitis c. Technical report, Robert Koch-Institut.
- Selby, M. J., Choo, Q. L., Berger, K., Kuo, G., Glazer, E., Eckart, M., Lee, C., Chien, D., Kuo, C., und Houghton, M. (1993). Expression, identification and subcellular localization of the proteins encoded by the hepatitis c viral genome. *J Gen Virol*, 74 (Pt 6):1103–13.
- Selby, M. J., Glazer, E., Masiarz, F., und Houghton, M. (1994). Complex processing and protein:protein interactions in the e2:ns2 region of hcv. *Virology*, 204(1):114–22.
- Sharp, T. V., Schwemmle, M., Jeffrey, I., Laing, K., Mellor, H., Proud, C. G., Hille, K., und Clemens, M. J. (1993). Comparative analysis of the regulation of the interferon-inducible protein kinase pkr by epstein-barr virus rnas eber-1 and eber-2 and adenovirus vai rna. *Nucleic Acids Res*, 21(19):4483–90.
- Shepard, C. W., Finelli, L., und Alter, M. J. (2005). Global epidemiology of hepatitis c virus infection. *Lancet Infect Dis*, 5(9):558–67.
- Shih, C. M., Lo, S. J., Miyamura, T., Chen, S. Y., und Lee, Y. H. (1993). Suppression of hepatitis b virus expression and replication by hepatitis c virus core protein in huh-7 cells. *J Virol*, 67(10):5823–32.

- Shimakami, T., Hijikata, M., Luo, H., Ma, Y. Y., Kaneko, S., Shimotohno, K., und Murakami, S. (2004). Effect of interaction between hepatitis c virus ns5a and ns5b on hepatitis c virus rna replication with the hepatitis c virus replicon. *J Virol*, 78(6):2738–48.
- Shimizu, Y. K., Igarashi, H., Kiyoohara, T., Cabezon, T., Farci, P., Purcell, R. H., und Yoshikura, H. (1996). A hyperimmune serum against a synthetic peptide corresponding to the hypervariable region 1 of hepatitis c virus can prevent viral infection in cell cultures. *Virology*, 223(2):409–12.
- Shimizu, Y. K., Igarashi, H., Kiyoohara, T., Shapiro, M., Wong, D. C., Purcell, R. H., und Yoshikura, H. (1998). Infection of a chimpanzee with hepatitis c virus grown in cell culture. *J Gen Virol*, 79 (Pt 6):1383–6.
- Shirota, Y., Luo, H., Qin, W., Kaneko, S., Yamashita, T., Kobayashi, K., und Murakami, S. (2002). Hepatitis c virus (hcv) ns5a binds rna-dependent rna polymerase (rdrp) ns5b and modulates rna-dependent rna polymerase activity. *J Biol Chem*, 277(13):11149–55.
- Simmonds, P. (2004). Genetic diversity and evolution of hepatitis c virus—15 years on. *J Gen Virol*, 85(Pt 11):3173–88.
- Simmonds, P., Alberti, A., Alter, H. J., Bonino, F., Bradley, D. W., Brechot, C., Brouwer, J. T., Chan, S. W., Chayama, K., Chen, D. S., und et al. (1994). A proposed system for the nomenclature of hepatitis c viral genotypes. *Hepatology*, 19(5):1321–4.
- Smith, D. B., Mellor, J., Jarvis, L. M., Davidson, F., Kolberg, J., Urdea, M., Yap, P. L., und Simmonds, P. (1995). Variation of the hepatitis c virus 5' non-coding region: implications for secondary structure, virus detection and typing. the international hcv collaborative study group. *J Gen Virol*, 76 (Pt 7):1749–61.
- Smith, J. A., Schmeichel, S. C., Williams, B. R., Silverman, R. H., und Schiff, L. A. (2005). Involvement of the interferon-regulated antiviral proteins pkr and rnase l in reovirus-induced shutoff of cellular translation. *J Virol*, 79(4):2240–50.
- Song, J., Nagano-Fujii, M., Wang, F., Florese, R., Fujita, T., Ishido, S., und Hotta, H. (2000). Nuclear localization and intramolecular cleavage of n-terminally deleted ns5a protein of hepatitis c virus. *Virus Res*, 69(2):109–17.
- Srinivas, R. V., Ray, R. B., Meyer, K., und Ray, R. (1996). Hepatitis c virus core protein inhibits human immunodeficiency virus type 1 replication. *Virus Res*, 45(2):87–92.
- Stark, G. R., Kerr, I. M., Williams, B. R., Silverman, R. H., und Schreiber, R. D. (1998). How cells respond to interferons. *Annu Rev Biochem*, 67:227–64.
- Sumpter, R., Loo, Y. M., Foy, E., Li, K., Yoneyama, M., Fujita, T., Lemon, S. M., und Gale, M., J. (2005). Regulating intracellular antiviral defense and permissiveness to hepatitis c virus rna replication through a cellular rna helicase, rig-i. *J Virol*, 79(5):2689–99.
- Sun, J., Li, K., Shata, M. T., und Chan, T. S. (2004). The immunologic basis for hepatitis c infection. *Curr Opin Gastroenterol*, 20(6):598–602.
- Suzuki, F., Arase, Y., Suzuki, Y., Tsubota, A., Akuta, N., Hosaka, T., Someya, T., Kobayashi, M., Saitoh, S., Ikeda, K., Kobayashi, M., Matsuda, M., Takagi, K., Satoh,

- J., und Kumada, H. (2004). Single nucleotide polymorphism of the mxa gene promoter influences the response to interferon monotherapy in patients with hepatitis c viral infection. *J Viral Hepat*, 11(3):271–6.
- Symons, J. A., Alcami, A., und Smith, G. L. (1995). Vaccinia virus encodes a soluble type i interferon receptor of novel structure and broad species specificity. *Cell*, 81(4):551–60.
- Takahashi, K., Kishimoto, S., Yoshizawa, H., Okamoto, H., Yoshikawa, A., und Mishiro, S. (1992). p26 protein and 33-nm particle associated with nucleocapsid of hepatitis c virus recovered from the circulation of infected hosts. *Virology*, 191(1):431–4.
- Talon, J., Horvath, C. M., Polley, R., Basler, C. F., Muster, T., Palese, P., und Garcia-Sastre, A. (2000). Activation of interferon regulatory factor 3 is inhibited by the influenza a virus ns1 protein. *J Virol*, 74(17):7989–96.
- Tan, S. L. und Katze, M. G. (1999). The emerging role of the interferon-induced pkr protein kinase as an apoptotic effector: a new face of death? *J Interferon Cytokine Res*, 19(6):543–54.
- Tan, S. L. und Katze, M. G. (2001). How hepatitis c virus counteracts the interferon response: the jury is still out on ns5a. *Virology*, 284(1):1–12.
- Tan, S. L., Nakao, H., He, Y., Vijaysri, S., Neddermann, P., Jacobs, B. L., Mayer, B. J., und Katze, M. G. (1999). Ns5a, a nonstructural protein of hepatitis c virus, binds growth factor receptor-bound protein 2 adaptor protein in a src homology 3 domain/ligand-dependent manner and perturbs mitogenic signaling. *Proc Natl Acad Sci U S A*, 96(10):5533–8.
- Taniguchi, S., Okamoto, H., Sakamoto, M., Kojima, M., Tsuda, F., Tanaka, T., Mune-kata, E., Muchmore, E. E., Peterson, D. A., und Mishiro, S. (1993). A structurally flexible and antigenically variable n-terminal domain of the hepatitis c virus e2/ns1 protein: implication for an escape from antibody. *Virology*, 195(1):297–301.
- Taniguchi, T. und Takaoka, A. (2002). The interferon-alpha/beta system in antiviral responses: a multimodal machinery of gene regulation by the irf family of transcription factors. *Curr Opin Immunol*, 14(1):111–6.
- Taylor, D. R. (2000). Hepatitis c virus: evasion of the interferon-induced antiviral response. *J Mol Med*, 78(4):182–90.
- Taylor, D. R., Shi, S. T., Romano, P. R., Barber, G. N., und Lai, M. M. (1999). Inhibition of the interferon-inducible protein kinase pkr by hcv e2 protein. *Science*, 285(5424):107–10.
- Tellinghuisen, T. L., Marcotrigiano, J., Gorbalyena, A. E., und Rice, C. M. (2004). The ns5a protein of hepatitis c virus is a zinc metalloprotein. *J Biol Chem*, 279(47):48576–87.
- Thompson, J. D., Higgins, D. G., und Gibson, T. J. (1994). Clustal w: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res*, 22(22):4673–80.

- Thomson, M. und Liang, T. J. (2000). Molecular biology of hepatitis c virus. In Liang, T. und Hoofnagle, J., editors, *Hepatitis C*, Biomedical Research Reports. Academic Press, London.
- Tsukiyama-Kohara, K., Iizuka, N., Kohara, M., und Nomoto, A. (1992). Internal ribosome entry site within hepatitis c virus rna. *J Virol*, 66(3):1476–83.
- van den Broek, M. F., Muller, U., Huang, S., Zinkernagel, R. M., und Aguet, M. (1995). Immune defence in mice lacking type i and/or type ii interferon receptors. *Immunol Rev*, 148:5–18.
- Vona, G., Tuveri, R., Delpuech, O., Vallet, A., Canioni, D., Ballardini, G., Baptiste Trabut, J., Le Bail, B., Nalpas, B., Carnot, F., Pol, S., Brechot, C., und Thiers, V. (2004). Intrahepatic hepatitis c virus rna quantification in microdissected hepatocytes. *J Hepatol*, 40(4):682–8.
- Wakita, T., Pietschmann, T., Kato, T., Date, T., Miyamoto, M., Zhao, Z., Murthy, K., Habermann, A., Krausslich, H. G., Mizokami, M., Bartenschlager, R., und Liang, T. J. (2005). Production of infectious hepatitis c virus in tissue culture from a cloned viral genome. *Nat Med*.
- Wang, C., Sarnow, P., und Siddiqui, A. (1993). Translation of human hepatitis c virus rna in cultured cells is mediated by an internal ribosome-binding mechanism. *J Virol*, 67(6):3338–44.
- Wang, J. T., Sheu, J. C., Lin, J. T., Wang, T. H., und Chen, D. S. (1992). Detection of replicative form of hepatitis c virus rna in peripheral blood mononuclear cells. *J Infect Dis*, 166(5):1167–9.
- Watashi, K., Hijikata, M., Hosaka, M., Yamaji, M., und Shimotohno, K. (2003). Cyclosporin a suppresses replication of hepatitis c virus genome in cultured hepatocytes. *Hepatology*, 38(5):1282–8.
- Weihua, X., Ramanujam, S., Lindner, D. J., Kudaravalli, R. D., Freund, R., und Kalvakolanu, D. V. (1998). The polyoma virus t antigen interferes with interferon-inducible gene expression. *Proc Natl Acad Sci U S A*, 95(3):1085–90.
- Westaway, E. G., Mackenzie, J. M., Kenney, M. T., Jones, M. K., und Khromykh, A. A. (1997). Ultrastructure of kunjin virus-infected cells: colocalization of ns1 and ns3 with double-stranded rna, and of ns2b with ns3, in virus-induced membrane structures. *J Virol*, 71(9):6650–61.
- Wiese, M., Berr, F., Lafrenz, M., Porst, H., und Oesen, U. (2000). Low frequency of cirrhosis in a hepatitis c (genotype 1b) single-source outbreak in germany: a 20-year multicenter study. *Hepatology*, 32(1):91–6.
- Winston, L. A. und Hunter, T. (1996). Intracellular signalling: putting jaks on the kinase map. *Curr Biol*, 6(6):668–71.
- Wormald, S. und Hilton, D. J. (2004). Inhibitors of cytokine signal transduction. *J Biol Chem*, 279(2):821–4.
- Xie, Z. C., Rieu-Boj, J. I., Lasarte, J. J., Guillen, J., Su, J. H., Civeira, M. P., und Prieto, J. (1998). Transmission of hepatitis c virus infection to tree shrews. *Virology*, 244(2):513–20.

- Xu, Z., Choi, J., Lu, W., und Ou, J. H. (2003). Hepatitis c virus f protein is a short-lived protein associated with the endoplasmic reticulum. *J Virol*, 77(2):1578–83.
- Yao, N., Hesson, T., Cable, M., Hong, Z., Kwong, A. D., Le, H. V., und Weber, P. C. (1997). Structure of the hepatitis c virus rna helicase domain. *Nat Struct Biol*, 4(6):463–7.
- Yasui, K., Wakita, T., Tsukiyama-Kohara, K., Funahashi, S. I., Ichikawa, M., Kajita, T., Moradpour, D., Wands, J. R., und Kohara, M. (1998). The native form and maturation process of hepatitis c virus core protein. *J Virol*, 72(7):6048–55.
- Yatsuhashi, H., Fujino, T., Matsumoto, T., Inoue, O., Koga, M., und Yano, M. (1999). Immunohistochemical analysis of hepatic interferon alpha-beta receptor level: relationship between receptor expression and response to interferon therapy in patients with chronic hepatitis c. *J Hepatol*, 30(6):995–1003.
- Yokoyama, S., Ohishi, N., Shamoto, M., Watanabe, Y., und Yagi, K. (1997). Isolation and expression of rat interferon beta gene and growth-inhibitory effect of its expression on rat glioma cells. *Biochem Biophys Res Commun*, 232(3):698–701.
- Yoneyama, M., Kikuchi, M., Natsukawa, T., Shinobu, N., Imaizumi, T., Miyagishi, M., Taira, K., Akira, S., und Fujita, T. (2004). The rna helicase rig-i has an essential function in double-stranded rna-induced innate antiviral responses. *Nat Immunol*, 5(7):730–7.
- Yoneyama, M., Suhara, W., Fukuhara, Y., Fukuda, M., Nishida, E., und Fujita, T. (1998). Direct triggering of the type i interferon system by virus infection: activation of a transcription factor complex containing irf-3 and cbp/p300. *Embo J*, 17(4):1087–95.
- Yoneyama, M., Suhara, W., Fukuhara, Y., Sato, M., Ozato, K., und Fujita, T. (1996). Autocrine amplification of type i interferon gene expression mediated by interferon stimulated gene factor 3 (isgf3). *J Biochem (Tokyo)*, 120(1):160–9.
- Yoshida, H., Kato, N., Shiratori, Y., Otsuka, M., Maeda, S., Kato, J., und Omata, M. (2001). Hepatitis c virus core protein activates nuclear factor kappa b-dependent signaling through tumor necrosis factor receptor-associated factor. *J Biol Chem*, 276(19):16399–405.
- Young, D. F., Didcock, L., Goodbourn, S., und Randall, R. E. (2000). Paramyxoviridae use distinct virus-specific mechanisms to circumvent the interferon response. *Virology*, 269(2):383–90.
- Zech, B., Kurtenbach, A., Krieger, N., Strand, D., Blencke, S., Morbitzer, M., Salassidis, K., Cotten, M., Wissing, J., Obert, S., Bartenschlager, R., Herget, T., und Daub, H. (2003). Identification and characterization of amphiphysin ii as a novel cellular interaction partner of the hepatitis c virus ns5a protein. *J Gen Virol*, 84(Pt 3):555–60.
- Zeuzem, S. (2004). Standard therapie der akuten und chronischen hepatitis c. *Z Gastroenterol*, 42:714–719.
- Zeuzem, S., Hultcrantz, R., Bourliere, M., Goeser, T., Marcellin, P., Sanchez-Tapias, J., Sarrazin, C., Harvey, J., Brass, C., und Albrecht, J. (2004). Peginterferon alfa-2b plus ribavirin for treatment of chronic hepatitis c in previously untreated patients infected with hcv genotypes 2 or 3. *J Hepatol*, 40(6):993–9.

- Zeuzem, S., Lee, J. H., und Roth, W. K. (1997). Mutations in the nonstructural 5a gene of european hepatitis c virus isolates and response to interferon alfa. *Hepatology*, 25(3):740–4.
- Zeuzem, S., Schmidt, J. M., Lee, J. H., von Wagner, M., Teuber, G., und Roth, W. K. (1998). Hepatitis c virus dynamics in vivo: effect of ribavirin and interferon alfa on viral turnover. *Hepatology*, 28(1):245–52.
- Zeuzem, S., Teuber, G., Lee, J. H., Ruster, B., und Roth, W. K. (1996). Risk factors for the transmission of hepatitis c. *J Hepatol*, 24(2 Suppl):3–10.
- Zibert, A., Schreier, E., und Roggendorf, M. (1995). Antibodies in human sera specific to hypervariable region 1 of hepatitis c virus can block viral attachment. *Virology*, 208(2):653–61.
- Zimmermann, A., Trilling, M., Wagner, M., Wilborn, M., Bubic, I., Jonjic, S., Koszynowski, U., und Hengel, H. (2005). A cytomegaloviral protein reveals a dual role for stat2 in ifn-gamma signaling and antiviral responses. *J Exp Med*, 201(10):1543–53.
- Zimring, J. C., Goodbourn, S., und Offermann, M. K. (1998). Human herpesvirus 8 encodes an interferon regulatory factor (irf) homolog that represses irf-1-mediated transcription. *J Virol*, 72(1):701–7.

