

8 Literaturverzeichnis

Aarestrup, F.M., Jensen, N.E. (1997):

Prevalence and duration of intramammary infection in Danish heifers during the peripartum period.

J. Dairy Sci. 80, 307–312

Aarestrup, F.M., Wegener, H.C., Rosdahl, V.T., Jensen, N.E. (1995):

Staphylococcal and other bacterial species associated with intramammary infections in Danish dairy herds.

Acta vet. Scand. 36, 475–487

Arbeitsgemeinschaft Deutscher Rinderzüchter, ADR (1991):

Rinderproduktion in der Bundesrepublik Deutschland 1990, Bonn

Arbeitsgemeinschaft Deutscher Rinderzüchter, ADR (1996):

Rinderproduktion in der Bundesrepublik Deutschland 1995, Bonn

Arbeitsgemeinschaft Deutscher Rinderzüchter, ADR (1999):

Rinderproduktion in der Bundesrepublik Deutschland 1998, Bonn

Barkema, H.W., Schukken, Y.H., Lam, T.J.G.M., Galligan, D.T., Beiboer, M.L., Brand, A. (1997):

Estimation of interdependence among quarters of the bovine udder with subclinical mastitis and implications for analysis.

J. Dairy Sci. 80, 1592–1599

Barkema, H.W., Schukken, Y.H., Lam, T.J.G.M., Beiboer, M.L., Wilmink, H., Benedictus, G., Brand, A. (1998):

Incidence of clinical mastitis in dairy herds grouped in three categories by bulk milk somatic cell counts.

J. Dairy Sci. 81, 411–419

Barkema, H.W., Schukken, Y.H., Lam, T.J.G.M., Beiboer, M.L., Benedictus, G., Brand, A. (1999):

Management practices associated with the incidence rate of clinical mastitis.

J. Dairy Sci. 82, 1643–1654

Barker, A.R., Schrick, F.N., Lewis, M.J., Dowlen, H.H., Oliver, S.P. (1998):

Influence of clinical mastitis during early lactation on reproductive performance of Jersey cows.

J. Dairy Sci. 81, 1285–1290

Bartlett, P.C., van Wijk, J., Wilson, D.J., Green, C.D., Miller, G.Y., Majewski, G.A., Heider, L.E. (1991):

Temporal patterns of lost milk production following clinical mastitis in a large Michigan Holstein herd.

J. Dairy Sci. 74, 1561–1572

Bartlett, P.C., Miller, G.Y., Lance, S.E., Heider, L.E. (1992):

Managerial determinants of intramammary coliform and environmental streptococci infections in Ohio dairy herds.

J. Dairy Sci. 75, 1241–1252

Barto, P.B., Bush, L.J., Adam, G.D. (1982):

Feeding milk containing *Staphylococcus aureus* to calves.

J. Dairy Sci. 65, 271–274

Bascom, S.S., Young, A.J. (1998):

A summary of the reasons why farmers cull cows.

J. Dairy Sci. 81, 2299–2305

Bauman, D.E., Currie, W.B. (1980):

Partitioning of nutrients during pregnancy and lactation: a review of mechanisms involving homeostasis and homeorhesis.

J. Dairy Sci. 63, 1514–1529

Beaudeau, F., Frankena, K., Fourichon, C., Seegers, H., Faye, B., Noordhuizen, J.P.T.M. (1994):

Associations between health disorders of French dairy cows and early and late culling within the lactation.

Prev. Vet. Med. 19, 213–231

Beaudeau, F., Ducrocq, V., Fourichon, C., Seegers, H. (1995):

Effect of disease on length of productive life of French Holstein dairy cows assessed by survival analysis.

J. Dairy Sci. 78, 103-117

Bendixen, P.H., Vilson, B., Ekesbo, I. (1988):

Disease frequencies in dairy cows in Sweden. V. Mastitis.

Report of the Swedish University of Agricultural Sciences, Faculty of Veterinary Medicine, Department of Animal Hygiene

Bergmann, J. (1998):

Einfluß der negativen Energiebilanz bei Hochleistungskühen im Puerperium auf die Sensibilität der Hypophyse für GnRH und auf die Veränderung unterschiedlicher indirekter Stoffwechselfparameter.

Dissertation Freie Universität Berlin, Fachbereich Veterinärmedizin

Binde, M., Bakke, H. (1984):

Relationships between teat characteristics and udder health – a field survey.

Nord. Vet.-Med. 36, 111–116

Birgersson, A., Jonsson, P., Holmberg, O. (1992):

Species identification and some characteristics of coagulase-negative staphylococci isolated from bovine udders.

Vet. Microbiol. 31, 181–189

Bleckmann, E., Hoedemaker, M. (1996):

Möglichkeiten und Grenzen der bakteriologischen Untersuchung von Milchproben in der Tierarztpraxis.

Prakt. Tierarzt 77, collegium veterinarium XXVI, 22–23

Blobel, H., Schließer, T. (1994):

Handbuch der bakteriellen Infektionen bei Tieren. Band II/1 Staphylokokken-Infektionen und –Enterotoxine.

Gustav Fischer Verlag, Jena, Stuttgart, S. 133 ff.

Bramley, A.J. (1982):

Sources of *Streptococcus uberis* in the dairy herd. I. Isolation from bovine faeces and from straw bedding of cattle.

J. Dairy Res. 49, 369–373

Bramley, A.J. (1985):

The sources of mastitis pathogens for a dairy herd and their control.

Kieler Milchwirtschaftliche Forschungsberichte 37, 375–385

Bramley, A.J., Godinho, K.S., Grindal, R.J. (1981):

Evidence of penetration of the bovine teat duct by *Escherichia coli* in the interval between milkings.

J. Dairy Res. 48, 379–386

Braun, U., Forrer, R., Furer, W., Lutz, H. (1991):

Selenium and vitamin E in blood sera of cows from farms with increased incidence of disease.

Vet. Rec. 128, 543–547

Brentrup, H. (1998):

Euterinfektionen bei Erstkalbinnen - Vorkommen, Bedeutung, prophylaktische Maßnahmen.

Tierärztl. Umsch. 53, 15–18

Buhatel, T., Vesa, S., Csegezi, E. (1983):

I. Occurrence of mastitis in large herds using machine milking. II. Zootechnical factors in mammary gland diseases and reproductive disorders.

Proceedings of the 8th Romanian symposium on the breeding, management and diseases of ruminants. Pathology section. Cluj-Napoca, Romania, Institutul Agronomic Dr. Petru Groza, 241–252

Bunch, K.J., Heneghan, D.J.S., Hibbitt, K.G., Rowlands, G.J. (1984):

Genetic influences on clinical mastitis and its relationship with milk yield, season, and stage of lactation.

Livest. Prod. Sci. 11, 91–104

Butler, W.R., Smith, R.D. (1989):

Interrelationships between energy balance and postpartum reproductive function in dairy cattle.

J. Dairy Sci. 72, 767–783

Cai, T.-Q., Weston, P.G., Lund, L.A., Brodie, B., McKenna, D.J., Wagner, W.C. (1994):

Association between neutrophil functions and periparturient disorders in cows.

Am. J. Vet. Res. 55, 934–943

Canfield, R.W., Butler, W.R. (1990):

Energy balance and pulsatile LH secretion in early postpartum dairy cattle.

Domest. Anim. Endocrinol. 7, 323–330

Canfield, R.W., Butler, W.R. (1991):

Energy balance, first ovulation and the effects of naloxone on LH secretion in early postpartum dairy cows.

J. Anim. Sci. 69, 740–746

Cobo-Abreu, R., Martin, S.W., Willoughby, R.A., Ston, J.B. (1979):

The association between disease, production and culling in a university dairy herd.

Can. vet. J. 20, 191–195

Coleman, D.A., Thayne, W.V., Dailey, R.A. (1985):

Factors affecting reproductive performance of dairy cows.

J. Dairy Sci. 68, 1793–1803

Correa, M.T., Curtis, C.R., Erb, H.N., Scarlett, J.M., Smith, R.D. (1990):

An ecological analysis of risk factors for postpartum disorders of Holstein-Friesian cows from thirty-two New York farms.

J. Dairy Sci. 73, 1515–1524

Cullen, G.A., Hebert, C.N. (1967):

Some ecological observations on micro-organisms inhabiting bovine skin, teat canals and milk.

Br. vet. J. 123, 14–25

Curtis, C.R., Erb, H.N., Sniffen, C.J., Smith, R.D., Kronfeld, D.S. (1985):

Path analysis of dry period nutrition, postpartum metabolic and reproductive disorders, and mastitis in Holstein cows.

J. Dairy Sci. 68, 2347–2360

de Kruif, A., Mansfeld, R., Hoedemaker, M. (1998):

Tierärztliche Bestandsbetreuung beim Milchrind. Ferdinand Enke Verlag, Stuttgart

Deutsche Veterinärmedizinische Gesellschaft, DVG, Fachgruppe Milchhygiene (1980):

Die subklinische Mastitis des Rindes und Vorschläge für bundeseinheitliche Richtlinien zu ihrer Bekämpfung.

DVG, Fachgruppe Milchhygiene des Arbeitsgebietes Lebensmittelhygiene, Kiel

Deutz, A., Obritzhauser, W. (1996):

Beitrag zur akuten Streptokokkenmastitis des Rindes.

Prakt. Tierarzt 77, 406–412

Dirksen, G., Liebich, H.-G., Mayer, E. (1985):

Funktionelle Bedeutung adaptiver Veränderungen der Pansenschleimhaut von Wiederkäuern.
Bericht des 16. Kongresses der Deutschen Veterinärmedizinischen Gesellschaft.
Bad Nauheim, 341–345

Distl, O., Wurm, A., Glibotic, A., Brem, G., Kräusslich, H. (1989):

Analysis of relationships between veterinary recorded production diseases and milk
production in dairy cows.
Livest. Prod. Sci. 23, 67–78

Döpfer, D., Schukken, Y.H., Metzner, R.M., Petersen, D. (1993):

Betreuungsstrategien zur Sanierung von Milchviehbetrieben mit hohem Zellgehalt in der
Tankmilch.
Prakt. Tierarzt 74, collegium veterinarium XXIV, 70–72

Doggweiler, R., Hess, E. (1983):

Zellgehalt in der Milch ungeschädigter Euter.
Milchwissenschaft 38, 5–8

Dohoo, I.R., Martin, S.W. (1984a):

Disease, production and culling in Holstein-Friesian cows. IV. Effects of disease on
production.
Prev. Vet. Med. 2, 755–770

Dohoo, I.R., Martin, S.W. (1984b):

Disease, production and culling in Holstein-Friesian cows. V. Survivorship.
Prev. Vet. Med. 2, 771–784

Edinger, D., Tenhagen, B.-A., Heuwieser, W., Kalbe, P., Klünder, P., Baumgärtner, B.
(1998):

Effect of early puerperal mastitis in primiparous cows on production, somatic cell scores and
culling.
Proceedings of the 10th International Conference on Production Diseases in Farm Animals,
Utrecht, Wageningen Pers, Niederlande, 368

Edinger, D., Tenhagen, B.-A., Heuwieser, W., Kalbe, P., Klünder, P., Baumgärtner, B. (1999):

Einfluß des Zitzendippens ante partum auf die mikrobielle Beschaffenheit des Erstgemelks bei Erstkalbinnen und die Mastitisinzidenz in der ersten Woche post partum.

Kurzfassung der Vorträge der Milchkonferenz 1999 der Deutschen Gesellschaft für Milchwissenschaft, 23. und 24. September, Kiel

Edler, B., Hoedemaker, M., Drumm, B. (1995):

Vergleichende Untersuchungen zur Mastitis bei Erstkalbinnen und Kühen im peripartalen Zeitraum, 1. Mitteilung.

Prakt. Tierarzt 76, 855–867

Elbers, A.R.W., Miltenburg, J.D., de Lange, D., Crauwels, A.P.P., Barkema, H.W., Schukken, Y.H. (1998):

Risk factors for clinical mastitis in a random sample of dairy herds from the southern part of the Netherlands.

J. Dairy Sci. 81, 420–426

Erb, H.N., Martin, S.W., Ison, N., Swaminathan, S. (1981):

Interrelationships between production and reproductive diseases in Holstein cows. Conditional relationships between production and disease.

J. Dairy Sci. 64, 272–281

Erb, H.N., Smith, R.D., Oltenacu, P.A., Guard, C.L., Hillman, R.B., Powers, P.A., Smith, M.C., White, M.E. (1985):

Path model of reproductive disorders and performance, milk fever, mastitis, milk yield, and culling in Holstein cows.

J. Dairy Sci. 68, 3337–3349

Erskine, R.J., Eberhart, R.J., Hutchinson, L.J., Spencer, S.B., Campbell, M.A. (1988):

Incidence and types of clinical mastitis in dairy herds with high and low somatic cell counts.

J. Am. Vet. Med. Assoc. 192, 761–765

Esslemont, R.J., Kossaibati, M.A. (1997):

Culling in 50 dairy herds in England.

Vet. Rec. 140, 36–39

Faull, W.B., Walton, J.R., Bramley, A.J., Hughes, J.W. (1983):

Mastitis in a large, zero-grazed dairy herd.

Vet. Rec. 113, 415–420

Fenlon, D.R., Logue, D.N., Gunn, J., Wilson, J. (1995):

A study of mastitis bacteria and herd management practices to identify their relationship to high somatic cell counts in bulk tank milk.

Br. vet. J. 151, 17–25

Fox, L.K., Chester, S.T., Hallberg, J.W., Nickerson, S.C., Pankey, J.W., Weaver, L.D. (1995):

Survey of intramammary infections in dairy heifers at breeding age and first parturition.

J. Dairy Sci. 78, 1619–1628

Gonzalez, R.N., Jasper, D.E., Kronlund, N.C., Farver, T.B., Cullor, J.S., Bushnell, R.B., Dellinger, J.D. (1990):

Clinical mastitis in two California dairy herds participating in contagious mastitis control programs.

J. Dairy Sci. 73, 648–660

Grindal, R.J., Hillerton, J.E. (1991):

Influence of milk flow rate on new intramammary infection in dairy cows.

J. Dairy Res. 58, 263–268

Grindal, R.J., Walton, A.W., Hillerton, J.E. (1991):

Influence of milk flow rate and streak canal length on new intramammary infection in dairy cows.

J. Dairy Res. 58, 383–388

Gröhn, Y.T., Erb, H.N., McCulloch, C.E., Saloniemi, H.S. (1989):

Epidemiology of metabolic disorders in dairy cattle: association among host characteristics, disease, and production.

J. Dairy Sci. 72, 1876–1885

Gröhn, Y.T., Erb, H.N., McCulloch, C.E., Saloniemi, H.S. (1990):

Epidemiology of mammary gland disorders in multiparous Finnish Ayrshire cows.

Prev. Vet. Med. 25, 241–252

Grommers, F.J., van de Geer, D., in 't Veen, C.A.A. (1985):

Duration of bovine intramammary infections in commercial dairy herds.

Vet. Rec. 116, 581–584

Guidry, A.J., Paape, M.J., Pearson, R.E. (1976):

Effects of parturition and lactation on blood and milk cell concentrations, corticosteroids, and neutrophil phagozytosis in the cow.

Am. J. Vet. Res. 37, 1195–1200

Hamann, J., Reichmuth, J. (1990):

Exogene Einflüsse auf den Zellgehalt der Milch unter Berücksichtigung des Gesundheitszustandes der Milchdrüse.

Milchwissenschaft 45, 286–290

Hamann, J. (1992):

Mastitisbekämpfung auf der Grundlage zytologischer Befunde der Herdensammelmilch.

Kieler Milchwirtschaftliche Forschungsberichte 44, 327–338

Harman, J.L., Gröhn, Y.T., Erb, H.N., Casella, G. (1996):

Event-time analysis of the effect of season of parturition, parity, and concurrent disease on parturition-to-conception interval in dairy cows.

Am. J. Vet. Res. 57, 640–645

Harmon, R.J., Crist, W.L., Hemken, R.W., Langlois, B.E. (1986):

Prevalence of minor udder pathogens after intramammary dry treatment.

J. Dairy Sci. 69, 843–849

Harmon, R.J. (1994):

Symposium: mastitis and genetic evaluation for somatic cell count. Physiology of mastitis and factors affecting somatic cell counts.

J. Dairy Sci 77, 2103–2112

Harrison, R.O., Ford, S.P., Young, J.W., Conley, A.J., Freeman, A.E. (1990):

Increased milk production versus reproductive and energy status of high producing dairy cows.

J. Dairy Sci. 73, 2749–2758

Heinonen, M., Heinonen, K. (1989):

Retained placenta in cattle: the effect of treatment or nontreatment on puerperal diseases and subsequent fertility.

Acta vet. Scand. 30, 425–429

Hillers, J.K., Senger, P.L., Darlington, R.L., Fleming, W.N. (1984):

Effects of production, season, age of cow, days dry, and days in milk on conception to first service in large commercial dairy herds.

J. Dairy Sci. 67, 861–867

Hogan, J.S., White, D.G., Pankey, J.W. (1987):

Effects of teat dipping on intramammary infections by staphylococci other than *Staphylococcus aureus*.

J. Dairy Sci. 70, 873–879

Hogan, J.S., Smith, K.L., Todhunter, D.A., Schoenberger, P.S. (1988):

Rate of environmental mastitis in quarters infected with *Corynebacterium bovis* and *Staphylococcus* species.

J. Dairy Sci. 71, 2520–2525

Hogan, J.S., Smith, K.L., Hoblet, K.H., Todhunter, D.A., Schoenberger, P.S., Hueston, W.D., Pritchard, D.E., Bowman, G.L., Heider, L.E., Brockett, B.L., Conrad, H.R. (1989a):
Field survey of clinical mastitis in low somatic cell count herds.
J. Dairy Sci. 72, 1547–1556

Hogan, J.S., Smith, K.L., Hoblet, K.H., Todhunter, D.A., Schoenberger, P.S., Hueston, W.D., Pritchard, D.E., Bowman, G.L., Heider, L.E., Brockett, B.L., Conrad, H.R. (1989b):
Bacterial counts in bedding materials used on nine commercial dairies.
J. Dairy Sci. 72, 250–258

Honkanen-Buzalski, T., Griffin, T.K., Dodd, F.H. (1984):
Observations on *Corynebacterium bovis* infection of the bovine mammary gland. I. Natural infection.
J. Dairy Res. 51, 371–378

Huszenicza, G., Jánosi, S., Kulcsár, M., Kóródi, P., Dieleman, S.J., Bartyik, J., Rudas, P., Ribiczei-Szabó, P. (1998):
Gram-negative mastitis in early lactation may interfere with ovarian and certain endocrine functions and metabolism in dairy cows.
Reprod. Dom. Anim. 33, 147–153

International Dairy Federation (1987):
Bovine mastitis – definition and guidelines for diagnosis.
IDF Doc. 211

Jain, N.C. (1979):
Common mammary pathogens and factors in infection and mastitis.
J. Dairy Sci. 62, 128–134

Jarp, J. (1991):
Classification of coagulase-negative staphylococci isolated from bovine clinical and subclinical mastitis.
Vet. Microbiol. 27, 151–158

Jonsson, P., Olsson, S.O., Olofson, A.S., Holmberg, O., Funke, H. (1991):

Bacteriological investigations of clinical mastitis in heifers in Sweden.

J. Dairy Res. 58, 179–185

Jorstad, A., Farver, T.B., Riemann, H. (1989):

Teat canal diameter and other cow factors with possible influence on somatic cell counts in cow milk.

Acta vet. Scand. 30, 239–245

Jukola, E., Hakkarainen, J., Saloniemi, H., Sankari, S. (1996):

Blood selenium, vitamin E, vitamin A, and β -carotene concentrations and udder health, fertility treatments, and fertility.

J. Dairy Sci. 79, 838–845

Kehrli, M.E. Jr., Kimura K., Goff J.P., Stabel J.R., Nonnecke B.J. (1999):

Periparturient immunosuppression in dairy cows: nutrition and lactation effects.

Proceedings of the 10th International Conference on Production Diseases in Farm Animals, Utrecht, Wageningen Pers, Niederlande, 41–55

Kimura, K., Goff, J.P., Kehrli, M.E. Jr., Harp, J.A. (1999):

Phenotype analysis of peripheral blood mononuclear cells in periparturient dairy cows.

J. Dairy Sci. 82, 315–319

Kirchgeßner, M. (1987):

Tierernährung: Leitfaden für Studium, Beratung und Praxis.

7. Auflage, DLG-Verlag, Frankfurt, S. 284

Klarmann, D. (1997):

Antibiotika-Resistenzen wichtiger Infektionserreger 1996 in Weser-Ems.

Dtsch. tierärztl. Wochenschr. 104, 325–335

Klug, F., Franz, H., Baumung, A. (1988):

Beziehungen zwischen Gesundheit und Leistung bei Jungkühen.

Tierzucht 42, 556–558

Koldewey, E., Emanuelson, U., Janson, L. (1999):

Relation of milk production loss to milk somatic cell count.

Acta vet. Scand. 40, 47–56

Kossaibati, M.A., Esslemont, R.J. (1997):

The costs of production diseases in dairy herds in England.

Vet. J. 154, 41–51

Krabisch, P., Gangl, A. (1999):

Zur aktuellen Resistenzsituation von Cefoperazon – Auswertung einer deutschlandweiten Multicenterstudie.

Vortragzusammenfassung Pfizer Mastitis Symposium, BPT-Kongress in Nürnberg, 10–23

Labernia, J., Lopez-Gatius, F., Santolaria, P., Hanzen, C., Laurent, Y., Houtain, J.Y. (1998):

Influence of calving season on the interactions among reproductive disorders of dairy cows.

Animal Science 67, 387–393

Labohm, R., Götz, E., Luhofer, G., Hess, R.G., Bostedt, H. (1998a):

Factors influencing the somatic milk-cell-count in dairy cows. 1. Influence of bacteriological findings, stage and number of lactation.

Milchwissenschaft 53, 63–66

Labohm, R., Götz, E., Luhofer, G., Hess, R.G. (1998b):

Diagnostik des Mastitisrisikos: Interpretation von Zellzahlbefunden.

Prakt. Tierarzt 79, collegium veterinarium XXVIII, 82–85

Lacy-Hulbert, S.J., Hillerton, J.E. (1995):

Physical characteristics of the bovine teat canal and their influence on susceptibility to streptococcal infections.

J. Dairy Res. 62, 395–404

Laevens, H., Deluyker, H., Schukken, Y.H., de Meulemeester, L., Vandermeersch, R., de Muëlenaere, E., de Kruif, A. (1997):

Influence of parity and stage of lactation on the somatic cell count in bacteriologically negative dairy cows.

J. Dairy Sci. 80, 3219–3226

Lancelot R., Faye B., Lescourret F. (1997):

Factors affecting the distribution of clinical mastitis among udder quarters in French dairy cows.

Vet. Res. 28, 45–53

Landeskontrollverband Schleswig-Holstein, LKV (1996):

Die Leistungsergebnisse des Kontrolljahres 1995.

Landeskontrollverband Schleswig-Holstein e.V., Kiel

Landeskontrollverband Schleswig-Holstein, LKV (1998):

Die Leistungsergebnisse des Kontrolljahres 1997.

Landeskontrollverband Schleswig-Holstein e.V., Kiel

Landeskontrollverband Schleswig-Holstein, LKV (1999):

Die Leistungsergebnisse des Kontrolljahres 1998.

Landeskontrollverband Schleswig-Holstein e.V., Kiel

Lange, J., Bleckmann, E. (1999):

Erregerspektrum und Antibiotikaresistenz bei katarrhalischen Mastitiden des Rindes.

Prakt. Tierarzt 80, 807–812

Lescourret, F., Coulon, J.B., Faye, B. (1995):

Predictive model of mastitis occurrence in the dairy cow.

J. Dairy Sci. 78, 2167–2177

Le Van, P.L., Eberhart, R.J., Kesler, E.M. (1985):

Effects of natural intrammary *Corynebacterium bovis* infection on milk yield and composition.

J. Dairy Sci. 68, 3329–3336

Lewis, G.S. (1997):

Symposium: Health problems of the postpartum cow - Uterine health and disorders.

J. Dairy Sci. 80, 984–994

Linde, C., Holmberg, O., Åström, G. (1980):

The interference between coagulase negative staphylococci and *Corynebacterium bovis* and the common udder pathogens in the lactating cow.

Nord. Vet.-Med. 32, 552–558

Lotthammer, K.-H. (1987):

Gesundheitsüberwachung in Milchviehbeständen.

Aktuelle Themen der Tierernährung und Veredelungswirtschaft. Zusammenfassung der Vorträge der wissenschaftlichen Tagung, Cuxhaven, 79–93

Lotthammer, K.-H., Klarmann, D. (1999):

Auswertungen von Resistenzbestimmungen in einem Gebiet mit intensiver Tierproduktion.

Tierärztl. Prax. 27, 324–328

Lotthammer, K.-H., Wittkowski, G. (1994):

Fruchtbarkeit und Gesundheit der Rinder.

Verlag Eugen Ulmer, Stuttgart

Lucey, S., Rowlands, G.J. (1984):

The association between clinical mastitis and milk yield in dairy cows.

Anim. Prod. 39, 165–175

Lucey, S., Rowlands, G.J., Russell, A.M. (1986):

Short-term associations between disease and milk yield of dairy cows.

J. Dairy Res. 53, 7–15

Lucy, M.C., Beck, J., Staples, C.R., Head, H.H., de la Sota, R.L., Thatcher, W.W. (1992):

Follicular dynamics, plasma metabolites, hormones and insulin-like growth factor I (IGF-I) in lactating cows with positive or negative energy balance during the preovulatory period.

Reprod. Nutr. Dev. 32, 331–341

Luhofe, G., Klawonn, W., Labohm, R., Heß, R.G. (1996):

Amtlicher Tierarzt für Eutergesundheit und Milchhygiene: Arbeitsergebnisse in Rheinland-Pfalz.

Tierärztl. Prax. 24, 459–466

Mallard, B.A., Dekkers, J.C., Ireland, M.J., Leslie, K.E.; Sharif, S., Vankampen, C.L., Wagter, L., Wilkie, B.N. (1998):

Alteration in immune responsiveness during the peripartum period and its ramification on dairy cow and calf health.

J. Dairy Sci. 81, 585–595

Markusfeld, O. (1987):

Periparturient traits in seven high dairy herds. Incidence rates, association with parity, and interrelationships among traits.

J. Dairy Sci. 70, 158–166

Martin, G., Bergmann, A. (1993):

Untersuchungen über koagulasenegative Staphylokokken aus Eutersekretproben von Rindern auf dem Gebiet der östlichen Bundesländer.

Monatsh. Veterinärmed. 48, 559–565

Matos, J.S., White, D.G., Harmon, R.J., Langlois, B.E. (1991):

Isolation of *Staphylococcus aureus* from sites other than the lactating mammary gland.

J. Dairy Sci. 74, 1544–1549

Matthews, K.R., Harmon, R.J., Langlois, B.E. (1991):

Effect of naturally occurring coagulase-negative staphylococci infections on new infections by mastitis pathogens in the bovine.

J. Dairy Sci. 74, 1855–1859

Matthews, K.R., Harmon, R.J., Langlois, B.E. (1992):

Prevalence of *Staphylococcus* species during the periparturient period in primiparous and multiparous cows.

J. Dairy Sci. 75, 1835–1839

Meaney, W.J. (1981):

Mastitis levels in spring-calving dairy heifers.

Ir. Vet. J. 35, 205–209

Metzner, M., Mansfeld, R. (1992):

Tierärztliche Betreuung von Milcherzeugerbetrieben, Teil 2: Die Beurteilung von Fruchtbarkeitsparametern, Möglichkeiten und Grenzen.

Prakt. Tierarzt, 72, 800–814

Michel, G., Rausch, B. (1988):

Veränderungen von Zitzenmaßen des Rindereuters im Verlaufe von mehreren Laktationen.

Mh. Vet.-Med. 43, 337–339

Miettinen, P.V. (1990):

Metabolic balance and reproductive performance in Finnish dairy cows.

Zentralbl. Veterinärmed. A. 37, 417–424

Miettinen, P.V., Rainio, V.A., Kukkonen, S.A., Setälä, J.J. (1991):

Finnish dairy cows: energy balance and seasonal variation in fertility.

Zentralbl. Veterinärmed. A. 38, 427–432

Miltenburg, J.D., de Lange, D., Crauwels, A.P.P., Bongers, J.H., Tielen, M.J.M., Schukken, Y.H., Elbers, A.R.W. (1996):

Incidence of clinical mastitis in a random sample of dairy herds in the southern Netherlands.

Vet. Rec. 139, 204–207

Moore, D.A., Cullor, J.S., Bondurant, R.H., Sischo, W.M. (1991):

Preliminary field evidence for the association of clinical mastitis with altered interestrus intervals in dairy cattle.

Theriogenology 36, 257–265

Munch-Petersen, E. (1970):

Mastitis in bovine primiparae.

Vet. Rec. 87, 568–574

Myllys, V. (1995):

Staphylococci in heifer mastitis before and after parturition.

J. Dairy Res. 62, 51–60

Myllys, V., Rautula, H. (1995):

Characterization of clinical mastitis in primiparous heifers.

J. Dairy Sci. 78, 538–545

Myllys, V., Honkanen-Buzalski, T., Huovinen, P., Sandholm, M., Nurmi, E. (1994):

Association of changes in the bacterial ecology of bovine mastitis with changes in the use of milking machines and antibacterial drugs.

Acta vet. Scand. 35, 363–369

Myllys, V., Asplund, K., Brofeldt, E., Hirvelä-Koski, V., Honkanen-Buzalski, T., Junttila, J., Kulkas, L., Myllykangas, O., Niskanen, M., Saloniemi, H., Sandholm, M., Saranpää, T. (1998):

Bovine mastitis in Finland in 1988 and 1995 – Changes in prevalence and antimicrobial resistance.

Acta vet. Scand. 39, 119–126

Natzke, R.P. (1978):

Change in udder health and overmilking.

J. Dairy Sci. 61, 233–238

Natzke, R.P. (1981):

Elements of mastitis control.

J. Dairy Sci. 64, 1431–1442

Natzke, R.P., Everett, R.W., Guthrie, R.S., Keown, J.F., Meek, A.M., Merrill, W.G., Roberts, S.J., Schmidt, G.H. (1972):

Mastitis control program: effect on milk production.

J. Dairy Sci. 55, 1256–1260

Ndiweni, N, Field, T.R., Williams, M.R., Booth, J.M., Finch, J.M. (1991):

Studies on the incidence of clinical mastitis and blood levels of vitamin E and selenium in dairy herds in England.

Vet. Rec. 129, 86–88

Nickerson, S.C., Boddie, R.L. (1994):

Effect of naturally occurring coagulase-negative staphylococcal infections on experimental challenge with major mastitis pathogens.

J. Dairy Sci. 77, 2526–2536

Oliver, S.P., Mitchell, B.A. (1983):

Intramammary infections in primigravid heifers near parturition.

J. Dairy Sci. 66, 1180–1183

Oliver, S.P., Sordillo, L.M. (1989):

Approaches to the manipulation of mammary gland involution.

J. Dairy Sci. 72, 1647–1664

Oltenacu, P.A., Ekesbo, I. (1994):

Epidemiological study of clinical mastitis in dairy cattle.

Vet. Res. 25, 208–212

Oltenacu, P.A., Frick, A., Lindhé, B. (1990):

Epidemiological study of several clinical diseases, reproductive performance and culling in primiparous Swedish cattle.

Prev. Vet. Med. 9, 59–74

Opsomer, G., Grohn, Y.T., Hertl, J., Coryn, M., Deluyker, H., de Kruif, A. (2000):

Risk factors for post partum ovarian dysfunction in high producing dairy cows in Belgium: a field study.

Theriogenology 53, 841–857

Paape, M.J., Pearson, R.E., Schultze, W.D. (1978):

Variation among cows in the ability of milk to support phagocytosis and in the ability of polymorphonuclear leukocytes to phagocytose *Staphylococcus aureus*.

Am. J. Vet. Res. 39, 1907–1910

Pankey, J.W., Nickerson, S.C. (1985):

Effects of *Corynebacterium bovis* infections on susceptibility to major mastitis pathogens.

Kieler Milchwirtschaftliche Forschungsberichte 37, 385–389

Pankey, J.W., Drechsler, P.A., Wildman, E.E. (1991):

Mastitis prevalence in primigravid heifers at parturition.

J. Dairy Sci. 74, 1550–1552

Peeler, E.J., Otte, M.J., Esslemont, R.J. (1994):

Inter-relationships of periparturient diseases in dairy cows.

Vet. Rec. 134, 129–132

Péretz, G., Bugnard, F., Philpot, J.M., Pluvillage, P. (1995):

Risk factors for dairy cow mastitis.

Proceedings of the third IDF International Mastitis Seminar, Tel-Aviv, Israel, S6, 72–73

Philpot, W.N. (1979):

Control of mastitis by hygiene and therapy.

J. Dairy Sci. 62, 168–176

Pösö, J., Mäntysaari, E.A. (1996):

Relationships between clinical mastitis, somatic cell score, and production for the first three lactations of Finnish Ayrshire.

J. Dairy Sci. 79, 1284–1291

Preisler, M.T., Weber, P.S.D., Tempelman, R.J., Erskine, R.J., Hunt, H., Burton, J.L. (2000):

Glucocorticoid receptor expression profiles in mononuclear leukocytes of periparturient Holstein cows.

J. Dairy Sci. 83, 38–47

Pyörälä, S., Syväjärvi, J. (1987):

Bovine acute mastitis. Part I. Clinical aspects and parameters of inflammation in mastitis caused by different pathogens.

J. Vet. Med. 34, 573–584

Radostits, O.M., Leslie, K.E., Fetrow, J. (1994):

Herd health: food, animal, production, medicine.

2. Auflage, W.B. Saunders Company, Philadelphia, USA, 141 ff.

Rainard, P., Poutrel, B. (1982):

Dynamics of nonclinical bovine intramammary infections with major and minor pathogens.

Am. J. Vet. Res. 43, 2143–2146

Rainard, P., Poutrel, B. (1988):

Effect of naturally occurring intramammary infections by minor pathogens on new infections by major pathogens in cattle.

Am. J. Vet. Res. 49, 327–329

Rainard, P., Ducelliez, M., Poutrel, B. (1990):

The contribution of intramammary infections by coagulase-negative staphylococci to the herd bulk milk somatic cell count.

Vet. Res. Commun. 14, 193–198

Rajala, P.J., Gröhn, Y.T. (1998):

Disease occurrence and risk factor analysis in Finnish Ayrshire cows.

Acta vet. Scand. 39, 1–13

Rajala-Schultz, P.J., Gröhn, Y.T., McCulloch, C.E., Guard, C.L. (1999):

Effects of clinical mastitis on milk yield in dairy cows.

J. Dairy Sci. 82, 1213–1220

Rasmussen, L.K., Nielsen, B.L., Pryce, J.E., Mottram, T.T., Veerkamp, R.F. (1999):

Risk factors associated with the incidence of ketosis in dairy cows.

Animal Science 68, 379–386

Reinsch, N., Jaitner, J., Dempfle, L. (1997):

Investigations on functional traits in Simmental. 2. Incidence of diseases in German Simmental cows.

Livest. Prod. Sci. 52, 87–96

Richter, J., Götze, R. (1993):

Tiergeburtschilfe

4. Auflage, Parey Verlag, Berlin, Hamburg, 129 ff.

Rinder-Report '99. Ergebnisse der Rinderspezialberatung in Schleswig-Holstein (1999):

Betriebswirtschaftliche Mitteilungen der Landwirtschaftskammer Schleswig-Holstein, 530/31

Roberson, J.R., Fox, L.K., Hancock, D.D., Gay, J.M., Besser, T.E. (1994a):

Coagulase-positive *Staphylococcus* intramammary infections in primiparous dairy cows.

J. Dairy. Sci. 77, 958–969

Roberson, J.R., Fox, L.K., Hancock, D.D., Gay, J.M., Besser, T.E. (1994b):

Ecology of *Staphylococcus aureus* isolated from various sites on dairy farms.

J. Dairy Sci. 77, 3354–3364

Roberson, J.R., Fox, L.K., Hancock, D.D., Gay, J.M., Besser, T.E. (1998):

Sources of intramammary infections from *Staphylococcus aureus* in dairy heifers at first parturition.

J. Dairy Sci. 81, 687–693

Rogers, G.W., Hargrove, G.L., Lawlor, T.J., Ebersole, J.L. (1991):

Correlations among linear type traits and somatic cell counts.

J. Dairy Sci. 74, 1087–1091

Ronningen, O., Reitan, A.D. (1990):

Influence of static and dynamic teat characteristics and milking time on udder health in Norwegian Red cattle.

J. Dairy Res. 57, 171–177

Rosenberger, G. (1990):

Die klinische Untersuchung des Rindes. 3. Auflage, Parey Verlag, Berlin, Hamburg, 525–543

Rulof, U. (1997):

Untersuchungen über Art und Häufigkeit intramammärer Infektionen bei Färsen ante und post partum in einem norddeutschen Hochzuchtgebiet und die Effizienz präpartaler antibiotischer Behandlungen.

Dissertation, Tierärztliche Hochschule Hannover

Rupp, R., Boichard, D. (1999):

Genetic parameters for clinical mastitis, somatic cell score, production, udder type traits, and milking ease in first lactation Holsteins.

J. Dairy Sci. 82, 2198–2204

Sargeant, J.M., Scott, H.M., Leslie, K.E., Ireland, M.J., Bashiri, A. (1998):

Clinical mastitis in dairy cattle in Ontario: frequency of occurrence and bacteriological isolates.

Can. Vet. J. 39, 33–38

SAS Institute Inc. (1995):

Logistic regression examples using the SAS system.

Version 6, First Edition, Cary, North Carolina, USA

SAS Institute Inc. (1996):

SAS/STAT Software: Changes and enhancement through release 6.12, Cary, North Carolina, USA

Schepers, A.J., Lam, T.J.G.M., Schukken, Y.H., Wilmink, J.B.M., Hanekamp, W.J.A. (1997):

Estimation of variance components for somatic cell counts to determine thresholds for uninfected quarters.

J. Dairy Sci. 80, 1833–1840

Schukken, Y.H., Erb, H.N., Smith, R.D. (1988):

The relationship between mastitis and retained placenta in a commercial population of Holstein dairy cows.

Prev. Vet. Med. 5, 181–190

Schukken, Y.H., Grommers, F.J., van de Geer, D., Brand, A. (1989a):

Incidence of clinical mastitis on farms with low somatic cell counts in bulk milk.

Vet. Rec. 125, 60–63

Schukken, Y.H., van de Geer, D., Grommers, F.J., Smit, J.A., Brand, A. (1989b):

Intramammary infections and risk factors for clinical mastitis in herds with low somatic cell counts in bulk milk.

Vet. Rec. 125, 393–396

Schukken, Y.H., Erb, H.N., Scarlett, J.M. (1989c):

A hospital-based study of the relationship between retained placenta and mastitis in dairy cows.

Cornell Vet. 79, 319–326

Schukken, Y.H., Grommers, F.J., van de Geer, D., Erb, H.N., Brand, A. (1990):

Risk factors for clinical mastitis in herds with a low bulk milk somatic cell count. 1. Data and risk factors for all cases.

J. Dairy Sci. 73, 3463–3471

Schukken, Y.H., Grommers, F.J., van de Geer, D., Erb, H.N., Brand, A. (1991):

Risk factors for clinical mastitis in herds with a low bulk milk somatic cell count. 2. Risk factors for *Escherichia coli* and *Staphylococcus aureus*.

J. Dairy Sci. 74, 826–832

Seykora, A.J., McDaniel, B.T. (1985):

Heritabilities of teat traits and their relationships with milk yield, somatic cell counts, and percent two-minute milk.

J. Dairy Sci. 69, 2670–2683

Seykora, A.J., McDaniel, B.T. (1986):

Genetics statistics and relationships of teat and udder traits, somatic cell count, and milk production.

J. Dairy Sci. 69, 2395–2407

Sheldrake, R.F., Hoare, R.J.T., McGregor, G.D. (1983):

Lactation stage, parity, and infection affecting somatic cells, electrical conductivity, and serum albumin in milk.

J. Dairy Sci. 66, 542–547

Shpigel, N.Y., Winkler, M., Ziv, G., Saran, A. (1998):

Clinical, bacteriological and epidemiological aspects of clinical mastitis in Israeli dairy herds.

Prev. Vet. Med. 35, 1–9

Sieber, R.L., Farnsworth, R.J. (1981):

Prevalence of chronic teat-end lesions and their relationship to intramammary infection in 22 herds of dairy cattle.

J. Am. Vet. Med. Assoc. 178, 1263–1267

Slettbakk, T., Asbjørn, J., Farver, T.B., Hird, D.W. (1990):

Impact of milking characteristics and teat morphology on somatic cell counts in first-lactation Norwegian cattle.

Prev. Vet. Med. 8, 253–267

Smith, R.E., Hagstad, H.V. (1985):

Infection of the bovine udder with coagulase-negative staphylococci.

Kieler Milchwirtschaftliche Forschungsberichte 37, 611–614

Smith, K.L., Hogan, J.S. (1993):

Environmental mastitis.

Vet. Clin. North. Am. Food. Anim. Pract. 9, 489–498

Smith, K.L., Hogan, J.S. (1995):

Epidemiology of mastitis.

Proceedings of the third IDF International Mastitis Seminar, Tel-Aviv, Israel, S6, 3–13

Sobiraj, A., Ostertag, H.-U., Peip, D., Bostedt, H., Kielwein, G. (1988):

Klinische und bakteriologische Untersuchungsbefunde zur Mastitishäufigkeit erstmalig laktierender Rinder intra und post partum.

Tierärztl. Prax. 16, 243–249

Sobiraj, A., Kron, A., Schollmeyer, U., Failing, K. (1997):

Bundesweite Untersuchung zur Erregerverteilung und in-vitro Resistenz euterpathogener Bakterien in der Milch von Kühen mit subklinischer Mastitis.

Tierärztl. Prax. 25, 108–115

Solbu, H. (1983):

Disease recording in Norwegian dairy cattle I. Disease incidences and non genetic effects on mastitis, ketosis and milk fever.

Z. Tierzüchtg. Züchtgsbiol. 100, 139–157

Sørensen, M.K., Jensen, J., Christensen, L.G. (2000):

Udder conformation and mastitis resistance in Danish first-lactation cows: heritabilities, genetic and environmental correlations.

Acta Agric. Scand., Sect. A., Animal Sci. 50, 72–82

Staples, C.R., Thatcher, W.W., Clark, J.H. (1990):

Relationship between ovarian activity and energy status during the early postpartum period of high producing dairy cows.

J. Dairy Sci. 73, 938–947

Stevenson, J.S., Call, E.P. (1988):

Reproductive disorders in the periparturient dairy cow.

J. Dairy Sci. 71, 2572–2583

Swanson, E.W., Poffenbarger, J.I. (1979):

Mammary gland development of dairy heifers during their first gestation.

J. Dairy Sci. 62, 702–714

Thomas, C.L., Vinson, W.E., Pearson, R.E., Dickinson, F.N., Johnson, L.P. (1984):

Relationships between linear type scores, objective type measures, and indicators of mastitis.

J. Dairy Sci. 67, 1281–1292

Timms, L.L., Schultz, L.H. (1987):

Dynamics and significance of coagulase-negative staphylococcal intramammary infections.

J. Dairy Sci. 70, 2648–2657

Todhunter, D.A., Smith, K.L., Hogan, J.S., Schoenberger, P.S. (1991):

Gram-negative bacterial infections of the mammary gland in cows.

Am. J. Vet. Res. 52, 184–188

Todhunter, D.A., Cantwell, L.L., Smith, K.L., Hoblet, K.H., Hogan, J.S. (1993):

Characteristics of coagulase-negative staphylococci isolated from bovine intramammary infections.

Vet. Microbiol. 34, 373–380

Todhunter, D.A., Smith, K.L., Hogan, J.S. (1995):

Environmental streptococcal intramammary infections of the bovine mammary gland.

J. Dairy Sci. 78, 2366–2374

Tolle, A., Heeschen, W., Hamann, J. (1977):

Grundlagen einer systematischen Bekämpfung der subklinischen Mastitis des Rindes.

Kieler Milchwirtschaftliche Forschungsberichte 29, 3–103

Trinidad, P., Nickerson, S.C., Alley, T.K. (1990):

Prevalence of intramammary infection and teat canal colonization in unbred and primigravid dairy heifers.

J. Dairy Sci. 73, 107–114

Trolldenier, H. (1996):

Resistenzentwicklungen von Infektionserregern landwirtschaftlicher Nutztiere in Deutschland (1990–1994) – ein Überblick.

Dtsch. tierärztl. Wochenschr. 103, 256–260

Trolldenier, H. (1999):

Zur Resistenzproblematik in der Veterinärmedizin – Übersicht aus bundesweit erfaßten Daten.

Tierärztl. Prax. Ausg. G. 27, 317–323

Tsolov, S., Dimitrov, M., Koleva, M. (1995):

The gynaecological state of cows in relation to their mastitis morbidity.

Proceedings of the third IDF International Mastitis Seminar, Tel-Aviv, Israel, S4, 78–79

Waage, S., Sviland, S., Ødegaard, S.A. (1998a):

Identification of risk factors for clinical mastitis in dairy heifers.

J. Dairy Sci. 81, 1275–1284

Waage, S., Sviland, S., Ødegaard, S.A. (1998b):

Risk factors for clinical heifer mastitis.

Proceedings of the XXth World Buiatrics Congress, Sydney, Australia, 6-10 July, Vol. I, 231–235

Waage, S., Mørk, T., Røros, A., Aasland, D., Hunshamar, A., Ødegaard, S.A. (1999):

Bacteria associated with clinical mastitis in dairy heifers.

J. Dairy Sci. 82, 712–719

Ward, G.-E., Schultz, L.H. (1972):

Relationship of somatic cells in quarter milk to type of bacteria and production.

J. Dairy Sci. 55, 1428–1431

Weiss, W.P., Hogan, J.S., Smith, K.L., Hoblet, K.H. (1990):

Relationships among selenium, vitamin E, and mammary gland health in commercial dairy herds.

J. Dairy Sci. 73, 381–390

Weiss, W.P., Hogan, J.S., Todhunter, D.A., Smith, K.L. (1997):

Effect of vitamin E supplementation in diets with a low concentration of selenium on mammary gland health of dairy cows.

J. Dairy Sci. 80, 1728–1737

Wendt, K., Bostedt, H., Mielke, H., Fuchs, H.W. (1994):

Euter- und Gesäugekrankheiten.

Gustav Fischer Verlag, Jena, Stuttgart

Wenninger, A., Distl, O. (1994):

Harnstoff- und Azetongehalt in der Milch als Indikatoren für ernährungsbedingte Fruchtbarkeitsstörungen der Milchkuh.

Dtsch. tierärztl. Wschr. 101, 152–157

Wheelock, J.V., Smith, A., Dodd, F.H., Lyster, R.L.J. (1967):

Change in the quantity and composition of mammary gland secretion in the between lactations. 1. The beginning of the dry period.

J. Dairy Res. 34, 1–7

White, D.G., Harmon, R.J., Matos, J.E.S., Langlois, B.E. (1989):

Isolation and identification of coagulase-negative *Staphylococcus* species from bovine body sites and streak canals of nulliparous heifers.

J. Dairy Sci. 72, 1886–1892

Wilson, D.J., Gonzalez, R.N., Das, H.H. (1997):

Bovine mastitis pathogens in New York and Pennsylvania: prevalence and effects on somatic cell count and milk production.

J. Dairy Sci. 80, 2592–2598

Young, C.W., Legates, J.E., Lecce, J.G. (1960):

Genetic and phenotypic relationships between clinical mastitis, laboratory criteria, and udder height.

J. Dairy Sci. 43, 54–62

Zdunczyk, S., Ahlers, D., Grunert, E. (1992):

Zur Frage der Beziehungen zwischen bovinen klinischen Mastitiden zum Zeitpunkt der Abkalbung und Retentio secundinarum.

Dtsch. tierärztl. Wschr. 99, 386–389

Zebracki, A., Lubieniecki, B. (1976):

Metritis-Mastitis-Hypogalactie-Syndrom bei Kühen.

Dtsch. tierärztl. Wschr. 83, 551–552

Zecconi, A., Piccinini, R., Casula, A., Zeponi, A. (1998):

Epidemiological study on new intramammary infections after calving in 7 dairy herds.

Proceedings of the XXth World Buiatrics Congress, Sydney, Australia, 6–10 July, Vol. I, 291–295

Zurek, E., Foxcroft, G.R., Kennelly, J.J. (1995):

Metabolic status and interval to first ovulation in postpartum dairy cows.

J. Dairy Sci. 78, 1909–1920