

Research Article

Blue-Violet Light Irradiation Dose Dependently Decreases Carotenoids in Human Skin, Which Indicates the Generation of Free Radicals

Staffan Vandersee, Marc Beyer, Juergen Lademann, and Maxim E. Darvin

Charité-Universitätsmedizin Berlin, Department of Dermatology, Venereology and Allergology, Charitéplatz 1, 10117 Berlin, Germany

Correspondence should be addressed to Maxim E. Darvin; maxim.darvin@charite.de

Received 9 December 2014; Accepted 27 January 2015

Academic Editor: Jing Yi

Copyright © 2015 Staffan Vandersee et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In contrast to ultraviolet and infrared irradiation, which are known to facilitate cutaneous photoaging, immunosuppression, or tumour emergence due to formation of free radicals and reactive oxygen species, potentially similar effects of visible light on the human skin are still poorly characterized. Using a blue-violet light irradiation source and aiming to characterize its potential influence on the antioxidant status of the human skin, the cutaneous carotenoid concentration was measured noninvasively in nine healthy volunteers using resonance Raman spectroscopy following irradiation. The dose-dependent significant degradation of carotenoids was measured to be 13.5% and 21.2% directly after irradiation at 50 J/cm² and 100 J/cm² ($P < 0.05$). The irradiation intensity was 100 mW/cm². This is above natural conditions; the achieved doses, though, are acquirable under natural conditions. The corresponding restoration lasted 2 and 24 hours, respectively. The degradation of cutaneous carotenoids indirectly shows the amount of generated free radicals and especially reactive oxygen species in human skin. In all volunteers the cutaneous carotenoid concentration dropped down in a manner similar to that caused by the infrared or ultraviolet irradiations, leading to the conclusion that also blue-violet light at high doses could represent a comparably adverse factor for human skin.

1. Introduction

The exposure of the body to electromagnetic irradiation is a ubiquitous and lifelong event with numerous beneficial and adverse effects on the individual. Natural sunlight is by far the most important irradiation source. Visible (VIS) light (380–760 nm) represents about 40% of the emitted energy that reaches the surface of the earth [1, 2]. Nevertheless, most biological effects are attributed to the ultraviolet (UV) spectrum (290–400 nm), the rate of which is influenced by the latitude and the ozone layer's condition. The numerous effects of UV light on human skin have been subject to thorough research for decades [3]. Despite lacking the potential to penetrate deeply into the skin, UVB (290–320 nm) and UVA (320–400 nm) light exhibit a broad range of beneficial as well as adverse effects. Most of the latter effects are based on UV-induced DNA-mutation and immunosuppression [4], causing photoaging [5] and especially carcinogenesis [6]. Thus, the first sunscreens developed focused on blocking

UVB though primarily for sunburn prevention. Then, after discovering its predominantly immunosuppressive effects, also the UVA range became interesting [7, 8]. Defective DNA or cell membranes, antioxidant depletion, proinflammatory effects, collagen, and elastin degradation represent the most important causes of UV-induced skin damage [9–11]. These effects are associated with the direct photochemical effects of UV and with indirect effects related to the action of free radicals and especially reactive oxygen species (ROS), generated subsequent to UV exposure [12–14].

With these effects well addressed, and potent means of protection being available, the assessment of infrared (760–3,000 nm) (IR) and VIS in the context of photodamage moved into focus. Since 1982 [15] scientific groups worldwide have been investigating the influence of IR on the skin showing its capability of ROS generation [16–19]. In 2006, the induction of ascorbate radicals by VIS could be shown *ex vivo* [20]. In 2008, Cho et al. proved that VIS and IR in conjunction induce skin damage *in vivo*, using the activation of matrix

metalloproteinases (MMPs) and procollagen I formation as surrogate markers for the cutaneous degradation [21]. In 2009, Zastrow et al. [22] presented the action spectrum of the formation of free radicals in the skin *ex vivo* in the range between 280 and 1.600 nm. The free radicals were shown to be generated in the whole spectral range with the maximum in the UV. In 2010, the induction of ROS in the skin subsequent to IR-A (760–1.440 nm) and NIR (760–3.000 nm) irradiation was described [23, 24]. Another explanation for these findings was proposed showing the possible formation of heat-shock radicals in the skin subsequent to IR irradiation [25, 26]. However, in 2014, the different pathways for IR-induced ROS and heat-shock-induced ROS in the skin were characterized, postulating the generation of free radicals including ROS in the human skin subsequent to IR exposure independent of the applied doses [2]. In 2012, the production of ROS, proinflammatory cytokines, and MMP-1 expression in human skin subsequent to VIS exposure was reported [27]. Moreover, as the ROS generation is much greater in living skin in comparison to excised skin, the *in vivo* measurements are more informative than *ex vivo* measurements in this regard [28].

Blue-violet light (380–495 nm) is widely used in medicine for treatment of acne [29–31], psoriasis [32, 33], atopic dermatitis [34], and neonatal jaundice [35] and for wound healing [36, 37]. Moreover, the skin exposure to blue light results in antibacterial [38–40], antimicrobial [36, 41], and anti-inflammatory [42] effects. The side effects of blue-light therapy of neonates [43], suppression of dendritic cell activation [44] and effect on human dermal fibroblasts [45, 46] were also reported. The toxic effect of blue light on the skin was shown to be related to the generation of nonenzymatic nitric oxide (NO) radicals [47–49]. Blue-light-induced NO generation was proposed to be effective for therapy of local or systemic hemodynamic disorders [49]. The cytotoxic effect of blue light was even proven to be effective for treatment of superficial skin carcinomas in humans [50]. Moreover, it could be speculated that blue-light-induced ROS are responsible for the antibacterial and antimicrobial effects mentioned above.

Several systems physiologically protect the skin from oxidative stress. In addition to enzymes such as catalase, glutathione peroxidase, or superoxide dismutase, this group comprises substances that cannot be synthesized by the human organism, such as vitamins (C and E), carotenoids, flavonoids, and phytoestrogens [51, 52]. The important protective role of carotenoids in neutralizing excessive free radicals could be shown *in vivo* [53–55] as could their potential to act as representative markers for the antioxidant status of the human epidermis [56, 57].

That latter feature makes them suitable for assessing the potential effects of VIS on the skin as well. In addition, carotenoids can be dietetically supplied [58] and have been described to contribute to an enhanced cutaneous UV protection [59, 60]. This and the fact that they seem to generally correlate with the individual stress level and lifestyle factors such as alcohol consumption, smoking, and current health stage have highlighted their importance in an even more general context [61–63].

TABLE 1: Volunteer's characteristics.

Parameters	Mean (range)
Number of volunteers	9
Age	39.8 (26–63)
Gender	7 females, 2 males
Skin type according to Fitzpatrick	Type II: 8, type III: 1
Smoker/nonsmoker	2/7
Vegetarian/nonvegetarian	1/8

Thus, in order to assess the dose-dependent influence that blue-violet light can potentially exert on the antioxidant status of the skin, we exposed nine healthy volunteers to different dosages of blue-violet light and measured the kinetics course of the cutaneous carotenoid concentration by means of resonance Raman spectroscopy over a period of 24 hours after irradiation.

2. Material and Methods

2.1. Volunteers. Nine healthy volunteers (see Table 1 for volunteers' characteristics) aged between 26 and 63 years with skin types II and III according to the Fitzpatrick classification [64] were included in the study. Exclusion criteria were any kind of cutaneous diseases, pregnancy, consumption of potentially phototoxic drugs of any kind, intensive sun exposure during the last 2 weeks, and hair existence on the inner forearm. The investigations were carried out in accordance with the ethical guidelines of the Declaration of Helsinki and had been approved by the Ethics Committee of the Charité-Universitätsmedizin Berlin before the study had started. All volunteers gave their informed written consent.

The volunteers were instructed not to utilize any skin care products on the forearm for at least 72 hours and not to bathe or shower for at least 4 hours prior to the beginning and during the experiments.

2.2. Source of Blue-Light Irradiation. The skin was irradiated with blue-violet light (80% in the range 380–495 nm, maxima at 440 nm), which is normally used for defining the skin treatment area for digital phototherapy using a skintrek PT3, Lumedtec GmbH, Lüneburg, Germany. The emission spectrum can be seen in Figure 1.

2.3. Resonance Raman Spectroscopy. The carotenoids were measured in the volunteers' skin noninvasively using resonance Raman spectroscopy [65, 66]. Based on the absorption properties of the carotenoids, the resonant excitation is achieved in the blue range of the spectra [66]. Thus, the 488 nm blue-light wavelength of an argon CW laser was used as a source of carotenoid excitation. Under the applied excitation conditions, the carotenoid Raman lines are resonantly enhanced and clearly detected in the fingerprint range of the spectra on the high fluorescence background. The intensity of the carbon-carbon double-bond stretch vibration of the conjugated backbone of carotenoid molecules (C=C) measured at 1525 cm^{-1} was analyzed. This was done due to




FIGURE 1: The emission spectrum of blue-light source of irradiation.

the fact that the C=C chemical bonds of the carotenoid molecules are responsible for their antioxidant properties when the free radicals are neutralized [60, 67, 68]. Thus, the reaction with free radicals induces the destruction of the carotenoids' C=C bonds that is attributed to the decrease of the 1525 cm^{-1} Raman peak intensity. Our group has previously described the Raman device utilized in the present study in detail [66]. Resonance Raman spectroscopy was chosen in consideration of the strong advantages over other noninvasive methods [69].

2.4. Temperature Measurement. The skin surface temperature was measured using a noncontact thermometer (Rytek Schlender Messtechnik, Rüttnick, Germany).

2.5. Measurement of Light Intensity. The intensity of blue light was measured with a power meter (Hydrosun Medizintechnik GmbH, Type HBM-1, Müllheim, Germany).

2.6. Study Design. Two areas of $2 \times 2\text{ cm}^2$ were marked on the left and right inner forearms of the volunteers. Afterwards, the carotenoid concentration was measured five times at different places within the marked areas and the mean values were calculated as initial concentration. Then the blue-violet light irradiation was conducted by administering 50 J/cm^2 on the one forearm first and subsequently 100 J/cm^2 on the other forearm. The left or right forearms were randomized from volunteer to volunteer. To achieve the intended doses, the light source was positioned at such a distance to the skin surface that the light intensity exactly amounted to 100 mW/cm^2 (individually controlled by means of a power meter). Thus, the irradiation time was calculated to be 500 sec. and 1000 sec. in order to achieve a dose of 50 J/cm^2 and 100 J/cm^2 , respectively. Note that this is far above the intensity of illumination present, when using the device for UV-therapy planimetry in the therapeutical setting, which is about 3.7 mW/cm^2 (dose does not exceed 0.25 J/cm^2). The carotenoid concentration was then measured directly, as well as 1, 2, and 24 hours after blue-violet light irradiation. To




FIGURE 2: The kinetic course of the cutaneous carotenoids degradation after the irradiation with blue light at doses of 50 J/cm^2 (mean + SD, solid line) and 100 J/cm^2 (mean - SD, dotted line) and the subsequent restoration. * shows the significant degradation in comparison to the initial level before irradiation ($P < 0.05$).

exclude the formation of heat-shock radicals, the skin was continuously cooled during the irradiation by a fan and the skin temperature remotely measured with a thermometer before, during, and after irradiation.

The applied irradiation doses were selected as they represent a dosage that is readily acquirable under natural conditions by sunbathing. The range between 380 and 495 nm (violet and blue range of the spectra) represents 80% of the energy emitted by the skintrek illumination source (see Figure 1). In this wavelength range, according to the currently active ASTM sun reference spectrum, the emitted average energy accounts for about 161 W/m^2 in the USA [70]. This value corresponds to sun exposure in southern Europe. This means that the acquired dosage after one hour of sunbathing at noon would amount to about 57 J/cm^2 , which is in the range of our experimental settings (50 J/cm^2 and 100 J/cm^2).

2.7. Statistical Analyses. Statistical analyses were done by one-way or two-way analysis of variance (ANOVA) as appropriate. All analyses were conducted with GraphPad Prism, 4.0 (GraphPad Software Inc., La Jolla). Differences were considered to be significant at $P < 0.05$.

3. Results and Discussion

Figure 2 displays the kinetic course of the cutaneous carotenoids degradation after the irradiation with blue-violet light at doses of 50 J/cm^2 and 100 J/cm^2 and the subsequent restoration. A statistically significant decrease of the carotenoid concentration after the application of both doses was observed. The obtained mean magnitude of carotenoid destruction was determined as the difference between the carotenoid values measured before and directly after blue-violet light irradiation. It is about twice as high at a dosage of 100 J/cm^2 compared to 50 J/cm^2 (21.2% versus 13.5%). The median skin surface temperature measured immediately after

irradiation was 36.3°C (range 34.9–37.8) for 50 J/cm² and 36.1°C (range 32.2–38.5) for 100 J/cm². The initial skin temperature measured before irradiation was 31.5°C on average. No volunteer uttered any inconvenient sensations or other problems during the entire experiment. Some volunteers reported they felt only a slight skin temperature increase when the respective skin areas were exposed to the blue-violet light.

The carotenoid degradation in human skin subsequent to irradiation with blue-violet light can be explained by, inter alia, the direct carotenoid destruction by blue-violet light absorption, the effect of heat-shock radicals generated due to the temperature increase, and the generation of free radicals including ROS due to the blue-violet light activation of mitochondrial activity. The two first possible reasons can be eliminated as carotenoids are very stable under the applied irradiation conditions (the dissociation energy of carotenoid molecules is higher than the applied irradiation energies) [71, 72]. The temperature starts to play an important role in the generation of heat-shock radicals in the skin only when exceeding the value of 39°C [2]. The obtained reduction of the cutaneous carotenoid concentration subsequent to irradiation of human skin with blue-violet light is fully in correlation with the results obtained for UV [73], IRA, and NIR [74] irradiation of human skin *in vivo*, which were associated with the generation of free radicals and especially ROS in the skin. Thus, the dose-dependent generation of blue-violet light-induced ROS in the skin can be established confirming the results obtained previously [22, 49].

When being irradiated at doses of 50 J/cm², the carotenoid concentration in the skin was already restored one hour after exposure. In contrast to this, the restoration of the carotenoid concentration to the initial level takes 24 hours after irradiation when being irradiated at doses of 100 J/cm². Still, 2 hours after irradiation at 100 J/cm², almost 50% of the degraded carotenoids were restored. The restoration of cutaneous carotenoids after the irradiation with blue-violet light shows dose-dependent kinetics. Higher irradiation doses give rise to the generation of a higher amount of ROS and, therefore, to the depletion of higher amount of carotenoids. Consequently, more time is required to restore the carotenoid level that existed prior to irradiation. Conversely, low irradiation doses destroy a lower amount of carotenoids, which reduces the time span for restoration. The restoration efficacy is expected to be influenced by carotenoid-rich nutrition that was shown in other studies [75–77].

To determine whether or not the reference blue excitation light used in the applied resonance Raman system may trigger the same effects in the skin as irradiation with blue-violet light, the applied irradiation doses should be compared. The power of the utilized reference excitation blue light was 10 mW. The diameter of the excitation beam was 6.5 mm in order to eliminate the influence of skin inhomogeneities on the Raman measurements [65]. The exposure time was 5 sec. Thus, the illumination dose of the reference blue light was equal to 0.15 J/cm² and this is substantially lower than the applied blue-violet light irradiation doses (0.15 J/cm² versus

50/100 J/cm²). Therefore, no detectable effect of any reference excitation blue light on human skin can be affirmed.

While in the past sunscreen research addressed the measuring technology in order to exactly specify not only the UVB but also the UVA spectral ranges as a prerequisite for developing new or improving existing products, Zastrow et al. [22] revealed in 2009 that approximately 50% of the free radicals, which are induced by solar radiation in the human skin, originate from visible and infrared light. Although the radical formation in the infrared spectral range has become a topic of intense international research, studies demonstrating that high amounts of free radicals are also induced by visible and infrared light are rare. The presented study well demonstrates that visible blue-violet light also induces free radicals in human skin *in vivo*. The generation of free radicals in the remaining visible spectrum is highly probable and, therefore, is expected. These findings have the potential to provide fresh impetus to the development of sunscreens, in particular with regard to transferring from UV protection to protection in the whole solar spectrum.

4. Conclusions

The irradiation of the human skin with blue-violet light results in a dose-dependent significant degradation of the epidermal antioxidants ($P < 0.05$) as was shown by *in vivo* measurements of the carotenoid concentration using resonance Raman spectroscopy. The mean magnitude of the carotenoid destruction was determined to be 13.5% after irradiation at 50 J/cm² and 21.2% after irradiation at 100 J/cm². Depending on the irradiation dose, the restoration time was measured to be 1 hour for the dose of 50 J/cm² and 24 hours for the dose of 100 J/cm². The increase of the irradiation dose is likely to extend also both the magnitude of carotenoid destruction and the restoration time.

Based on the results obtained previously by other scientific groups, it could be established that free radicals and most probably ROS are generated in the human skin subsequent to irradiation with blue-violet light *in vivo*.

The obtained results could be essential for the development of future protection strategies based not only on UV but also on protection across the whole spectrum of the light.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgment

The authors would like to thank the Foundation “Skin Physiology” of the Donor Association for German Science and Humanities for the financial support.

References

- [1] J. E. Frederick, H. E. Snell, and E. K. Haywood, “Solar ultraviolet-radiation at the earths surface,” *Photochemistry and Photobiology*, vol. 50, pp. 443–450, 1989.

- [2] M. Y. Akhalaya, G. V. Maksimov, A. B. Rubin, J. Lademann, and M. E. Darvin, "Molecular action mechanisms of solar infrared radiation and heat on human skin," *Ageing Research Reviews*, vol. 16, no. 1, pp. 1–11, 2014.
- [3] J. Krutmann, A. Morita, and J. H. Chung, "Sun exposure: what molecular photodermatology tells us about its good and bad sides," *Journal of Investigative Dermatology*, vol. 132, no. 3, pp. 976–984, 2012.
- [4] D. A. Kelly, A. R. Young, J. M. McGregor, P. T. Seed, C. S. Potten, and S. L. Walker, "Sensitivity to sunburn is associated with susceptibility to ultraviolet radiation-induced suppression of cutaneous cell-mediated immunity," *Journal of Experimental Medicine*, vol. 191, no. 3, pp. 561–566, 2000.
- [5] F. Poon, S. Kang, and A. L. Chien, "Mechanisms and treatments of photoaging," *Photodermatology, Photoimmunology & Photomedicine*, 2014.
- [6] F. R. de Gruij, "Skin cancer and solar UV radiation," *European Journal of Cancer*, vol. 35, no. 14, pp. 2003–2009, 1999.
- [7] G. Bens, "Sunscreens," *Advances in Experimental Medicine and Biology*, vol. 810, pp. 429–463, 2014.
- [8] D. D. Moyal and A. M. Fourtanier, "Broad-spectrum sunscreens provide better protection from solar ultraviolet-simulated radiation and natural sunlight-induced immunosuppression in human beings," *Journal of the American Academy of Dermatology*, vol. 58, no. 5, pp. S149–S154, 2008.
- [9] M. E. Darvin, H. Richter, S. Ahlberg et al., "Influence of sun exposure on the cutaneous collagen/elastin fibers and carotenoids: negative effects can be reduced by application of sunscreen," *Journal of Biophotonics*, vol. 7, no. 9, pp. 735–743, 2014.
- [10] J. C. Monboisse, G. Poulin, P. Braquet, A. Randoux, C. Ferradini, and J. P. Borel, "Effect of oxy radicals on several types of collagen," *International Journal of Tissue Reactions*, vol. 6, no. 5, pp. 385–390, 1984.
- [11] Y. Kawaguchi, H. Tanaka, T. Okada et al., "Effect of reactive oxygen species on the elastin mRNA expression in cultured human dermal fibroblasts," *Free Radical Biology and Medicine*, vol. 23, no. 1, pp. 162–165, 1997.
- [12] J. H. Epstein, "Photocarcinogenesis, skin cancer, and aging," *Journal of the American Academy of Dermatology*, vol. 9, no. 4, pp. 487–502, 1983.
- [13] J. Krutmann, "The role of UVA rays in skin aging," *European Journal of Dermatology*, vol. 11, no. 2, pp. 170–171, 2001.
- [14] B. Poljšak and R. Dahmane, "Free radicals and extrinsic skin aging," *Dermatology Research and Practice*, vol. 2012, Article ID 135206, 4 pages, 2012.
- [15] L. H. Kligman, "Intensification of ultraviolet-induced dermal damage by infrared radiation," *Archives of Dermatological Research*, vol. 272, no. 3-4, pp. 229–238, 1982.
- [16] S. M. Schieke, H. Stege, V. Kürten, S. Grether-Beck, H. Sies, and J. Krutmann, "Infrared-A radiation-induced matrix metalloproteinase 1 expression is mediated through extracellular signal-regulated kinase 1/2 activation in human dermal fibroblasts," *Journal of Investigative Dermatology*, vol. 119, no. 6, pp. 1323–1329, 2002.
- [17] M. E. Darvin, I. Gersonde, H. Albrecht, L. Zastrow, W. Sterry, and J. Lademann, "In vivo Raman spectroscopic analysis of the influence of IR radiation on the carotenoid antioxidant substances beta-carotene and lycopene in the human skin. Formation of free radicals," *Laser Physics Letters*, vol. 4, no. 4, pp. 318–321, 2007.
- [18] P. Schroeder, J. Haendeler, and J. Krutmann, "The role of near infrared radiation in photoaging of the skin," *Experimental Gerontology*, vol. 43, no. 7, pp. 629–632, 2008.
- [19] P. Schroeder, J. Lademann, M. E. Darvin et al., "Infrared radiation-induced matrix metalloproteinase in human skin: implications for protection," *Journal of Investigative Dermatology*, vol. 128, no. 10, pp. 2491–2497, 2008.
- [20] R. Haywood, "Relevance of sunscreen application method, visible light and sunlight intensity to free-radical protection: a study of *ex vivo* human skin," *Photochemistry and Photobiology*, vol. 82, no. 4, pp. 1123–1131, 2006.
- [21] S. Cho, M. J. Lee, M. S. Kim et al., "Infrared plus visible light and heat from natural sunlight participate in the expression of MMPs and type I procollagen as well as infiltration of inflammatory cell in human skin in vivo," *Journal of Dermatological Science*, vol. 50, no. 2, pp. 123–133, 2008.
- [22] L. Zastrow, N. Groth, F. Klein et al., "The missing link—light-induced (280–1,600 nm) free radical formation in human skin," *Skin Pharmacology and Physiology*, vol. 22, no. 1, pp. 31–44, 2009.
- [23] M. E. Darvin, S. Haag, M. Meinke, L. Zastrow, W. Sterry, and J. Lademann, "Radical production by infrared A irradiation in human tissue," *Skin Pharmacology and Physiology*, vol. 23, no. 1, pp. 40–46, 2010.
- [24] M. E. Darvin, S. F. Haag, J. Lademann, L. Zastrow, W. Sterry, and M. C. Meinke, "Formation of free radicals in human skin during irradiation with infrared light," *Journal of Investigative Dermatology*, vol. 130, no. 2, pp. 629–631, 2010.
- [25] H. Piazena and D. K. Kelleher, "Effects of infrared-a irradiation on skin: discrepancies in published data highlight the need for an exact consideration of physical and photobiological laws and appropriate experimental settings," *Photochemistry and Photobiology*, vol. 86, no. 3, pp. 687–705, 2010.
- [26] T. Jung, A. Höhn, H. Piazena, and T. Grune, "Effects of water-filtered infrared A irradiation on human fibroblasts," *Free Radical Biology and Medicine*, vol. 48, no. 1, pp. 153–160, 2010.
- [27] F. Liebel, S. Kaur, E. Ruvolo, N. Kollias, and M. D. Southall, "Irradiation of skin with visible light induces reactive oxygen species and matrix-degrading enzymes," *Journal of Investigative Dermatology*, vol. 132, no. 7, pp. 1901–1907, 2012.
- [28] M. C. Meinke, R. Müller, A. Bechtel et al., "Evaluation of carotenoids and reactive oxygen species (ROS) in human skin after UV-irradiation: a critical comparison between in vivo and ex vivo investigations," *Experimental Dermatology*, 2014.
- [29] R. G. Wheeland and S. Dhawan, "Evaluation of self-treatment of mild-to-moderate facial acne with a blue light treatment system," *Journal of Drugs in Dermatology*, vol. 10, no. 6, pp. 596–602, 2011.
- [30] M. H. Gold, W. Sensing, and J. A. Biron, "Clinical efficacy of home-use blue-light therapy for mild-to moderate acne," *Journal of Cosmetic and Laser Therapy*, vol. 13, no. 6, pp. 308–314, 2011.
- [31] H. H. Kwon, J. B. Lee, J. Y. Yoon et al., "The clinical and histological effect of home-use, combination blue-red LED phototherapy for mild-to-moderate acne vulgaris in Korean patients: a double-blind, randomized controlled trial," *British Journal of Dermatology*, vol. 168, no. 5, pp. 1088–1094, 2013.
- [32] A. Weinstabl, S. Hoff-Lesch, H. F. Merk, and V. von Felbert, "Prospective randomized study on the efficacy of blue light in the treatment of psoriasis vulgaris," *Dermatology*, vol. 223, no. 3, pp. 251–259, 2011.
- [33] M. M. Kleinpenning, M. E. Otero, P. E. J. van Erp, M. J. P. Gerritsen, and P. C. M. van de Kerkhof, "Efficacy of blue

- light vs. red light in the treatment of psoriasis: a double-blind, randomized comparative study," *Journal of the European Academy of Dermatology and Venereology*, vol. 26, no. 2, pp. 219–225, 2012.
- [34] D. Becker, E. Langer, M. Seemann et al., "Clinical efficacy of blue light full body irradiation as treatment option for severe atopic dermatitis," *PLoS ONE*, vol. 6, no. 6, Article ID e20566, 2011.
- [35] J. F. Ennever, "Blue light, green light, white light, more light: treatment of neonatal jaundice," *Clinics in Perinatology*, vol. 17, no. 2, pp. 467–481, 1990.
- [36] Y. Zhang, Y. Zhu, A. Gupta et al., "Antimicrobial blue light therapy for multidrug-resistant *Acinetobacter baumannii* infection in a mouse burn model: implications for prophylaxis and treatment of combat-related wound infections," *Journal of Infectious Diseases*, vol. 209, no. 12, pp. 1963–1971, 2014.
- [37] N. Adamskaya, P. Dungal, R. Mittermayr et al., "Light therapy by blue LED improves wound healing in an excision model in rats," *Injury*, vol. 42, no. 9, pp. 917–921, 2011.
- [38] J. S. Guffey and J. Wilborn, "Effects of combined 405-nm and 880-nm light on *Staphylococcus aureus* and *Pseudomonas aeruginosa* in vitro," *Photomedicine and Laser Surgery*, vol. 24, no. 6, pp. 680–683, 2006.
- [39] T. Dai, A. Gupta, Y.-Y. Huang et al., "Blue light eliminates community-acquired methicillin-resistant *Staphylococcus aureus* in infected mouse skin abrasions," *Photomedicine and Laser Surgery*, vol. 31, no. 11, pp. 531–538, 2013.
- [40] T. G. S. Denis, T. Dai, and M. R. Hamblin, "Killing bacterial spores with blue light: when innate resistance meets the power of light," *Photochemistry and Photobiology*, vol. 89, no. 1, pp. 2–4, 2013.
- [41] R. Yin, T. Dai, P. Avci et al., "Light based anti-infectives: ultraviolet C irradiation, photodynamic therapy, blue light, and beyond," *Current Opinion in Pharmacology*, vol. 13, no. 5, pp. 731–762, 2013.
- [42] E. Shnitkind, E. Yaping, S. Geen, A. R. Shalita, and W.-L. Lee, "Anti-inflammatory properties of narrow-band blue light," *Journal of Drugs in Dermatology*, vol. 5, no. 7, pp. 605–610, 2006.
- [43] J. Oláh, E. Tóth-Molnár, L. Kemény, and Z. Csoma, "Long-term hazards of neonatal blue-light phototherapy," *British Journal of Dermatology*, vol. 169, no. 2, pp. 243–249, 2013.
- [44] M. R. Fischer, M. Abel, S. Lopez Kostka, B. Rudolph, D. Becker, and E. von Stebut, "Blue light irradiation suppresses dendritic cells activation in vitro," *Experimental Dermatology*, vol. 22, no. 8, pp. 558–560, 2013.
- [45] C. Opländer, S. Hidding, F. B. Werners, M. Born, N. Pallua, and C. V. Suschek, "Effects of blue light irradiation on human dermal fibroblasts," *Journal of Photochemistry and Photobiology B: Biology*, vol. 103, no. 2, pp. 118–125, 2011.
- [46] S. Bonatti, B. Hochman, V. M. Tucci-Viegas et al., "In vitro effect of 470 nm LED (light emitting diode) in keloid fibroblasts," *Acta Cirurgica Brasileira*, vol. 26, no. 1, pp. 25–30, 2011.
- [47] R. Mittermayr, A. Osipov, C. Piskernik et al., "Blue laser light increases perfusion of a skin flap via release of nitric oxide from hemoglobin," *Molecular Medicine*, vol. 13, no. 1-2, pp. 22–29, 2007.
- [48] J. Liebmann, M. Born, and V. Kolb-Bachofen, "Blue-light irradiation regulates proliferation and differentiation in human skin cells," *Journal of Investigative Dermatology*, vol. 130, no. 1, pp. 259–269, 2010.
- [49] C. Opländer, A. Deck, C. M. Volkmar et al., "Mechanism and biological relevance of blue-light (420–453 nm)-induced nonenzymatic nitric oxide generation from photolabile nitric oxide derivatives in human skin in vitro and in vivo," *Free Radical Biology and Medicine*, vol. 65, pp. 1363–1377, 2013.
- [50] A. Sparsa, K. Faucher, V. Sol et al., "Blue light is phototoxic for B16f10 murine melanoma and bovine endothelial cell lines by direct cytotoxic effect," *Anticancer Research*, vol. 30, no. 1, pp. 143–148, 2010.
- [51] M. Darvin, L. Zastrow, W. Sterry, and J. Lademann, "Effect of supplemented and topically applied antioxidant substances on human tissue," *Skin Pharmacology and Physiology*, vol. 19, no. 5, pp. 238–247, 2006.
- [52] J. J. Thiele, C. Schroeter, S. N. Hsieh, M. Podda, and L. Packer, "The antioxidant network of the stratum corneum," *Current Problems in Dermatology*, vol. 29, pp. 26–42, 2001.
- [53] J. Lademann, S. Schanzer, M. Meinke, W. Sterry, and M. E. Darvin, "Interaction between carotenoids and free radicals in human skin," *Skin Pharmacology and Physiology*, vol. 24, no. 5, pp. 238–244, 2011.
- [54] N. I. Krinsky, "Carotenoids as antioxidants," *Nutrition*, vol. 17, no. 10, pp. 815–817, 2001.
- [55] M. E. Darvin, J. W. Fluhr, S. Schanzer et al., "Dermal carotenoid level and kinetics after topical and systemic administration of antioxidants: enrichment strategies in a controlled in vivo study," *Journal of Dermatological Science*, vol. 64, no. 1, pp. 53–58, 2011.
- [56] S. F. Haag, B. Taskoparan, M. E. Darvin et al., "Determination of the antioxidative capacity of the skin in vivo using resonance Raman and electron paramagnetic resonance spectroscopy," *Experimental Dermatology*, vol. 20, no. 6, pp. 483–487, 2011.
- [57] M. C. Meinke, A. Friedrich, K. Tscherch et al., "Influence of dietary carotenoids on radical scavenging capacity of the skin and skin lipids," *European Journal of Pharmaceutics and Biopharmaceutics*, vol. 84, no. 2, pp. 365–373, 2013.
- [58] J. Lademann, A. Patzelt, S. Schanzer et al., "Uptake of antioxidants by natural nutrition and supplementation: pros and cons from the dermatological point of view," *Skin Pharmacology and Physiology*, vol. 24, no. 5, pp. 269–273, 2011.
- [59] E. Fernández-García, "Skin protection against UV light by dietary antioxidants," *Food & Function*, vol. 5, no. 9, pp. 1994–2003, 2014.
- [60] W. Stahl and H. Sies, "Photoprotection by dietary carotenoids: concept, mechanisms, evidence and future development," *Molecular Nutrition and Food Research*, vol. 56, no. 2, pp. 287–295, 2012.
- [61] M. E. Darvin, A. Patzelt, F. Knorr, U. Blume-Peytavi, W. Sterry, and J. Lademann, "One-year study on the variation of carotenoid antioxidant substances in living human skin: influence of dietary supplementation and stress factors," *Journal of Biomedical Optics*, vol. 13, no. 4, Article ID 044028, 2008.
- [62] J. Lademann, W. Köcher, R. Yu et al., "Cutaneous carotenoids: the mirror of lifestyle?" *Skin Pharmacology and Physiology*, vol. 27, no. 4, pp. 201–207, 2014.
- [63] M. E. Darvin, W. Sterry, J. Lademann, and A. Patzelt, "Alcohol consumption decreases the protection efficiency of the antioxidant network and increases the risk of sunburn in human skin," *Skin Pharmacology and Physiology*, vol. 26, no. 1, pp. 45–51, 2013.
- [64] T. B. Fitzpatrick, "The validity and practicality of sun-reactive skin types I through VI," *Archives of Dermatology*, vol. 124, no. 6, pp. 869–871, 1988.
- [65] M. E. Darvin, I. Gersonde, S. Ey et al., "Noninvasive detection of beta-carotene and lycopene in human skin using raman spectroscopy," *Laser Physics*, vol. 14, no. 2, pp. 231–233, 2004.

- [66] M. E. Darvin, I. Gersonde, H. Albrecht, S. A. Gonchukov, W. Sterry, and J. Lademann, "Determination of beta carotene and lycopene concentrations in human skin using resonance Raman spectroscopy," *Laser Physics*, vol. 15, no. 2, pp. 295–299, 2005.
- [67] N. I. Krinsky and E. J. Johnson, "Carotenoid actions and their relation to health and disease," *Molecular Aspects of Medicine*, vol. 26, no. 6, pp. 459–516, 2005.
- [68] M. E. Darvin, W. Sterry, J. Lademann, and T. Vergou, "The role of carotenoids in human skin," *Molecules*, vol. 16, no. 12, pp. 10491–10506, 2011.
- [69] M. E. Darvin, M. C. Meinke, W. Sterry, and J. Lademann, "Optical methods for noninvasive determination of carotenoids in human and animal skin," *Journal of Biomedical Optics*, vol. 18, no. 6, Article ID 61230, 2013.
- [70] ASTM, "Standard tables for reference solar spectral irradiances: direct normal and hemispherical on 37° tilted surface," Tech. Rep. G173-03 A, ASTM International, Conshohocken, Pa, USA, 2012.
- [71] J. S. Hurst, J. E. Centreras, W. G. Siems, and F. J. G. M. van Kuijk, "Oxidation of carotenoids by heat and tobacco smoke," *BioFactors*, vol. 20, no. 1, pp. 23–35, 2004.
- [72] C. S. Boon, D. J. McClements, J. Weiss, and E. A. Decker, "Factors influencing the chemical stability of carotenoids in foods," *Critical Reviews in Food Science and Nutrition*, vol. 50, no. 6, pp. 515–532, 2010.
- [73] M. E. Darvin, I. Gersonde, H. Albrecht, W. Sterry, and J. Lademann, "In vivo Raman spectroscopic analysis of the influence of UV radiation on carotenoid antioxidant substance degradation of the human skin," *Laser Physics*, vol. 16, no. 5, pp. 833–837, 2006.
- [74] M. E. Darvin, A. Patzelt, M. Meinke, W. Sterry, and J. Lademann, "Influence of two different IR radiators on the antioxidative potential of the human skin," *Laser Physics Letters*, vol. 6, no. 3, pp. 229–234, 2009.
- [75] M. E. Darvin, I. Gersonde, H. Albrecht, W. Sterry, and J. Lademann, "Resonance Raman spectroscopy for the detection of carotenoids in foodstuffs. Influence of the nutrition on the antioxidative potential of the skin," *Laser Physics Letters*, vol. 4, no. 6, pp. 452–456, 2007.
- [76] S. Jung, M. E. Darvin, H.-S. Chung et al., "Antioxidants in asian-korean and caucasian skin: the influence of nutrition and stress," *Skin Pharmacology and Physiology*, vol. 27, no. 6, pp. 293–302, 2014.
- [77] S. T. Mayne, B. Cartmel, S. Scarmo et al., "Noninvasive assessment of dermal carotenoids as a biomarker of fruit and vegetable intake," *American Journal of Clinical Nutrition*, vol. 92, no. 4, pp. 794–800, 2010.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

