

**2,3-Bis(1-methylimidazol-2-yl)quinoxaline (bmiq), a new ligand with decoupled electron transfer and metal coordination sites: the very different redox behaviour of isoelectronic complexes with [PtCl<sub>2</sub>] and [AuCl<sub>2</sub>]<sup>+</sup>†**Ece Bulak,<sup>a</sup> Tereza Varnali,<sup>b</sup> Brigitte Schwederski,<sup>a</sup> Biprajit Sarkar,<sup>a</sup> Ingo Hartenbach,<sup>a</sup> Jan Fiedler<sup>c</sup> and Wolfgang Kaim<sup>\*a</sup>

Received 24th September 2010, Accepted 9th December 2010

DOI: 10.1039/c0dt01282c

The new, potentially ambidentate heterocyclic ligand 2,3-bis(1-methylimidazol-2-yl)quinoxaline (bmiq) was obtained from 2,3-bis(1-methylimidazol-2-yl)glyoxal and 1,2-diaminobenzene. Its coordination to PtCl<sub>2</sub> and to the isoelectronic [AuCl<sub>2</sub>]<sup>+</sup> in [AuCl<sub>2</sub>(bmiq)](AuCl<sub>4</sub>) occurs *via* the imine N donors of the imidazolyl groups, leading to the formation of seven-membered chelate rings with boat conformation. According to the spectroelectrochemistry (UV-vis-NIR, EPR), the reversible electron addition to the [PtCl<sub>2</sub>(bmiq)] and the free ligand takes place in the (non-coordinated) quinoxaline part of the molecule, similarly as for related complexes of dipyrido[3,2-*a*:2',3'-*c*]phenazines (dppz), 2,3-bis(2-pyridyl)quinoxalines (bpq) and 2,3-bis(dialkylphosphino)quinoxalines (QuinoxP). DFT calculations confirm the experimental results (structures, spectroscopy) and also point to the coordination potential of the quinoxaline N atoms. The electron addition to [AuCl<sub>2</sub>(bmiq)]<sup>+</sup> takes place not at the ligand but at the metal site, according to experimental and DFT results.

**Introduction**

The coordination compounds of ligands in which the metal binding location and the electron transfer site are spatially and electronically separated have raised attention in various areas. The molecule dipyrido[3,2-*a*:2',3'-*c*]phenazine (dppz, Scheme 1) and its many derivatives were thus shown by various techniques to exhibit such a situation<sup>1</sup> and the ability of certain complexes of dppz to exhibit environmentally and, especially, intercalation dependent luminescence made them popular as “molecular light switches”.<sup>2</sup> Specifically, the chelate coordination to form a five-membered ring

occurs at the  $\alpha$ -diimine site of the 1,10-phenanthroline part of the molecule, whereas the rather facile electron addition takes place mainly in the quinoxaline moiety.<sup>1a</sup>

The 2,3-bis(dialkylphosphino)quinoxaline ligands (QuinoxP, Scheme 1), although developed for the purpose of stereoselective organic catalysis with an oxidatively stable diphosphine,<sup>3</sup> were shown to exhibit a similar separation of chelate coordination activity (at the diphosphine site) and electron transfer (at the quinoxaline heterocycle).<sup>4</sup>

A related type of ligand is based on the commercially available 2,3-bis(2-pyridyl)quinoxaline (dpq).<sup>5</sup>

We have now obtained another such system, 2,3-bis(1-methylimidazol-2-yl)quinoxaline (bmiq, Scheme 2), by reacting 1,2-diaminobenzene (*o*-phenylenediamine) with the known 2,3-bis(1-methylimidazol-2-yl)glyoxal.<sup>6</sup>


In addition to using the biorelevant imidazole substituents, reminiscent of the histidine side chain,<sup>7,8</sup> we were interested in probing the possibility of chelation *via* both imidazol-2-yl imine

<sup>a</sup>Institut für Anorganische Chemie, Universität Stuttgart, Pfaffenwaldring 55, D-70550 Stuttgart, Germany. E-mail: kaim@iac.uni-stuttgart.de

<sup>b</sup>Chemistry Department, Bogazici University, Bebek, Istanbul 34342, Turkey

<sup>c</sup>J. Heyrovský Institute of Physical Chemistry, v.v.i., Academy of Sciences of the Czech Republic, Dolejškova 3, CZ-18223 Prague, Czech Republic

† CCDC reference numbers 792125–792127. For crystallographic data in CIF or other electronic format see DOI: 10.1039/c0dt01282c



Scheme 1



**Table 1** Crystallographic data

	bmiq	[PtCl <sub>2</sub> (bmiq)]	[AuCl <sub>2</sub> (bmiq)](AuCl <sub>4</sub> )
empirical formula	C <sub>16</sub> H <sub>14</sub> N <sub>6</sub>	C <sub>16</sub> H <sub>14</sub> Cl <sub>2</sub> N <sub>6</sub> Pt	C <sub>16</sub> H <sub>14</sub> Au <sub>2</sub> Cl <sub>6</sub> N <sub>6</sub>
FW	290.33	556.32	896.97
Space group	<i>P</i> 2 <sub>1</sub> / <i>n</i>	<i>P</i> 2 <sub>1</sub> / <i>c</i>	<i>P</i> 2 <sub>1</sub> / <i>n</i>
<i>a</i> /Å	9.0875(5)	14.5272(7)	9.7732(2)
<i>b</i> /Å	10.3843(7)	9.5752(4)	24.6103(5)
<i>c</i> /Å	14.7640(8)	13.5829(7)	10.1615(2)
$\beta$ /°	105.664(3)	101.724(2)	107.0950(10)
<i>V</i> /Å <sup>3</sup>	1341.50(14)	1849.98(12)	2336.08(8)
<i>Z</i>	4	4	4
<i>D<sub>c</sub></i> /g cm <sup>-3</sup>	1.438	1.997	2.550
$\mu$ /mm <sup>-1</sup>	0.09	7.89	13.25
<i>R</i> <sub>int</sub>	0.105	0.044	0.047
<i>R</i> <sub>1</sub>	0.090	0.060	0.047
w <i>R</i> <sub>2</sub>	0.163	0.110	0.071
Goof	1.079	1.057	1.021
largest difference peak/hole/e Å <sup>-3</sup>	0.382/-0.343	3.143/-3.196	1.840/-1.926

[PtCl<sub>2</sub>(bmiq)], colourless platelet (0.7 × 0.6 × 0.2 mm) and [AuCl<sub>2</sub>(bmiq)](AuCl<sub>4</sub>), reddish platelet (2.2 × 1.2 × 0.8 mm). A NONIUS Kappa CCD diffractometer was used at 100 K, additional crystallographic information is given in Table 1. The structures were solved using direct methods with refinement by full-matrix least squares of *F*<sup>2</sup>, employing the program system SHELXL 97<sup>20</sup> in connection with a numerical absorption correction in case of bmiq and an empirical correction for [PtCl<sub>2</sub>(bmiq)] and [AuCl<sub>2</sub>(bmiq)](AuCl<sub>4</sub>). All non-hydrogen atoms were refined anisotropically.

### DFT calculations


Density functional theory (DFT) calculations were carried out using the program package GAUSSIAN.<sup>21</sup> The hybrid functional B3LYP (UB3LYP for radical anions) with the LANL2DZ basis set was employed.<sup>6c</sup> The ligand, the complexes and their one-electron reduced forms were subjected to calculations. Also, PtCl<sub>2</sub> complexes yielding a 7-membered (A) and a 5-membered (B) chelate ring were studied (see Scheme 3).

## Results and discussion


### Synthesis and structure

The new heterocycle bmiq has been obtained *via* a standard synthetic condensation procedure for preparing quinoxalines,<sup>19</sup> using the previously introduced<sup>6</sup> 2,3-bis(1-methylimidazol-2-yl)glyoxal. The compound was characterised by single-crystal X-ray crystal diffraction (Table 1), the molecules in the crystal studied exhibit a pronounced, unsymmetrical twisting between the planes of the quinoxaline  $\pi$ -acceptor system and the  $\pi$  electron donating methylimidazolyl substituents (Fig. 1, Table 2).

The complexation of bmiq with PtCl<sub>2</sub> involving the [PtCl<sub>2</sub>(dmsO)<sub>2</sub>] precursor gave a single mononuclear product, [PtCl<sub>2</sub>(bmiq)], characterised by <sup>1</sup>H-NMR and single-crystal X-ray crystallography as the symmetrically methylimidazolyl-chelated coordination compound (Fig. 2). The resulting seven-membered ring, involving both imine functions of the imidazoles, is not planar but exhibits a typical<sup>6</sup> boat conformation (Table 2). The dihedral



**Fig. 1** The molecular structure of bmiq in the crystal (ellipsoid probability: 95%).



**Fig. 2** The molecular structure of [PtCl<sub>2</sub>(bmiq)] in the crystal (ellipsoid probability: 95%).


angles between the quinoxaline and the imidazolyl rings amount to about 69°.

A similar chelation and conformation was found for [AuCl<sub>2</sub>(bmiq)]<sup>+</sup>, which was obtained as the tetrachloroaurate salt from the reaction of bmiq with KAuCl<sub>4</sub> (Fig. 3, Table 2).

Significant intermolecular interactions were not observed for the compounds presented here, presumably due to the non-planar molecular structures with severely twisted methylimidazolyl substituents. The structural properties of bmiq and its complexes could be well reproduced by DFT calculations (Table 2). This method suggested a higher energy for the five-membered chelate alternative B (Scheme 3) over the experimentally observed seven-membered chelate form A by 12.42 kcal mol<sup>-1</sup> in the case of the PtCl<sub>2</sub> complex.

**Table 2** Experimental and calculated bond lengths and angles for compounds

Bond lengths/Å	bmiq (exp)	bmiq (DFT)	bmiq <sup>-</sup> (DFT)	[PtCl <sub>2</sub> (bmiq)] (exp)	[PtCl <sub>2</sub> (bmiq)] (DFT)	[PtCl <sub>2</sub> (bmiq)] <sup>-</sup> (DFT)	[AuCl <sub>2</sub> (bmiq)] <sup>+</sup> (exp)	[AuCl <sub>2</sub> (bmiq)] <sup>+</sup> (DFT)	[AuCl <sub>2</sub> (bmiq)] <sup>+</sup> (DFT)
N1–C1	1.321(3)	1.348	1.358	1.349(8)	1.355	1.360	1.332(7)	1.359	1.392
N1–C3	1.377(3)	1.386	1.391	1.364(8)	1.387	1.392	1.374(7)	1.390	1.386
N2–C1	1.364(3)	1.389	1.398	1.346(8)	1.385	1.387	1.353(7)	1.374	1.388
N2–C2	1.373(3)	1.390	1.397	1.371(8)	1.391	1.396	1.374(7)	1.393	1.392
C5–C10	1.439(3)	1.457	1.413	1.432(9)	1.457	1.418	1.421(7)	1.450	1.458
C5–N6	1.324(3)	1.340	1.395	1.327(7)	1.341	1.379	1.322(7)	1.342	1.344
C10–N5	1.322(3)	1.339	1.370	1.322(8)	1.342	1.379	1.300(7)	1.342	1.344
M–Cl1				2.300(2)	2.405	2.429	2.268(1)	2.383	2.504
M–Cl2				2.308(2)	2.405	2.429	2.257(1)	2.383	2.504
M–N1				2.005(5)	2.029	2.030	2.015(4)	2.058	2.228
M–N3				1.991(5)	2.029	2.030	2.014(4)	2.058	2.228
Bond angles/°									
N3–M–Cl1				89.5(2)	88.57	89.16	90.8(1)	89.96	88.82
N3–M–Cl2				177.3(2)	177.98	177.27	178.2(1)	178.63	172.36
N1–M–N3				88.4(2)	90.12	89.61	88.5(2)	89.50	85.43
N1–M–Cl2				89.9(2)	88.57	89.16	90.2(1)	89.96	89.96
Cl2–M–Cl1				92.2(1)	92.70	91.97	90.47(5)	90.54	96.36
Dihedral angle/°									
C1N1C3C2N2/ C6N3C7C8N4	48.91	61.31	76.21	69.03	66.97	64.80	71.15	64.97	61.53

**Fig. 3** The molecular structure of [AuCl<sub>2</sub>(bmiq)]<sup>+</sup> in the crystal of [AuCl<sub>2</sub>(bmiq)][AuCl<sub>4</sub>] (ellipsoid probability: 95%).

Although PtCl<sub>2</sub> compounds in particular can be light emitting on excitation,<sup>10</sup> the complex of bmiq was found to be non-luminescent in the solid or in CH<sub>2</sub>Cl<sub>2</sub> solution, probably due to non-radiative decay involving the non-rigid methylimidazolyliquinoline functions. The compound [AuCl<sub>2</sub>(bmiq)](AuCl<sub>4</sub>) showed emission at 440 nm after excitation at 300 nm in dichloromethane.


### One-electron reduction

The one-electron character of the primary reduction processes in the free ligand bmiq and in the complexes was checked coulometrically and, in the case of reversible steps, was confirmed by the nearly 60 mV difference between the cathodic peak and

**Table 3** Electrochemical data of compounds

compound	solvent <sup>a</sup>	T/K	redox potential <sup>b</sup> (V vs. Fc <sup>+0</sup> )
bmiq	CH <sub>2</sub> Cl <sub>2</sub>	293	-2.31 [70] <sup>c</sup>
[PtCl <sub>2</sub> (bmiq)]	CH <sub>2</sub> Cl <sub>2</sub>	293	-1.62 [80] <sup>c</sup>
[AuCl <sub>2</sub> (bmiq)][AuCl <sub>4</sub> ]	C <sub>3</sub> H <sub>7</sub> CN	203	<sup>d</sup>

<sup>a</sup> 0.1 mol dm<sup>-3</sup> Bu<sub>4</sub>NPF<sub>6</sub> as electrolyte. <sup>b</sup> Half-wave potentials measured at 100 mV s<sup>-1</sup> scan rate; [peak potential differences/mV] <sup>c</sup> *i*<sub>pc</sub>/*i*<sub>pa</sub> ≈ 1.0. <sup>d</sup> Irreversible reduction, interfering waves around 0.0 V from the two Au<sup>III</sup> species.

**Fig. 4** The cyclic voltammogram of [PtCl<sub>2</sub>(bmiq)] in CH<sub>2</sub>Cl<sub>2</sub>/0.1 mol dm<sup>-3</sup> Bu<sub>4</sub>NPF<sub>6</sub> (100 mV s<sup>-1</sup> scan rate).

the anodic counter-peak in cyclic voltammetry (Fig. 4). Whereas bmiq and its PtCl<sub>2</sub> complex could be reduced reversibly to the persistent radical anions bmiq<sup>-</sup> and [PtCl<sub>2</sub>(bmiq)]<sup>-</sup> under ambient conditions, the gold(III) species [AuCl<sub>2</sub>(bmiq)](AuCl<sub>4</sub>) showed only irreversible electron uptake, even at -70 °C in *n*-butyronitrile. The inability to detect an *in situ* EPR signal leaves the formation of a persistent species [AuCl<sub>2</sub>(bmiq)]<sup>-</sup> in question. The complexes are more easily reduced than the free ligand (Table 3), in agreement with the changed structure and with the






Fig. 5 ESR spectrum of  $\text{bmqi}^{\bullet-}$  obtained by reduction of  $\text{bmqi}$  with K in THF at 293 K.




Fig. 6 ESR spectrum of electrochemically generated  $[\text{PtCl}_2(\text{bmqi})]^{\bullet-}$  at 293 K in  $\text{CH}_2\text{Cl}_2/0.1 \text{ mol dm}^{-3} \text{ Bu}_4\text{NPF}_6$ .

$\sigma$ -acceptor functions of the dichlorometal fragments. Both  $\text{bmqi}^{\bullet-}$  and  $[\text{PtCl}_2(\text{bmqi})]^{\bullet-}$  show hyperfine-split EPR signals (Fig. 5 and 6), the spectra being dominated by 1:2:3:2:1 quintet coupling from two equivalent  $^{14}\text{N}$  atoms, of 6.2 G and 6.0 G for  $\text{bmqi}^{\bullet-}$  and  $[\text{PtCl}_2(\text{bmqi})]^{\bullet-}$ , respectively ( $^{195}\text{Pt}$  coupling estimated at  $\leq 10$  G). These values indicate<sup>22</sup> spin concentration in the pyrazine ring of the quinoxaline part of the molecule, an assumption which is supported by DFT spin density calculations (Fig. 7 and 8) and also by the structural changes calculated for the radical anions  $\text{bmqi}^{\bullet-}$  and  $[\text{PtCl}_2(\text{bmqi})]^{\bullet-}$  (Table 2). The major changes on reduction occur in the pyrazine ring, while the coordination is mainly affected by the added charge. It is concluded, therefore, that the electronic structure of  $\text{bmqi}$  and its complexes is similar to that of the corresponding  $\text{dppz}$ ,  $\text{bpq}$  and Quinox-P ligands, *i.e.* the electron transfer site (quinoxaline) is spatially and electronically separated from the metal binding site<sup>1a,4</sup> (here: the imidazole rings).




Fig. 7 DFT calculated spin densities for  $\text{bmqi}^{\bullet-}$ .

Although some additional hyperfine splitting is visible in the EPR spectra (Fig. 5 and 6), the total number of theoretical lines




Fig. 8 DFT calculated spin densities for  $[\text{PtCl}_2(\text{bmqi})]^{\bullet-}$ .

(70 875) from 8 different nuclei ( $5 \times ^1\text{H}$ ,  $3 \times ^{14}\text{N}$ ) in  $\text{bmqi}^{\bullet-}$  prevented us from further analysis and assignment of the smaller ( $\leq 1$  G) EPR coupling constants.

The isotropic  $g$  values of  $\text{bmqi}^{\bullet-}$  and  $[\text{PtCl}_2(\text{bmqi})]^{\bullet-}$  differ only slightly (2.0034 and 2.0004), which confirms the hypothesis on the separation of spin localisation and metal binding. The dichloroplatinum(II) complexes of normal  $\alpha$ -diimines (where electron accommodation and coordination are not separated) show a much more pronounced shift of the isotropic  $g$  and a considerable  $g$  anisotropy.<sup>23</sup>

The inability to detect a neutral paramagnetic species  $[\text{AuCl}_2(\text{bmqi})]^{\bullet}$  by EPR could be due to its instability even at  $-70$  °C or to very rapid EPR relaxation with concomitant line broadening of a genuine gold(II) compound. True  $\text{Au}^{\text{II}}$  compounds are still elusive even if several kinds of assumed gold(II) species with extensive spin delocalisation to S and similar donor ligands were reported.<sup>24</sup> In agreement with these results, the DFT calculations predict considerable structural changes around the metal after reduction of  $[\text{AuCl}_2(\text{bmqi})]^{\bullet}$  (Table 2).

The UV-vis spectroelectrochemical reduction of the neutral precursors to  $\text{bmqi}^{\bullet-}$  and  $[\text{PtCl}_2(\text{bmqi})]^{\bullet-}$  is illustrated in Fig. 9 and 10. The similarity confirms the comparable site of the electron addition, *i.e.* the quinoxaline part of  $\text{bmqi}$ , with its typical quinoxaline radical absorption of around 600 nm.<sup>4,25</sup>





Fig. 9 UV-vis spectroelectrochemical monitoring of the reduction of  $\text{bmqi}$  in  $\text{CH}_3\text{CN}/0.1 \text{ mol dm}^{-3} \text{ Bu}_4\text{NPF}_6$ .




**Fig. 10** UV-vis spectroelectrochemical reduction of  $[\text{PtCl}_2(\text{bmiq})]$  in  $\text{CH}_3\text{CN}/0.1 \text{ mol dm}^{-3} \text{ Bu}_4\text{NPF}_6$ .

The distinct differences between the isoelectronic platinum(II) and gold(III) complexes are best explained using DFT calculated MO energies and diagrams (Table 4, Fig. 11). While the absolute energies are dependant on the (different) charges, the *relative* orbital ordering reveals a significant dichotomy: the positively charged gold system exhibits much more stabilised metal-based orbitals, leading to a LUMO involving mainly Au–Cl and Au–N  $\sigma$  bonds. The occupation of such MOs generally causes chemical reactivity such as chloride dissociation.<sup>26</sup> In contrast, the dichloroplatinum analogue has the unoccupied metal-chloride

**Table 4** DFT calculated orbital energies of complexes

Orbitals	Orbital energies (DFT)	
	$[\text{PtCl}_2(\text{bmiq})]$	$[\text{AuCl}_2(\text{bmiq})]^+$
$\pi^*$	–0.117, –0.088, –0.034	–0.224, –0.202, –0.149
$\sigma^*(\text{M–Cl})$	–0.028	–0.263
HOMO	–0.198	–0.373



**Fig. 11** The DFT-calculated frontier orbital representations of the one-electron reduced gold and platinum complexes.

$\sigma$  bond orbitals lying above the bmiq ligand-located  $\pi^*$  MOs (Table 4), which results in the observed reversible formation of a persistent anion radical complex.

Summarising, we could add a new member to the series (Scheme 1) of quinoxaline-based chelate ligands with separated metal binding and electron transfer sites. The ligand bmiq provides imidazole-imine N donors for coordination and could thus model the binding by two histidine side chains. Moreover, we could analyse the differences between the isoelectronic  $[\text{PtCl}_2]$  and  $[\text{AuCl}_2]^+$  complexes experimentally (by cyclic voltammetry, EPR and UV-vis spectroelectrochemistry) and by DFT. The relative positioning of the unoccupied metal-chloride vs. ligand  $\pi^*$  orbitals determines the presence or absence of reactivity on one-electron reduction of the  $[\text{MCl}_2(\text{bmiq})]^n$ .

## Acknowledgements

This work was supported by Deutsche Forschungsgemeinschaft and Fonds der Chemischen Industrie.

## References

- (a) J. Fees, W. Kaim, M. Moscherosch, W. Matheis, J. Klima, M. Krejci and S. Zalis, *Inorg. Chem.*, 1993, **32**, 166; (b) J. Fees, M. Ketterle, A. Klein, J. Fiedler and W. Kaim, *J. Chem. Soc., Dalton Trans.*, 1999, 2595; (c) A. Klein, T. Scheiring and W. Kaim, *Z. Anorg. Allg. Chem.*, 1999, **625**, 1177; (d) P. J. Walsh, K. C. Gordon, N. J. Lundin and A. G. Blackman, *J. Phys. Chem.*, 2005, **109**, 5933; (e) S. L. Howell, K. C. Gordon, M. R. Waterland, K. H. Leung and D. L. Phillips, *J. Phys. Chem. A*, 2006, **110**, 11194; (f) W. Chen, C. Turro, L. A. Friedman, J. K. Barton and N. J. Turro, *J. Phys. Chem. B*, 1997, **101**, 6995.
- (a) J. C. Chambron, J. P. Sauvage, E. Amouyal and P. Koffi, *New J. Chem.*, 1985, **9**, 527; (b) F. Westerlund, F. Pierard, M. P. Eng, B. Nördén and P. Lincoln, *J. Phys. Chem. B*, 2005, **109**, 17327; (c) N. Gupta, N. Grover, G. A. Neyhart, W. Liang, P. Singh and H. H. Thorp, *Angew. Chem.*, 1992, **104**, 1058, (*Angew. Chem., Int. Ed. Engl.*, 1992, **31**, 1048); (d) J. K. Barton, *Science*, 1986, **233**, 727.
- (a) T. Imamoto, K. Sugita and K. Yoshida, *J. Am. Chem. Soc.*, 2005, **127**, 11934; (b) T. Imamoto, A. Kumada and K. Yoshida, *Chem. Lett.*, 2007, **36**, 500.
- A. K. Das, E. Bulak, B. Sarkar, F. Lissner, Th. Schleid, M. Niemeyer, J. Fiedler and W. Kaim, *Organometallics*, 2008, **27**, 218.
- (a) S. E. Page, A. Flood and K. C. Gordon, *J. Chem. Soc., Dalton Trans.*, 2002, 1180; (b) J. Granifo, M. E. Vargas, M. T. Garland and R. Baggio, *Inorg. Chim. Acta*, 2000, **305**, 143; (c) J. Granifo, M. E. Vargas, H. Rocha, M. T. Garland and R. Baggio, *Inorg. Chim. Acta*, 2001, **321**, 209; (d) M. A. Ivanov, G. O. Finagenova, M. V. Puzyk and K. P. Balashev, *Opt. Spectrosc.*, 2006, **100**, 367.
- (a) A. Knödler, M. Wanner, J. Fiedler and W. Kaim, *J. Chem. Soc., Dalton Trans.*, 2002, 3079; (b) O. Sarper, E. Bulak, W. Kaim and T. Varnali, *THEOCHEM*, 2006, **773**, 35; (c) E. Bulak, M. Leboschka, B. Schwederski, O. Sarper, T. Varnali, J. Fiedler, F. Lissner, Th. Schleid and W. Kaim, *Inorg. Chem.*, 2007, **46**, 5562.
- I. Sóvágó, K. Várnagy and K. Ósz, *Comments Inorg. Chem.*, 2002, **23**, 149.
- S. Abuskhuna, M. McCann, J. Briody, M. Devereux and V. McKee, *Polyhedron*, 2004, **23**, 1731.
- (a) B. Lippert, (ed.), *Cisplatin*, Wiley-VCH, Weinheim, 1999; (b) C. Mock, I. Puscasu, M. J. Rauterkus, G. Tallen, J. E. A. Wolff and B. Krebs, *Inorg. Chim. Acta*, 2001, **319**, 109; (c) S. Roy, K. D. Hagen, D. P. U. Maheswari, M. Lutz, A. L. Spek, J. Reedijk and G. P. van Wezel, *Chem. Med. Chem.*, 2008, **3**, 1427.
- (a) W. B. Connick, V. M. Miskowski, V. H. Houlding and H. B. Gray, *Inorg. Chem.*, 2000, **39**, 2585 and literature cited; (b) K. E. Dungey, B. D. Thompson, N. A. P. Kane-Maguire and L. L. Wright, *Inorg. Chem.*, 2000, **39**, 5192; (c) W. Kaim, A. Dogan, M. Wanner, A. Klein, I. Tiritiris, Th. Schleid, D. J. Stufkens, T. L. Snoeck, E. J. L. McInnes, J. Fiedler and S. Zalis, *Inorg. Chem.*, 2002, **41**, 4139.

- 11 (a) M. Kato, *Bull. Chem. Soc. Jpn.*, 2007, **80**, 287; (b) R. McGuire, Jr., M. H. Wilson, J. J. Nash, P. E. Fanwick and D. R. McMillin, *Inorg. Chem.*, 2008, **47**, 2946; (c) C. Vogler, B. Schwederski, A. Klein and W. Kaim, *J. Organomet. Chem.*, 1992, **436**, 367; (d) M. Kato, S. Kishi, Y. Wakamatsu, Y. Sugi, Y. Osamura, T. Koshiyama and M. Hasegawa, *Chem. Lett.*, 2005, **34**, 1368; (e) M. Kato, A. Omura, A. Toshikawa, S. Kishi and Y. Sugimoto, *Angew. Chem.*, 2002, **114**, 3315, (*Angew. Chem., Int. Ed.*, 2002, **41**, 3183).
- 12 (a) P. Du, J. Schneider, F. Li, W. Zhao, U. Patel, F. N. Castellano and R. Eisenberg, *J. Am. Chem. Soc.*, 2008, **130**, 5056; (b) S. Rau, D. Walther and J. G. Vos, *Dalton Trans.*, 2007, 915.
- 13 (a) W. B. Connick, R. E. Marsh, W. P. Schaefer and H. B. Gray, *Inorg. Chem.*, 1997, **36**, 913; (b) V. M. Textor and H. R. Oswald, *Z. Anorg. Allg. Chem.*, 1974, **407**, 244; (c) R. H. Herber, M. Croft, M. J. Coyer, B. Bilash and A. Sahiner, *Inorg. Chem.*, 1994, **33**, 2422; (d) W. B. Connick, L. M. Henling, R. E. Marsh and H. B. Gray, *Inorg. Chem.*, 1996, **35**, 6261.
- 14 (a) E. A. M. Geary, K. L. McCall, A. Turner, P. R. Murray, E. J. L. McInnes, L. A. Jack, L. J. Yellowlees and N. Robertson, *Dalton Trans.*, 2008, 3701; (b) P. S. Braterman, J.-I. Song, C. Vogler and W. Kaim, *Inorg. Chem.*, 1992, **31**, 222; (c) W. Kaim and A. Klein, *Organometallics*, 1995, **14**, 1176; (d) W. Kaim, B. Schwederski, A. Dogan, J. Fiedler, C. J. Kuehl and P. J. Stang, *Inorg. Chem.*, 2002, **41**, 4025.
- 15 (a) F. Basolo, and R. G. Pearson, *Mechanisms of Inorganic Reactions, A Study of Metal Complexes in Solution*, 2nd ed., Wiley, New York, 1967; (b) M. L. Tobe, and J. Burgess, *Inorganic Reaction Mechanisms*, Addison Wesley Longman, Harlow, England, 1999.
- 16 (a) R. A. Periana, D. J. Taube, S. Gamble, H. Taube, T. Satoh and H. Fujii, *Science*, 1998, **280**, 560; (b) A. E. Shilov, and G. B. Shul'pin, *Activation and Catalytic Reactions of Saturated Hydrocarbons*, Kluwer, Dordrecht, 2000.
- 17 (a) S. R. Seidel and P. J. Stang, *Acc. Chem. Res.*, 2002, **35**, 972; (b) W.-Y. Sun, M. Yoshizawa, T. Kusukawa and M. Fujita, *Curr. Opin. Chem. Biol.*, 2002, **6**, 757.
- 18 (a) M. Krejčík, M. Danek and F. Hartl, *J. Electroanal. Chem.*, 1991, **317**, 179; (b) W. Kaim, S. Ernst and V. Kasack, *J. Am. Chem. Soc.*, 1990, **112**, 173.
- 19 P. Anzenbacher, Jr., A. C. Try, H. Miyaji, K. Jursíková, V. M. Lynch, M. Marquez and J. L. Sessler, *J. Am. Chem. Soc.*, 2000, **122**, 10268.
- 20 G. M. Sheldrick, *SHELXL-97, Program for refinement of crystal structures*, University of Göttingen, Germany, 1997.
- 21 M. J. Frisch *et al.*, *Gaussian 98, Revision A.3*, Gaussian, Inc., Pittsburgh, PA, 1998.
- 22 W. Kaim, *Chem. Ber.*, 1982, **115**, 910.
- 23 (a) A. Dogan, B. Sarkar, A. Klein, F. Lissner, Th. Schleid, J. Fiedler, S. Zálíš, V. K. Jain and W. Kaim, *Inorg. Chem.*, 2004, **43**, 5973; (b) W. Kaim, A. Dogan, M. Wanner, A. Klein, I. Tiritiris, T. Schleid, D. J. Stufkens, T. L. Snoeck, E. J. L. McInnes, J. Fiedler and S. Zálíš, *Inorg. Chem.*, 2002, **41**, 4139; (c) A. Klein, S. Hasenzahl, W. Kaim and J. Fiedler, *Organometallics*, 1998, **17**, 3532; (d) W. Kaim and A. Klein, *Organometallics*, 1995, **14**, 1176.
- 24 (a) L. Ihlo, M. Kampf, R. Böttcher and R. Kirmse, *Z. Naturforsch.*, 2002, **57b**, 171; (b) M. Kampf, R.-M. Olk and R. Kirmse, *Z. Anorg. Allg. Chem.*, 2002, **628**, 34; (c) J. L. Shaw, J. Wolowska, D. Collison, J. A. K. Howard, E. J. L. McInnes, J. McMaster, A. J. Blake, C. Wilson and M. Schröder, *J. Am. Chem. Soc.*, 2006, **128**, 13827.
- 25 F. Hilgers, W. Kaim, A. Schulz and S. Zálíš, *J. Chem. Soc., Perkin Trans. 2*, 1994, 135.
- 26 (a) W. Kaim, R. Reinhardt, E. Waldhör and J. Fiedler, *J. Organomet. Chem.*, 1996, **524**, 195; (b) W. Kaim, in *New Trends in Molecular Electrochemistry*, Hrsg. A. J. L. Pombeiro, Fontis Media, Lausanne, 2004, p. 127–151.