

6. References:

- Berrettini WH, Vuoristo J, Ferraro TN, Buono RJ, Wildenauer D, Ala-Kokko L (1998): Human G(olf) gene polymorphisms and vulnerability to bipolar disorder. *Psychiatr Genet* 8:235-8.
- Buck KJ, Finn DA (2001): Genetic factors in addiction: QTL mapping and candidate gene studies implicate GABAergic genes in alcohol and barbiturate withdrawal in mice. *Addiction* 96:139-49.
- Buck KJ, Metten P, Belknap JK, Crabbe JC (1997): Quantitative trait loci involved in genetic predisposition to acute alcohol withdrawal in mice. *J Neurosci* 17:3946-55.
- Buono RJ, Lohoff FW, Sander T, et al (2004): Association between variation in the human KCNJ10 potassium ion channel gene and seizure susceptibility. *Epilepsy Res* 58:175-83.
- Chioza B, Osei-Lah A, Wilkie H, et al (2002): Suggestive evidence for association of two potassium channel genes with different idiopathic generalised epilepsy syndromes. *Epilepsy Res* 52:107-16.
- De Fusco M, Marconi R, Silvestri L, et al (2003): Haploinsufficiency of ATP1A2 encoding the Na⁺/K⁺ pump alpha2 subunit associated with familial hemiplegic migraine type 2. *Nat Genet* 33:192-6.
- Dietrich WF, Miller J, Steen R, et al (1996): A comprehensive genetic map of the mouse genome. *Nature* 380:149-52.
- Ferraro TN, Buono RJ (2006): Polygenic epilepsy. *Adv Neurol* 97:389-98.
- Ferraro TN, Golden GT, Smith GG, et al (2001): Quantitative genetic study of maximal electroshock seizure threshold in mice: evidence for a major seizure susceptibility locus on distal chromosome 1. *Genomics* 75:35-42.
- Ferraro TN, Golden GT, Smith GG, et al (2004): Fine mapping of a seizure susceptibility locus on mouse Chromosome 1: nomination of Kcnj10 as a causative gene. *Mamm Genome* 15:239-51.
- Ferraro TN, Golden GT, Smith GG, et al (1997): Mapping murine loci for seizure response to kainic acid. *Mamm Genome* 8:200-8.
- Ferraro TN, Golden GT, Smith GG, et al (1999): Mapping loci for pentylenetetrazol-induced seizure susceptibility in mice. *J Neurosci* 19:6733-9.

- Ferraro TN, Golden GT, Snyder R, et al (1998a): Genetic influences on electrical seizure threshold. *Brain Research* 813:207-210.
- Ferraro TN, Golden, G. T., Dahl, J.P., Smith, Schwebel, C. L., MacDonald, R., Lohoff, F. W., Berrettini, W. H. and Buono, R. J. (2007): Bacterial artificial chromosome-mediated gene transfer confirms effect of a quantitative trait locus for seizure susceptibility in mice. *Epilepsia* in press.
- Ferraro TN, Schill JF, Ballas C, et al (1998b): Genotyping microsatellite polymorphisms by agarose gel electrophoresis with ethidium bromide staining: application to quantitative trait loci analysis of seizure susceptibility in mice. *Psychiatr Genet* 8:227-33.
- Hall CS (1947): Genetic differences in fatal audiogenic seizures between two inbred strains of house mice. *J.Hered.* 38:2-6.
- Jallon P, Latour P (2005): Epidemiology of idiopathic generalized epilepsies. *Epilepsia* 46:10-14.
- Jansen LA, Uhlmann EJ, Crino PB, Gutmann DH, Wong M (2005): Epileptogenesis and Reduced Inward Rectifier Potassium Current in Tuberous Sclerosis Complex-1-Deficient Astrocytes. *Epilepsia* 46:1871-1880.
- Kaunisto MA, Harno H, Vanmolkot KR, et al (2004): A novel missense ATP1A2 mutation in a Finnish family with familial hemiplegic migraine type 2. *Neurogenetics* 5:141-6.
- Lenzen KP, Heils A, Lorenz S, et al (2005): Supportive evidence for an allelic association of the human KCNJ10 potassium channel gene with idiopathic generalized epilepsy. *Epilepsy Res* 63:113-8.
- Mulley JC, Scheffer IE, Harkin LA, Berkovic SF, Dibbens LM (2005): Susceptibility genes for complex epilepsy. *Hum Mol Genet* 14 Spec No. 2:R243-9.
- Noebels JL (2003): Exploring new gene discoveries in idiopathic generalized epilepsy. *Epilepsia* 44 Suppl 2:16-21.
- Ottman R (2005): Analysis of genetically complex epilepsies. *Epilepsia* 46:7-14.
- Sander T, Berlin W, Gscheidel N, Wendel B, Janz D, Hoehe MR (2000): Genetic variation of the human mu-opioid receptor and susceptibility to idiopathic absence epilepsy. *Epilepsy Res* 39:57-61.
- Schlesinger K, Boggan WO, Griek BJ (1968): Pharmacogenetic correlates of pentylenetetrazol and electroconvulsive seizure thresholds in mice. *Psychopharmacologia* 13:181-8.

- Schroder W, Hinterkeuser S, Seifert G, et al (2000): Functional and molecular properties of human astrocytes in acute hippocampal slices obtained from patients with temporal lobe epilepsy. *Epilepsia* 41 Suppl 6:S181-4.
- Segall L, Scanzano R, Kaunisto MA, et al (2004): Kinetic alterations due to a missense mutation in the Na,K-ATPase alpha 2 subunit cause familial hemiplegic migraine type 2. *J Biol Chem*.
- Shang L, Lucchese CJ, Haider S, Tucker SJ (2005): Functional characterisation of missense variations in the Kir4.1 potassium channel (KCNJ10) associated with seizure susceptibility. *Molecular Brain Research* 139:178-183.
- Spadaro M, Ursu S, Lehmann-Horn F, et al (2004): A G301R Na(+)/K(+)-ATPase mutation causes familial hemiplegic migraine type 2 with cerebellar signs. *Neurogenetics*.
- Swoboda KJ, Kanavakis E, Xaidara A, et al (2004): Alternating hemiplegia of childhood or familial hemiplegic migraine? A novel ATP1A2 mutation. *Ann Neurol* 55:884-7.
- Turnbull J, Lohi H, Kearney JA, et al (2005): Sacred disease secrets revealed: the genetics of human epilepsy. *Hum. Mol. Genet.* 14:2491-2500.
- Upton N, Stratton S (2003): Recent developments from genetic mouse models of seizures. *Current Opinion in Pharmacology* 3:19-26.
- Vanmolkot KR, Kors EE, Hottenga JJ, et al (2003): Novel mutations in the Na⁺, K⁺-ATPase pump gene ATP1A2 associated with familial hemiplegic migraine and benign familial infantile convulsions. *Ann Neurol* 54:360-6.
- Wickenden A (2002): K(+) channels as therapeutic drug targets. *Pharmacol Ther* 94:157-82.