

7. Bibliography:

Research Publications, peer reviewed:

- Buono RJ, Ferraro TN, O'Connor MJ, Sperling MR, Abbey M, Finanger E, **Lohoff F**, Mulholland N, Berrettini WH. Lack of association between temporal lobe epilepsy and a novel polymorphism in the alpha 2 subunit gene (ATP1A2) of the sodium potassium transporting ATPase. *Am J Med Genet.* 2000 Feb;96(1):79-83.
- Buono RJ, Ferraro TN, O'Connor MJ, Sperling MR, Ryan SG, Scattergood T, Mulholland N, Gilmore J, **Lohoff FW**, Berrettini WH. Lack of Association Between an Interleukin 1 Beta (IL-1beta) Gene Variation and Refractory Temporal Lobe Epilepsy. *Epilepsia.* 2001 Jun;42(6):782-4.
- Ferraro TN, Golden GT, Smith GG, Longman RL, Snyder RL, DeMuth D, Szpilzak I, Mulholland N, Eng E, **Lohoff FW**, Buono RJ, Berrettini WH. Quantitative genetic study of maximal electroshock seizure threshold in mice: evidence for a major seizure susceptibility locus on distal chromosome 1. *Genomics.* 2001 Jul;75(1-3):35-42.
- Buono RJ, **Lohoff FW**, Sander T, Sperling MR, O'Connor MJ, Dlugos DJ, Ryan SG, Golden GT, Zhao H, Scattergood TM, Berrettini WH, Ferraro TN. Association between variation in the human KCNJ10 potassium ion channel gene and seizure susceptibility. *Epilepsy Res.* 2004 Feb;58(2-3):175-83.
- Ferraro TN, Golden GT, Smith GG, Martin JF, **Lohoff FW**, Gieringer TA, Zamboni D, Schwebel CL, Press DM, Kratzer SO, Zhao H, Berrettini WH, Buono RJ. Fine mapping of a seizure susceptibility locus on mouse Chromosome 1: nomination of Kcnj10 as a causative gene. *Mamm Genome.* 2004 Apr;15(4):239-51.
- Dahl JP, Doyle GA, Buono RJ, Ferraro TN, **Lohoff FW**, Oslin DW, Berrettini WH. Lack of Association Between Single Nucleotide Polymorphisms in the Corticotropin Releasing Hormone Receptor 1 (CRHR1) Gene and Alcohol Dependence. *J Psychiatr Res.* 2005 Sep;39(5):475-9. Epub 2005 Jan 20.

Lenzen K, Heils A, Lorenz S, A. Hempelmann A, F. **Lohoff FW**, Schmitz B, Sander T. Supportive evidence for an allelic association of the human *KCNJ10* potassium channel gene and idiopathic generalized epilepsy. *Epilepsy Res.* 2005 Feb;63(2-3):113-8.

Lohoff FW, Ferraro TN, McNabb L, Schwebel C, Dahl JP, Buono RJ, Berrettini WH. No association between variations in the neuronal nicotinic acetylcholine receptor alpha 2 subunit gene (*CHRNA2*) and bipolar disorder. *Psychiatry Res.* 2005 Jun 30;135(3):171-7.

Lohoff FW, Ferraro TN, Sander T, Zhao H, Dahl JP, Berrettini WH, Buono RJ. No association between common variations in the human alpha 2 subunit gene (*ATP1A2*) of the sodium potassium transporting ATPase and idiopathic generalized epilepsy. *Neurosci Lett.* 2005 Jul 1;382(1-2):33-8.

Lohoff FW, Ferraro TN, Dahl JP, Hildebrandt MA, O'Connor MJ, Sperling MR, Berrettini WH, Buono RJ. Lack of association between variations in the brain-derived neurotrophic factor (*BDNF*) gene and temporal lobe epilepsy. *Epilepsy Res.* 2005 Aug-Sep;66(1-3):59-62.

Dahl JP, Weller AE, Kampman KM, Oslin DW, **Lohoff FW**, Ferraro TN, O'Brien CP and WH Berrettini. Confirmation of the Association Between a Polymorphism in the Promoter Region of the Prodynorphin Gene and Cocaine Dependence. *Am J Med Genet B Neuropsychiatr Genet.* 2005 Nov 5;139(1):106-8.

Lohoff FW, Sander T, Ferraro TN, Dahl JP, Gallinat J, Berrettini WH. Confirmation of Association between the Val66Met Variation in the *BDNF* Gene and Bipolar Disorder. *Am J Med Genet B Neuropsychiatr Genet.* 2005 Nov 5;139(1):51-3.

Dahl JP, Kampman KM, Oslin DW, Weller AE, **Lohoff FW**, Ferraro TN, O'Brien CP, Berrettini WH. Association of a Polymorphism in the *Homer1* Gene with Cocaine Dependence in an African American Population. *Psychiatr Genet.* 2005 Dec;15(4):277-283.

Lohoff FW, Berrettini WH. Lack of association between variations in the melanocortin 5 receptor gene (*MC5R*) and bipolar disorder. *Psychiatr Genet.* 2005 Dec;15(4):255-8.

- Weller AE, Dahl JP, **Lohoff FW**, Ferraro TN, and WH Berrettini. Analysis of Variations in the NAPG Gene on Chromosome 18p11 in Bipolar Disorder. *Psychiatr Genet*. 2006 Feb;16(1):3-8.
- Dahl JP, Cubells JF, Ray R, Weller AE, **Lohoff FW**, Ferraro TN, Oslin DW, Kampman KM, Tang Y, Gelernter J, Kranzler HR, O'Brien CP and WH Berrettini. Analysis of Variations in the Tryptophan Hydroxylase-2 (TPH2) Gene in Cocaine Dependence. *Addict Biol*. 2006 Mar;11(1):76-83
- Weller AE, Dahl JP, **Lohoff FW**, Kampman KM, Dackis C, Ferraro TN, O'Brien CP and WH Berrettini. No Association Between Polymorphisms in the Par-4 gene and Cocaine Dependence. *Psychiatr Genet*. 2006 Oct;16(5):193-196.
- Lohoff FW**, Dahl JP, Ferraro TN, Arnold SE, Gallinat J, Sander T, Berrettini WH. Variations in the vesicular monoamine transporter 1 gene (VMAT1 / SLC18A1) are associated with bipolar I disorder. *Neuropsychopharmacology*. 2006 Dec;31(12):2739-47. Epub 2006 Aug 23.
- Lang UE, Sander T, **Lohoff FW**, Hellweg R, Bajbouj M, Winterer G, Gallinat J. Association of the met66 allele of brain-derived neurotrophic factor (BDNF) with smoking. *Psychopharmacology (Berl.)*, 2006 Dec 21;
- Ferraro TN, Golden GT, Dahl JP, Smith GG, Schwebel CG, MacDonald R, **Lohoff FW**, Berrettini WH and Buono RJ. Analysis of a Quantitative Trait Locus for Seizure Susceptibility in Mice Using Bacterial Artificial Chromosome-Mediated Gene Transfer. *Epilepsia*. 2007 May 23;
- Ferraro TN, Smith GG, Schwebel CL, **Lohoff FW**, Furlong P, Berrettini WH, Buono RJ. Quantitative Trait Locus for Seizure Susceptibility on Mouse Chromosome 5 Confirmed with Reciprocal Congenic Strains. *Physiol Genomics*. 2007 Aug 14; [Epub ahead of print]
- Lohoff FW**, Bloch PJ, Weller AE, Ferraro TN, Berrettini WH. Association analysis of the pituitary adenylate cyclase-activating polypeptide (PACAP/ADCYAP1) gene in bipolar disorder. *Psychiatric Genetics*, 2007. in press

Research Publications, non-peer reviewed:

Lohoff FW. Psychiatry training in light of uncertainty. American Journal of Psychiatry, Resident journal, November 2006: Volume 1, Issue 2

Abstracts:

Buono, R., **Lohoff, F.**, Ferraro, T., Sperling, M., O'Connor, M., Ryan, S., Sander, T., Berrettini, W. A novel polymorphism in the human KIR4.1 gene: implications for general seizure susceptibility. *Epilepsia* 41 (Suppl 7):73,2000.

Lohoff FW, Buono RJ, Ferraro TN, Sperling MR, O'Connor MJ, Ryan SG, Sander T, Berrettini WH. Variation in the human potassium channel gene Kir 4.1 and susceptibility to epilepsy. (presented at the 11th European Student Conference, Berlin, Germany, 2000)

Lohoff FW, Buono RJ, Ferraro TN, Sperling MR, O'Connor MJ, Ryan SG, Sander T, Berrettini WH. Association analysis of a novel polymorphism in the human potassium channel gene Kir 4.1 and susceptibility to epilepsy. (presented at the 4th European Congress on Epileptology, Florence, Italy, 2000)

T.N. Ferraro, G.T. Golden, R.J. Buono, G. Smith, D. DiVirgilio, D. Demuth, I. Szpilzak, D. Zamboni, **F.W. Lohoff**, W.H. Berrettini. Seizure-related locus on mouse chromosome 1: Refined mapping with reciprocal congenic strains. (Abstract Neuroscience meeting 2001)

RJ Buono, **FW Lohoff**, GT Golden, G Smith, MJ O'Connor, MR Sperling,, WH Berrettini and TN Ferraro. KCN10: A Putative Seizure Susceptibility Gene. (Abstract Epilepsy Society meeting 2001)

Buono, R., **Lohoff, F.**, Golden, G., Smith, G., O'Connor, M., Sperling, M., Ferraro, T. KCN10: A Putative Seizure-Susceptibility Gene. *Epilepsia* 43 (Suppl 7) 210, 2001.

Ferraro TN, Golden GT, Smith GG, Martin JF, **Lohoff FW**, Geringer TA, Kratzer SO, Press DM, Berrettini WH, Buono RJ. Seizure sensitivity in inbred mouse strains: relationship to kcnj10 polymorphism. (American Epilepsy Society meeting 2002)

Ferraro TN, Golden GT, Smith GG, Martin JF, **Lohoff FW**, Geringer TA, Kratzer SO, Press DM, Berrettini WH, Buono RJ. Potassium ion channel gene polymorphism and seizure sensitivity in inbred mouse strains. (Abstract Neuroscience meeting 2002)

R.J. Buono; **F.W. Lohoff**, T. Sander; M.R. Sperling; M.J. O'Connor; D.J. Dlugos; S.G. Ryan; G.T. Golden; H. Zhao; W.H. Berrettini; T.N. Ferraro. fine mapping the human kcnj10 locus: identification of a seizure susceptibility allele. (Abstract Neuroscience meeting 2002)

Lohoff FW, Ferraro TN, Sander T, Mulholland N, Sperling MR, O'Connor MJ, Ryan SG, Berrettini WH, Buono RJ. Variations in the human alpha 2 subunit gene (ATP1A2) of the sodium potassium transporting ATPase and susceptibility to idiopathic generalized epilepsy. (Slide presentation, abstract, Neuroscience meeting 2003)

Lohoff FW, Sander T, Gallinat J, Berrettini WH. Confirmation Of Association Between The Val66Met Variation In The BDNF Gene And Bipolar Disorder (2004). Biol Psychiatry 55: 149-242.

Lohoff FW, Sander T, Gallinat J, Berrettini WH. Association between the Val66Met variation in the BDNF gene and bipolar disorder: a replication study. (American Psychiatric Association meeting 5/3/2004, New York, NY)

Lohoff FW, Sander T, Gallinat J, Berrettini WH. Variations in the BDNF gene and susceptibility to bipolar disorder. (Society for Neuroscience meeting 10/23/2004, Washington DC)

Lohoff FW, Sander T, Gallinat J, Berrettini WH. Variations in the brain-derived neurotrophic factor (BDNF) gene and susceptibility to bipolar I disorder. (American Society for Human Genetics meeting, Toronto, 10/29/2004)

Lohoff FW, Ferraro TN, McNabb L, Schwebel C, Dahl JP, Buono RJ, Berrettini WH. No association between variations in the neuronal nicotinic acetylcholine receptor alpha 2 subunit gene (CHRNA2) and bipolar disorder. (2005). Biol Psychiatry 57: S159-S177.

Thomas N Ferraro, Gregory T Golden, John P Dahl, George G Smith, Candice L Schwebel, James F Martin, Ross MacDonald, **Falk W Lohoff**, Wade H Berrettini, Russell J Buono. Bacterial Artificial Chromosome (BAC) Transgenesis Confirms Effect of a Quantitative Trait Locus (QTL) for Seizure

Susceptibility in Mice. (Society for Neuroscience meeting, 11/12/2005, Washington DC)

Russell J Buono, Gregory T Golden, John P Dahl, George G Smith, Candice L Schwebel, James F Martin, Ross MacDonald, **Falk W Lohoff**, Wade H Berrettini, Thomas N Ferraro. Bacterial artificial chromosome (bac) transgenesis confirms effect of a quantitative trait locus for seizure susceptibility in mice: further evidence that coding variation in *kcnj10* influences seizure threshold. (American Epilepsy Society Meeting, 11/13/2005, Washington DC)

FW Lohoff, JP Dahl, TN Ferraro, J Gallinat, T Sander, WH Berrettini. Polymorphisms in the VMAT1 gene are associated with bipolar I disorder. (Society for Neuroscience meeting 11/14/2005, Washington DC)

T.N. Ferraro, G.T. Golden, G.G. Smith, J.F. Martin, R. MacDonald, A. Gupta, J. Scena, J. Picasso, C. Schwebel, **F.W. Lohoff**, J.P. Dahl, W.H. Berrettini, R.J. Buono: Quantitative Trait Loci (QTL) Mapping of Anticonvulsant Response to Valproic Acid (VPA). *Epilepsia* 46 Suppl. 8 (Abst 3.031) 2005

R.J. Buono, G.T. Golden, J.P. Dahl, G.G. Smith, C.L. Schwebel, J.F. Martin, R. MacDonald, **F.W. Lohoff**, W.H. Berrettini, T.N. Ferraro: Bacterial Artificial Chromosome (BAC) Transgenesis Confirms Effect of a Quantitative Trait Locus for Seizure Susceptibility in Mice: Further Evidence That Coding Variation in *KCNJ10* Influences Seizure Threshold. *Epilepsia* 46 Suppl. 8 (Abst 3.026) 2005.

FW Lohoff, JP Dahl, TN Ferraro, J Gallinat, T Sander, WH Berrettini. Variations in the vesicular monoamine transporter 1 gene (*VMAT1* / *SLC18A1*) are associated with bipolar I disorder. (International Society for Psychiatric Genetics meeting 10/17/2005, Boston, MA)

FW Lohoff, JP Dahl, TN Ferraro, J Gallinat, T Sander, WH Berrettini. Association of polymorphisms in the vesicular monoamine transporter 1 gene (*VMAT1* / *SLC18A1*) and bipolar I disorder. (2006). *Biol Psychiatry* 59: 1-90.

T.N. Ferraro, G.T. Golden, G.G. Smith, C.L. Schwebel, F.W. Lohoff, W.H. Berrettini, RJ Buono: Chromosome 5 quantitative trait locus (QTL) for seizure susceptibility in C57BL/6 and DBA/2 mice confirmed using reciprocal congenic strains. Society for Neuroscience meeting, 10/15/2006, Atlanta, GA

FW Lohoff, AE Weller, JP Dahl, TN Ferraro, J Gallinat, T Sander, SJ Kanen, SE Arnold, RE Gur, WH Berrettini. Association between polymorphism in the vesicular monoamine transporter 1 gene (VMAT1 / SLC18A1) on chromosome 8p and schizophrenia. (Society for Neuroscience meeting 10/16 /2006, Atlanta, GA)

T.N. Ferraro, G.T. Golden, G.G. Smith, C.L. Schwebel, **F.W. Lohoff**, W.H. Berrettini, R.J. Buono: Chromosome 5 Quantitative Trait Locus (QTL) for Seizure Susceptibility in C57BL/6 and DBA/2 Mice Confirmed Using Reciprocal Congenic Strains. *Epilepsia* 47 Suppl. 2006.

FW Lohoff, AE Weller, JP Dahl, TN Ferraro, J Gallinat, T Sander, SJ Kanen, SE Arnold, RE Gur, WH Berrettini. Variations in the vesicular monoamine transporter 1 gene (VMAT1 / SLC18A1) on chromosome 8p are associated with schizophrenia. (International Society for Psychiatric Genetic, 10/31/2006, Cagliari, Italy)

Lohoff FW, Bloch PJ, Weller AE, Ferraro TN, Berrettini WH. Shared bipolar/schizophrenia susceptibility genes on chromosome 8p. (American Psychiatric Association / APIRE meeting 5/19/2007, San Diego, CA)

Lohoff FW, Bloch PJ, Weller AE, Ferraro TN, Berrettini WH. Association analysis of the pituitary adenylate cyclase-activating polypeptide (PACAP/ADCYAP1) gene in bipolar disorder. (2007). *Biol Psychiatry* 61: S178-S265.

Lohoff FW, Bloch PJ, Weller AE, Ferraro TN, Berrettini WH. Polymorphisms in the pituitary adenylate cyclase-activating polypeptide (PACAP/ADCYAP1) gene and susceptibility to bipolar disorder. World Congress of Psychiatric Genetics 2007, New York, NY, USA, 10/9/07

Lohoff FW, Bloch PJ, Weller AE, Ferraro TN, Berrettini WH. Analysis of variation in the pituitary adenylate cyclase-activating polypeptide (PACAP/ADCYAP1) gene and susceptibility to bipolar disorder. American Society of Human Genetics Meeting 2007, San Diego, CA, USA. 10/25/07

Editorials, Reviews, Chapters, including participation in committee reports:

Lohoff FW and Berrettini WH. Genetics of Mood Disorders. In: Neurobiology of Mental Illness by Eric J. Nestler (Author), Dennis S. Charney (Editor)
Paperback: 1272 pages, Publisher: Oxford University Press, USA; 4
edition , 2007