G. ANHANG

G.1. Abkürzungsverzeichnis

AG	Antigen
AK	Antikörper
APACHE	Acute Physiology and Chronic Health Evaluation
APP	Akute Phase Protein
ARDS	Adult Respiratory Distress Syndrome = akutes Lungenversagen
AU	Arbeitsunfall
C	Komplementfaktor
CD	Cluster of Differentation
EC	Endothelial Cells = Endothelzellen
EDTA	Ethylene Diamine Tetra-Acetate = Äthylendiamintetraessigsäure
EGF	Epidermal Growth Factor
EK	Erythrozytenkonzentrat
ELAM	Endothelial Leukocyte Adhesion Molecule = E-Selektin = CD62E
ELISA	Enzyme Linked Immuno Sorbent Assay
FITC	Fluoresceine Isothio Cyanat
FS	Forward Light Scatter
GCS	Glasgow Coma Score
gmp 140	Granule Membrane Protein = P-Selektin = CD 62P
GOT	Glutamat - Oxalazetat - Transaminase
HDL	High Density Lipoprotein
HEV	High Endothelial Venule
HIV	Human Immundeficiency Virus

HRP	Horse Raddish Peroxidase = Meerrettichperoxidase
ICAM	Intercellular Adhesion Molecule
IgG	Immunglobulin
ISS	Injury Severity Score
LECAM	Leukocyte Adhesion Molecule (=CD62L)
Mo	Makrophagen
MODS	Multiple Organ Dysfunction Syndrom = Multiples Ordandysfunktionssyndro
MOF	Multiple Organ Failure = Multiorganversagen
NK-Zellen	Natürliche Killerzellen
PEEP	Positiver Endexpiratorischer Beatmungsdruck
PMN	Polymorphkernige Zellen = Granulozyten
PTS	Polytraumaschlüssel
RD 1	Rhodamin 1
SAPS	Simplified Acute Physiology Score
SHT	Schädel-Hirn-Trauma
sLECAM	soluble LECAM = lösliches CD62L
SS	Side Light Scatter
TNF-α	Tumor Nekrose Faktor-α
ZNS	Zentralnervensystem

G.2.Literaturverzeichnis

Adams D, Shaw S. Leukocyte-endothelial interactions and regulation of leukocyte migration. Lancet 1994; 343:831-836.

Armstead VE, Minchenko AG, Campbell B, et al. P-Selectin is up-regulated in vital organs during murine traumatic shock. Faseb. J. 1997, 11:1271-1279.

Bajorath J, Aruffo A. A template for generation and comparison of threedimensional selectin models. Biochemical and Biophysical Research Communications 1995; 216(3):1018-1023.

Baker C, Oppenheimer L, Stephens B, et al. Epidemiology of Trauma Deaths. The American Journal of Surgery 1980; 140:144-150.

Baker SP, O'Neill B. The Injury Severity Score: an update. The Journal of Trauma 1976; 16:882-885.

Baue AE. Progress in Trauma Care through understanding the Cell Biology of Injury. In: Faist E, Meakins J, Schildberg FW, eds. Host Defense Dysfunction in Trauma, Shock and Sepsis. Berlin, Heidelberg, New York: Springer Verlag, 1993:3-14.

Baue AE. The Horror Autotoxicus and Multiple Organ Failure. Arch Surg 1992; 127:1451-1462.

Beal AL, Cerra FB. Multiple Organ Failure Syndrome in the 1990s. JAMA 1994; 271:226-233.

Berlin C, Berg EL, Briskin MJ, et al. □4□7 Integrin Mediates Lymphocyte Binding to the Mucosal Vascular Addressin MAdCAM-1. Cell 1993; 74:185-195.

Bernard GR, Artigas A, Brigham KL et al. The Consensus Committee (1994) Report of the American-European consensus conference on ARDS: definitions, mechanisms, relevant outcomes and clinical trial coordination. Intensive Care Medicine 1994; 20:225-232.

Blann AD, Nadar SK, Lip GY. The adhesion molecule P-selectin and cardiovascular disease. European heart journal 2003; 24:2166-2179.

Boldt J, Wollbruck M, Kuhn D, et al. Do plasma levels of circulating soluble adhesion molecules differ between surviving and nonsurviving critically ill patients? Chest 1995; 107:787-792.

Brandley BK, Swiedler SJ, Robbins PW. Carbohydrate Ligands of the LEC Cell Adhesion Molecules. Cell 1990; 63:861-863.

Carlos T, Harlan J. Leukocyte-Endothelial Adhesion Molecules. Blood 1994; 84(7):2068-2101.

Cellner C, Patey N, Fromont-Hankard G, et al. In situ endothelial cell adhesion molecule expression in ulcerative colitis. E-Selectin in situ expression correlates with clinical, endoscopical, histological activity and outcome. European Journal of Gastroenterological Hepatology 1997; 9:1197-1203.

Cocks RA, Chan TY. Alteration in leukocyted adhesion molecule expression following minor, moderate and major trauma. Eur J Emeg Med 1997, 4(4):193-195.

Cowley HC, Heney D, Gearing AJ, et al. Increased circulating adhesion molecule concentrations in patients with the systemic inflammatory response syndrome: A prospective cohort study. Critical Care Medicine 1994; 22:651-657.

De-Bandt M, Meyer D, Hakim J, et al. Antibodies to proteinase-3 mediate expression of intercellular adhesion molecule-1 (ICAM-1, CD54). British Journal of Rheumatology 1997; 36:839-846.

Dietrich HJ, Peter K. The Intensive Care Unit of Tomorrow. In: Faist E, Meakins J, Schildberg FW, eds. Host Defense Dysfunction in Trauma, Shock and Sepsis. Berlin, Heidelberg, New York: Springer

Verlag, 1993:31-35.

Donelly SC, Robertson C.Mediators, mechanisms and mortality in major trauma. Resuscitation 1994; 28:87-92.

Donnelly SC, Haslett C, Dransfield I, et al. Role of selectins in development of adult respiratory distress syndrome. Lancet 1994; 344:215-219.

Dorinsky P, Gadek J. Multiple Organ Failure. Clinics in Chest Medicine 1990; 11/4:581-591.

Ertel W, Trentz O. Polytrauma und Multiorgan-Dysfunktionssyndrom (MODS). Definition–Pathophysiologie-Therapie. Zentralblatt für Chirurgie 1994; 119(3):159-167.

Faist E, Baue AE, Dittmer H, et al. Multiple Organ Failure in Polytrauma Patients. The Journal of Trauma 1983; 23/9:775-787.

Fassbender K, Schminke U, Ries S, et al. Endothelial-derived adhesion molecules in bacterial meningitis: association to cytokine release and intrathecal leukocyte recruitment. Journal of Neuroimmunology 1997; 74:130-134.

Friedl HP, Trentz O. Polytrauma: Definition - Schock - Multiorganversagen. Unfallchirurgie 1992; 18:64-68.

Fujimi S, Ogura H, Tanaka H, et al. Increased production of leukocyte microparticles with enhanced expression of adhesion molecules from activated polymorphonuclear leukocytes in severely injured patients. J Trauma 2003; 54(1):114-9.

Gando S, Kameue T, Matsuda N, et al. Combined activation of coagulation and inflammation haas an important role in multiple organ dysfunction and poor outcome after severe trauma. Thromb Haemost 2002; 88(6):943-9.

Gearing AJ, Neman W. Circulating adhesion molecules in disease. Immunology today 1993; 14:506-512.

Goldfarb RD, Georgakopoulos N, Achkter I, et al. E-Selectin levels elevate prior to organ injury during bacteremia induced by peritonitis in the pig. Critical Care Medicine 1996; 24: A91.

Goris RJ, te Boekhorst TP, Nuytinck JK, et al. Multiple Organ Failure - generalized autodestructive inflammation?. Archives of Surgery 1985; 120:1109-1115.

Grob PJ, Holch M, Brunner W. Posttraumatisches/ postoperatives Immundefektsyndrom. Schweizerische Medizinische Wochenschrift 1987; 117/13:471-480.

Guillou PJ. Biological variation in the development of sepsis after surgery and trauma. Lancet 1993; 342:217-220.

Holmes K, Fowlkes BJ. Current Protocols in Immunology, New York: John Wiley and Sons Inc. 1992.

Johnston GI, Cook RG, McEver RP. Cloning of GMP-140, a Granule Membrane Protein of Platelets and Endothelium: Sequence Similarity to Proteins Involved in Cell Adhesion and Inflammation. Cell 1989; 56:1033-1044.

Kayal S, Jais JP, Aguini N, et al. Elevated circulating E-Selectin, Intercellular adhesion molecule 1, and von Willebrand Factor in patients with severe infection. Am. J. Respir. Crit. Care Med. 1998; 157:776-784.

Kishimoto TK, Warnok A, Jutila MA, et al. Antibodies gainst human neutrophil LECAM-1 and endothelial cell ELAM-1 inhibit a common CD18-independent adhesion pathway in vitro. Blood 1991; 78:805-811.

Knaus WA, Draper EA, Wagner DP, et al. APACHE II: A severity of disease classification system.

Critical Care Medicine 1985; 13:818-829.

Kobayashi H, Wang Y, Puri P. Increased levels of circulating endothelial leukocyte adhesion molecule 1 (ELAM-1) in children with reflux nephropathy. Europ. Urol.1997; 31:343-346.

Kvasnicka J, Briza J, Krska Z, et al. Increase of soluble cytoadhesive molecules sE-selectin and sIcam-1 and hyperfibrinogenaemia in patients with Polytrauma without septicaemia. Sb Lek. 1998; 99(2):93-6.

Larsen R. Trauma und Polytrauma. In: Larsen R, Ed. Anästhesie, 4th edn. München, Wien, Baltimore: Urban & Schwarzenberg1994:916-932.

Lasky LA, Singer MS, Dowbenko D, et al. An Endothelial Ligand for L-Selectin Is a Novel Mucin-like Molecule. Cell 1992; 69:927-938.

Law MM, Cryer HG, Abraham E. Elevated levels of soluble ICAM-1 correlate with the development of multiple organ failure in severely injured trauma patients. The Journal of Trauma 1994; 37:100-110.

Lawrence MB, Springer TA. Neutrophils roll on E-Selectin. The Journal of Immunology 1993; 151:6338-6346.

Le Gall JR, Lemeshow S, Saulnier S. A new Simplified Acute Physiology Score (SAPS II) based on a European/ North American multicenter study. JAMA 1993; 270:2957-2963.

Macey MG, McCarthy DA. Quantitation of Adhesion Molecules and other Function-Associated Antigens on Human Peripheral Blood Leukocytes. Cytometry 1993; 14:898-908.

Maekawa K, Futami S, Nishida M, et al. Effects of trauma and sepsis on solubile L-selectin and cell surface expression of L-selectin and CD11b. J. Trauma 1998; 44:460-467.

Malik AB, Wang Y. Cellular mechanisms of leukocyte adhesion. In Schlag G, Redl H, eds. Pathophysiology of shock, sepsis and organ failure. Berlin, Heidelberg, New York: Springer 1993:61-74.

Marcus BC, Hynes KL, Gewertz BL. Loss of endothelial barrier function requires neutrophil adhesion. Surgery 1997; 122:420-426.

Mazrer CA, Friess H, Kretschmann B. Over-Expression of ICAM-1, VCAM-1 and ELAM-1 might influence tumor progression in colorectal cancer. Int. J. Cancer 1998; 79:76-81.

McEver R. Leukocyte Interactions Mediated by Selectins. Thrombosis and Haemostasis 1991; 66:80-87.

McEver R. Selectins: Novel Receptors that Mediate Leukocyte Adhesion During Inflammation. Thrombosis and Haemostasis 1991; 65:223-228.

Miller S, Miller C, Trunkey D. The Immune Consequences of Trauma. Surgical Clinics of North America 1982; 62:167-179.

Müller JC, Buhrer C, Kiening K, et al. Decreased soluble adhesion molecule L-selectin plasma concentrations after major trauma. J Trauma 1998, 45(4): 705-708.

Nieuwenhoven EJ, Lefering R, Neugebauer E, Goris RJA. Clinical relevance of sepsis scores. Langenbeck´s Archives of Surgery 1998; 383:11-14.

Nuytinck HKS, Offermans XJ, Kubat K, et al. Whole-Body Inflammation in Trauma Patients-an autopsy study. Arch Surg 1988; 123:1519-1524.

Pan J, Fu G, Lin X, et al. The expression of L-selectin at transcription and translation level in hemorrhagic shock rat. Zhonghua Nei Ke Za Zhi 22002; 41(7): 472-5.

Parks DR, Herzenberg DA, Herzenberg LA. Flow Cytometry and Fluorescence-Activated Cell Sorting. In: Paul WE, Ed. Fundamental Immunology, 2nd edn. New York: Raven Press Ltd. 1989:781-802.

Peitzmann AB, Harbrecht BG, Udekwu AO et al. Hemorrhagic Shock. Current Problems in Surgery 1995; 32:967-1002.

Rainer TH, Ng MH, Lam NY, et al. Role of monocyte L-selectin in the development of post-traumattic organ failure. Resuscitation 2001; 51(2):139-49.

Redl H, Nikolai A, Kneidinger R, et al. Endothelial and leukocyte activation in experimental polytrauma and sepsis. Behring Institut Mitteilungen 1993; 92:218- 228.

Redl H, Schlag G, Dinges HP, et al. Leukocyte-Endothelial Interactions in Trauma and Sepsis. In: Faist E, Meakins J, Schildberg FW, eds. Host Defense Dysfunction in Trauma, Shock and Sepsis. Berlin, Heidelberg, New York. Springer 1993:277-286.

Regel G, Pape HC, Pohlemann T, et al. Scores als Entscheidungshilfe. Unfallchirurg 1994; 97:211-216.

Regel G, Sturm JA, Pape HC. Das Multiorganversagen (MOV) Ausdruck eines generalisierten Zellschadens aller Organe nach schwerem Trauma. Unfallchirurg 1991; 94:487-497.

Ridings PC, Bloomfield GL, Halloway S, t al. Sepsis-Induced Acute Lung Injury Is Attenuated by Selectin Blochade Following the Onset of Sepsis. Archives of Surgery 1995; 130:1199-1208.

Rohr U. sCD14, TNFα, Interleukin-6, sICAM-1 und sE-Selektin im septischen Geschehen. Dissertation 1998, Humboldt Universität, Berlin.

Roumen RH, Redl H, Schlag G, Goris RJ et al. Inflammatory mediators in relation to the development of multiple organ failure in patients after severe blunt trauma. Critical Care Medicine 1995; 23/3:474-480.

Roumen RM, Hendriks T, Goris JA. Cytokine patterns in patients after major vascular surgery, hemorrhagic shock, and severe blunt trauma - relation with subsequent ARDS and MOF. Annals of Surgery 1993; 218:769-776.

Sako D, Chang XJ, Barone KM, et al. Expression Cloning of a Functional Glycoprotein Ligand for P-Selectin. Cell 1993; 75:1179-1186.

Seidelin JB, Nielsen OH, Strom J. Soluble L-selectin levels predict survival in sepsis. Intensive care medicine 2002; 28:1613-1618.

Sessler CN, Windsor AC, Schwartz M, et al. Circulating ICAM-1 is increased in septic shock. Am J Respir Crit Care Med 1995; 151:1420-1427.

Shahgasempour S, Woodroffe SB, Garnett HM. Alterations in the expression of ELAM-1, ICAM-1 and VCAM-1 after in-vitro infection of endothelial cells with a clinical isolate of human cytomegalovirus. Microbiological Immunology 1997; 41:21-129.

Shaw S. Leukocyte Differentiation Database. 5th Workshop 1995.

Shen J, T-To SS, Schrieber L, et al. Early E-Selectin, VCAM-1, ICAM-1 and late major histocompatibility complex antigen induction on human endothelial cells by flavivirus and comodulation of adhesion molecule expression by immune cytokines. Journal of Virology 1997; 71:9323-9332.

Siemiatkowski A, Rogowski F, Borkowski J, et al. Soluble selectin profiles associated with severe trauma. Arch Immunol Ther Exp (Warsz) 2001; 49(4): 317-24.

Simon SI, Rochon YP, Lynam EB, et al. □2-Integrin and L-Selectin are obligatory receptors in

neutrophil aggregation. Blood 1993, 82:1097-1106.

Simons RK, Hoyt DB, Winchell RM, et al. Elevated Selectin levels afeer severe trauma: a marker for sepsis and organ failure and a potential target for immunomodulatory therapy. Journal of trauma 1996; 41:653-662.

Smith CW. Leukocyte - endothelial cell interactions. Seminars in Hematology 1993; 30/4:45-55.

Smith CW. Principles of immune recognition. In: Rich R, Ed. Clinical Immunology - Principles and Practice. St. Louis: Mosby-Year Book Inc. 1996:180-191

Spertini O, Schleiffenbaum B, White-Owen C, et al. ELISA for quantitation of L-selectin shed from leukocytes in vivo. Journal Immunol. Methods 1992; 156:115-123.

Springer TA. Adhesion receptors of the immune system. Nature 1990; 346:425-434.

Stengel D, Bauwens K, Keh D, Gerlach H, Ekkernkamp A, Tauber R, Kerner T. Prognostic value of an early soluble L-selectin (sCD62L) assay for risk assessment in blunt multiple trauma: a metaanalysis. Clin Chem 2005; 51(1):16-24.

Svoboda P, Kantorova I, Ochmann J. Dynamics of interleukin 1, 2, 6 and tumor necrosis factor alpha in multiple trauma patients. Journal of Trauma 1994; 36:336-340.

Tözeren A, Ley K. How do selectins mediate leukozyte rolling in venules? Biophysical Journal 1992; 63:700-709.

Van Griensven M, Barkhausen T, Hildebrand F, et al. L-selectin shows time and gender dependency in association with MODS. Injury 2004; 35(11):1087-95.

Ward PA, Mulligan MS, Warren JS. Neutrophils, Cytokines, Oxygen Radicals, and Lung Injury. In: Faist, Meakins, Schildberg, Eds. Host Defense Dysfunction in Trauma, Shock and Sepsis. Berlin, Heidelberg, New York: Springer 1993:177-180.

G.3.Lebenslauf

Aus Datenschutzgründen wird der Lebenslauf in der elektronischen Fassung nicht veröffentlicht

G.4.Erklärung

Ich, Stefan Höfler, erkläre, dass ich die vorgelegte Dissertationsschrift mit dem Thema:

„Lösliche und zellgebundene Selektine im frühen Verlauf nach Polytraumatisierung“

selbst verfasst und keine anderen als die angegebenen Quellen und Hilfsmittel benutzt, ohne die (unzulässige) Hilfe Dritter verfasst und auch in Teilen keine Kopien anderer Arbeiten dargestellt habe.

G.5.Danksagung

Für die Unterstützung und Ermöglichung dieser Arbeit möchte ich Herrn Prof. Dr. H. Gerlach danken.

Herrn Dr. D. Keh und Herrn PD Dr. T. Kerner danke ich für ihre Betreuung während der Studie.

Ein ganz besonderer Dank gebührt dem Kollegen Dr. med. O. Ahlers für seine stets kompetente und überaus liebenswerte Unterstützung.