

- ABBASI, I.; BRANZBURG, A.; CAMPOS-PONCE, M.; ABDEL HAFEZ, S.K.; RAOUL, F.; CRAIG, P.H.S. and HAMBURGER, J. (2003):
Copro-diagnosis of *Echinococcus granulosus* infection in dogs by amplification of a newly identified repeated DNA sequence.
Am. J. Trop. Med. Hyg. 69, 324-330
- ALLAN, J.C.; CRAIG, P.S.; GARCIA NOVAL, J.; MENCOS, F.; LIU, D.; WANG, Y.; WEN, H.; ZHOU, P.; STRINGER, R.; ROGAN, M. and ZEHYALE, E. (1992):
Coproantigen detection for immunodiagnosis of echinococcosis and humans.
Parasitology 104, 347-355
- ALLARD, A.; GIRONES, R.; JUTO, R. and WADELL, G. (1990):
Polymerase chain reaction for detection of adenoviruses in stool samples.
J. Clin. Microbiol. 28, 2659-2667
- ALTHER, P. (1996):
Beitrag zur Epidemiologie und Diagnose der *Echinococcus multilocularis*-Infektion bei Endwirten.
Vet. Med. Diss., Universität Zürich.
- ALTINTAS, N. (2003):
Past to present: echinococcosis in Turkey.
Acta Trop. 85, 105-112
- AMMAN, R. and ECKERT, J. (1995):
Clinical diagnosis and treatment of echinococcosis in humans.
In: THOMPSON, R.C.A. and LYMBERRY, A.J. (eds): *Echinococcus and hydatid disease*.
CAB International, Wallingford, pp. 411-463
- AUER, H.; PICHER, O. and ASPÖCK, H. (1986):
Erfahrungen bei der Serodiagnostik der Echinokokkosen mittels ELISA.
Mitt. Österr. Ges. Tropenmed. Parasit. 8, 17-22
- BABOS, S. and NEMETH, I. (1962):
Zur Frage bei der Serodiagnostik der Echinococcosis. (in ungarisch).
Magyar Allatorvosok Lapja 17, 58-60
- BAI, Y.; CHENG, N.; JIANG, C.; WANG, Q. and DARUENG, C. (2002):
Survey on cystic echinococcosis in Tibetans, West China.
Acta Trop. 82, 381-385

BARDONNET, K.; BENCHIKH-ELFEGOUN, M.C.; BART, J.M.; HARRAGA, S.; HANNACHE, N.; HADDAD, S.; DUMON, H.; VUITTON, D.A. and PIARROUX, R. (2003):

Cystic echinococcosis in Algeria: cattle act as reservoirs of a sheep strain and may contribute to human contamination.

Vet. Parasitol. 116, 35-44

BARONET, D.; WALTNER-TOEWS, D.; CRAIG, P.S. and JOSHI, D.D. (1994):

Echinococcus granulosus infections in the dogs of Kathmandu, Nepal.

Ann. Trop. Med. Parasitol. 88, 485-492

BIAVA, M.F.; DAO, A. and FORTIER, B. (2001):

Laboratory diagnosis of cystic hydatid disease.

World. J. Surg. 25, 10-14

BOCH, J. and SUPPERER, R. (1983):

Veterinärmedizinische Parasitologie. 3. Auflage, Verlag Paul Parey, Berlin, pp. 39-53

BRETAGNE, S.; GUILLOU, J.P.; MORAND, M. and HOUIN, R. (1993):

Detection of *Echinococcus multilocularis* DNA in fox faeces using DNA amplification.

Parasitology 106, 193-199

BREYER, I.; GEORGIEVA, D.; KURDOVA, R and GOTTSSTEIN, B (2004):

Echinococcus granulosus strain typing in Bulgaria: the G1 genotype is predominant in intermediate and definitive wild hosts.

Parasitol. Res. 93, 127-130

BUDKE, C.M.; CAMPOS-PONCE, M.; WANG, Q. and TORGERSON, P.R. (2005):

A canine purgation study and risk factor analysis for echinococcosis in a high endemic region of the Tibetan plateau.

Vet. Parasitol. 127, 49-55

BÜHLER, A. (1972):

Untersuchungen über den Echinokokkenbefall bei Schlachttieren in der Schweiz und die Übertragbarkeit larvaler Echinokokken auf Labortiere.

Vet. Med. Diss., Universität Zürich

CABRERA, M.; CANOVA, S.; ROSENVZVIT, M. and GUARNERA, E. (2002):

Identification of *Echinococcus granulosus* eggs.

Diagn. Microbiol. Infect. Dis. 44, 29-34

CARMONA, C.; PERDOMA, R.; CARBO, A.; ALVAREZ, C.; MONTI, J.; GRAUERT, R.; STERN, D.; PERERA, G.; LLOYD, S.; BZINI, R.; GEMMELL, M.A. and YARZABAL, L. (1998):

Risk factors associated with human cystic echinococcosis in Florida, Uruguay: results of a mass screening study using ultrasound and serology.

Am. J. Trop. Med. Hyg. 58, 599-605

CHORDI, A.; GONZALEZ-CASTRO, J. and TORMO, J. (1962):
Aportacion al estudio de les helmenthiasis intestinales en los perros.II. Resultados de la prueba de flocculation con "Benthid" y de la de fijacion de complemento en perros con *Echinococcus granulosus*.
Rev. Iber. Parasitol. 22, 285-290

CHRISTOFI, G.; DEPLAZES, P.; CHRISTOFI, N.; TANNER, I.; ECONOMIDES, P. and ECKERT, J. (2002):
Screening of dogs for *Echinococcus granulosus* coproantigen in a low endemic situation in Cyprus.
Vet. Parasitol. 104, 299-306

COHEN, H.; PAOLILLO, E.; BONIFACINO, R.; BOTTA, B.; PARADA, L.; CABRERA, P.; SNOWDEN, K.; GASSER, R.; TESSIER, R.; DIBARBOURE, L.; WEN, H.; ALLAN, J.C.; SOTO DE ALFARO, H.; ROGAN, M.T. and CRAIG, P.S. (1998):
Human cystic echinococcosis in a Uruguayan community: a sonographic, serologic and epidemiological study.
Am. J. Trop. Med. Hyg. 59, 620-627

CRAIG, P.S. and RICKARD, M.D. (1981):
Studies on the specific immunodiagnosis of larval cestode infections of cattle and sheep using antigens purified by affinity chromatography in an enzyme-linked immunosorbent assay (ELISA).
Int. J. Parasitol. 11, 441-449

CRAIG, P.S.; HOCKING, R.E.; MITCHELL, G.F. and RICKARD, M.D. (1981):
Murine hybridoma-derived antibodies in the processing of antigens for the immunodiagnosis of hydatid (*Echinococcus granulosus*) infection in sheep.
Parasitology 83, 303-317

CRAIG, P.S.; MACPHERSON, C.N.L. and NELSON, G.S. (1986):
The identification of eggs of *Echinococcus* by immunofluorescence using a specific anti-oncospherical monoclonal antibody.
Am. J. Trop. Med. Hyg. 35, 152-158

CRAIG, P.S.; MACPHERSON, C.N.L.; WATSON-JONES, D.L. and NELSON, G.S. (1988):
Immunodetection of *Echinococcus* eggs from naturally infected dogs and from environmental contamination sites in settlements in Turkana, Kenya.
Trans. Roy. Soc. Trop. Med. Hyg. 82, 268-274

CRAIG, P.S. (1993):
Immunodiagnosis of *Echinococcus granulosus*. In: ANDERSEN, F.L. (eds.): Compendium on cystic echinococcosis with special reference to Xinjiang, China Provo, Brigham Young University, USA, pp. 85-118

CRAIG, P.S.; GASSER, R.B.; PARADA, L.; CABRERA, P.; PARIETTI, S.; BORGUES, C.; ACUTTIS, A.; AGULLA, J.; SNOWDEN, K. and PAOLILLO, E. (1995):
Diagnosis of canine echinococcosis: comparision of coproantigen and serum antibody tests with arecoline purgation in Uruguay.
Vet. Parasitol. 56, 293-301

CRAIG, P.S. (1997):
Echinococcus granulosus : immunodiagnosis and vaccination, a perspective.
Parassitologia 39, 345-347

CRAIG, P.S.; GIRAUDOUX, P.; SHI, D.; BARTHOLOMOT, B.; BARNISH, G.; DELATTRE, P.; QUERRE, J.P.; HARRAGA, S.; BAO, G.; WANG, Y.; LU, F.; ITO, A. and VUITTON, D.A. (2000):
An epidemiological and ecological study of human alveolar echinococcosis transmission in south Gansu, China.
Acta Trop. 77, 167-177

CRAIG, P.S.; ROGAN, M.T. and CAMPOS-PONCE, M. (2003):
Echinococcosis: disease, detection and transmission.
Parasitology 127, 5-20

D'ALESSANDRO, A.; RAUSCH, R.L.; CUELLO. C. and ARISTIZABAL, N. (1979):
Echinococcus vogeli in man, with a review of polycystic hydatid disease in Columbia and neighbouring countries.
Am. J. Trop. Med. Hyg. 28, 303-317

DE RYCKE, P.H. and DE COOMAN, E.P. (1968):
The evagination of *echinococcus granulosus* scolices from lung cysts of bovin origin.
Z. Parasitenkd. 30, 171-175

DEPLAZES, P. and ECKERT, J. (1988):
Untersuchungen zur Infektion des Hundes mit *Taenia hydatigena*.
Schweiz Arch. Tierheilk. 130, 289-306

DEPLAZES, P.; GOTTSSTEIN, B.; STINGELIN, Y. and ECKERT, J. (1990):
Detection of *Taenia hydatigena* copro-antigens by ELISA in dogs.
Vet. Parasitol. 36, 91-103

DEPLAZES, P.; GOTTSSTEIN, B.; ECKERT, J.; JENKINS, D.J.; EWALD, D. and JIMENEZ-PALAZIOS, S. (1992):
Detection of Echinococcus coproantigens by enzyme-linked immunosorbent assay in dogs, dingoes and foxes.
Parasitol. Res. 78, 303-308

DEPLAZES, P.; JIMENEZ-PALACIOS, S.; GOTTSSTEIN, B.; SKAGGS, J. and ECKERT, J. (1994):
Detection of Echinococcus coproantigens in stray dogs of northern Spain.
Appl. Parasitol. 35, 297-301

DEPLAZES, P. and ECKERT, J. (1996):
Diagnosis of the *Echinococcus multilocularis* infection in final hosts.
Appl. Parasitol. 37, 245-252

DEPLAZES, P.; ALTHER, P.; TANNER, I.; THOMPSON, R.C.A. and ECKERT, J. (1999): *Echinococcus multilocularis* coproantigen detection by enzyme-linked immunosorbent assay in fox, dog and cat populations.
J. Parasitol. 85, 115-121

DINKEL, A.; von NICKISCH-ROSENEGK, M.; BILGER, B.; MERLI, M.; LUCIUS, R. and ROMIG, TH. (1998):
Detection of *Echinococcus multilocularis* in the definitive host: Coprodiagnosis by PCR as an alternative to necropsy.
J. Clin. Microbiol. 36, 1871-1876

DINKEL, A.; NJOROGE, E.M.; ZIMMERMANN, A.; WÄLZ, M.; ZEYHLE, E.; ELMAHDI, I.E.; MECKENSTEDT, U. and ROMIG, T. (2004):
A PCR system for detection of species and genotypes of the *Echinococcus granulosus*-complex, with reference to the epidemiological situation in Eastern Africa.
Int. J. Parasitol. 34, 645-653

DOWLING, P.; ABO-SHEHADA, M. and TORGERSON, P.R. (2000):
Risk factors associated with human cystic echinococcosis in Jordan: result of a case -control study.
Ann. Trop. Med. Parasitol. 94, 69-75

ECKERT, J. (1981):
Echinokokkose.
Berl. Münch. Tierärztl. Wochenschr. 94, 369-378

ECKERT, J.; THOMPSON, R.C.A.; MICHAEL, S.A., KUMARATILAKE, L.M. and EL-SAWAH, H.M. (1989):
Echinococcus granulosus of camel origin: development in dogs and parasite morphology.
Parasitol. Res. 75, 536-544

ECKERT, J.; DEPLAZES, P.; EWALD, D. and GOTTSSTEIN, B. (1991):
Parasitologische und immunologische Methoden zum Nachweis von *Echinococcus multilocularis* bei Füchsen.
Mitt. Österr. Ges. Tropmed. Parasit. 13, 25-30

ECKERT, J.; THOMPSON, R.C.A.; LYMBERRY, A.J.; PAWLOWSKI, Z.S.; GOTTSSTEIN, B. and MORGAN, U.M. (1993):
Further evidence for the occurrence of a distinct strain of *Echinococcus granulosus* in European pigs.
Parasitol. Res. 79, 42-48

ECKERT, J. (1996):
Der "gefährlichste Fuchsbandwurm" (*Echinococcus multilocularis*) und die alveolare Echinokokkose des Menschen in Mitteleuropa.
Berl. Münch. Tierärztl. Wochenschr. 109, 202-210

ECKERT, J. and THOMPSON, R.C.A. (1996):
Intraspecific variation of *Echinococcus granulosus* and related species.
Acta Trop. 64, 19-34

ECKERT, J. (1997):
Epidemiology of *Echinococcus multilocularis* and *Echinococcus granulosus* in Central Europe.
Parassitologia 39, 337-344

ECKERT, J. (1998):
Alveolar echinococcosis (*Echinococcus multilocularis*) and other forms of echinococcosis (*Echinococcus oligarthus* and *Echinococcus vogeli*). In: PALMER, S.R; SOULSBY, E.J.L. and SIMPSON, D.I.H. (eds.): Zoonoses. Oxford: Oxford University Press, pp. 689-716

ECKERT, J. and DEPLAZES, P. (1999):
Alveolar echinococcosis in humans: the current situation in Central Europe and the need for countermeasures.
Parasitol. Today 15, 315-319

ECKERT, J. (2000):
Helminthosen von Hund und Katze. In: ROMMEL, M.; ECKERT, J.; KUTZER, E.; KÖRTING, W. and SCHNIEDER, T. (eds.): Veterinärmedizinische Parasitologie. 5 vollst. Neubearb. Aufl.-Berlin, Parey, pp. 527-569

ECKERT, J.; SCHANTZ, P.M.; GASSER, R.B.; TORGERSON, P.R.; BESSONOV, A.S.; MOVSESSIAN, S.O.; THAKUR, A.; GRIMM, F. and NIKOGOSSIAN, M.A. (2001 a):
Geographic distribution and prevalence. In: ECKERT, J.; GEMMELL, M.A.; MESLIN, F-X. and PAWLOWSKI, Z. (eds.): WHO/OIE Manual on echinococcosis in humans and animals: a public health problem of global concern. Office International des Epizooties, Paris, pp. 100-142

ECKERT, J.; DEPLAZES, P.; CRAIG, P.S.; GEMMELL, M.A.; GOTTTSTEIN, B.; HEATH, D.; JENKINS, D.J.; KAMIYA, M. and LIGHTOWLERS, M. (2001 b):
Echinococcosis in animals: clinical aspects, diagnosis and treatment. In: ECKERT, J.; GEMMELL, M.A.; MESLIN F-X, PAWLOWSKI, Z. (eds.): WHO/OIE Manual on echinococcosis in humans and animals: a public health problem of global concern. Office International des Epizooties, Paris, pp. 72-99

ECONOMIDES, P. (1998):
Experience gained and evaluation of the echinococcosis/hydatidosis eradication programme in Cyprus. In: Proceedings in the International Conference on Veterinary Medicine and Human Health in the 21st Century, Kuwait, October 19-21

ECONOMIDES, P. and THRASOU, K. (1999):
Echinococcosis/hydatidosis and programmes for its control in the Mediterranean countries.
Arch. Int. Hidatidol. 33, 63-83

ELEZI, B. (1992):
Ekinokokoza paresore dhe dytesore ne Shqiperi.
Shtepia Botuese e Librit Universitar, Universiteti Tirane

EL-SHEHABI, F.S.; KAMHAWI, S.A.; SCHANTZ, P.M.; CRAIG, P.S. and ABDEL-HAFEZ, S.K. (2000):

Diagnosis of canine echinococcosis: Comparison of coproantigen detection with necropsy in stray dogs and red foxes from Northern Jordan.

Parasitology 7, 83-90

EUZEBY, J. (1991):

The epidemiology of hydatidosis with special reference to the Mediterranean area.

Parassitologia 33, 25-39

FARIAS, L.N.; MALGOR, L.; CASSARAVILLA, C.; BRAGANCA, C. and DE LA RUE, M.L. (2004):

Echinococcosis in Southern Brazil: efforts toward implementation of a control program in Santana do Livramento, Rio Grande do Sul.

Rev. Inst. Med trop. S. Paulo 46, 153-156

FRANK, W. (1982):

Biologie und Epidemiologie des *Echinococcus granulosus* und *Echinococcus multilocularis*.

Aktuelle Probl. Chir. Orthop. 23, 12-25

FROSCH, P.M.; MUHLSCHLEGEGL, F.; SYGULLA, L.; HARTMANN, M. and FROSCH, M. (1994):

Identification of cDNA clone from the larval stage of *Echinococcus granulosus* with homologies to the *E. multilocularis* antigen.

Parasitol. Res. 80, 703-705

GASSER, R.B.; LIGHTLOWERS, M.W.; RICKARD, M.D.; LYFORD, R.A. and DAWKINS, H.J.S. (1990):

Serological screening of farm dogs for *Echinococcus granulosus* infection in an endemic region.

Aust. Vet. J. 67, 145-147

GASSER, R.B.; JENKINS, D.J.; PAOLILLO, E.; PARADA, L.; CABRERA, P. and CRAIG, P.S. (1993):

Serum antibodies in canine Echinococcosis.

Int. J. Parasitol. 23, 579-586

GEMMELL, M.A. and JOHNSTONE, P.D. (1976):

Factors regulating tapeworm populations: dispersion of eggs of *Taenia hydatigena* on pasture.

Ann. Trop. Med. Parasitol. 70, 431-434

GEMMELL, M.A. (1978):

The effect of weather on tapeworm eggs and its epidemiological implications.

Weather and Parasitic Animal Disease; World Meteorological Organization Technical Note 159, 83-94

GEMMELL, M.A.; LAWSON, J.R.; ROBERT, M.G.; KERIN, B.R. and MASON, C.C. (1986):

Population dynamics in echinococcosis and cysticercosis: comparison of the response of

Echinococcus granulosus, *Taenia hydatigena* and *Taenia ovis* to control.

Parasitology 93, 357-369

GEMMELL, M.A.; LAWSON, J.R. and ROBERTS, M.G. (1986):
Population dynamics in echinococcosis and cysticercosis, biological parameters of *Echinococcus granulosus* in dogs and sheep.
Parasitology 92, 599-620

GEMMELL, M.A.; LAWSON, J.R. and ROBERTS, M.G. (1987):
Population dynamics in echinococcosis and cysticercosis. Evaluation of the biological parameters of *Taenia hydatigena* and *Taenia ovis* and comparison with those of *Echinococcus granulosus*.
Parasitology 94, 161-180

GEMMELL, M.A. (1990):
Australasian contributions to an understanding of the epidemiology and control of hydatid disease caused by *Echinococcus granulosus*: past, present and future.
Int. J. Parasitol. 20, 431-458

GOTTSTEIN, B.; ECKERT, J. and FEY, R. (1983):
Serological differentiation between *Echinococcus granulosus* and *Echinococcus multilocularis* infection in man.
Z. Parasitenkd. 69, 347-356

GOTTSTEIN, B. and MOWATT, M.R. (1991):
Sequencing and characterization of an *Echinococcus multilocularis* DNA probe and its use in the polymerase chain reaction.
Mol. Biochem. Parasitol. 81, 183-194

GOTTSTEIN, B. and REICHEN, J. (1996):
Echinococcosis/hydatidosis.
In: COOK, G.C. (eds): Mansons Tropical Diseases, London, Saunders, pp. 1486-1508

GOTTSTEIN, B.; SAUCY, F.; DEPLAZES, P.; REICHEN, J.; DEMIERRE, G.; ZÜRCHER, CH.; BUSATO, A. and PUGIN, P. (2001):
Is a high prevalence of *Echinococcus multilocularis* in wild and domestic animals associated with increased disease incidence in humans?
Emerging Infectious Diseases 7, 408-412

GREEN, E.L. (1986):
Immunological detection of parasite antigen in faeces.
Parasitol. Today 3, 144-151

GUARNERA, E.A.; ZANZOTTERA, E.M.; PEREYRA, H. and FRANCO, A.J. (2000):
Ultrasonographic diagnosis of ovine cystic echinococcosis.
Vet. Radiol. Ultras. 42, 352-354

HAHN, E. (1987):
Verbreitung von *Echinococcus granulosus* (1984/85) bei Schlachtrindern an drei Schlachthöfen in Deutschland und Rückübertragungsversuche zur Klärung der Wirtsspezifität.
Vet. Med. Diss., FU-Berlin

HARANDI, M.F.; HOBBS, R.P.; ADAMS, P.J.; MOBEDI, I.; MORGAN-RYAN, U.M. and THOMPSON, R.C. (2002):
Molecular and morphological characterization of *Echinococcus granulosus* of human and animal origin in Iran.
Parasitology 125, 367-373

HÖRCHNER, F.; HAHN, E.; HUWER, M. and SANFT, S. (1986):
Occurrence of hydatid cysts in the Federal Republic of Germany.
XI. International Congress of Infectious and Parasitic Diseases 20.7.-26.7.1986 München

HOXHA, F.T. (2003):
Trajtimi bashkehoore i cistes ekonokokut te melcise ne Kliniken e Kirurgjise.
Punim Magjistrature, Universiteti Prishtine

HÜBNER, K.H. and METZGER, H.O.F. (1982):
Computertomographische Diagnostik der Echinokokkose.
Aktuel. Probl. Chir. Orthop. 23, 75-79

IBRAHEM, M.M.; CRAIG, P.S.; MCVIE, A.; ERSFELD, K. and ROGAN, M.T. (1996):
Echinococcus granulosus antigen B and seroreactivity in natural ovine hydatidosis.
Res. Vet. Sci. 61, 102-106

JENKINS, D.J. and RICKARD, M.D. (1986):
Specific antibody responses in dogs experimentally infected with *Echinococcus granulosus*.
Am. J. Trop. Med. Hyg. 35, 345-349

JENKINS, D.J.; GASSER, R.B.; ZEYHLE, E.; ROMIG, T. and MACPHERSON, C.N.L. (1990): Assessment of a serological test for the detection of *Echinococcus granulosus* infection in dogs in Kenya.
Acta Trop. 47, 245-248

JENKINS, D.J.; FRASER, A.; BRADSHAW, H. and CRAIG, P.H.S. (2000):
Detection of *Echinococcus granulosus* coproantigens in Australian canids with natural or experimental infection.
J. Parasitol. 86, 140-145

JENS, H. and DÖLLE, W. (1982):
Klinik und Differentialdiagnose der Echinokokkose des Menschen.
Aktuel. Probl. Chir. Orthop. 23, 46-51

KAGAN, I.G. (1968):
A review of serological tests for the diagnosis of hydatid disease.
Bulletin of WHO 39, 25-37

KATIC-RADIVOJEVIC, S. and POPOVIC, I. (1997):
Echinococcosis in Yugoslavia (Montenegro): prevalence of *E. granulosus* in dogs and hydatidosis in sheep.
Arch. Int. Hidatidol. 32, 294

KITTELBERGER, R.; REICHEL, M.P.; JENNER, J.; HEATH, D.D.; LIGHTOWLERS, M.W.; MORO, P.; IBRAHEM, M.M.; CRAIG, P.S. and O'KEEFE, J.S. (2002): Evaluation of three enzyme-linked immunosorbent assays (ELISA) for the detection of serum antibodies in sheep infected with *Echinococcus granulosus*.
Vet. Parasitol. 110, 57-76

KOLAROVA, L. (1999):
Echinococcus multilocularis: new epidemiological insights in central and eastern Europe.
Helminthologia 36, 193-200

KUMARATILAKE, L.M. and THOMPSON, R.C.A. (1982)
A review of the taxonomy and specification of the genus Echinococcus Rudolphi.
Z. Parasitenkd. 68, 121-146

KUMARATILAKE, L.M.; THOMPSON, R.C.A. and ECKERT, J. (1986):
Echinococcus granulosus of equine origin from different countries possess uniform morphological characters.
Int. J. Parasitol. 16, 529-540

LAHMAR, S.; KILANI, M.; TORGERSON, P.R. (2001):
Frequency distribution of *Echinococcus granulosus* and other helminths in stray dogs in Tunisia.
Ann. Trop. Med. Parasitol. 95, 69-76

LAHMAR, S.; DEBBEK, H.; ZHANG, L.H.; McMANUS, D.P.; SOUSSI, A; CHELLY, S. and TORGERSON, P.R. (2004):
Transmission dynamics of the *Echinococcus granulosus* sheep-dog strain (G1 genotype) in camels in Tunisia.
Vet. Parasitol. 121, 151-156

LARRIEU, E.J.; COSTA, M.T.; DEL CARPIO, M.; MOGUILLANSKY, S.; BIANCHI, G. and YADON, Z.E. (2002):
A case control study of the risk factors for cystic echinococcosis among the children of Rio Negro province, Argentina.
Ann. Trop. Med. Parasitol. 96, 43-52

LAVIKAINEN, A.; LEHTINEN, M.J.; MERI, T.; HIRVELÄ-KOSKI, V. and MERI, S. (2003):
Molecular genetic characterization of the Fennoscandian cervid strain, a new genotypic group (G 10) of *Echinococcus granulosus*.
Parasitology 127, 207-215

LAWSON, J.R. and GEMMEL, M.A. (1983):
Hydatidosis and Cysticercosis: The dynamics of transmission.
Adv. Parasitol. 22, 261-297

LAWSON, J.R. and GEMMELL, M.A. (1985):
The potential role of blowflies in the transmission of taeniid tapeworm eggs.
Parasitology 91, 129-143

LIGHTLOWERS, M.W.; RICKARD, M.D.; HONEY, R.D.; OBENDORF, D.L. and MITCHELL, G.F. (1984):

Serological diagnosis of *Echinococcus granulosus* infection in sheep using cyst fluid antigen processed by antibody affinity chromatography.

Aust. Vet. J. 61, 101-108

LOPERA, L.; MORO, P.L.; CHAVEZ, A.; MONTES, G.; GONZALEZ, A. and GILMAN, R.H. (2003):

Field evaluation of a coproantigen enzyme-linked immunosorbent assay for diagnosis of canine echinococcosis in a rural Andean village in Peru.

Vet. Parasitol. 117, 37-42

LORENZINI, R. and RUGGIERI, A. (1986):

Distribution of echinococcosis/hydatidosis in Italy.

Parassitologia 34, 45-51

LYMBERY, A.J.; THOMPSON, R.C.A. and HOBBS, R.P. (1990):

Genetic diversity and genetic differentiation in *Echinococcus granulosus* (Batsch, 1786) from domestic and sylvatic hosts on the mainland of Australia.

Parasitology 101, 283-289

MACHNICKA, Z. and KRAWCZUK, S. (1988):

Hymenolepis diminuta antigen: detection of feces in rats.

Bull. Pol. Acad. Science, Biological Science 35, 103-106

MACPHERSON, C.N.L. (1983):

An active intermediate host role for man in the lifecycle of *Echinococcus granulosus* in Turkana, Kenya.

Am. J. Trop. Med. Hyg. 32, 397-404

MACPHERSON, C.N.L. and CRAIG, P.S. (2000):

Dogs and cestode zoonoses. In: MACPHERSON, C.N.L.; MESLIN, F.X. and WANDELER, A.I., (eds.): Dogs, Zoonoses and Public Health. CABI Publishing, Wallingford, UK, pp. 177-211

MALGOR, R.; NONAKA, N.; BASMADJIAN, I.; SAKAI, H.; CARAMBULA, B.; OKU, Y.; CARMONA, C. and KAMIYA, M. (1997):

Coproantigen Detection in Dogs Experimentally and Naturally Infected with *Echinococcus granulosus* by a Monoclonal Antibody-based Enzyme-linked Immunosorbent Assay.

Int. J. Parasitol. 27, 1605-1612

MARTINEZ-FERNANDEZ, A.R. (1986):

The epidemiology of hydatidosis in Spain. COMAC Epidemiology. Workshop on Subspecies Related Incidence of Echinococcosis in Europe. 11-12 October, Athens/Greece.

MATHIS, A.; DEPLAZES, P. and ECKERT, J. (1996):

An improved test system for PCR-based specific detection of *Echinococcus multilocularis* eggs.

J. Helminthol. 70, 219-222

MATOFF, K. (1968):

Neues von der Biologie des *Echinococcus granulosus* und der Echinococcose.
Angew. Parasitol. 9, 65-73

MAXSON, A.D.; WACHIRA, T.M.; ZEHYLE, E.E.; FINE, A.; MWANGI, T.W. and SMITH, G. (1996):

The use of ultrasound to study the prevalence of hydatid cysts in the right lung and liver of sheep and goats in Turkana, Kenya.

Int. J. Parasitol. 26, 1335-1338

McMANUS, D.P. (1981):

A biochemical study of adult and cystic stages of *Echinococcus granulosus* of human and animal origin from Kenya.

J. Helminthol. 55, 21-27

McMANUS, D.P. and MACPHERSON, C.N.L. (1984):

Strain characterization in the hydatid organism *Echinococcus granulosus*: current status and new perspectives.

Ann. Trop. Med. Parasitol. 78, 193-198

MONNIER, PH.; CLIQUET, F.; AUBERT, M. and BRETAGNE, S. (1996):

Improvement of a polymerase chain reaction assay for the detection of *Echinococcus multilocularis* DNA in faecal samples of foxes.

Vet. Parasitol. 67, 185-195

MORO, P.L.; BONIFACIO, N.; GILMAN, R.H.; LOPERA, L.; SILVA, B.; TAKUMOTO, R.; VERASTEGUI, M. and CABRERA, L. (1999):

Field diagnosis of *Echinococcus granulosus* infection among intermediate and definitive hosts in an endemic focus of human cystic echinococcosis.

Trans. Roy. Soc. Trop. Med. Hyg. 93, 611-615

MOVSESIJAN, M. and MLADENOVIC, Z. (1971):

The possibility of using different development stages of *E. granulosus* for detection of specific antibodies against their parasite.

Veterinarski Glasnik 25, 159-163

MÜLLER-GRAF, C.D.M. (1995):

A coprological survey of the intestinal parasites of wild lions (*Panthera leo*) in the Serengeti and the Ngorongoro Crater, Tanzania, East Africa.

J. Parasitol. 8, 812- 814

MWAMBETE, K.D.; PONCE-GORDO, F. and CUESTA-BANDERA, C. (2004):

Genetic identification and host range of the Spanish strains of *Echinococcus granulosus*.

Acta Trop. 91, 87-93

NELSON, G.S. (1972):

Human behaviour in the transmission of parasitic diseases. In: CANNING, E.U. and WRIGHT, C.A. (eds.): Behavioural Aspects of Parasite Transmission, Zoological Journal of the Linnean Society, 51, supplement No.1. London: Academic Press, pp. 109-122

NIETSCHE, W. (1985):
Echinokokkose: Parasitologie, Klinik, Diagnostik, Therapie, Prophylaxe.
Therapiewoche 35, 1547-1554

OKU, Y.; MALGOR, R.; BENAVIDEZ, U.; CARMONA, C. and KAMIYA, H. (2004):
Control program against hydatidosis and the decreased prevalence in Uruguay.
International Congress Series 1267, Japan 28-29 October, 98-104

PALMER, S.R.; BIFFIN, A.H.; CRAIG, P.S. and WALTERS, T.M. (1996):
Control of hydatid disease in Wales.
Br. Med. J. 312, 674-675

PAWLOWSKI, Z.; MROZEWICZ, B.; STEFANIAK, J.; SCHANTZ, P.; WILSON, M.; ECKERT, J.; JAQUIER, P.; HAREMSKI, T.; NOWOSIELSKI, J. and ZIETA, B. (1993):
Echinococcus granulosus pig strain from Poland has a low infectivity to humans.
Am. J. Trop. Med. Hyg. 49 (Supp.), 342-343

PETROVIC, Z. (1979):
Der heutige Stand der Taeniasis und Zystizerkose sowie der Echinococcose und Hydatidose
in Jugoslawien.
Parasit. Hung. 12, 37-39

PIRSCHEL, J. (1982):
Sonographie der Echinokokkose.
Aktuel. Probl. Chir. Orthop. 23, 61-68

RAOUL, F.; DEPLAZES, P.; NONAKA, N.; PIARROUX, R.; VUITTON, D.A. and GIRAUDOUX, P. (2001):
Assessment of the epidemiological status of *Echinococcus multilocularis* in foxes in France
using ELISA coprotest on fox faeces collected in the field.
Int. J. Parasitol. 31, 1579-1588

RAUSCH, R.L. (1967):
On the ecology and distribution of *Echinococcus spp.* (Cestoda: Taenidae), and characteristics
of their development in the intermediate host.
Annu. Parasitol. 42, 19-63

RAUSCH, R.L. and BERNSTEIN, J.J. (1972):
Echinococcus vogeli spp. (Cestoda : Taeniidae) from the bushdog *Speothos venaticus* (Lund).
Z. Tropenmed. Parasitol. 23, 25-34

RAUSCH, R.L.; RAUSCH, V.R. and D'ALESSANDRO, A. (1978):
Discrimination of the larval stages of *Echinococcus oligarthrus* (Diesing 1863).
Am. J. Trop. Med. Hyg. 27, 1195-1202

RAUSCH, R.L.; FAY, F.H. and WILLIAMSON, F.S. (1990):
The ecology of *Echinococcus multilocularis* (Cestoda: Taeniidae) on St. Lawrence Island,
Alaska. II. Helminth populations in the definitive host.
Ann. Parasitol. Hum. Comp. 65, 131-140

RAUSCH, R.L. (1995):

Life cycle patterns and geographic distribution of *Echinococcus* species. In: THOMPSON, R.C.A. (eds.): The Biology of *Echinococcus* and Hydatid Disease. George Allen and Unwin, London, pp. 44-80

SADAKA, H.A.; KHALIFA, A.M.; ELDEIN, S.Z.; TAHIA, K. and ELDEIN, K.M. (2002):

Urinary antigen detection for diagnosis of hydatid disease.

J. Egypt. Soc. Parasitol. 32, 69-78

SAKAI, H.; MALGOR, R.; BASMADJIAN, I.; GALLARDO, R.; CARMONA, C.; SATO, H.; OKU, Y. and KAMIYA, M. (1995):

An enzyme-linked immunosorbent assay (ELISA) for the detection of *Echinococcus granulosus* coproantigens in dogs.

Jap. Jour. Parasitol. 44, 453-461

SAKASHITA, M.; SAKAI, H.; KOHNO, H.; OOI, H.K.; OKU, Y.; YAGI, M.; ITO, M. and KAMIYA, M. (1995):

Detection of *Echinococcus multilocularis* coproantigens in experimentally infected dogs using murine monoclonal antibody against adult worms.

Jap. Jour. Parasitol. 44, 413-420

SAMPIO SILVA, M.L.; ROMBERT, P.; VEIGA, S.G.; MORAES, D. and SCHURIAN, E. (1986):

Epidemiologie et mesures de controle dans L'echinococcosse-hydatidose au Portugal. COMAC Epidemiology. Workshop on Subspecies Related Incidence of Echinococcosis in Europe. 19-21 October, Athens/Greece.

SCHANTZ, P.M. and SCHWABE, C.W. (1969):

Worldwide status of hydatid disease control.

J. Am. Vet. Assoc. 155, 2104-2121

SCHANTZ, P.M. and COLLI, C. (1973):

Echinococcus oligarthrus (Diesing 1863) from Geoffroy's Cat (*Felis geoffroyi* D'Orbigny y Gervais) in temperate South Amerika.

J. Parasitol. 59, 1138-114

SCHANTZ, P.P.; CHAI, J.; CRAIG, P.S.; ECKERT, J.; JENKINS, D.J.; MACPHERSON, C.N.L. and THAKUR, A. (1995):

Epidemiology and Control of Hydatid Disease. In: THOMSON, R.C.A. and LYMBERY, A.J.(eds.): *Echinococcus* and Hydatid Disease, CAB International, Wallingford, pp. 233- 331

SCOTT, J.C.; STEFANIAK, J.; PAWLOWSKI, Z.S. and McMANUS, D.P. (1997):

Molecular genetic analysis of human hydatid cases from Poland: identification of a new genotypic group of *Echinococcus granulosus*.

Parasitology 114, 37-43

SHAIKENOV, B.S; RYSMUKHAMBETOVA, A.T.; MASSENOV, B.; DEPLAZES, P.; MATHIS, A. and TORGERSON, P.R. (2004):

Short Report: The use of a polymerase chain reaction to detect *Echinococcus granulosus* (G1 strain) eggs in soil samples.

Am. J. Trop. Med. Hyg. 71(4), 441-443

SIDRANSKI, D.; TOKINO, T.; HAMILTON, S.R.; KINZLER, K.W.; LEVIN, B.; FROST, P. and VOGELSTEIN, B. (1992):

Identification of ras oncogene mutations in the stool of patients with curable colorectal tumors.

Science 256, 102-104

SILES-LUCAS, M. and GOTTSSTEIN, B. (2001):

Molecular tools for the diagnosis of cystic and alveolar echinococcosis.

Trop. Med. Int. Health 6, 463-475

SILES-LUCAS, M.; FELLEISEN, R.; CUESTA-BANDERA, C.; GOTTSSTEIN, B. and ECKERT, J. (1994):

Comparative genetic analysis of Swiss and Spanish isolates of *Echinococcus granulosus* by Southern hybridisation and Random Amplified Polymorphic DNA technique.

Appl. Parasitol. 35, 107-117

SMITH, G.C.; GANGADHARAN, B.; TAYLOR, Z.; LAURENSEN, K.; BRADSHAW, H.; HIDE, G.; HUGHES, J.M.; DINKEL, A.; ROMIG, T. and CRAIG, P.S. (2003):

Prevalence of zoonotic important parasites in the red fox (*Vulpes vulpes*) in Great Britain.

Vet. Parasitol. 118, 133-142

SMYTH, J.D. and DAVIES, Z. (1974):

Occurrence of physiological strains of *Echinococcus granulosus* demonstrated by in vitro culture of protoscolices from sheep and horse hydatid cysts.

Int. J. Parasitol. 4, 443-445

SOILEV, CH. and BOEVA, V. (1982):

Zur Epidemiologie der Echinococcosis bei Haustieren und Menschen in der VR Bulgarien.

Angew. Parasitol. 23, 129-134

SOUSA, O.E. and THATCHER, V.E. (1969):

Observations of the life-cycle of *Echinococcus oligarthrus* (Diesing 1863) in the Republic of Panama.

Ann. Trop. Med. Parasitol. 63, 165-175

STEFANIC, S.; SHAIKENOV, B.; DEPLAZES, P.; DINKEL, A.; TORGERSON, P. and MATHIS, A. (2004):

Polymerase chain reaction for detection of patent infections of *Echinococcus granulosus* ("sheep-strain") in naturally infected dogs.

Parasitol. Res. 92, 347-351

STIGER, C.; HEGGLIN, G.; SCHWARZENBACH, G.; MATHIS, A. and DEPLAZES, P. (2002):

Spatial and temporal aspects of urban transmission of *E.multilocularis*.

Parasitology 124, 631-640

STÖSSEL, T. (1989):

Literaturübersicht zur Häufigkeit und geographischen Verbreitung der Echinococcosis bei Menschen und Tieren in Ländern der EG und EFTA.

Vet. Med. Diss., Universität Zürich

SVOBODOVA, V. and LENSKA, B. (2002):

Echinococcosis in dogs in the Czech Republic.

Acta Vet. Brno 71, 347-350

THENEVET, P.S.; JENSEN, O.; MELLADO, I.; TORRECILLAS, C.; RASO, S.; FLORES, M.E.; MINVIELLI, M.C. and BASUALDO, J.A. (2003):

Presence and persistence of intestinal parasites in canine fecal material collected from the environment in the Province of Chubut, Argentine Patagonia.

Vet. Parasitol. 117, 263-269

THOMPSON, R.C.A (1979):

Biology and speciation of *Echinococcus granulosus*.

Aust. Vet. J. 55, 93-98

THOMPSON, R.C.A. and KUMARATILAKE, L.M. (1982):

Intraspecific variation in *Echinococcus granulosus*: The Australian situation and perspectives for the future.

Trans. Roy. Soc. Trop. Med. Hyg. 76, 13-16

THOMPSON, R.C.; KUMARATILAKE, A. and ECKERT, J. (1984):

Observations on *Echinococcus granulosus* of cattle origin in Switzerland.

Int. J. Parasitol. 14, 283-291

THOMPSON, R.C.A. (1986):

Biology and systematics of Echinococcus. In: THOMPSON, R.C.A. (eds.): The biology of Echinococcus and hydatid disease. George Allen & Unwin, London, pp. 5-43

THOMPSON, R.C.A. and LYMBERRY, A.J. (1988):

The nature, extent and significance of variation within the genus Echinococcus.

Adv. Parasitol. 27, 209-259

THOMPSON, R.C. and LYMBERRY, A.J.(1996):

Genetic variability in parasites and host-parasite interactions.

Parasitology 112, 7-22

THOMPSON, R.C.A. and McMANUS, D.P. (2002):

Towards a taxonomic revision of the genus Echinococcus.

Trends in Parasitology 18, 452-457

TODOROV, T. and BOEVA, B. (1999):
Human echinococcosis in Bulgaria: a comparative epidemiological analysis.
Bulletin of the WHO 77, 110-118

TORGERSON, P.R.; SHAIKENOV, B.S.; RYSMUKHAMBETOVA, A.T.; USSENBAYEV, A.E.; ABDYBEKOVA, A.M. and BURTISURNOV, K.K. (2003):
Modelling the transmission dynamics of *Echinococcus granulosus* in dogs in rural Kazakhstan.
Parasitology 126, 417-424

VAN DER GIESSEN, J.W.B.; ROMBOUT, Y.; FRANCHIMONT, H.; LIMPER, L. and HOMAN, W. (1999):
Detection of *Echinococcus multilocularis* in foxes in the Netherlands.
Vet. Parasitol. 82, 49-57

VEIT, P.; BILGER, B.; SCHAD, V.; SCHÄFER, J.; FRANK, W. and LUCIUS, R. (1995):
Influence of environmental factors on the infectivity of *Echinococcus multilocularis* eggs.
Parasitology 110, 79-86

VISCIDI, R.; LAUGHON, B.E.; HANVANICH, M.; BARTLET, J.G. and YOLKEN, R.H. (1984):
Improved enzyme immunoassays for detection of antigens in fecal specimens. Investigation
and correction of interfering factors.
J. Immunol. Methods 67, 129-143

VUITTON, D.A. (2002):
Host immunogenetics and role in epidemiology of larval cestodes. In: CRAIG, P.S. and PAWLOWSKI, Z. (eds.): Cestode Zoonoses, Echinococcosis and Cysticercosis. An Emergent
and Global Problem Nato Science Series, Vol. 341, Amsterdam: IOS Press, pp. 183-194

WACHIRA, T.M.; MACPHERSON, C.N.L. and GATHUMA, J.M. (1991):
Release and survival of *Echinococcus granulosus* eggs in different environments in Turkana,
and their possible impact on the incidence of hydatidosis in man and livestock.
J. Helminthol. 65, 55-61

WALTHER, E. (1982):
Konventionelle Röntgendiagnostik der Echinokokkose.
Aktuel. Probl. Chir. Orthop. 23, 61-68

WANG, Q.; QIU, J.; SCHANTZ, P.; HE, J.; ITO, A. and LIU, F. (2001):
Investigation of risk factor for development of human hydatidosis among households raising
livestock in Tibetan areas of western Sichuan Province.
Chin. J. Parasitol. Parasitic Disease 19, 93-96

WIKERHAUSER, T. and BRGLEZ, J. (1984):
An attempt to infect calves with *Echinococcus granulosus* of porcine origine. The fourth
European multicolloquium of parasitology. Oktober 1984, Izmir.

WORBES, H. (1992):

Zum Vorkommen von *Echinococcus granulosus* und *E. multilocularis* in Thüringen.
Angew. Parasitol. 33, 193-204

World Health Organization (1981):

FAO/UNEP/WHO guidelines on echinococcosis/hydatidosis surveillance, prevention and control.

Wrld. Hlth. Org. Geneva, Zwitzerland

WYN-JONES, G. and CLARKSON, M.J. (1984):

Radiologic detection of ovine hydatidosis.

Vet. Radiology 25, 182-186

YONG, W.K.; HEATH, D.D. and VAN KNAPEL, F. (1984):

Comparison of cestode antigens in an enzyme-linked immunosorbent assay for the diagnosis of *Echinococcus granulosus*, *T. hydatigena* and *T. ovis* infections in sheep.

Res. Vet. Sci. 36, 24-31