

Literaturverzeichnis

- [1] S. Tremmel. *Design, Synthese und Stabilität von beta-Faltblatt Modellpeptiden*. Dissertation, Martin-Luther-Universität, 2005.
- [2] B. Rost. Did evolution leap to create the protein universe? *Curr Opin Struct Biol*, 12(3):409–416, 2002.
- [3] L. Holm, C. Sander. Dictionary of recurrent domains in protein structures. *Proteins*, 33(1):88–96, 1998.
- [4] M. Ramirez-Alvarado, T. Kortemme, F. J. Blanco, L. Serrano. Beta-hairpin and beta-sheet formation in designed linear peptides. *Bioorg Med Chem*, 7(1):93–103, 1999.
- [5] F. A. Syud, H. E. Stanger, S. H. Gellman. Interstrand side chain–side chain interactions in a designed beta-hairpin: significance of both lateral and diagonal pairings. *J Am Chem Soc*, 123(36):8667–8677, 2001.
- [6] C. L. Nesloney, J. W. Kelly. Progress towards understanding beta-sheet structure. *Bioorg Med Chem*, 4(6):739–766, 1996.
- [7] M. Ramirez-Alvarado, F. J. Blanco, L. Serrano. De novo design and structural analysis of a model beta-hairpin peptide system. *Nat Struct Biol*, 3(7):604–612, 1996.
- [8] J. F. Espinosa, V. Munoz, S. H. Gellman. Interplay between hydrophobic cluster and loop propensity in beta-hairpin formation. *J Mol Biol*, 306(3):397–402, 2001.
- [9] B. Ciani, M. Jourdan, M. S. Searle. Stabilization of beta-hairpin peptides by salt bridges: role of preorganization in the energetic contribution of weak interactions. *J Am Chem Soc*, 125(30):9038–9047, 2003.
- [10] A. G. Street, S. L. Mayo. Intrinsic beta-sheet propensities result from van der waals interactions between side chains and the local backbone. *Proc Natl Acad Sci U S A*, 96(16):9074–9076, 1999.

- [11] S. R. Griffiths-Jones, G. J. Sharman, A. J. Maynard, M. S. Searle. Modulation of intrinsic phi,psi propensities of amino acids by neighbouring residues in the coil regions of protein structures: Nmr analysis and dissection of a beta-hairpin peptide. *J Mol Biol*, 284(5):1597–1609, 1998.
- [12] C. M. Santiveri, M. Rico, M. A. Jimenez, M. T. Pastor, E. Perez-Paya. Insights into the determinants of beta-sheet stability: 1h and 13c nmr conformational investigation of three-stranded antiparallel beta-sheet-forming peptides. *J Pept Res*, 61(4):177–188, 2003.
- [13] R. S. Prajapati, M. Sirajuddin, V. Durani, S. Sreeramulu, R. Varadarajan. Contribution of cation-pi interactions to protein stability. *Biochemistry*, 45(50):15000–15010, 2006.
- [14] J. F. Espinosa, F. A. Syud, S. H. Gellman. Analysis of the factors that stabilize a designed two-stranded antiparallel beta-sheet. *Protein Sci*, 11(6):1492–1505, 2002.
- [15] A. K. Dunker, Z. Obradovic. The protein trinity–linking function and disorder. *Nat Biotechnol*, 19(9):805–806, 2001.
- [16] L. M. Iakoucheva, C. J. Brown, J. D. Lawson, Z. Obradovic, A. K. Dunker. Intrinsic disorder in cell-signaling and cancer-associated proteins. *J Mol Biol*, 323(3):573–584, 2002.
- [17] D. Tobi, I. Bahar. Structural changes involved in protein binding correlate with intrinsic motions of proteins in the unbound state. *Proc Natl Acad Sci U S A*, 102(52):18908–18913, 2005.
- [18] J. M. Thornton, A. E. Todd, D. Milburn, N. Borkakoti, C. A. Orengo. From structure to function: approaches and limitations. *Nat Struct Biol*, 7 Suppl:991–994, 2000.
- [19] M. Jager, H. Nguyen, J. C. Crane, J. W. Kelly, M. Gruebele. The folding mechanism of a beta-sheet: the ww domain. *J Mol Biol*, 311(2):373–393, 2001.
- [20] X. Jiang, J. Kowalski, J. W. Kelly. Increasing protein stability using a rational approach combining sequence homology and structural alignment: Stabilizing the ww domain. *Protein Sci*, 10(7):1454–1465, 2001.
- [21] C. Sigurdson, M. Polymenidou, A. Aguzzi. Reconstructing prions: fibril assembly from simple yeast to complex mammals. *Neurodegener Dis*, 2(1):1–5, 2005.
- [22] J. Ziegler, C. Viehrig, S. Geimer, P. Rosch, S. Schwarzinger. Putative aggregation initiation sites in prion protein. *FEBS Lett*, 580(8):2033–2040, 2006.

- [23] L. M. Miller, Q. Wang, T. P. Telivala, R. J. Smith, A. Lanzirotti, J. Miklossy. Synchrotron-based infrared and x-ray imaging shows focalized accumulation of cu and zn co-localized with beta-amyloid deposits in alzheimer's disease. *J Struct Biol*, 155(1):30–37, 2006.
- [24] S. Lesne, M. T. Koh, L. Kotilinek, R. Kayed, C. G. Glabe, A. Yang, M. Gallagher, K. H. Ashe. A specific amyloid-beta protein assembly in the brain impairs memory. *Nature*, 440(7082):352–357, 2006.
- [25] M. R. Hayden, S. C. Tyagi, M. M. Kerklo, M. R. Nicolls. Type 2 diabetes mellitus as a conformational disease. *Jop*, 6(4):287–302, 2005.
- [26] C. M. Cowan, L. A. Raymond. Selective neuronal degeneration in huntington's disease. *Curr Top Dev Biol*, 75:25–71, 2006.
- [27] M. Gross. Proteins that convert from alpha helix to beta sheet: implications for folding and disease. *Curr Protein Pept Sci*, 1(4):339–347, 2000.
- [28] Y. Takahashi, A. Ueno, H. Mihara. Mutational analysis of designed peptides that undergo structural transition from alpha helix to beta sheet and amyloid fibril formation. *Structure*, 8(9):915–925, 2000.
- [29] M. Lopez de la Paz, L. Serrano. Sequence determinants of amyloid fibril formation. *Proc Natl Acad Sci U S A*, 101(1):87–92, 2004.
- [30] R. Srinivasan, E. M. Jones, K. Liu, J. Ghiso, R. E. Marchant, M. G. Zagorski. ph-dependent amyloid and protofibril formation by the abri peptide of familial british dementia. *J Mol Biol*, 333(5):1003–1023, 2003.
- [31] K. Giri, N. P. Bhattacharyya, S. Basak. ph-dependent self-assembly of polyalanine peptides. *Biophys J*, 92(1):293–302, 2007.
- [32] T. Wang, R. C. Wade. Implicit solvent models for flexible protein-protein docking by molecular dynamics simulation. *Proteins*, 50(1):158–169, 2003.
- [33] K. Chen, J. Maley, P. H. Yu. Potential implications of endogenous aldehydes in beta-amyloid misfolding, oligomerization and fibrillogenesis. *J Neurochem*, 2006.
- [34] E. Malolepsza, M. Boniecki, A. Kolinski, L. Piela. Theoretical model of prion propagation: a misfolded protein induces misfolding. *Proc Natl Acad Sci U S A*, 102(22):7835–7840, 2005.
- [35] D. M. Walsh, I. Klyubin, J. V. Fadeeva, W. K. Cullen, R. Anwyl, M. S. Wolfe, M. J. Rowan, D. J. Selkoe. Naturally secreted oligomers of amyloid beta protein potent-

- ly inhibit hippocampal long-term potentiation in vivo. *Nature*, 416(6880):535–539, 2002.
- [36] D. M. Walsh, D. J. Selkoe. Deciphering the molecular basis of memory failure in alzheimer's disease. *Neuron*, 44(1):181–193, 2004.
- [37] R. L. Baldwin. Alpha-helix formation by peptides of defined sequence. *Biophys Chem*, 55(1-2):127–135, 1995.
- [38] E. Lacroix, A. R. Viguera, L. Serrano. Elucidating the folding problem of alpha-helices: local motifs, long-range electrostatics, ionic-strength dependence and prediction of nmr parameters. *J Mol Biol*, 284(1):173–191, 1998.
- [39] C. M. Santiveri, J. Santoro, M. Rico, M. A. Jimenez. Factors involved in the stability of isolated beta-sheets: Turn sequence, beta-sheet twisting, and hydrophobic surface burial. *Protein Sci*, 13(4):1134–1147, 2004.
- [40] M. Lopez De La Paz, K. Goldie, J. Zurdo, E. Lacroix, C. M. Dobson, A. Hoenger, L. Serrano. De novo designed peptide-based amyloid fibrils. *Proc Natl Acad Sci U S A*, 99(25):16052–16057, 2002.
- [41] M. T. Pastor, A. Esteras-Chopo, M. Lopez de la Paz. Design of model systems for amyloid formation: lessons for prediction and inhibition. *Curr Opin Struct Biol*, 15(1):57–63, 2005.
- [42] E. Lacroix, T. Kortemme, M. Lopez de la Paz, L. Serrano. The design of linear peptides that fold as monomeric beta-sheet structures. *Curr Opin Struct Biol*, 9(4):487–493, 1999.
- [43] L. J. Ball, R. Kuhne, J. Schneider-Mergener, H. Oschkinat. Recognition of proline-rich motifs by protein-protein-interaction domains. *Angew Chem Int Ed Engl*, 44(19):2852–2869, 2005.
- [44] S. S. Li. Specificity and versatility of sh3 and other proline-recognition domains: structural basis and implications for cellular signal transduction. *Biochem J*, 390(Pt 3):641–653, 2005.
- [45] M. Sudol. Structure and function of the ww domain. *Prog Biophys Mol Biol*, 65(1-2):113–132, 1996.
- [46] M. Sudol, H. I. Chen, C. Bougeret, A. Einbond, P. Bork. Characterization of a novel protein-binding module—the ww domain. *FEBS Lett*, 369(1):67–71, 1995.

- [47] S. Wiesner, G. Stier, M. Sattler, M. J. Macias. Solution structure and ligand recognition of the ww domain pair of the yeast splicing factor prp40. *J Mol Biol*, 324(4):807–822, 2002.
- [48] O. Y. Fedoroff, S. A. Townson, A. P. Golovanov, M. Baron, J. M. Avis. The structure and dynamics of tandem ww domains in a negative regulator of notch signaling, suppressor of deltex. *J Biol Chem*, 279(33):34991–5000, 2004.
- [49] P. Bork, M. Sudol. The ww domain: a signalling site in dystrophin? *Trends Biochem Sci*, 19(12):531–533, 1994.
- [50] H. I. Chen, M. Sudol. The ww domain of yes-associated protein binds a proline-rich ligand that differs from the consensus established for src homology 3-binding modules. *Proc Natl Acad Sci U S A*, 92(17):7819–7823, 1995.
- [51] M. J. Macias, M. Hyvonen, E. Baraldi, J. Schultz, M. Sudol, M. Saraste, H. Oschkinat. Structure of the ww domain of a kinase-associated protein complexed with a proline-rich peptide. *Nature*, 382(6592):646–649, 1996.
- [52] E. K. Koepf, H. M. Petrassi, M. Sudol, J. W. Kelly. Ww: An isolated three-stranded antiparallel beta-sheet domain that unfolds and refolds reversibly; evidence for a structured hydrophobic cluster in urea and gdnhcl and a disordered thermal unfolded state. *Protein Sci*, 8(4):841–853, 1999.
- [53] V. Kanelis, D. Rotin, J. D. Forman-Kay. Solution structure of a nedd4 ww domain-enac peptide complex. *Nat Struct Biol*, 8(5):407–412, 2001.
- [54] M. A. Verdecia, M. E. Bowman, K. P. Lu, T. Hunter, J. P. Noel. Structural basis for phosphoserine-proline recognition by group iv ww domains. *Nat Struct Biol*, 7(8):639–643, 2000.
- [55] M. J. Macias, V. Gervais, C. Civera, H. Oschkinat. Structural analysis of ww domains and design of a ww prototype. *Nat Struct Biol*, 7(5):375–379, 2000.
- [56] B. Aghazadeh, M. K. Rosen. Ligand recognition by sh3 and ww domains: the role of n-alkylation in ppii helices. *Chem Biol*, 6(9):R241–6, 1999.
- [57] B. K. Kay, M. P. Williamson, M. Sudol. The importance of being proline: the interaction of proline-rich motifs in signaling proteins with their cognate domains. *Faseb J*, 14(2):231–241, 2000.
- [58] A. Zarrinpar, R. P. Bhattacharyya, W. A. Lim. The structure and function of proline recognition domains. *Sci STKE*, 2003(179):RE8, 2003.

- [59] T. P. Creamer. Left-handed polyproline ii helix formation is (very) locally driven. *Proteins*, 33(2):218–226, 1998.
- [60] B. Bochicchio, A. M. Tamburro. Polyproline ii structure in proteins: identification by chiroptical spectroscopies, stability, and functions. *Chirality*, 14(10):782–792, 2002.
- [61] L. Otte, U. Wiedemann, B. Schlegel, J. R. Pires, M. Beyermann, P. Schmieder, G. Krause, R. Volkmer-Engert, J. Schneider-Mergener, H. Oschkinat. Ww domain sequence activity relationships identified using ligand recognition propensities of 42 ww domains. *Protein Sci*, 12(3):491–500, 2003.
- [62] F. Toepert, J. R. Pires, C. Landgraf, H. Oschkinat, J. Schneider-Mergener. Synthesis of an array comprising 837 variants of the hyap ww protein domain. *Angew Chem Int Ed Engl*, 40(5):897–900, 2001.
- [63] M. B. Yaffe, M. Schutkowski, M. Shen, X. Z. Zhou, P. T. Stukenberg, J. U. Rahfeld, J. Xu, J. Kuang, M. W. Kirschner, G. Fischer, L. C. Cantley, K. P. Lu. Sequence-specific and phosphorylation-dependent proline isomerization: a potential mitotic regulatory mechanism. *Science*, 278(5345):1957–1960, 1997.
- [64] M. Shen, P. T. Stukenberg, M. W. Kirschner, K. P. Lu. The essential mitotic peptidyl-prolyl isomerase pin1 binds and regulates mitosis-specific phosphoproteins. *Genes Dev*, 12(5):706–720, 1998.
- [65] K. P. Lu, Y. C. Liou, I. Vincent. Proline-directed phosphorylation and isomerization in mitotic regulation and in alzheimer's disease. *Bioessays*, 25(2):174–181, 2003.
- [66] N. Abovich, M. Rosbash. Cross-intron bridging interactions in the yeast commitment complex are conserved in mammals. *Cell*, 89(3):403–412, 1997.
- [67] M. T. Bedford, R. Reed, P. Leder. Ww domain-mediated interactions reveal a spliceosome-associated protein that binds a third class of proline-rich motif: the proline glycine and methionine-rich motif. *Proc Natl Acad Sci U S A*, 95(18):10602–10607, 1998.
- [68] N. R. Gavva, R. Gavva, K. Ermekova, M. Sudol, C. J. Shen. Interaction of ww domains with hematopoietic transcription factor p45/nf-e2 and rna polymerase ii. *J Biol Chem*, 272(39):24105–24108, 1997.
- [69] A. Komuro, M. Saeki, S. Kato. Association of two nuclear proteins, npw38 and npwbp, via the interaction between the ww domain and a novel proline-rich motif containing glycine and arginine. *J Biol Chem*, 274(51):36513–36519, 1999.

- [70] R. Yagi, L. F. Chen, K. Shigesada, Y. Murakami, Y. Ito. A ww domain-containing yes-associated protein (yap) is a novel transcriptional co-activator. *Embo J*, 18(9):2551–2562, 1999.
- [71] M. Sudol, K. Sliwa, T. Russo. Functions of ww domains in the nucleus. *FEBS Lett*, 490(3):190–195, 2001.
- [72] A. Komuro, M. Nagai, N. E. Navin, M. Sudol. Ww domain-containing protein yap associates with erbB-4 and acts as a co-transcriptional activator for the carboxyl-terminal fragment of erbB-4 that translocates to the nucleus. *J Biol Chem*, 278(35):33334–33341, 2003.
- [73] S. Rentschler, H. Linn, K. Deininger, M. T. Bedford, X. Espanol, M. Sudol. The ww domain of dystrophin requires ef-hands region to interact with beta-dystroglycan. *Biol Chem*, 380(4):431–442, 1999.
- [74] J. L. Ilsley, M. Sudol, S. J. Winder. The ww domain: linking cell signalling to the membrane cytoskeleton. *Cell Signal*, 14(3):183–189, 2002.
- [75] O. Staub, S. Dho, P. Henry, J. Correa, T. Ishikawa, J. McGlade, D. Rotin. Ww domains of nedd4 bind to the proline-rich py motifs in the epithelial na⁺ channel deleted in liddle's syndrome. *Embo J*, 15(10):2371–2380, 1996.
- [76] L. Schild, Y. Lu, I. Gautschi, E. Schneeberger, R. P. Lifton, B. C. Rossier. Identification of a py motif in the epithelial na channel subunits as a target sequence for mutations causing channel activation found in liddle syndrome. *Embo J*, 15(10):2381–2387, 1996.
- [77] T. Russo, R. Faraonio, G. Minopoli, P. De Candia, S. De Renzis, N. Zambrano. Fe65 and the protein network centered around the cytosolic domain of the alzheimer's beta-amyloid precursor protein. *FEBS Lett*, 434(1-2):1–7, 1998.
- [78] P. J. Lu, G. Wulf, X. Z. Zhou, P. Davies, K. P. Lu. The prolyl isomerase pin1 restores the function of alzheimer-associated phosphorylated tau protein. *Nature*, 399(6738):784–788, 1999.
- [79] L. A. Passani, M. T. Bedford, P. W. Faber, K. M. McGinnis, A. H. Sharp, J. F. Gusella, J. P. Vonsattel, M. E. MacDonald. Huntington's ww domain partners in huntington's disease post-mortem brain fulfill genetic criteria for direct involvement in huntington's disease pathogenesis. *Hum Mol Genet*, 9(14):2175–2182, 2000.
- [80] M. Sudol. From src homology domains to other signaling modules: proposal of the 'protein recognition code'. *Oncogene*, 17(11 Reviews):1469–1474, 1998.

- [81] A. K. Bednarek, K. J. Laflin, R. L. Daniel, Q. Liao, K. A. Hawkins, C. M. Aldaz. Wwox, a novel ww domain-containing protein mapping to human chromosome 16q23.3-24.1, a region frequently affected in breast cancer. *Cancer Res*, 60(8):2140–2145, 2000.
- [82] H. Lubs, F. E. Abidi, R. Echeverri, L. Holloway, A. Meindl, R. E. Stevenson, C. E. Schwartz. Golabi-ito-hall syndrome results from a missense mutation in the ww domain of the pqbp1 gene. *J Med Genet*, 43(6):e30, 2006.
- [83] J. Przezdziak, S. Tremmel, I. Kretzschmar, M. Beyermann, M. Bienert, R. Volkmer-Engert. Probing the ligand-binding specificity and analyzing the folding state of spot-synthesized fbp28 ww domain variants. *Chembiochem*, 7(5):780–788, 2006.
- [84] M. Petrovich, A. L. Jonsson, N. Ferguson, V. Daggett, A. R. Fersht. Phi-analysis at the experimental limits: Mechanism of beta-hairpin formation. *J Mol Biol*, 2006.
- [85] Y. W. Chen, A. R. Fersht, K. Henrick. Contribution of buried hydrogen bonds to protein stability. the crystal structures of two barnase mutants. *J Mol Biol*, 234(4):1158–1170, 1993.
- [86] K. Takano, Y. Yamagata, M. Kubota, J. Funahashi, S. Fujii, K. Yutani. Contribution of hydrogen bonds to the conformational stability of human lysozyme: calorimetry and x-ray analysis of six ser → ala mutants. *Biochemistry*, 38(20):6623–6629, 1999.
- [87] C. N. Pace, G. Horn, E. J. Hebert, J. Bechert, K. Shaw, L. Urbanikova, J. M. Scholtz, J. Sevcik. Tyrosine hydrogen bonds make a large contribution to protein stability. *J Mol Biol*, 312(2):393–404, 2001.
- [88] G. Bulaj, D. P. Goldenberg. Mutational analysis of hydrogen bonding residues in the bpti folding pathway. *J Mol Biol*, 313(3):639–656, 2001.
- [89] S. K. Holmgren, K. M. Taylor, L. E. Bretscher, R. T. Raines. Code for collagen's stability deciphered. *Nature*, 392(6677):666–667, 1998.
- [90] R. Kaul, A. R. Angeles, M. Jager, E. T. Powers, J. W. Kelly. Incorporating beta-turns and a turn mimetic out of context in loop 1 of the ww domain affords cooperatively folded beta-sheets. *J Am Chem Soc*, 123(22):5206–5212, 2001.
- [91] C. Renner, S. Alefelder, J. H. Bae, N. Budisa, R. Huber, L. Moroder. Fluoroprolines as tools for protein design and engineering. *Angew Chem Int Ed Engl*, 40(5):923–925, 2001.

- [92] Y. Tang, G. Ghirlanda, N. Vaidehi, J. Kua, D. T. Mainz, Iii Wa Goddard, W. F. DeGrado, D. A. Tirrell. Stabilization of coiled-coil peptide domains by introduction of trifluoroleucine. *Biochemistry*, 40(9):2790–2796, 2001.
- [93] N. C. Yoder, K. Kumar. Fluorinated amino acids in protein design and engineering. *Chem Soc Rev*, 31(6):335–341, 2002.
- [94] N. Ferguson, J. R. Pires, F. Toepert, C. M. Johnson, Y. P. Pan, R. Volkmer-Engert, J. Schneider-Mergener, V. Daggett, H. Oschkinat, A. Fersht. Using flexible loop mimetics to extend phi-value analysis to secondary structure interactions. *Proc Natl Acad Sci U S A*, 98(23):13008–13013, 2001.
- [95] S. Deechongkit, P. E. Dawson, J. W. Kelly. Toward assessing the position-dependent contributions of backbone hydrogen bonding to beta-sheet folding thermodynamics employing amide-to-ester perturbations. *J Am Chem Soc*, 126(51):16762–16771, 2004.
- [96] S. Deechongkit, H. Nguyen, E. T. Powers, P. E. Dawson, M. Gruebele, J. W. Kelly. Context-dependent contributions of backbone hydrogen bonding to beta-sheet folding energetics. *Nature*, 430(6995):101–105, 2004.
- [97] S. Deechongkit, S. L. You, J. W. Kelly. Synthesis of all nineteen appropriately protected chiral alpha-hydroxy acid equivalents of the alpha-amino acids for boc solid-phase depsipeptide synthesis. *Org Lett*, 6(4):497–500, 2004.
- [98] S. Deechongkit, H. Nguyen, M. Jager, E. T. Powers, M. Gruebele, J. W. Kelly. beta-sheet folding mechanisms from perturbation energetics. *Curr Opin Struct Biol*, 16(1):94–101, 2006.
- [99] Jr. Pearce, K. H., M. H. Ultsch, R. F. Kelley, A. M. de Vos, J. A. Wells. Structural and mutational analysis of affinity-inert contact residues at the growth hormone-receptor interface. *Biochemistry*, 35(32):10300–10307, 1996.
- [100] E. Di Cera. Site-specific thermodynamics: Understanding cooperativity in molecular recognition. *Chem Rev*, 98(4):1563–1592, 1998.
- [101] B. Holst, S. Zoffmann, C. E. Elling, S. A. Hjorth, T. W. Schwartz. Steric hindrance mutagenesis versus alanine scan in mapping of ligand binding sites in the tachykinin nk1 receptor. *Mol Pharmacol*, 53(1):166–175, 1998.
- [102] Ronald Frank. Spot-synthesis: an easy technique for the positionally addressable, parallel chemical synthesis on a membrane support. *Tetrahedron*, 48(42):9217–9232, 1992.

- [103] H. Wenschuh, R. Volkmer-Engert, M. Schmidt, M. Schulz, J. Schneider-Mergener, U. Reineke. Coherent membrane supports for parallel microsynthesis and screening of bioactive peptides. *Biopolymers*, 55(3):188–206, 2000.
- [104] U. Reineke, R. Volkmer-Engert, J. Schneider-Mergener. Applications of peptide arrays prepared by the spot-technology. *Curr Opin Biotechnol*, 12(1):59–64, 2001.
- [105] R. Frank. The spot-synthesis technique. synthetic peptide arrays on membrane supports—principles and applications. *J Immunol Methods*, 267(1):13–26, 2002.
- [106] R. Frank, H. Overwin. Spot synthesis. epitope analysis with arrays of synthetic peptides prepared on cellulose membranes. *Methods Mol Biol*, 66:149–169, 1996.
- [107] A. Kramer, T. Keitel, K. Winkler, W. Stocklein, W. Hohne, J. Schneider-Mergener. Molecular basis for the binding promiscuity of an anti-p24 (hiv-1) monoclonal antibody. *Cell*, 91(6):799–809, 1997.
- [108] A. Kramer, U. Reineke, L. Dong, B. Hoffmann, U. Hoffmuller, D. Winkler, R. Volkmer-Engert, J. Schneider-Mergener. Spot synthesis: observations and optimizations. *J Pept Res*, 54(4):319–327, 1999.
- [109] C. Landgraf, S. Panni, L. Montecchi-Palazzi, L. Castagnoli, J. Schneider-Mergener, R. Volkmer-Engert, G. Cesareni. Protein interaction networks by proteome peptide scanning. *PLoS Biol*, 2(1):E14, 2004.
- [110] P. Boisguerin, R. Leben, B. Ay, G. Radziwill, K. Moelling, L. Dong, R. Volkmer-Engert. An improved method for the synthesis of cellulose membrane-bound peptides with free c termini is useful for pdz domain binding studies. *Chem Biol*, 11(4):449–459, 2004.
- [111] A. A. Weiser, M. Or-Guil, V. Tapia, A. Leichsenring, J. Schuchhardt, C. Frommel, R. Volkmer-Engert. Spot synthesis: reliability of array-based measurement of peptide binding affinity. *Anal Biochem*, 342(2):300–311, 2005.
- [112] S. Bhargava, K. Licha, T. Knaute, B. Ebert, A. Becker, C. Grotzinger, C. Hessenius, B. Wiedenmann, J. Schneider-Mergener, R. Volkmer-Engert. A complete substitutional analysis of vip for better tumor imaging properties. *J Mol Recognit*, 15(3):145–153, 2002.
- [113] R. B. Merrifield. Solid phase peptide synthesis. i. the synthesis of a tetrapeptide. *J. Am. Chem. Soc.*, 85(14):2149–2154, 1963.
- [114] N. Sewald, H.-D. Jakubke. *Peptides: Chemistry and Biology*. WILEY-VCH Verlag GmbH, Weinheim, 2002.

- [115] R.C. Asherton, E.; Sheppard. *Solid Phase Peptide Synthesis, A Practical Approach.* IRL Press, Oxford, 1989.
- [116] C. Sune, T. Hayashi, Y. Liu, W. S. Lane, R. A. Young, M. A. Garcia-Blanco. Ca150, a nuclear protein associated with the rna polymerase ii holoenzyme, is involved in tat-activated human immunodeficiency virus type 1 transcription. *Mol Cell Biol*, 17(10):6029–6039, 1997.
- [117] K. T. Lin, R. M. Lu, W. Y. Tarn. The ww domain-containing proteins interact with the early spliceosome and participate in pre-mrna splicing in vivo. *Mol Cell Biol*, 24(20):9176–9185, 2004.
- [118] M. Sanchez-Alvarez, A. C. Goldstrohm, M. A. Garcia-Blanco, C. Sune. Human transcription elongation factor ca150 localizes to splicing factor-rich nuclear speckles and assembles transcription and splicing components into complexes through its amino and carboxyl regions. *Mol Cell Biol*, 26(13):4998–5014, 2006.
- [119] L. Otte. *Charakterisierung von WW-Domänen-Ligand-Wechselwirkungen mit Hilfe synthetischer Protein- und Peptidbibliotheken.* Dissertation, Humboldt-Universität, 2004.
- [120] A. C. Goldstrohm, T. R. Albrecht, C. Sune, M. T. Bedford, M. A. Garcia-Blanco. The transcription elongation factor ca150 interacts with rna polymerase ii and the pre-mrna splicing factor sf1. *Mol Cell Biol*, 21(22):7617–7628, 2001.
- [121] B. Chattopadhyay, S. Ghosh, P. K. Gangopadhyay, S. K. Das, T. Roy, K. K. Sinha, D. K. Jha, S. C. Mukherjee, A. Chakraborty, B. S. Singhal, A. K. Bhattacharya, N. P. Bhattacharyya. Modulation of age at onset in huntington's disease and spinocerebellar ataxia type 2 patients originated from eastern india. *Neurosci Lett*, 345(2):93–96, 2003.
- [122] S. Holbert, I. Denghien, T. Kiechle, A. Rosenblatt, C. Wellington, M. R. Hayden, R. L. Margolis, C. A. Ross, J. Dausset, R. J. Ferrante, C. Neri. The gln-ala repeat transcriptional activator ca150 interacts with huntingtin: neuropathologic and genetic evidence for a role in huntington's disease pathogenesis. *Proc Natl Acad Sci U S A*, 98(4):1811–1816, 2001.
- [123] N. Ferguson, J. Becker, H. Tidow, S. Tremmel, T. D. Sharpe, G. Krause, J. Flinders, M. Petrovich, J. Berriman, H. Oschkinat, A. R. Fersht. General structural motifs of amyloid protofilaments. *Proc Natl Acad Sci U S A*, 103(44):16248–16253, 2006.
- [124] A. Y. Hung, M. Sheng. Pdz domains: structural modules for protein complex assembly. *J Biol Chem*, 277(8):5699–5702, 2002.

- [125] F. Bauer, H. Sticht. A proline to glycine mutation in the lck sh3-domain affects conformational sampling and increases ligand binding affinity. *FEBS Lett.*, 2007.
- [126] N. Ferguson, J. Berriman, M. Petrovich, T. D. Sharpe, J. T. Finch, A. R. Fersht. Rapid amyloid fiber formation from the fast-folding ww domain fbp28. *Proc Natl Acad Sci U S A*, 100(17):9814–9819, 2003.
- [127] S. Tremmel, M. Beyermann, H. Oschkinat, M. Bienert, D. Naumann, H. Fabian. (13)c-labeled tyrosine residues as local ir probes for monitoring conformational changes in peptides and proteins. *Angew Chem Int Ed Engl.*, 44(29):4631–4635, 2005.
- [128] R. Doelling, M. Beyermann, J. Haenel, F. Kernchen, E. Krause, P. Franke, M. Brudel, M. Bienert. Piperidine-mediated side product formation for asp(obut)-containing peptides. *J. Chem. Soc., Chem. Commun.*, 7:853 – 854, 1994.
- [129] M. Mutter, A. Nefzi, T. Sato, X. Sun, F. Wahl, T. Wohr. Pseudo-prolines (psi pro) for accessing inaccessible peptides. *Pept Res.*, 8(3):145–153, 1995.
- [130] Stanley C. Gill, Peter H. von Hippel. Calculation of protein extinction coefficients from amino acid sequence data. *Analytical Biochemistry*, 182(2):319–326, 1989.
- [131] C. Krittanai, W. C. Johnson. Correcting the circular dichroism spectra of peptides for contributions of absorbing side chains. *Anal Biochem*, 253(1):57–64, 1997.
- [132] L. Swint, A. D. Robertson. Thermodynamics of unfolding for turkey ovomucoid third domain: thermal and chemical denaturation. *Protein Sci.*, 2(12):2037–2049, 1993.
- [133] N. G. Bandur, K. Harms, U. Koert. First stereoselective synthesis of a tyr-tyr e-alkene isostere. *Synlett*, (1):99–102, 2007.
- [134] H. Wenschuh, H. Gausepohl, L. Germeroth, M. Ulbricht, H. Matuschewski, A. Kramer, R. Volkmer-Engert, N. Heine, T. Ast, D. Scharn, J. Schneider-Mergener. *Combinatorial Chemistry, A Practical Approach*. Oxford University Press, Oxford, 2000.
- [135] M. T. Bedford, D. C. Chan, P. Leder. Fbp ww domains and the abl sh3 domain bind to a specific class of proline-rich ligands. *Embo J.*, 16(9):2376–2383, 1997.
- [136] D. C. Chan, M. T. Bedford, P. Leder. Formin binding proteins bear wwp/ww domains that bind proline-rich peptides and functionally resemble sh3 domains. *Embo J.*, 15(5):1045–1054, 1996.

- [137] M. Sudol, T. Hunter. New wrinkles for an old domain. *Cell*, 103(7):1001–1004, 2000.
- [138] H. Hu, J. Columbus, Y. Zhang, D. Wu, L. Lian, S. Yang, J. Goodwin, C. Luczak, M. Carter, L. Chen, M. James, R. Davis, M. Sudol, J. Rodwell, J. J. Herrero. A map of ww domain family interactions. *Proteomics*, 4(3):643–655, 2004.
- [139] A. A. Kortt, M. Lah, G. W. Oddie, C. L. Gruen, J. E. Burns, L. A. Pearce, J. L. Atwell, A. J. McCoy, G. J. Howlett, D. W. Metzger, R. G. Webster, P. J. Hudson. Single-chain fv fragments of anti-neuraminidase antibody nc10 containing five- and ten-residue linkers form dimers and with zero-residue linker a trimer. *Protein Eng*, 10(4):423–433, 1997.
- [140] H. Nguyen, M. Jager, A. Moretto, M. Gruebele, J. W. Kelly. Tuning the free-energy landscape of a ww domain by temperature, mutation, and truncation. *Proc Natl Acad Sci U S A*, 100(7):3948–3953, 2003.
- [141] J. R. Pires, F. Taha-Nejad, F. Toepert, T. Ast, U. Hoffmuller, J. Schneider-Mergener, R. Kuhne, M. J. Macias, H. Oschkinat. Solution structures of the yap65 ww domain and the variant l30 k in complex with the peptides gtppppytvg, n-(n-octyl)-gpppy and plppy and the application of peptide libraries reveal a minimal binding epitope. *J Mol Biol*, 314(5):1147–1156, 2001.
- [142] U. Wiedemann, P. Boisguerin, R. Leben, D. Leitner, G. Krause, K. Moelling, R. Volkmer-Engert, H. Oschkinat. Quantification of pdz domain specificity, prediction of ligand affinity and rational design of super-binding peptides. *J Mol Biol*, 343(3):703–718, 2004.

An dieser Stelle möchte ich allen danken, die zum Gelingen dieser Arbeit beigetragen haben.

Zuerst möchte ich meiner Familie danken, die mich immer unterstützt hat, besonders in den letzten schwierigen Jahren. Weiterhin gilt mein Dank Fabian, der mich immer selbstlos in allen Belangen begleitet hat.

Mein Dank gilt Prof. H. Oschkinat, für Fragen immer ein offenes Ohr zu gehabt zu haben.

Ich danke herzlich Dr. R. Volkmer für die Möglichkeit der Durchführung dieser Doktorarbeit, für die gute Betreuung, die fachlichen Ratschläge und fruchtbaren Diskussionen.

Besonders herzlich möchte ich mich bei Ines Kretzschmar bedanken, die mich von Anfang an in allen Fragen selbstlos unterstützt und mich in diesem Projekt hilfreich begleitet hat. Auch allen anderen Mitgliedern (auch ehemaligen) der Arbeitsgruppe "Molekulare Bibliotheken" möchte ich für fachliche und moralische Unterstützung danken: Dr. B. Ay, Dr. P. Boisguerin, J. Bongartz, Dr. C. Dröseler, Z. Fidan, A. Hayungs, M. Hovestädt, I. Jäger, C. Landgraf, K. Landgraf, R. Leben, A. Leichsenring, C. Mahrenholz, J. Müller, Dr. L. Otte, Dr. M. Portwich, Dr. A. Schlosser, Dr. R. Stigler, M. Streitz, V. Tapia und A. Weiser.

Ich bedanke mich bei allen Mitgliedern der Forschergruppe 475, besonders bei N. Bandur und Prof. R. Seckler. Weiterhin gilt mein Dank den Mitgliedern des FMP-Berlin, besonders Dr. S. Tremmel, H. Holtmann, H. Nikolenko und Dr. S. Keller.

Zuletzt danke ich allen anderen, die mich in jeder Hinsicht unterstützt haben.