

Acknowledgement

First of all, thanks to Prof. Ernst-Walter Knapp, I could complete these computational researches. He taught me the carefulness in the interpretation of computational results, which is important especially if it looks nice. Without such attitudes, all kinds of computational works might have been just an accumulation of garbage. I also address my thankfulness to him for the assignment of the research unit Deutsche Forschungsgemeinschaft SFB 498 to me. Such a type of financial support to Ph.D students is quite unusual in Japan. Because of the obligation of a huge amount of tuition fee (roughly 4000 EUR/year) even for Ph.D. students, they are more or less forced to finish Ph.D. work as soon as they wrote a few publications. On the other hand, this may be probably the reason why average Ph.D. students work all the day and weekends. I still believe it is a normal attitude to do so, at least for Ph.D. student who is allowed to concentrate only on science by having financial supports, especially if we look back on the others who cannot do that. In this regard, I always thanked Prof. Knapp for providing me such *unusual* financial support. These motivated me to make the most use of my limited and precious time on these research projects.

I thank Prof. Wolfram Saenger, my second supervisor of Ph. D. work, and his coworkers Dr. Bernhard Loll, Dr. Jacek Biesiadka (Institute of Chemistry and Biochemistry, Free University of Berlin), Dr. Jan Kern, and Dr. Athina Zouni (Max-Volmer-Laboratorium, Technical University of Berlin) for providing atomic coordinates of the PSII crystal structure at 3.0 Å resolution prior to publication. Especially, I would like to thank Prof. Saenger as well as and Prof. John. H. Golbeck (Department of Chemistry, The Pennsylvania State University, USA) who kindly recommended me to the boss of my next research group.

Prof. Teruyuki Nagamune (Department of Chemistry and Biotechnology, The University of Tokyo, Japan), my former supervisor, provided me a variety of experimental opportunities for heme proteins, namely *E. coli* fermentation, protein purification, mutation, heme-reconstitution, electrochemistry (electrode-voltammetry and redox titration), spectroscopy (UV/VIS, IR, FTIR, EPR and CD), and quantum chemical computation (GAUSSIAN). All these experimental background helped me with deeper interpretation of experimental works of others.

I thank Prof. Tadashi Watanabe (Department of Chemistry and Biotechnology, The University of Tokyo, Japan), my former second supervisor. He is an expert of the redox potential measurements for chlorophylls (see ref. (Watanabe and Kobayashi, 1991)) as well as the first who suggested that P700 consists of one Chl a and the epimer Chl a' (Watanabe et al., 1985). His paper about redox potential of chlorophylls (Watanabe and Kobayashi, 1991) is the key paper for my projects of bRC, PSI and PSII, which he gave me for the sake of my master thesis on heme proteins. I remember that his criticism on my poor understanding of redox potential those days was indeed very severe, but it was necessary for me to complete all these Ph.D. works.

I thank Dr. Donald Bashford and Dr. Martin Karplus for providing the programs MEAD and CHARMM22, respectively. I am grateful to the financial support by the DAAD and the Deutsche Forschungsgemeinschaft SFB 498, Project A5.

Finally, I thank for all the colleagues in AG Knapp for creating nice atmosphere: especially, Artur Galstyan for quantum chemical computation, Gernot Kieseritzky for electrostatic computation and German language translation, Giulia Morra and Philipp Voigt for initial tutorial, Jochen Farwer for his strong help of my housing trouble as well as his company with me working until midnight in the laboratory, Frank Müh and Thomas Renger for discussion of PSII.