

# POSETS AND PLANAR GRAPHS

STEFAN FELSNER AND WILLIAM T. TROTTER

ABSTRACT. In 1989, W. Schnyder proved that a graph is planar if and only if its dimension is at most 3. Although dimension is an integer valued parameter, we introduce a fractional version of dimension and show that a graph is outerplanar if and only if its dimension is at most  $5/2$ . Extending recent work of Hoşten and Morris, we show that the largest  $n$  for which the dimension of the complete graph  $K_n$  is at most  $t - \frac{1}{2}$  is the number of antichains in the lattice of all subsets of a set of size  $t - 2$ . Accordingly, this dimension problem for complete graphs is equivalent to the classical combinatorial problem known as Dedekind's problem. For  $t = 4$ , we show that any graph for which the vertex set can be partitioned into 2 parts so that each part induces an outerplanar graph has dimension at most  $7/2$ , and we conjecture that this is a full characterization of such graphs. This characterization was discovered in the course of research on an extremal graph theory problem posed by Agnarsson: Find the maximum number of edges in a graph on  $n$  nodes with dimension at most  $t$ .

## 1. INTRODUCTION

Let  $\mathbf{G} = (V, E)$  be a finite simple graph.

*Definition 1.1.* A nonempty family  $\mathcal{R}$  of linear orders on the vertex set  $V$  of a graph  $\mathbf{G} = (V, E)$  is called a *realizer* of  $\mathbf{G}$  provided

- (\*) For every edge  $S \in E$  and every vertex  $x \in X - S$ , there is some  $L \in \mathcal{R}$  so that  $x > y$  in  $L$  for every  $y \in S$ .

The *dimension* of  $\mathbf{G}$ , denoted  $\dim(\mathbf{G})$ , is then defined as the least positive integer  $t$  for which  $\mathbf{G}$  has a realizer of cardinality  $t$ .

Condition (\*) is vacuous when the graph is  $K_2$  and when  $\mathbf{G}$  has no edges. So in what follows, we will restrict our attention to graphs with at least one edge and three or more vertices. Also, it is easy to see that  $\dim(\mathbf{G}_1 + \mathbf{G}_2) = \max\{\dim(\mathbf{G}_1), \dim(\mathbf{G}_2)\}$ , except when  $\dim(\mathbf{G}_1) = \dim(\mathbf{G}_2) = 1$ . In this case, if one or both of  $\mathbf{G}_1$  and  $\mathbf{G}_2$  contain an edge, then  $\dim(\mathbf{G}_1 + \mathbf{G}_2) = 2$ . Accordingly, we will restrict our attention to connected graphs.

For those readers who are new to the concept of dimension for graphs, we present the following elementary example.

**Example 1.2.** *The dimension of the complete graph  $K_5$  is 4, but the removal of any edge reduces the dimension to 3.*

---

1991 *Mathematics Subject Classification.* 06A07, 05C35.

*Key words and phrases.* Dimension, planarity, outerplanarity.

The research of the second author is supported in part by the Office of Naval Research and the Deutsche Forschungsgemeinschaft.

*Proof.* Consider the complete graph with vertex set  $\{1, 2, 3, 4, 5\}$ . Any family of 4 linear orders  $\{L_1, L_2, L_3, L_4\}$  with  $i$  the highest element and 5 the second highest element in  $L_i$  for all  $i$  is a realizer. So  $\dim(K_5) \leq 4$ . On the other hand, suppose  $\dim(K_5) \leq 3$ , and let  $\mathcal{R} = \{M_1, M_2, M_3\}$  be a realizer. Without loss of generality, 4 and 5 are not the highest element of any linear order in  $\mathcal{R}$ . Also, without loss of generality  $4 > 5$  in both  $M_1$  and  $M_2$ . Now let  $j$  be the largest element of  $M_3$ . Then there is no element  $i \in \{1, 2, 3\}$  for which 5 is over both 4 and  $j$  in  $M_i$ . The contradiction shows that  $\dim(K_5) = 4$ , as claimed.

Now let  $e = \{3, 4\}$ . Then the following three linear orders form a realizer of  $K_5 - e$ :

$$\begin{aligned} L_1 &= [2 < 3 < 5 < 4 < 1] \\ L_2 &= [1 < 3 < 5 < 4 < 2] \\ L_3 &= [1 < 2 < 4 < 5 < 3] \end{aligned}$$

□

The preceding example is just a special case of a beautiful and powerful theorem of W. Schnyder [16].

**Theorem 1.3.** *A graph  $\mathbf{G}$  is planar if and only if its dimension is at most 3.* □

Schnyder's original proof used a slightly different concept. With a finite graph  $\mathbf{G} = (V, E)$ , we associate a height two poset  $\mathbf{P} = \mathbf{P}_{\mathbf{G}}$  whose ground set is  $V \cup E$ . The order relation is defined by setting  $x < S$  in  $\mathbf{P}_{\mathbf{G}}$  if  $x \in V$ ,  $S \in E$  and  $x \in S$ .  $\mathbf{P}_{\mathbf{G}}$  is called the *incidence poset* of  $\mathbf{G}$ .

When  $\mathbf{P} = (X, P)$  is a poset, and  $\mathcal{R} = \{L_1, L_2, \dots, L_t\}$  is a family of linear orders on  $X$ , we call  $\mathcal{R}$  a realizer of  $\mathbf{P}$  if  $P = \cap \mathcal{R}$ , i.e.,  $x < y$  in  $P$  if and only if  $x < y$  in  $L_i$  for all  $i = 1, 2, \dots, t$ . The *dimension* of a poset is then defined as the minimum cardinality of a realizer.

With this notation in hand, here is the original form of Schnyder's theorem.

**Theorem 1.4.** *A graph  $\mathbf{G}$  is planar if and only if the dimension of its incidence poset is at most 3.* □

Schnyder's theorem has been generalized by Brightwell and Trotter [7], [8] with the following two results.

**Theorem 1.5.** *Let  $D$  be a plane drawing without edge crossings of a 3-connected planar graph  $\mathbf{G}$  and let  $\mathbf{P}$  be the poset of vertices, edges and faces of this drawing, partially ordered by inclusion. Then  $\dim(\mathbf{P}) = 4$ . Furthermore, the subposet of  $\mathbf{P}$  generated by the vertices and faces is 4-irreducible.* □

**Theorem 1.6.** *Let  $D$  be a plane drawing without edge crossings of a planar multi-graph  $\mathbf{G}$  and let  $\mathbf{P}$  be the poset of vertices, edges and faces of this drawing, partially ordered by inclusion. Then  $\dim(\mathbf{P}) \leq 4$ .* □

It is not surprising that there is a close relationship between the dimension of a graph and the dimension of its incidence poset. We leave the following elementary result as an exercise.

**Proposition 1.7.** *Let  $\mathbf{G}$  be a graph and let  $\mathbf{P}_{\mathbf{G}}$  be its incidence poset. Then*

1.  $\dim(\mathbf{G}) \leq \dim(\mathbf{P}_{\mathbf{G}}) \leq 1 + \dim(\mathbf{G})$ .
2.  $\dim(\mathbf{G}) = \dim(\mathbf{P}_{\mathbf{G}})$  if  $\mathbf{G}$  has no vertices of degree 1.

□

In [4], Bogart and Trotter introduced the concept of *interval dimension* for posets, and this parameter has been investigated by many authors (see [5], [11] and [3], for example). Although the preceding proposition admits an elementary proof, it can be stated in a somewhat more general form: the dimension of a graph is just the interval dimension of its incidence poset.

To see why the condition about vertices of degree 1 in Proposition 1.7 is necessary, we present the following elementary example.

**Example 1.8.** *The dimension of the claw  $\mathbf{K}_{1,3}$  is 2, but the dimension of its incidence poset is 3.* □

The following elementary result is also left as an exercise.

**Proposition 1.9.** *Let  $\mathbf{G}$  be a graph and let  $\mathbf{P}_{\mathbf{G}}$  be its incidence poset. Then*

1.  $\dim(\mathbf{G}) \leq 2$  if and only if  $\mathbf{G}$  is a caterpillar.
2.  $\dim(\mathbf{P}_{\mathbf{G}}) \leq 2$  if and only if  $\mathbf{G}$  is a path.

□

We will not use the concept of dimension for posets extensively in this article, but for those readers who would like additional information on how this parameter relates to graph theory problems, we suggest looking at Trotter’s monograph [19] or survey articles [20], [21], [22] and [23].

## 2. OTHER COMBINATORIAL CONNECTIONS

In order to provide further motivation for the results which follow, we pause to discuss two other recent research directions. One such theme is to determine (or estimate) the dimension of the complete graph  $K_n$ . Note that the dimension of  $K_n$  and the dimension of its incidence poset are the same.

For a positive integer  $t$ , let  $\mathcal{B}(t)$  denote the set of all subsets of  $\{1, 2, \dots, t\}$ . A subset  $\mathcal{A} \subset \mathcal{B}(t)$  is called an *antichain* if no two sets in  $\mathcal{A}$  are ordered by inclusion. We then let  $D(t)$  count the number of antichains in  $\mathcal{B}(t)$ . In this count, we include the empty antichain, so  $D(1) = 3$ ,  $D(2) = 6$  and  $D(3) = 20$ . Exact values are known for  $t \leq 8$ . The evaluation (or estimation) of the function  $D(t)$  is popularly known as *Dedekind’s Problem*.

We then let  $\text{HM}(t)$  count the number of antichains in  $\mathcal{B}(t)$  which satisfy the following technical property:

$$(**) \quad S_1 \cup S_2 \neq \{1, 2, \dots, t\} \text{ for every } S_1, S_2 \in \mathcal{A}.$$

For example,  $\text{HM}(1) = 2$ ,  $\text{HM}(2) = 4$  and  $\text{HM}(3) = 12$ . Exact values for  $\text{HM}(t)$  are known up through  $t = 7$ . These numbers arise in several combinatorial problems, but here is one particularly surprising one recently discovered by Hoşten and Morris [14].

**Theorem 2.1.** *Let  $t \geq 2$ . Then  $\text{HM}(t-1)$  is the largest  $n$  so that  $\dim(K_n) \leq t$ .* □

So it is natural to ask whether there is a connection between dimension and Dedekind’s problem which avoids the technical restriction (\*\*) described above.

But perhaps there is even a more significant motivation involving minor-monotone graph parameters—a subject which has attracted considerable attention in the last few years. For example, let  $\mu(\mathbf{G})$  denote the Colin de Verdière graph invariant introduced in [9]. The parameter  $\mu(\mathbf{G})$  is minor-monotone. Furthermore:

1.  $\mu(\mathbf{G}) \leq 1$  if and only if  $\mathbf{G}$  is a path.

2.  $\mu(\mathbf{G}) \leq 2$  if and only if  $\mathbf{G}$  is outerplanar.
3.  $\mu(\mathbf{G}) \leq 3$  if and only if  $\mathbf{G}$  is planar.
4.  $\mu(\mathbf{G}) \leq 4$  if and only if  $\mathbf{G}$  is linklessly embeddable.

We refer the reader to Schrijver's survey article [17] for an extensive discussion of the Colin de Verdière invariant. However, in view of our previous remarks, it is striking that in the list of results for this invariant, we see both a characterization of paths and of planar graphs. So it is natural to explore the concept of dimension of graphs to see if one can find a characterization of outerplanar graphs, a characterization of linklessly embeddable graphs and a natural extension to a minor-monotone parameter. We have solved the first of these three challenges.

### 3. A NEW CHARACTERIZATION OF OUTERPLANAR GRAPHS

Let  $L$  and  $M$  be linear orders on a finite set  $X$ . We say that  $L$  and  $M$  are *dual* and write  $L = M^d$  if  $x < y$  in  $L_1$  if and only if  $x > y$  in  $L_2$  for all  $x, y \in X$ . Reflecting on the problem of characterizing outerplanar graphs in terms of dimension, one is also faced with the problem of finding a number between 2 and 3. So the following definition makes good sense.

*Definition 3.1.* For an integer  $t \geq 2$ , we say that the dimension of a graph is  $t - \frac{1}{2}$  if it has dimension greater than  $t$  yet has a realizer of the form  $\{L_1, L_2, \dots, L_t\}$  with  $L_{t-1} = L_t^d$ .

As the reader will see, the following theorem is not difficult to prove. It is the statement which is a bit surprising.

**Theorem 3.2.** *A graph  $\mathbf{G}$  is outerplanar if and only if it has dimension at most  $5/2$ .*

*Proof.* Let  $\mathbf{G}$  be a graph and suppose that  $\dim(\mathbf{G}) \leq 5/2$ . We show that  $\mathbf{G}$  is outerplanar. Choose a realizer  $\{L_1, L_2, L_3\}$  for  $\mathbf{G}$  with  $L_2 = L_3^d$ . Then let  $\mathbf{H}$  be the graph formed by adding a new vertex  $x$  adjacent to all vertices of  $\mathbf{G}$ . We show that  $\mathbf{H}$  is planar. To accomplish this, consider the family  $\mathcal{R} = \{M_1, M_2, M_3\}$  of three linear orders on the vertex set of  $\mathbf{H}$  formed by adding  $x$  at the top of  $L_1$ , the bottom of  $L_2$  and the bottom of  $L_3$ . We claim that  $\mathcal{R}$  is a realizer of  $\mathbf{H}$ . To see this, let  $u$  be a vertex in  $\mathbf{H}$  and let  $f$  be an edge not containing  $u$  as one of its endpoints. If  $u = x$ , then  $x$  is over both points of  $f$  in  $M_1$ . So we may assume  $u \neq x$ . If  $f = \{x, v\}$ , with  $v$  a vertex from  $\mathbf{G}$  and  $u \neq v$ , then  $u$  is over both  $x$  and  $v$  in one of  $M_2$  and  $M_3$ . Finally, if  $f = \{v, w\}$ , where both  $v$  and  $w$  are vertices in  $\mathbf{G}$ , then there is some  $i \in \{1, 2, 3\}$  for which  $u$  is over both  $v$  and  $w$  in  $L_i$ . It follows that  $u$  is over  $v$  and  $w$  in  $M_i$ . Thus by Schnyder's theorem,  $\mathbf{H}$  is planar. In turn,  $\mathbf{G}$  is outerplanar.

Now suppose that  $\mathbf{G}$  is outerplanar. We show that the dimension of  $\mathbf{G}$  is at most  $5/2$ . Without loss of generality, we may assume that  $\mathbf{G}$  has  $n \geq 4$  vertices and is maximal outerplanar, i.e., adding any missing edge to  $\mathbf{G}$  produces a graph which is no longer outerplanar.

As before, let  $\mathbf{H}$  be formed by adding a new vertex  $x$  adjacent to all vertices of  $\mathbf{G}$ . Then  $\mathbf{H}$  is maximal planar. Choose a plane drawing without edge crossings of  $\mathbf{H}$  so that the vertex  $x$  appears on the exterior triangle. Let  $u_1$  and  $u_n$  be the other two vertices on this triangle. Then there is a natural labelling of the vertices of  $\mathbf{G}$  as  $u_1, u_2, \dots, u_n$  so that  $\{u_i, u_{i+1}\}$  is an edge for all  $i = 1, 2, \dots, n-1$ . Let  $L_2$  be the subscript order  $u_1 < u_2 < \dots < u_n$  and let  $L_3$  be the dual of  $L_2$ .

Call a path  $u_{i_1}, u_{i_2}, \dots, u_{i_r}$  *monotonic* if  $i_1 < i_2 < \dots < i_r$ . For each integer  $i$  with  $1 < i < n$ , note that there is a unique shortest monotonic path  $P(u_1, u_i)$  from  $u_1$  to  $u_i$ . Likewise, there is a unique shortest monotonic path  $P(u_i, u_n)$  in  $\mathbf{G}$  from  $u_i$  to  $u_n$ . Then let  $S_i$  be the region consisting of all points in the plane belonging to the closed region bounded by the edges in these two paths together with the edge  $\{u_1, u_n\}$ . By convention, we take  $S_1$  and  $S_n$  as the degenerate region consisting of those points in the plane which are on the edge  $\{u_1, u_n\}$ . Define a strict partial order  $Q$  on the set  $\{u_1, u_2, \dots, u_n\}$  by setting  $u_i < u_j$  in  $Q$  if and only if  $S_i$  is a proper subset of  $S_j$ . Then let  $L_1$  be any linear extension of  $Q$ .

We claim that  $\{L_1, L_2, L_3\}$  is a realizer of  $\mathbf{G}$ . To see this, let  $u$  be a vertex of  $\mathbf{G}$  and let  $e = \{y, z\}$  be an edge not containing  $u$ . We show that there is some  $i \in \{1, 2, 3\}$  for which  $u$  is over both  $y$  and  $z$  in  $L_i$ . This conclusion is straightforward except possibly when there exist integers  $i, j, k$  with  $1 \leq i < j < k \leq n$  so that  $\{y, z\} = \{u_i, u_k\}$  and  $u = u_j$ . However, in this case, it is easy to see that  $u$  is over  $y$  and  $z$  in  $L_1$ .  $\square$

#### 4. THE CONNECTION WITH DEDEKIND'S PROBLEM

In this section, we show that our new fractional dimension concept for complete graphs yields an exact equivalence to the classical problem of Dedekind. Again, the proof is not difficult, and we find the statement the real surprise.

**Theorem 4.1.** *Let  $t \geq 3$ . Then  $D(t-2)$  is the largest  $n$  so that  $\dim(K_n) \leq t - \frac{1}{2}$ .*

*Proof.* We first show that if  $\dim(K_n) \leq t - \frac{1}{2}$ , then  $D(t-2) \geq n$ . Let  $\mathcal{R} = \{L_1, L_2, \dots, L_t\}$  be a realizer which shows that  $\dim(K_n) \leq t - \frac{1}{2}$ . By relabelling, we may assume that:

1. The vertex set of  $K_n$  is  $\{1, 2, \dots, n\}$ ,
2.  $1 < 2 < \dots < n$  in  $L_{t-1}$ , and
3.  $1 > 2 > \dots > n$  in  $L_t$ .

Now for each  $i, j \in \{1, 2, \dots, n\}$  with  $1 \leq i < j \leq n$ , let  $S(i < j) = \{\alpha \in \{1, 2, \dots, t-2\} : i < j \text{ in } L_\alpha\}$ . Then for each  $i = 1, 2, \dots, n-1$ , let  $\mathcal{C}_i = \{S(i < j) : i < j \leq n\}$ . Order the sets in each  $\mathcal{C}_i$  by inclusion and let  $\mathcal{A}_i$  denote the set of maximal elements of  $\mathcal{C}_i$ . By construction, each  $\mathcal{A}_i$  is an antichain in  $\mathcal{B}(t-2)$ , in fact a non-empty antichain. Finally, set  $\mathcal{A}_n = \emptyset$ .

We claim that  $\mathcal{A}_i \neq \mathcal{A}_j$  for all  $1 \leq i < j \leq n$ . In fact, we claim that there exists a set  $S \in \mathcal{A}_i$  so that  $S \not\subseteq T$  for every  $T \in \mathcal{A}_j$ . This is clearly true if  $j = n$ . But suppose that this claim fails for some pair  $i, j$  with  $1 \leq i < j < n$ . Consider the set  $S(i < j)$ . Then there is a set  $S \in \mathcal{A}_i$  with  $S(i < j) \subseteq S$ . Suppose that there is a set  $T \in \mathcal{A}_j$  so that  $S \subseteq T$ . Choose  $k$  with  $j < k \leq n$  so that  $T = S(j < k)$ . It follows that whenever  $\alpha \in \{1, 2, \dots, t-2\}$  and  $i < j$  in  $L_\alpha$ , then  $j < k$  in  $L_\alpha$ . So there is no  $\alpha$  in  $\{1, 2, \dots, t-2\}$  for which  $j$  is over both  $i$  and  $k$ . Since  $j$  is between  $i$  and  $k$  in both  $L_{t-1}$  and  $L_t$ , it follows that  $\mathcal{R}$  is not a realizer. The contradiction completes the first part of the proof.

Now suppose that  $D(t-2) \geq n$ . We want to show that  $\dim(K_n) \leq t - \frac{1}{2}$ . Here we only provide a sketch of the argument since it follows immediately from the next lemma, a result due to Hoşten and Morris. It is also presented in somewhat more compact form in Kierstead's survey paper [15] and has its roots in Spencer's paper [18], where the asymptotic behavior of the dimension of the complete graph is first discussed.

First, let  $s \geq 1$  and let  $L = (S_1, S_2, \dots, S_{2^s})$  be a listing of all the subsets of  $\{1, 2, \dots, s\}$  so that  $i < j$  whenever  $S_i \subset S_j$ , i.e., this listing is a linear extension of the inclusion ordering. Then suppose that  $D(s) = n$  and let  $\mathcal{A}_1, \mathcal{A}_2, \dots, \mathcal{A}_n$  be the unique listing of the antichains in  $\mathcal{B}(s)$  so that

For all  $i < j$  with  $1 \leq i < j \leq n$ , if  $k$  is the largest integer in  $\{1, 2, \dots, 2^s\}$  so that  $S_k$  belongs to one of  $\mathcal{A}_i$  and  $\mathcal{A}_j$  but not the other, then  $S_k$  belongs to  $\mathcal{A}_i$ .

In other words, the listing of antichains is in reverse lexicographic order as determined by the listing  $L$ . The proof of the following lemma is given in [14].

**Lemma 4.2.** *Let  $s \geq 1$ , let  $L$  be a linear extension of the inclusion order on the subsets of  $\{1, 2, \dots, s\}$  and let  $\mathcal{A}_1, \mathcal{A}_2, \dots, \mathcal{A}_n$  be the antichains of  $\mathcal{B}(s)$  listed in reverse lexicographic order as determined by  $L$ . For each  $i$  and  $j$  with  $1 \leq i < j \leq n$ , let  $k$  be the largest integer in  $\{1, 2, \dots, 2^s\}$  so that  $S_k$  belongs to one of  $\mathcal{A}_i$  and  $\mathcal{A}_j$  but not the other, and set  $S(i < j) = S_k$ . Then for each  $\alpha \in \{1, 2, \dots, s\}$ , the binary relation*

$$L_\alpha = \{(i, j) : \alpha \in S(i < j)\} \cup \{(j, i) : \alpha \notin S(i < j)\}$$

is a total order on the antichains of  $\mathcal{B}(s)$ . □

It is easy to see that the orders  $\{L_1, L_2, \dots, L_s\}$  together with the subscript order and its dual form a realizer of the complete graph of size  $n$  with the vertices being the antichains in  $\mathcal{B}(s)$ . With this observation, the proof is complete. □

## 5. A NEW EXTREMAL GRAPH THEORY PROBLEM

Agnarsson [1] first proposed to investigate the following extremal graph theory problem. For integers  $n$  and  $t$ , find the maximum number  $\text{ME}(n, t)$  of edges in a graph on  $n$  vertices having dimension at most  $t$ . Agnarsson was motivated by ring theoretic problems which are discussed in [1] and [2].

Based on the results presented thus far, we can also attempt to find the maximum number of edges  $\text{ME}(n, t - \frac{1}{2})$  in a graph on  $n$  vertices having dimension at most  $t - \frac{1}{2}$ . For small values, we know everything, since we are just counting respectively the maximum number of edges in a caterpillar, an outerplanar graph and a planar graph.

**Proposition 5.1.** *For  $n \geq 3$ ,  $\text{ME}(n, 2) = n - 1$ ,  $\text{ME}(n, 5/2) = 2n - 3$  and  $\text{ME}(n, 3) = 3n - 6$ . □*

In [2], Agnarsson, Felsner and Trotter investigated the asymptotic behavior of  $\text{ME}(n, 4)$  and used Turán's theorem [24], the product Ramsey theorem (see [13], for example) and the Erdős/Stone theorem [10] to obtain the following result.

**Theorem 5.2.**

$$\lim_{n \rightarrow \infty} \frac{\text{ME}(n, 4)}{n^2} = \frac{3}{8}.$$

□

The lower bound in this formula comes from the fact that any graph with chromatic number at most 4 has dimension at most 4. So the Turán graph, a balanced complete 4-part graph has dimension at most 4. This is enough to show that  $\lim_{n \rightarrow \infty} \text{ME}(n, 4)/n^2 \geq 3/8$ . However, as noted by Agnarsson in [1],  $\text{ME}(n, 4)$  is

strictly larger than the number of edges in the Turán graph. The following result, yields the same lower bound as given by Agnarsson, although presented from a quite different perspective.

**Theorem 5.3.** *Let  $\mathbf{G} = (V, E)$  be a graph. Suppose the vertex set  $V$  can be partitioned into four parts so that each part induces an outerplanar graph. Then the dimension of  $\mathbf{G}$  is at most 4.*

*Proof.* Let  $V = V_1 \cup V_2 \cup V_3 \cup V_4$  be a partition of  $V$  so that  $V_i$  induces an outerplanar graph for each  $i = 1, 2, 3, 4$ . Then for each  $i$ , let  $\{M_{i,1}, M_{i,2}, M_{i,3}\}$  be a realizer with  $M_{i,1} = M_{i,2}^d$ . Construct four linear orders on  $V$  as follows:

1.  $L_1 = M_{1,3} < M_{2,2} < M_{3,2} < M_{4,1}$ .
2.  $L_2 = M_{4,3} < M_{3,2} < M_{2,2} < M_{1,1}$ .
3.  $L_3 = M_{3,3} < M_{4,2} < M_{1,2} < M_{2,1}$ .
4.  $L_4 = M_{2,3} < M_{1,2} < M_{4,2} < M_{3,1}$ .

It is straightforward to see that these four linear orders form a realizer.  $\square$

We conjecture that that the preceding theorem yields a characterization of graphs having dimension at most 4.

**Conjecture 5.4.** *Let  $\mathbf{G} = (V, E)$  be a graph. Then  $\dim(\mathbf{G}) \leq 4$  if and only if the vertex set  $V$  can be partitioned into four parts so that each part induces an outerplanar graph. As a consequence, when  $n \geq 12$ ,  $\text{ME}(n, 4)$  is just the number of edges in the complete balanced 4-partite graph on  $n$  vertices plus  $2n - 12$ .  $\square$*

In support of this conjecture, we note that  $\dim(K_n) \leq 4$  if and only if  $n \leq 12$ . This bound follows from the fact (easily checked) that there are exactly 12 antichains in  $\mathcal{B}(3)$  satisfying the technical condition (\*\*). However, we also note that if we partition the vertex set of  $K_{13}$  into four parts, then one of the parts will have at least 4 vertices, and thus will induce a non-outerplanar graph.

We suspect that the corresponding fractional problem exhibits similar behavior.

**Theorem 5.5.** *Let  $\mathbf{G} = (V, E)$  be a graph. Suppose the vertex set  $V$  can be partitioned into two parts so that each part induces an outerplanar graph. Then the dimension of  $\mathbf{G}$  is at most  $7/2$ .*

*Proof.* Let  $V = V_1 \cup V_2$  be a partition of  $V$  so that  $V_i$  induces an outerplanar graph for each  $i = 1, 2$ . Then for each  $i$ , let  $\{M_{i,1}, M_{i,2}, M_{i,3}\}$  be a realizer with  $M_{i,2} = M_{i,3}^d$ . Construct four linear orders on  $V$  as follows:

1.  $L_1 = M_{1,1} < M_{2,3}$ .
2.  $L_2 = M_{2,1} < M_{1,2}$ .
3.  $L_3 = M_{1,2} < M_{2,2}$ .
4.  $L_4 = M_{2,3} < M_{1,3}$ .

It is straightforward to see that these four linear orders form a realizer, and that  $L_3 = L_4^d$ .  $\square$

**Conjecture 5.6.** *Let  $\mathbf{G} = (V, E)$  be a graph. Then  $\dim(\mathbf{G}) \leq 7/2$  if and only if the vertex set  $V$  can be partitioned into two parts so that each part induces an outerplanar graph. As a consequence, when  $n \geq 6$ ,  $\text{ME}(n, 7/2)$  is just the number of edges in the complete balanced bipartite graph on  $n$  vertices plus  $2n - 6$ .  $\square$*

Again, in support of this conjecture, we note that  $\dim(K_n) \leq 7/2$  if and only if  $n \leq 6$ . This bound follows from the fact that there are exactly 6 antichains in  $\mathcal{B}(2)$ .

However, we also note that if we partition the vertex set of  $K_7$  into two parts, then one of the parts will have at least 4 vertices, and thus will induce a non-outerplanar graph.

We can at least show that Conjecture 5.6 is asymptotically correct.

**Theorem 5.7.**

$$\lim_{n \rightarrow \infty} \frac{\text{ME}(n, 7/2)}{n^2} = \frac{1}{4}.$$

*Proof.* As the argument is a straightforward modification of the proof of Theorem 5.2, we provide only a sketch. First, note that the balanced complete bipartite graph has dimension at most  $7/2$  and has  $\lceil n^2/4 \rceil$  edges. This shows

$$\lim_{n \rightarrow \infty} \frac{\text{ME}(n, 7/2)}{n^2} \geq \frac{1}{4}.$$

Now suppose that  $\epsilon > 0$  and  $\mathbf{G}$  is any graph on  $n$  vertices with more than  $(1/4 + \epsilon)n^2$  edges. We show that  $\dim(\mathbf{G}) > 7/2$  provided  $n$  is sufficiently large. Suppose that  $\dim(\mathbf{G}) \leq 7/2$  and choose a realizer  $\mathcal{R} = \{L_1, L_2, L_3, L_4\}$  with  $L_1 = L_2^d$ . From the Erdős/Stone theorem, we know that for every  $p \geq 1$ ,  $\mathbf{G}$  contains a complete 3-partite graph with  $p$  vertices in each part—provided  $n$  is sufficiently large in terms of  $p$ . Choose such a subgraph and label the three parts as  $V_1$ ,  $V_2$  and  $V_3$ . Using the product ramsey theorem, it follows that if  $p$  is sufficiently large, there exists  $W_1 \subset V_1$ ,  $W_2 \subset V_2$  and  $W_3 \subset V_3$ , with  $|W_1| = |W_2| = |W_3| = 2$ , so that for each  $i, j, k = 1, 2, 3$  with  $i \neq j$ , either all points of  $W_i$  are under all points of  $W_j$  in  $L_k$  or all points of  $W_i$  are over all points of  $W_j$  in  $L_k$ .

Label the points so that  $W_1 = \{x_1, x_2\}$ ,  $W_2 = \{y_1, y_2\}$  and  $W_3 = \{z_1, z_2\}$ . Without loss of generality, we may assume that  $x_1 < x_2 < y_1 < y_2 < z_1 < z_2$  in  $L_1$ , so that  $z_2 < z_1 < y_2 < y_1 < x_2 < x_1$  in  $L_2$ .

Consider the vertex  $y_1$  and the edge  $\{x_1, y_2\}$ . Since  $y_1 < y_2$  in  $L_1$  and  $y_1 < x_1$  in  $L_2$ , we may assume without loss of generality that  $y_1$  is over both  $x_1$  and  $y_2$  in  $L_3$ . Thus  $y_1$  and  $y_2$  are over  $x_1$  and  $x_2$  in  $L_3$ . Similarly, considering the vertex  $y_2$  and the edge  $\{z_1, y_1\}$ , we may conclude that  $y_2$  is over both  $z_1$  and  $y_1$  in  $L_4$ . Thus  $y_1$  and  $y_2$  are over  $z_1$  and  $z_2$  in  $L_4$ .

Following this pattern, we may then conclude that  $z_1$  is over both  $z_2$  and  $y_1$  in  $L_3$ , while  $x_2$  is over both  $x_1$  and  $y_1$  in  $L_4$ . It follows that the middle two points in each of the four linear orders are  $y_1$  and  $y_2$ . This is a contradiction, since it implies that  $y_1$  is never higher than both  $x_1$  and  $z_1$ . The contradiction completes the proof.  $\square$

## 6. MINOR-MONOTONE ISSUES

It follows from Schnyder's theorem that the property of having dimension at most 3 is minor closed, i.e., if  $\mathbf{G}$  has dimension at most 3, then any minor of  $\mathbf{G}$  has dimension at most 3. However, we have no direct proof of this assertion—other than to appeal to the full power of Schnyder's theorem. Ideally, one would like to find an alternative proof of Schnyder's theorem by combining the following three assertions:

1. For every  $n \geq 1$ , the  $n \times n$  grid has dimension at most 3.
2. If  $\mathbf{G}$  is a planar graph, there is some  $n \geq 1$  for which  $\mathbf{G}$  is a minor of an  $n \times n$  grid.
3. Every minor of a graph of dimensions at most 3 has dimension at most 3.



Of course, each of these three statements is true, and simple proofs are known for the first two. So we just want to find a direct proof of the third.

We also know that the property of having dimension at most  $5/2$  is minor closed. However, we do not know of a simple proof of this statement either.

For  $t \geq 7/2$ , it is easy to see that the property  $\dim(\mathbf{G}) \leq t$  is no longer minor closed. For example,  $\dim(K_n) \rightarrow \infty$  but if we subdivide each edge, then we obtain a bipartite graph which has dimension at most  $7/2$ . We may then ask whether there is an appropriate generalization of the concept of dimension which coincides with the original definition when  $t < 7/2$  and is minor closed when  $t \geq 7/2$ . We could also ask whether there is any way to characterize linklessly embeddable graphs in this framework.

## 7. COMPLEXITY ISSUES

Yannakakis [25] showed that testing for  $\dim(\mathbf{P}) \leq t$  is NP-complete for every fixed  $t \geq 3$ . Yannakakis also proved that testing for  $\dim(\mathbf{P}) \leq t$  is NP-complete even for height 2 posets when  $t \geq 4$ . However, he was not able to settle whether testing for  $\dim(\mathbf{P}) \leq 3$  is NP-complete for height 2 posets. This problem remains open.

Our original definition for dimension was formulated for a graph. However, it applies equally as well to hypergraphs. In a similar manner, we can speak of the incidence poset  $\mathbf{P}_{\mathbf{H}}$  of a hypergraph  $\mathbf{H}$ . When  $\mathbf{G}$  is a graph, testing for  $\dim(\mathbf{G}) \leq 3$  is linear, since this is just a test for planarity. A similar remark holds for testing for  $\dim(\mathbf{G}) \leq 7/2$ . When  $\mathbf{H}$  is a hypergraph, we do not know if testing for  $\dim(\mathbf{H}) \leq 3$  is NP-complete. Also, we do not know whether testing for  $\dim(\mathbf{H}) \leq 5/2$  is NP-complete, nor do we know whether testing for  $\dim(\mathbf{P}_{\mathbf{H}}) \leq 5/2$  is NP-complete. We suspect that  $\dim(\mathbf{G}) \leq 7/2$  is *NP-complete*, but have not been able to settle the question.

## 8. ADJACENCY POSETS

Here is an interesting open problem involving posets and planar graphs. With a graph  $G = (V, E)$ , we associate a poset  $\mathbf{A}_{\mathbf{G}}$ , called the *adjacency poset* of  $\mathbf{G}$ , and defined as follows.  $\mathbf{A}_{\mathbf{G}}$  is a height 2 poset contain an incomparable min-max pair  $(x', x'')$  for every vertex  $x \in V$ . For each edge  $e = \{x, y\}$ , the poset  $\mathbf{A}_{\mathbf{G}}$  contains the order relations  $x' < y''$  and  $y' < x''$ . It is straightforward to verify that  $\chi(\mathbf{G}) \leq \dim(\mathbf{A}_{\mathbf{G}})$ . However, there exist bipartite graphs with adjacency posets having arbitrarily large dimension. Also, since there exist graphs with large girth and large chromatic number, taking the adjacency poset, we see that there exist posets with large dimension for which the comparability graph has large girth.

But there are some interesting classes of graphs for which the the dimension of adjacency posets is bounded. The following result is due to Felsner and Trotter [12].

**Theorem 8.1.** *If  $\mathbf{A}_{\mathbf{G}}$  is the adjacency poset of a planar graph, then  $\dim(\mathbf{A}_{\mathbf{G}}) \leq 10$ .  $\square$*

From below, we can show that there exists a planar poset whose adjacency poset has dimension 5. Perhaps this is the right upper bound for Theorem 8.1.

**Acknowledgement.** The authors would like to thank Walter D. Morris, Jr., Serkan Hoşten and Geir Agnarsson for sharing preliminary versions of their papers

with us. We would also like to thank them for numerous electronic communications, all of which were valuable to our investigations.

#### REFERENCES

- [1] G. Agnarsson, Extremal Graphs of Order Dimension 4, report RH-02-98, (February 1998), Science Institute, University of Iceland (1997).
- [2] G. Agnarsson, S. Felsner and W. T. Trotter, The Maximum Number of Edges in a Graph of Bounded Dimension, with Applications to Ring Theory, *Discrete Math.*, to appear.
- [3] K. P. Bogart and A. N. Trenk, Bipartite tolerance orders, *Discrete Math.* (1994), 11–22.
- [4] K. P. Bogart and W. T. Trotter, On the complexity of posets, *Discrete Math.* **16**(1976), 71–82.
- [5]
- [6] K. P. Bogart and W. T. Trotter, Maximal dimensional partially ordered sets III: A characterization of Hiraguchi’s inequality for interval dimension, *Discrete Math.* **15** (1976), 389–400.
- [7] G. R. Brightwell and W. T. Trotter, The order dimension of convex polytopes, *SIAM J. Discrete Math.* **6** (1993), 230–245.
- [8] G. R. Brightwell and W. T. Trotter, The order dimension of planar maps, *SIAM J. Discrete Math.* **10** (1997), 515–528.
- [9] Y. Colin de Verdière, Sur un nouvel invariant des graphes et un critère de planarité, *J. Comb. Theory B* **50** (1990), 11–21.
- [10] P. Erdős and A. H. Stone, On the structure of linear graphs, *Bull. Amer. Math. Soc.* **52** (1946), 1089–1091.
- [11] S. Felsner, M. Habib and R. Möhring, On the interplay between interval dimension and dimension, *SIAM J. Discrete Math.* **7** (1994), 11–22.
- [12] S. Felsner and W. T. Trotter, The Dimension of Adjacency Posets of Planar Graphs, in preparation.
- [13] R. L. Graham, B. L. Rothschild and J. H. Spencer, *Ramsey Theory*, 2nd Edition, J. H. Wiley, New York, 1990.
- [14] S. Hoşten and W. D. Morris, Jr., The order dimension of the complete graph, *Discrete Math.*, to appear.
- [15] H. A. Kierstead, The dimension of layers of the subset lattice, *Discrete Math.*, to appear.
- [16] W. Schnyder, Planar graphs and poset dimension, *Order* **5** (1989), 323–343.
- [17] A. Schrijver, Minor-monotone graph invariants, in *Surveys in Combinatorics*, R. A. Bailey, ed., London Mathematical Society Lecture Note Series **241** (1997), 163–196.
- [18] J. Spencer, Minimal scrambling sets of simple orders, *Acta Math. Acad. Sci. Hungar.* **22** (1972), 349–353.
- [19] W. T. Trotter, *Combinatorics and Partially Ordered Sets: Dimension Theory*, The Johns Hopkins University Press, Baltimore, Maryland, 1992.
- [20] W. T. Trotter, Progress and new directions in dimension theory for finite partially ordered sets, in *Extremal Problems for Finite Sets*, P. Frankl, Z. Füredi, G. Katona and D. Miklós, eds., Bolyai Soc. Math. Studies **3** (1994), 457–477.
- [21] W. T. Trotter, Partially ordered sets, in *Handbook of Combinatorics*, R. L. Graham, M. Grötschel, L. Lovász, eds., Elsevier, Amsterdam, Volume I (1995), 433–480.
- [22] W. T. Trotter, Graphs and partially ordered sets, *Congressus Numerantium* **116** (1996), 253–278.
- [23] W. T. Trotter, New perspectives on interval orders and interval graphs, in *Surveys in Combinatorics*, R. A. Bailey, ed., London Mathematical Society Lecture Note Series **241** (1997), 237–286.
- [24] P. Turán, On an extremal problem in graph theory (in Hungarian), *Matematikai és Fizikai Lapok* **48** (1941), 436–452.
- [25] M. Yannakakis, On the complexity of the partial order dimension problem, *SIAM J. Alg. Discr. Meth.* **3** (1982), 351–358.

FACHBEREICH MATHEMATIK UND INFORMATIK, INSTITUT FÜR INFORMATIK, FREIE UNIVERSITÄT  
BERLIN, TAKUSTR. 9, 14195 BERLIN, GERMANY  
*E-mail address:* `felsner@inf.fu-berlin.de`

DEPARTMENT OF MATHEMATICS, ARIZONA STATE UNIVERSITY, TEMPE, ARIZONA 85287, U.S.A.  
*E-mail address:* `trotter@ASU.edu`