

7. LITERATURVERZEICHNIS

ABU AL-SOUD, W. and P. RADSTRÖM (1998):

Capacity of nine thermostable DNA polymerases to mediate DNA amplification in the presence of PCR-inhibiting samples.

Appl. Environ. Microbiol. **64**, 3748-3753

ALEKSEEV, A.N., H.V. DUBININA, L.P. ANTYKOVA, T.I. DZHIVANYAN,

S.G.T. RIJPKEMA, N. VERBEEK-DE KRUIF, and M. CINCO (1998):

Tick-borne borreliosis pathogen identification in *Ixodes* ticks (Acarina, Ixodidae) collected in St. Petersburg and Kalingrad baltic regions of Russia.

J. Med. Entomol. **35**, 136-142

ANDERSON, J.F. (1991):

Epizootiology of Lyme borreliosis.

Scand. J. Inf. Dis. Suppl. **77**, 23-34

ANGULO, A.B. (1986):

Lyme disease in cats.

Southw. Vet. **37**, 108-109

ARTHUR, D.R. (1963):

British Ticks, Butterworths, London

ASSOUS, M.V., D. POSTIC, G. PAUL, P. NEVOT and G. BARANTON (1994):

Individualisation of two new genomic groups among American *Borrelia burgdorferi* sensu lato strains.

FEMS Microbiol. Lett. **121**, 93-98

AZUMA, Y., K. KAWAMURA, H. ISOGAI, E. ISOGAI (1993):

Neurologic abnormalities in two dogs suspected Lyme disease.

Microbiol. Immunol. **37**, 325-329

BARBOUR, A.G. (1984):

Isolation and cultivation of Lyme disease spirochetes.

In: Steere, A.C., S.E. Malawista, J.E. Craft, D.K. Fischer, M. Garcia-Blanco (Hrsg.),
Lyme Disease, First Int. Symp., Yale J. Biol. Med., Connecticut, 71-75

BARK, S. (1986):

Zur Diagnose und Verbreitung der einheimischen Zecken-Borreliose beim Tier.

Vet. med. Diss., Tierärztl. Fakultät München

BASTA, J., J. PLCH, D. HULINKA, M. DANIEL (1999):

Incidence of *Borrelia garinii* and *Borrelia afzelii* in *Ixodes ricinus* ticks in an urban environment, Prague, Czech Republic, between 1995 and 1998.

Eur. J. Clin. Microbiol. Infect. Dis. **18**, 515-517

BAUERFEIND, R., U. KREIS, R. WEIß, L.H. WIELER, and G. BALJER (1998):

Detection of *Borrelia burgdorferi* in urine specimens from dogs by a nested polymerase chain reaction.

Zent.bl. Bakt. **287**, 347-361

BEICHEL, E., T.N. PETNEY, D. HASSSLER, M. BRUCKNER, M. MAIWALD (1996):

Tick infestation patterns and prevalence of *Borrelia burgdorferi* in ticks collected at a veterinary clinic in Germany.

Vet. Parasitol. **65**, 147-155

BERGLUND, J. and R. EITREM (1993):

Tick-borne borreliosis in the archipelago of southern Sweden.

Scand. J. Infect. Dis. **25**, 67-72

BERGMANN, J., A. LIEBISCH und K. POHLMEYER (1992):

Zum Vorkommen der einheimischen Borreliose bei Zecken, Wild- und Haustieren in einem niedersächsischen Moor.

Vet. **7**, 12-15

BERGSTRÖM, S., B. OLSEN, N. BURMANN, L. GOTHEFORS, T.G.T. JAENSON, M. JONSSON, H.A. MEJLON (1992):

Molecular characterization of *Borrelia burgdorferi* isolated from *Ixodes ricinus* in northern Sweden.

Scand. J. Infect. Dis. **24**, 181-188

BERNASCONI, M.V., C. VALSANGIACOMO, T. BALMELLI, O. PETER, and J.C. PIFFARETTI (1997):

Tick zoonoses in the southern part of Switzerland (Canton Ticino): occurrence of *Borrelia burgdorferi* sensu lato and *Rickettsia* sp..

Eur. J. Epidemiol. **13**, 209-215

BIGL, S., L. MÜLLER, G. PÖNITZ, C. MICKEL, B.-M. KLAPPER (1999):

Untersuchungen zur Epidemiologie der Borreliose im Freistaat Sachsen 1997.

Bundesgesundhbl. **42**: 219-225

BOSLER, E.M., D.P. COHEN, T.L. SCHULZE, C. OLSEN, W. BERNARD, B. LISSMANN (1988):

Host responses to *Borrelia burgdorferi* in dogs and horses.

Ann. N.Y. Ac. Sci. **539**, 221-233

BRAND, A. (1990):

Vergleichende seroepidemiologische Untersuchung von Rindern auf Zeckenborreliose in der Südheide und im Weserbergland.

Vet. med. Diss., Tierärztl. Hochschule Hannover

BREITSCHWERDT, E.B., W.L. NICHOLSON, A.R. KIEHL, C. STEERS, D.J. MEUTEN, and J.F. LEVINE (1994):

Natural infections with *Borrelia* spirochetes in two dogs from Florida.

J. Clin. Microbiol. **32**, 352-357

BREM, S., E. GÖBEL, U. BUSCH, D. KAHLAU, A. SCHÖNBERG, H. KOPP, and P. MEYER (1999a):

Nonmotile „rigid“ spiral forms observed during cultivation of *Bacillus* sp. - similar to atypical forms of *Borrelia burgdorferi* sensu lato.

Zentr.bl. Bakt. **289**, 744

BREM, S., E. GÖBEL, U. BUSCH, D. KAHLAU, J. HEIDRICH, A. SCHÖNBERG, H. KOPP, P. MEYER (1999b):

Nonmotile „rigid“ spiral shapes similar to atypical *Borrelia burgdorferi* sensu lato forms – observed during cultivation of different bacteria.

VIII. Int. Conf. Lyme Borreliosis Emerging Tick-Borne Dis., 20.-24.07.99 in München,
Abstr. P 363

BROWNING, A., S.D. CARTER, A. BARNES, C. MAY, D. BENNETT (1993):

Lameness associated with *Borrelia burgdorferi* infection in the horse.

Vet. Rec. **132**, 610-611

BURGDORFER, W., A.G. BARBOUR, S.F. HAYES, J.L. BENACH, E. GRUNWALDT, and J.P. DAVIS (1982):

Lyme disease- a tick-borne spirochelosis?

Sci. **216**, 1317-1319

BURGESS, E.C. (1986):

Experimental inoculation of dogs with *Borrelia burgdorferi*.

Zbl. Bakt. Hyg. (A) **263**, 49-54

BURGESS, E.C. (1988):

Borrelia burgdorferi infection in Wisconsin horses and cows.

Ann. N.Y. Ac. Sci. **539**, 235-243

BURGESS, E.C. (1991):

The role of wild mammals in the transmission of *Borrelia burgdorferi*.

Bull. Soc. Vect. Ecol. **16**, 50-58

BURGESS, E.C. (1992):

Experimentally induced infection of cats with *Borrelia burgdorferi*.

Am. J. Vet. Res. **53**, 1507-1511

BURGESS, E.C. and M. MATTISON (1987):

Encephalitis associated with *Borrelia burgdorferi* infection in a horse.

JAVMA **191**, 1457-1458

BURGESS, E.C., D. GILLETTE and J.P. PICKETT (1986):

Arthritis and panuveitis as manifestation of *Borrelia burgdorferi* infection in a Wisconsin pony.

JAVMA **189**, 1340-1342

BURGESS, E.C., A. GENDRON-FITZPATRICK and W.O. WRIGHT (1987):

Arthritis and systemic diseases caused by *Borrelia burgdorferi* infection in a cow.

JAVMA **191**, 1468-1470

BURMEISTER, G., C. HÜLBE, G. SCHÖTTLER, K. FIEDLER, H. STEINBRINK, L.W. SCHRÖDER, S. SOMMER (1998):

Ergebnisse 6-jähriger Zeckenuntersuchungen auf Borrelien und FSME-Viren in Mecklenburg-Vorpommern.

18. Tagung der DVG, 24.-28.03.98 in Dresden, Abstr. V 15

CARTER, S.D., C. MAY, A. BARNES, D. BENNETT (1994):

Borrelia burgdorferi infection in UK horses.

Equine Vet. J. **26**, 187-190

CERRI, D., F. FARINA, E. ANDREANI, R. NUVOLOLONI, A. PEDRINI (1994):

Experimental infection of dogs with *Borrelia burgdorferi*.

Res. Vet. Sci. **57**, 256-258

CINCO, M., R. MURGIA, S. BONIN, D. PADOVAN, G. STANTA (1996):

Detection of three species of *Borrelia burgdorferi* in *I. ricinus* in northern Italy by polymerase chain reaction.

Alpe Adria Microbiol. J. **5**, 253-260

COHEN, D., E.M. BOSLER, W. BERNARD, D. MEIRS, R. EISNER and T. SCHULZE (1988):

Epidemiologic studies of Lyme disease in horses and their public health significance.

Ann. N.Y. Ac. Sci. **539**, 244-257

COHEN, N.D., F.C. HECK, B. HEIM, D.M. FLAD, E.M. BOSLER, and D. COHEN (1992):

Seroprevalence of antibodies to *Borrelia burgdorferi* in a population of horses in Texas.

JAVMA **201**, 1030-1034

COONS, A.H., H.J. CREECH and R.N. JONES (1941):

Immunological properties of an antibody containing a fluorescent group.

Proc. Soc. Exp. Biol. Med. **47**, 200-202

CORNELY, M. und U. SCHULTZ (1992):

Zur Zeckenfauna Ostdeutschlands.

Ang. Paras. **33**, 173-183

CRAINE, N.G., S.E. RANDOLPH, P.A. NUTTALL (1995):

Seasonal variation in the role of grey squirrels as hosts of *Ixodes ricinus*, the tick vector of the Lyme disease spirochete, in a British woodland.

Fol. Parasit. **42**, 73-80

CUTLER, S., P. PHIPPS, M. CRANWELL, and D. WRIGHT (1990):

Borrelia burgdorferi antibodies in British cattle.

Proc. IV. Int. Conf. Lyme Borreliosis, 18.-21. Juni 1990 in Stockholm, P 43

DE BOER, R., K.E. HOVIUS, M.K.E. NOHLMANS, and J.S. GRAY (1993):

The woodmouse (*Apodemus sylvaticus*) as a reservoir of tick-transmitted spirochetes (*Borrelia burgdorferi*) in the Netherlands.

Zbl. Bakt. **279**, 404-416

DEDIE, K. (1989):

Das Erythema chronicum migrans, eine Schildzecken-Borreliose.

Tierärztl. Umsch. **44**, 480-482

DEMAERSCHALCK, I., A.B. MESSAOUD, M. DE KESEL, B. HOYOIS, Y. LOBET,

P. HOET, G. BIGAIGNON, A. BOLLEN, and E. GODFROID (1995):

Simultaneous presence of different *Borrelia burgdorferi* genospecies in biological fluids of Lyme disease patients.

J. Clin. Microbiol. **33**, 602-608

DENNIS, D.T. (1997):

Epidemiology of Lyme Borreliosis.

Proc. 2nd Int. Symp. Lyme Disease, 27.-28. Oktober 1997 in Japan, 121-164

DOBY, J.M., S. CHEVRIER, A. COUATARMANACH (1987):

Spirochetose a tiques par *Borrelia burgdorferi* chez le cheval en Bretagne.

Bull. Soc. Fr. Parasit. **2**, 5

DOBY, J.M., C. LEMBLE, G. BIGAIGNON, M. KREMER, C. ROLAND,

M.C. LAMBERT (1990):

Borrelia burgdorferi, agent des spirochetes a tiques (maladie de Lyme et autres formes cliniques) chez *Ixodes ricinus* (Acari: Ixodidae) en Alsace.

Bull. Soc. Fr. Parasit. **8**, 339-350

DOBY, J.M., C. BETREMIEUX, J. BARRAT, C. ROLLAND (1991a):

Tickborne spirochetosis due to *Borrelia burgdorferi* in wild carnivores in France: Results of the serological examination of 372 foxes.

Bull. Soc. Path. Exot. **84**, 46-53

DOBY, J.M., C. BETREMIEUX, C. ROLLAND, J. BARRAT (1991b):

Are large forest mammals reservoirs of *Borrelia burgdorferi*, the agent of Lyme disease? Serological survey of 543 deer and boar.

Rec. Med. Vet. **167**, 55-61

DORN, W., U. JACOBI und C. FLÜGEL (1995):

Zum Vorkommen von *Borrelia burgdorferi* in freilandgefangenen *Ixodes-ricinus*-Entwicklungsstadien in Thüringen.

In: Süss, J. (Hrsg.), Durch Zecken übertragbare Erkrankungen / FSME und Lyme-Borreliose, 3. Potsdamer Symp., 11. März 1995, Weller, Schriesheim, 162-174

DRESSLER, F. (1998):

Probleme der Standardisierung in der Diagnostik der Lyme Borreliose.

In: Talaska, T. (Hrsg.), Für die Praxis: Lyme Borreliose, Chromik Offsetdruck, Brieskow-Finkenheerd, ISBN 3-00-002363-1, 68-69

EIFFERT, H., A. OHLENBUSCH, H.J. CHRISTEN, R. THOMSEN, A.

SPIELMANN, F.R. MATUSCHKA (1995):

Nondifferentiation between Lyme disease spirochetes from vector ticks and human cerebrospinal fluid.

J. Infect. Dis. **171**, 476-479

ENG, T.R., M.L. WILSON, A. SPIELMANN, and C.C. LASTAVICA (1988):

Greater risk of *Borrelia burgdorferi* infection in dogs than in people.

J. Infect. Dis. **158**, 1410-1411

ENGVALL, E. and P. PERLMANN (1972):

Enzyme-linked immunosorbent assay, ELISA.

J. Imm. **109**, 129-135

ESTRADA-PEÑA, A., J.A. OTEO, R. ESTRADA-PEÑA, C. GORTAZAR, J.J. OSACAR, J.A. MORENO, and J. CASTELLA (1995):

Borrelia burgdorferi sensu lato in ticks (Acari: Ixodidae) from two different foci in Spain.

Exp. Appl. Acarol. **19**, 173-180

FEIR, D., C.R. SANTANELLO, B.W. LI, C.S. XIE, E. MASTERS, R. MARCONI, and G. WEIL (1994):

Evidence supporting the presence of *Borrelia burgdorferi* in Missouri.

Am. J. Trop. Med. Hyg. **51**, 475-482

FINGERLE; V., U. HAUSER, G. LIEGL, B. PETKO, V. PREAC-MURSIC, and B. WILSKE (1995):

Expression of outer surface proteins A and C of *Borrelia burgdorferi* in *Ixodes ricinus*.

J. Clin. Microbiol. **33**, 1867-1869

FINKBEINER-WEBER, B. (1996):

Die Rolle des Igels und seiner Ektoparasiten in der Epidemiologie der Lyme-Borreliose.

Vet. med. Diss., Tierärztl. Hochschule Hannover

FISCHER, A. und G. LEUTERER (1992):

Borrelia-burgdorferi assoziierte Lymphadenitis purulenta bei einem Hund.

Kleint. Prax. **37**, 13-16

FRESE, S. (1997):

Untersuchungen zum klinischen Verlauf der Lyme-Borreliose beim Hund nach experimenteller Infektion durch den Holzbock *Ixodes ricinus*.

Vet. med. Diss., FU Berlin

FRIDRIKSDOTTIR, V., G. OVERNES and S. STUEN (1992):

Suspected Lyme borreliosis in sheep.

Vet. Rec. **130**, 323-324

FUKUNAGA, M. (1997):

Vector tick and *Borrelia*.

Proc. 2nd Int. Symp. Lyme Disease, 27.-28. Oktober 1997 in Japan, 96-105

GERN, L., Z. ZHU, A. AESCHLIMANN (1990):

Development of *Borrelia burgdorferi* in *Ixodes ricinus* females during blood feeding.

Ann. Paras. Hum. **65**, 89-93

GERN, L., E. ROUVINEZ, L.N. TOUTOUNGI, and E. GODFROID (1997):

Transmission cycles of *Borrelia burgdorferi* sensu lato involving *Ixodes ricinus* and/or *I. hexagonus* and the European hedgehog, *Erinaceus europaeus*, in suburban areas in Switzerland.

Fol. Parasit. **44**, 309-314

GERN, L., A. ESTRADA-PEÑA, F. FRANSEN, J.S. GRAY, T.G.T. JAENSON, F. JONGEJAN, O. KAHL, E. KORENBERG, R. MEHL, and P.A. NUTTALL (1998):

European reservoir hosts of *Borrelia burgdorferi* sensu lato.

Zent.bl. Bakt. **287**, 196-204

GEUE, L. (1993):

Seroprävalenzuntersuchungen zum Auftreten von Antikörpern gegen *Borrelia burgdorferi* s.l. beim Rotfuchs (*Vulpes vulpes*) im Land Brandenburg.
unveröffentlicht

GIBSON, M.D., C.R. YOUNG, M.T. OMRAN, J. EDWARDS, K. PALMA, L. RUSSELL, J. RAWLINGS (1993):

Borrelia burgdorferi infection in cats.

JAVMA **202**, 237-239

GILL, J.S., R.G. MCLEAN, D.F. NEITZEL, and R.C. JOHNSON (1993):

Serologic analysis of white-tailed deer sera for antibodies to *Borrelia burgdorferi* by enzyme-linked immunosorbent assay and western immunoblotting.

J. Clin. Microbiol. **31**, 318-322

GILOT, B., B. DEGEILH, J. PICHOT, B. DOCHE, and C. GUIGUEN (1996):

Prevalence of *Borrelia burgdorferi* (sensu lato) in *Ixodes ricinus* (L.) populations in France, according to a phytoecological zoning of the territory.

Eur. J. Epidem. **12**, 395-401

GOLUBIC, D., S. RIJPKEMA, I. KVAKAN, T. REZEK, V. SKVORC, and E. RUZIC (1995):

Serological evidence for Lyme borreliosis in northwest Croatia.

Vet. Arh. **65**, 127-133

GOLUBIC, D., S. RIJPKEMA, N. TKALEC-MAKOVEC and E. RUZIC (1998):

Epidemiologic, ecologic and clinical characteristics of Lyme borreliosis in northwest Croatia.

Acta med. Croat. **52**, 7-13

GRAUER, G.F., E.C. BURGESS, A.C. COOLEY, and J.H. HAGEE (1988):

Renal lesions associated with *Borrelia burgdorferi* infection in a dog.

JAVMA **193**, 237-239

GRAY, J.S., O. KAHL, C. JANETZKI-MITTMANN, J. STEIN and E. GUY (1994):

Acquisition of *Borrelia burgdorferi* by *Ixodes ricinus* ticks fed on the European hedgehog, *Erinaceus europeans* L.

Exp. Appl. Acarol. **18**, 485-491

GRAY, J.S., O. KAHL, C. JANETZKI, J. STEIN and E. GUY (1995):

The spatial distribution of *Borrelia burgdorferi*-infected *Ixodes ricinus* in the Connemara region of county Galway, Ireland.

Exp. Appl. Acarol. **19**, 163-172

GREENE, R.T., R.L. WALKER, W.L. NICHOLSON, H.W. HEIDNER, J.F. LEVINE, E.C. BURGESS, M. WYAND, E.B. BREITSCHWERDT, and H.A. BERKHOFF (1988):

Immunoblot analysis of immunoglobulin G response to the Lyme disease agent (*Borrelia burgdorferi*) in experimentally and naturally exposed dogs.

J. Clin. Microbiol. **26**, 648-653

GREINER, M., D. SOHR and P. GÖBEL (1995):

A modified ROC analysis for the selection of cut-off values and the definition of intermediate results of serodiagnostic tests.

J. Immunol. Meth. **185**, 123-132

GUBLER, D.J. (1999):

Laboratory diagnosis of Lyme disease: lessons learned and prospects for the future.
VIII. Int. Conf. Lyme Borreliosis Emerging Tick-Borne Dis., 20.-24.07.99 in München,
Abstr. 76

GUETHOFF, W. und T. WEINKE (1998):

Borreliosen in der Inneren Medizin - Klinik und Therapie.

In: Talaska, T. (Hrsg.), Für die Praxis: Lyme Borreliose, Chromik Offsetdruck,
Brieskow-Finkenheerd, ISBN 3-00-002363-1, 79-85

GUPTA, S. (1994):

Studies regarding incidence of ticks in relation to their role as vector of *Borrelia burgdorferi* in Brandenburg state (Germany).

Vet. med. Diss., FU Berlin

GUPTA, S.K., A. SCHÖNBERG, H.F. MATTHES, TH. HIEPE und H.J. RUSCHER (1994):

Zum Vorkommen von *Borrelia burgdorferi* in Schildzecken *Ixodes ricinus* (Acari: Ixodidae) im Land Brandenburg.

Z. ärztl. Fortbild. **88**, 97-100

GUPTA, S.K., A. SCHÖNBERG and TH. HIEPE (1995):

Prevalence of ticks in relation to their role as vector of *Borrelia burgdorferi* under autochthonous conditions.

Appl. Parasitol. **36**, 97-106

GUSTAFSON, R., T.G.T. JAENSON, A. GARDULF, H. MEJLON, B.

SVENUNGSSON (1995):

Prevalence of *Borrelia burgdorferi* sensu lato infection in *Ixodes ricinus* in Sweden.

Scand. J. Infect. Dis. **27**, 597-601

GUY, E.C. and R.G. FARQUHAR (1991):

Borrelia burgdorferi in urban parks.

Lanc. **338**, 253

HAYES, S.F. and W. BURGDORFER (1993):

Ultrastructure of *Borrelia burgdorferi*.

In: Weber, K. and W. Burgdorfer (Hrsg.), Aspects of Lyme Borreliosis, Springer, Berlin, Heidelberg, 29-43

HEIDRICH, J., A. SCHÖNBERG, S. STEUBER, K. NÖCKLER, P. SCHULZE, W.P.

VOIGT, and E. SCHEIN (1999):

Investigation of skin samples from red foxes (*Vulpes vulpes*) in eastern Brandenburg (Germany) for the detection of *Borrelia burgdorferi* s.l.

Zentr.bl. Bakt. **289**, 666-672

HERBES, R.G., N. VERBEEK-DE KRUIF, S.G.T. RIJPKEMA,

J.F.P. SCHELLEKENS (1995):

Survey for *Borrelia burgdorferi* among roe deer and ticks in three parts of the Netherlands.

Tijdschr. Diergeesk. **120**, 722-725

HJELLE, A. (1966):

Borrelia-like organism in the urine of lambs suffering from eczema facialis.

Nat. **212**, 856-857

HOVIND-HOGEN, K. (1984):

Ultrastructure of spirochetes isolated from *Ixodes ricinus* and *Ixodes dammini*.

In: Steere, A.C., S.E. Malawista, J.E. Craft, D.K. Fischer, M. Garcia-Blanco (Hrsg.), Lyme Disease, First Int. Symp., Yale J. Biol. Med., Connecticut, 93-98

HOVMARK, A., E. ASBRINK, O. SCHWAN, B. HEDERSTEDT,

D. CHRISTENSSON (1986):

Antibodies to *Borrelia* spirochetes in sera from Swedish cattle and sheep.

Acta Vet. Scand. **27**, 479-485

HOVMARK, A.; T.G.T. JAENSON, E. ASBRINK, A. FORSMAN, E. JANSSON

(1988):

First isolation of *Borrelia burgdorferi* from rodents collected in northern Europe.

Acta Path. Microbiol. Immunol. Scand. **96**, 917-920

HUBALEK, Z. and J. HALOUZKA (1998):

Prevalence rates of *Borrelia burgdorferi* sensu lato in host-seeking *Ixodes ricinus* ticks in Europe.

Parasit. Res. **84**, 167-172

HUBBARD, M.J., A.S. BAKER, and K.J. CANN (1998):

Distribution of *Borrelia burgdorferi* s.l. spirochete DNA in British ticks (Argasidae and Ixodidae) since the 19th century, assessed by PCR.

Med. Vet. Entomol. **12**, 89-97

HÜLBE, C., H. HERRMANN, S. OHEIM, L.W. SCHRÖDER und M. v. STENGLIN

(1995):

Untersuchungen zur Verbreitung von *Borrelia burgdorferi* in Mecklenburg-Vorpommern.

Hyg. Med. **20**, 345-350

HULINKA, D., P. BARTAK, J. HERCOGOVA, J. HANCIL, J. BASTA, J. SCHRAMLOVA (1994):

Electron microscopy of Langerhans cells and *Borrelia burgdorferi* in Lyme disease patients.

Zbl. Bakt. **280**, 348-359

HUMAIR, P.F., N. TURRIAN, A. AESCHLIMANN, and L. GERN (1993):

Borrelia burgdorferi in a focus of Lyme borreliosis: epizootiologic contribution of small mammals.

Fol. Parasit. **40**, 65-70

HUNFELD, K.P., P. OSCHMANN, R. KAISER, J. SCHULZE, V. BRADE (1998):

Diagnostik.

In: Oschmann, P., P. Kraiczy (Hrsg.), Lyme-Borreliose und Frühsommer-Meningoenzephalitis, UNI-MED, Bremen, 76-109

ISOGAI, E., H. ISOGAI, H. KAWABATA, T. MASUZAWA, Y. YANAGIHARA, K. KIMURA, T. SAKAI, Y. AZUMA, N. FUJII, and S. OHNO (1994):

Lyme disease in a wild fox (*Vulpes vulpes schrencki*) and in ticks.

J. Wildl. Dis. **30**, 439-444

JAENSON, T.G.T. and L. TÄLLEKLINT (1996):

Lyme borreliosis spirochetes in *Ixodes ricinus* (Acari: Ixodidae) and the varying hare on isolated islands in the Baltic Sea.

J. Med. Entomol. **33**, 339-343

JENEK, J. and A. GLAZACZOW (1996):

The evaluation of *Borrelia burgdorferi* sensu lato spirochetes distribution in *Ixodes ricinus* ticks collected in selected regions of Wielkopolska + regions by polymerase chain reaction (PCR) method.

Przegl. Epidem. **50**, 383-386

JOHNSON, R.C., F.W. HYDE and C.M. RUMPEL (1984a):

Taxonomy of the Lyme disease spirochetes.

In: Steere, A.C., S.E. Malawista, J.E. Craft, D.K. Fischer, M. Garcia-Blanco (Hrsg.), Lyme Disease, First Int. Symp., Yale J. Biol. Med., Connecticut, 79-87

JOHNSON, R.C., N. MAREK and C. KODNER (1984b):

Infection of Syrian hamsters with Lyme disease spirochetes.

J. Clin. Microbiol. **20**, 1099-1101

JOHNSON, S.E., G.C. KLEIN, G.P. SCHMID, G.S. BOWEN, J.C. FEELEY, and T. SCHULZE (1984c):

Lyme disease: a selective medium for isolation of the suspected etiological agent, a spirochete.

J. Clin. Microbiol. **19**, 81-82

KÄSBOHRER, A. und A. SCHÖNBERG (1990):

Serologische Untersuchungen zum Vorkommen von *Borrelia burgdorferi* bei Haustieren in Berlin (West).

Berl. Münch. Tierärztl. Ws. **103**, 374-378

KAHL, O. (1991):

Lyme Borreliosis - an ecological perspective of a tick-borne human disease.

Anz. Schädl. K. **64**, 45-55

KAHL,O. (1998):

Betrachtungen zur Ökologie der Lyme-Borreliose in der Region Berlin-Brandenburg und zum Infektionsrisiko bei Zeckenstich.

In: Talaska, T. (Hrsg.), Für die Praxis: Lyme Borreliose, Chromik Offsetdruck, Brieskow-Finkenheerd, ISBN 3-00-002363-1, 19-21

KAHL, O. und L. GEUE (1996):

Zur Rolle des Rotfuchses, *Vulpes vulpes*, als ein möglicher Reservoirwirt für *Borrelia burgdorferi* s.l. (Bb).

17. Tagung der DVG, 27.-29. März 1996 in München, P 57

KAHL, O., K. SCHMIDT, A. SCHÖNBERG, U. LAUKAMM-JOSTEN, W. KNÜLLE, and U. BIENZLE (1989):

Prevalence of *Borrelia burgdorferi* in *Ixodes ricinus* ticks in Berlin (West).

Zbl. Bakt. Hyg. (A) **270**, 434-440

KAHL, O., C. JANETZKI, J.S. GRAY, J. STEIN and R.J. BAUCH (1992):

Tick infection rates with *Borrelia*: *Ixodes ricinus* versus *Haemaphysalis concinna* and *Dermacentor reticulatus* in two locations in eastern Germany.

Med. Vet. Entomol. **6**, 363-366

KAHL, O., L. GERN, J.S. GRAY, E.C. GUY, F. JONGEJAN, F. KIRSTEIN, K. KURTENBACH, S.G.T. RIJPKEMA, and G. STANEK (1998):

Detection of *Borrelia burgdorferi* sensu lato in ticks: immunofluorescence assay versus polymerase chain reaction.

Zent.bl. Bakt. **287**, 205-210

KAUFMAN, A.C., C.E. GREENE, R.A. MC GRAW (1993):

Optimization of polymerase chain reaction for the detection of *Borrelia burgdorferi* in biological specimens.

J. Vet. Diagn. Invest. **5**, 548-554

KAWABATA, H., H. TASHIBU, K. YAMADA, T. MASUZAWA, and Y. YANAGIHARA (1994):

Polymerase chain reaction analysis of *Borrelia species* isolated in Japan.

Microbiol. Immunol. **38**, 591-598

KAZMIERCZAK, J.J. and E.C. BURGESS (1989):

Antibodies to *Borrelia* sp. in wild foxes and coyotes from Wisconsin and Minnesota.

J. Wildl. Dis. **25**, 108-111

KERSTEN, A., C. POITSCHEK, S. RAUCH, E. ABERER (1995):

Effects of penicillin, ceftriaxone, and doxycycline on morphology of *Borrelia burgdorferi*.

Antimicr. Ag. Chemother. **39**, 1127-1133

KIMURA, K., E. ISOGAI, H. ISOGAI, Y. KAMEWAKA, T. NISHIKAWA, N. ISHII, and N. FUJII (1995):

Detection of Lyme disease spirochetes in the skin of naturally infected wild sika deer (*Cervus nippon yesoensis*) by PCR.

Appl. Environ. Microbiol. **61**, 1641-1642

KIRSTEIN, F., S. RIJPKEMA, M. MOLKENBOER and J.S. GRAY (1997):

The distribution and prevalence of *B. burgdorferi* genomospecies in *Ixodes ricinus* ticks in Ireland.

Eur. J. Epidemiol. **13**, 67-72

KLICH, M., M.W. LANKESTER, KINGWAN WU, and K.W. WU (1996):

Spring migratory birds (Aves) extend the northern occurrence of blacklegged tick (Acari: Ixodidae).

J. Med. Entomol. **33**, 581-585

KOPP, A. (1990):

Natürliche und experimentelle Infektionen bei Haustieren mit durch Zecken übertragenen Borrelien.

Vet. med. Diss., Tierärztl. Hochschule Hannover

KRAICZY, P., G. ACKER, V. BRADE (1998):

Erregereigenschaften.

In: Oschmann, P., P. Kraiczy (Hrsg.), Lyme-Borreliose und Frühsommer-Meningoenzephalitis, UNI-MED, Bremen, 16-25

KRAMPITZ, H. und S. BARK (1987):

Zur Epidemiologie der *Ixodes*-Borreliose in Süddeutschland.

Immun. Infekt. **15**, 141-145

KREIS, U. (1996):

Nachweis der Infektion mit *Borrelia burgdorferi* beim Hund mittels Polymerase-Kettenreaktion und Kulturverfahren.

Vet. med. Diss., Justus-Liebig-Universität Gießen

KURTENBACH, K., M. PEACEY, S.G.T. RIJPKEMA, A.N. HOODLESS, P.A. NUTTALL, and S.E. RANDOLPH (1998):

Differential transmission of the genospecies of *Borrelia burgdorferi* sensu lato by game birds and small rodents in England.

Appl. Environ. Microbiol. **64**, 1169-1174

KURTENBACH, K., H.S. SEWELL, S.S. SCHÄFER, and S. DE MICHELIS (1999):

Biology, ecology and systematics of the genus *Borrelia*.

Zentr.bl. Bakt. **289**, 639-642

LANDBO, A.S. and P.T. FLÖNG (1992):

Borrelia burgdorferi infection in *Ixodes ricinus* from habitats in Denmark.

Med. Vet. Entomol. **6**, 165-167

LEBECH, A.M., O. CLEMMENSEN, and K. HANSEN (1995):

Comparison of in vitro culture, immunohistochemical staining, and PCR for detection of *Borrelia burgdorferi* in tissue from experimentally infected animals.

J. Clin. Microbiol. **33**, 2328-2333

LE FLECHE, A., D. POSTIC, K. GIRARDET, O. PETER, and G. BARANTON (1997):

Characterization of *Borrelia lusitaniae* sp. nov. by 16S ribosomal DNA sequence analysis.

Int. J. Syst. Bact. **47**, 921-925

LEUBA-GARCIA, S., M.D. KRAMER, R. WALLICH, and L. GERN (1994):

Characterization of *Borrelia burgdorferi* isolated from different organs of *Ixodes ricinus* ticks collected in nature.

Zbl. Bakt. **280**, 468-475

LEVY, S.A. and P.H. DURAY (1988):

Complete heart block in a dog seropositive for *Borrelia burgdorferi*.

J. Vet. Int. Med. **2**, 138-144

LIEBISCH, A. (1993a):

Biologie und Ökologie der Zecken.

In: Horst, H. (Hrsg.), Einheimische Zeckenborreliose (Lyme-Krankheit) bei Mensch und Tier, PERIMED-spitta, Nürnberg, 31-47

LIEBISCH, A. (1993b):

Zeckenborreliose bei Haustieren.

In: Horst, H. (Hrsg.), Einheimische Zeckenborreliose (Lyme-Krankheit) bei Mensch und Tier, PERIMED-spitta, Nürnberg, 164-187

LIEBISCH, G. (1997):

Der Nachweis von Borrelien bei Haus- und Wildtieren: Patienten oder Reservoir der Lyme-Borreliose?

Proc. 22. Kongreß der DVG, 8.-11. April 1997 in Bad Nauheim, P 40

LIEBISCH, G., L. HOFFMANN, F. PFEIFER und A. LIEBISCH (1996):

Zur Zeckenfauna des Rotfuchses und deren Bedeutung für die Lyme Borreliose.

17. Tagung der DVG, 27.-29. März 1996 in München, Abstr. P 77

LIEBISCH, G., B. SOHNS, and W. BAUTSCH (1998):

Detection and typing of *Borrelia burgdorferi* sensu lato in *Ixodes ricinus* ticks attached to human skin by PCR.

J. Clin. Microbiol. **36**, 3355-3358

LISSMANN, B.A., E.M. BOSLER, H. CAMAY, B.G. ORMISTON, and J.L. BENACH (1984):

Spirochete-associated (Lyme disease) in a dog.

JAVMA **185**, 219-220

LIVESLEY, M.A., D. CAREY, L.GERN and P.A. NUTTALL (1994):

Problems of isolating *Borrelia burgdorferi* from ticks collected in United Kingdom foci of Lyme disease.

Med. Vet. Entomol. **8**, 172-178

LOTTMANN, H., B. WILSKE, H. HERRMANN (1996):

Characterization of *Borrelia burgdorferi* sensu lato strains isolated from *Ixodes ricinus* in Mecklenburg-Vorpommern.

Med. Microbiol. Immunol. **184**, 181-184

MADIGAN, J. (1993):

Lyme disease in horses.

Large Anim. Vet. **4**, 22-25

MAGNARELLI, L.A., J.F. ANDERSON, A.B. SCHREIER, and C.M. FICKE (1987):

Clinical and serological studies of canine Borreliosis.

JAVMA **191**, 1089-1092

MAGNARELLI, L.A., J.F. ANDERSON, E. SHAW, J.E. POST, and F.C. PALKA (1988):

Borreliosis in equids in northeastern United States.

Am. J. Vet. Res. **49**, 359-362

MAGNARELLI, L.A., J.F. ANDERSON, H.R. LEVINE, S.A. LEVY (1990):

Tick parasitism and antibodies to *Borrelia burgdorferi* in cats.

JAVMA **197**, 63-66

MAGNARELLI, L.A., J. F. ANDERSON and K.C. III STAFFORD (1994):

Detection of *Borrelia burgdorferi* in urine of *Peromyscus leucopus* by inhibition enzyme-linked immunosorbent assay.

J. Clin. Microbiol. **32**, 777-782

MAGNARELLI, L.A., A. DENICOLA, K.C. III STAFFORD, and J.F. ANDERSON (1995):

Borrelia burgdorferi in an urban environment: white-tailed deer with infected ticks and antibodies.

J. Clin. Microbiol. **33**, 541-544

MAIWALD, M., C. STOCKINGER, D. HASSSLER, M. v. KNEBEL-DOEBERITZ, and H.-G. SONNTAG (1995):

Evaluation of the detection of *Borrelia burgdorferi* DNA in urine samples by polymerase chain reaction.

Infect. **23**, 173-177

MARTIN, P., G. BIGAIGNON, P. GILLON, A. THIRION, and A. FAIN (1990):

Frequence de *Borrelia burgdorferi* (maladie de Lyme) et repartition de son vecteur *Ixodes ricinus* (Acari: Ixodidae) dans le district Mosan en Belgique.

Bull. Soc. Fr. Parasit. **8**, 331-338

MASUZAWA, T. (1997):

Molecular epidemiology of Lyme *Borreliae* in the far east.

Proc. 2nd Int. Symp. Lyme Disease, 27.-28. Oktober 1997 in Japan, 63-81

MASUZAWA, T. and Y. YANAGIHARA (1998):

Die Erreger - Klassifikation, Vorkommen und klinische Relevanz.

In: Talaska, T. (Hrsg.), Für die Praxis: Lyme Borreliose, Chromik Offsetdruck, Brieskow-Finkenheerd, ISBN 3-00-002363-1, 11-18

MATEICKA, F., D. KOZAKOVA, P.A. ROSA, and E. KMETY (1997):

Identification of *Borrelia burgdorferi* sensu lato tick isolates from Slovakia by PCR typing with 16S rRNA primers.

Zbl. Bakt. **286**, 355-361

MATUSCHKA, F.R., P. FISCHER, M. HEILER, S. BLÜMCKE, and A. SPIELMAN (1992):

Stage associated risk of transmission of the Lyme disease spirochete by European *Ixodes* ticks.

Parasit. Res. **78**, 695-698

MATUSCHKA, F.R., M. HEILER, H. EIFFERT, P. FISCHER, H. LOTTER, and A. SPIELMANN (1993):

Diversionary role of hoofed game in the transmission of Lyme disease spirochetes.
Am. J. Trop. Med. Hyg. **48**, 693-699

MATUSCHKA, F.R., T.W. SCHINKEL, B. KLUG, A. SPIELMAN, and D. RICHTER (1998):

Failure of *Ixodes* ticks to inherit *Borrelia afzelii* infection.
Appl. Environ. Microbiol. **64**, 3089-3091

MAY, C., S.D. CARTER, A. BARNES, C. MCLEAN, D. BENNETT, A. COUTTS, C.K. GRANT (1994):

Borrelia burgdorferi infection in cats in the UK.
J. Small Anim. Pract. **35**, 517-520

MEHL, R., P. SANDVEN, L.R. BRAATHEN (1987):

The tick *Ixodes ricinus*, a vector of spirochosis (in Norwegian).
Tidsskr. Norske Laegeforen. **107**, 1642-1644

MEHLE, J. (1996):

Lyme borreliosis in animals.
Vet. Novice **22**, 179-183

MIYAMOTO, K., M. NAKAO, K. UCHIKAWA, and H. FUJITA (1992):

Prevalence of Lyme borreliosis spirochetes in ixodid ticks of Japan, with special reference to a new potential vector, *Ixodes ovatus* (Acari: Ixodidae).
J. Med. Entomol. **29**, 216-220

MÖRNER, A.P., P. OLSON, M. CARLSON, and M. OLERSBACKEN (1990):

Clinical and serological studies of borreliosis in Swedish dogs.
Proc. IV. Int. Conf. Lyme Borreliosis, 18.-21. Juni 1990 in Stockholm, P 40

MONIN, R., L. GERN and A. AESCHLIMANN (1987):

A study of the different modes of transmission of *Borrelia burgdorferi* by *Ixodes ricinus*.

Zbl. Bakt. Suppl. **18**, 14-20

MOTER, S.E., H. HOFMANN, R. WALLICH, M.M. SIMON, and M.D. KRAMER (1994):

Detection of *Borrelia burgdorferi* sensu lato in lessional skin of patients with erythema migrans and acrodermatitis chronica atrophicans by ospA-specific PCR.

J. Clin. Microbiol. **32**, 2980-2988

NADZAMOVA, D., L. CISLAKOVA, M. TRAVNICEK, B. PET'KO and A. STEFANICIKOVA (1999):

The occurrence of IgG antibodies against Lyme Borreliosis agent *Borrelia burgdorferi* in free living animals from east Slovakia.

Proc. 3rd. Int. Conf. Ticks and Tick-Borne Pathogens, Slovakia, B 4

NIEPOLD, J. (1990):

Untersuchungen von Rinderseren auf das simultane Vorkommen von Infektionen mit Borrelien (*Borrelia burgdorferi*) und Babesien (*Babesia divergens*).

Vet. med. Diss., Tierärztl. Hochschule Hannover

NORMAN, G.L., J.M. ANTIG, G. BIGAIGNON, W.R. HOGREFE (1996):

Serodiagnosis of Lyme borreliosis by *Borrelia burgdorferi* sensu stricto, *B. garinii*, and *B. afzelii* Western blots (immunoblots).

J. Clin. Microbiol. **34**, 1732-1738

NUTTALL, P., S. RANDOLPH, D. CAREY, N. CRAINE, A. LIVESLEY, L. GERN (1994):

The ecology of Lyme borreliosis in the UK.

In: Axford, J.S., D.H.E. Rees (Hrsg.), Lyme borreliosis, New York, Plenum Press 125-129

OHEIM, S. und H. HERRMANN (1994):

Untersuchungen zur Verbreitung von *Borrelia burgdorferi* im Raum Greifswald in der Region Vorpommern des Landes Mecklenburg-Vorpommern.

Z. f. Ä. Fb. **88**, 807-810

OLIVER, J.H. (1997):

Ecology of ticks and transmission of *Borrelia burgdorferi*.

Proc. 2nd Int. Symp. Lyme Disease, 27.-28. Oktober 1997 in Japan, 82-95

OLIVER, J.H., T.M. KOLLARS, F.W. CHANDLER, A.M. JAMES, E.J. MASTERS, R.S. LANE, and L.O. HUEY (1998):

First isolation and cultivation of *Borrelia burgdorferi* sensu lato from Missouri.

J. Clin. Microbiol. **36**, 1-5

PETER, O., A.G. BRETZ and D. BEE (1995):

Occurrence of different genospecies of *Borrelia burgdorferi* sensu lato in ixodid ticks of Valais, Switzerland.

Eur. J. Epidemiol. **11**, 463-467

PETNEY, T.N., D. HASSSLER, M. BRÜCKNER, and M. MAIWALD (1996):

Comparison of urinary bladder and ear biopsy samples for determining prevalence of *Borrelia burgdorferi* in rodents in central Europe.

J. Clin. Microbiol. **34**, 1310-1312

PFISTER, K., B. BIGLER, J. NESVADBA, L. GREEN, and A. AESCHLIMANN (1989):

Borrelia burgdorferi infections in dogs in Switzerland.

Zbl. Bakt. Suppl. **18**, 26-31

PICKEN, M.M., R.N. PICKEN, D. HAN, Y. CHENG, F. STRLE (1996):

Single-tube nested polymerase chain reaction assay based on flagellin gene sequences for detection of *Borrelia burgdorferi* sensu lato.

Eur. J. Clin. Microbiol. Infect. Dis. **15**, 489-498

POKORNY, P. (1990):

Borrelia sp. in *Ixodes ricinus* ticks in the Prague area (in Czech.).

Cesk. Epidem. Mikrob. Imun. **39**, 32-38

POKORNY, P. and S. ZAHRADKOVA (1990):

The occurrence of *Borreliae* in *Ixodes ricinus* ticks in the Brno area (in Czech.).

Cesk. Epidem. Mikrob. Imun. **39**, 166-170

PORSTMANN, T. (1998):

Labordiagnostik der Borrelioseninfektion.

In: Talaska, T. (Hrsg.), Symposium Lyme-Borreliose, 11. September 1998 in Gross Lindow, 35-38

POST, J.E., E.E. SHAW and S.D. WRIGHT (1988):

Suspected borreliosis in cattle.

Ann. N.Y. Ac. Sci. **539**, 488

PREAC-MURSIC, V. and B. WILSKE (1993):

Biology of *Borrelia burgdorferi*.

In: Weber, K., W. Burgdorfer (Hrsg.), Aspects of Lyme Borreliosis, Springer, Berlin, Heidelberg, 44-58

PREAC-MURSIC, V., B. WILSKE and G. SCHIERZ (1986):

European *Borrelia burgdorferi* isolated from humans and ticks: culture conditions and antibiotic susceptibility.

Zbl. Bakt. Hyg. (A) **263**, 112-118

PREAC-MURSIC, V., G. WANNER, S. REINHARDT, B. WILSKE, U. BUSCH, W. MARGET (1996):

Formation and cultivation of *Borrelia burgdorferi* spheroplast-l-form variants.

Infect. **24**, 218-226

PROKOPCAKOVA, H., J. PETERKOVA, B. PETKO (1992):

Examination of ticks for the presence of *Borrelia* sp. in the Kosice area (in Slovak.).

Cesk. Epidem. Mikrob. Imun. **41**, 236-239

QUESSADA, T., CH. NAVETAT, S. ARNAUD, H. LEUDET, J. PICHOT (1999):

Detection of *Borrelia burgdorferi* s.l. by PCR in questing *Ixodes ricinus* (Linne, 1758) from the region of Lyon (France).

VIII. Int. Conf. Lyme Borreliosis Emerging Tick-Borne Dis., 20.-24.07.99 in München, Abstr. P 243

RATH, P.M., B. IBERSHOFF, A. MOHNHAUPT, J. ALBIG, B. ELJASCHEWITSCH, D. JÜRGENS, I. HORBACH, F.J. FEHRENBACH (1996):

Seroprevalence of Lyme Borreliosis in forestry workers from Brandenburg, Germany.

Eur. J. Clin. Microbiol. Infect. Dis. **15**, 372-377

RIJPKEMA, S., J. NIEUWENHUIJS, F.F.J. FRANSSEN, F. JONGEJAN (1994):

Infection rates of *Borrelia burgdorferi* in different instars of *Ixodes ricinus* ticks from the Dutch North Sea island of Ameland.

Exp. Appl. Acarol. **18**, 531-542

RIJPKEMA, S.G.T., M.J.C.H. MOLKENBOER, L.M. SCHOUMLS, F. JONGEJAN, J.F.P. SCHELLEKENS (1995):

Simultaneous detection and genotyping of three genomic groups of *Borrelia burgdorferi* sensu lato in Dutch *Ixodes ricinus* by characterization of the amplified intergenic spacer region between 5S and 23S rRNA genes.

J. Clin. Microbiol. **33**, 3091-3095

RIJPKEMA, S.G.T., R.G. HERBES, N. VERBEEK-DE KRUIF, and J.F.P. SCHELLEKENS (1996):

Detection of four species of *Borrelia burgdorferi* sensu lato in *Ixodes ricinus* ticks collected from roe deer (*Capreolus capreolus*) in the Netherlands.

Epidemiol. Infect. **117**, 563-566

RINDER, R. (1998):

PCR-Sinn, Unsinn, Qualitätssicherung.

18. Tagung der DVG, 24.-28.03.98 in Dresden, Abstr. V 95

ROESSLER, D., H. EIFFERT, S. JAURIS-HEIPKE, G. LEHNERT, V. PREAC-MURSIC, J. TEEPE, T. SCHLOTT, E. SOUTSCHEK, AND B. WILSKE (1995):

Molecular and immunological characterization of the p 83/100 protein of various *Borrelia burgdorferi* sensu lato strains.

Med. Microbiol. Immunol. **184**, 23-32

ROESSLER, D., V. VASILIU, B. WILSKE (1996):

Development of OspA type specific PCR for characterization of *Borrelia burgdorferi* sensu lato.

J. Microbiol. Meth. **27**, 102, Abstr. 19

ROLFS, A., I. SCHULLER, U. FINCKH, I. WEBER-ROLFS (1992):

PCR Principles and reaction components.

In: Rolfs, A., I. Schuller, U. Finckh, I. Weber-Rolfs (Hrsg.), PCR: Clinical diagnostics and research, Springer, Berlin, Heidelberg, New York, London, Paris, Tokyo, Hong Kong, Barcelona, Budapest, 1-21

RUSSELL, R.C., S.L. DOGGETT, R. MUNRO, J. ELLIS, D. AVERY, C. HUNT and D. DICKESON (1994):

Lyme disease: a search for a causative agent in ticks in south-eastern Australia.

Epidemiol. Infect. **112**, 375-384

RUZIC-SABLJIC, E., F. STRLE, J. CIMPERMAN (1993):

The *Ixodes ricinus* tick as a vector of *Borrelia burgdorferi* in Slovenia.

Eur. J. Epidemiol. **9**, 396-400

SAINT GIRONS, I., L. GERN, J.S. GRAY, E.C. GUY, E. KORENBERG, P.A. NUTTALL, S.G.T. RIJPKEMA, A. SCHÖNBERG, G. STANEK, and D. POSTIC (1998):

Identification of *Borrelia burgdorferi* sensu lato species in Europe.

Zent.bl. Bakt. **287**, 190-195

SCHEMPP, C., H. BOCKLAGE, M. OWSIANOWSKI, R. LANGE, C.E. ORFANOS, H. GOLLNICK (1993):

In-vivo- und In-vitro-Nachweis einer Borrelieninfektion bei einer morpheaähnlichen Hautveränderung mit negativer Borrelien-Serologie.

Hautarzt **44**, 14-18

SCHÖFFEL, I., E. SCHEIN, U. WITTSTADT und J. HENTSCHE (1991):

Zur Parasitenfauna des Rotfuchses in Berlin (West).

Berl. Münch. Tierärztl. Ws. **104**, 153-157

SCHÖNBERG, A., C. CAMEY, O. KAHL, B. WILSKE, V. PREAC-MURSIC, and K. HOVIND-HOUGEN (1988):

First isolation of *Borrelia burgdorferi*, the agent of Lyme Borreliosis, from *Ixodes ricinus* (Acari: Ixodidae) in Berlin (West).

Zbl. Bakt. Hyg. (A) **268**, 487-494

SCHÖNBERG, A., K. SCHMIDT und A. KÄSBOHRER (1989):

Eine durch Zecken übertragbare Zoonose: Lyme-Borreliose.

Bundesgesundhbl. **32**, Sonderdruck, 190-193

SCHÖNBERG, A., B. HAHN-HEY, U. KÄMPE, K. SCHMIDT, and W.A. ELLIS (1992):

The isolation and identification of *Leptospira interrogans* serovar bratislava from a pig in Germany.

J. Vet. Med. **39**, 362-368

SCHÖNBERG, A., A. KÄSBOHRER, S. BREM und P.M. RATH (1994):

Further important details about immobile spiral microorganism in skin biopsies and cerebrospinal fluid from patients with Lyme Borreliosis.

VI. Int. Conf. Lyme Borreliosis, 19.-22. Juni 1994 in Bologna, Abstr. P 086 T

SCHÖNBERG, A., C. LOSER und S. GUPTA (1995):

Borrelia-burgdorferi-Isolate aus Zecken der neuen Bundesländer und ihre Reaktionen mit monoklonalen Antikörpern.

In: Süss, J. (Hrsg.), Durch Zecken übertragbare Erkrankungen / FSME und Lyme-Borreliose, 3. Potsdamer Symp., 11. März 1995, Weller, Schriesheim, 155-161

SCHÖNBERG, A., W. ERLER und B. JACOB (1996):

Vergleichende Untersuchungen von *Borrelia-burgdorferi*-Isolaten aus dem Berliner Raum als Beitrag zur serologischen Diagnostik.

Bundesgesundhbl. **39**, Sonderdruck, 332-336

SCHRÖCK, K. und A. SCHÖNBERG (1998):

Veterinärmedizinische Aspekte der Lyme-Borreliose.

In: Talaska, T. (Hrsg.), Für die Praxis: Lyme Borreliose, Chromik Offsetdruck, Brieskow-Finkenheerd, ISBN 3-00-002363-1, 34-38

SCHULZE, P., S. MACEDO, A. SCHÖNBERG und M. SCHWEBS (1995):

Morphologische Unterschiede von *Borrelia-burgdorferi*-Stämmen bei elektronenmikroskopischen Untersuchungen.

In: Süss, J. (Hrsg.), Durch Zecken übertragbare Erkrankungen / FSME und Lyme-Borreliose, 3. Potsdamer Symp., 11. März 1995, Weller, Schriesheim, 175-181

SCHWARTZ, I., S. VARDE, R.B. NADELMAN, G. P. WORMSER, and D. FISH (1997):

Inhibition of efficient polymerase chain reaction amplification of *Borrelia burgdorferi* DNA in blood-fed ticks.

Am. J. Trop. Med. Hyg. **56**, 339-342

SIMON, M.M., Y. BAUER, W. ZHONG, H. HOFMAN, and R. WALLICH (1999):

Lyme disease: Pathogenesis and vaccine development.

Zentr.bl. Bakt. **289**, 690-695

SIMPSON, W.J., M.E. SCHRUMPF, and T.G. SCHWAN (1990):

Reactivity of human Lyme borreliosis sera with a 39-kilodalton antigen specific to *Borrelia burgdorferi*.

J. Clin. Microbiol. **28**, 1329-1337

SINSKY, R.J. and J. PIESMAN (1989):

Ear punch biopsy method for detection and isolation of *Borrelia burgdorferi* from rodents.

J. Clin. Microbiol. **27**, 1723-1727

SKOTARCZAK, B., B. WODECKA and A. STACHOW (1998):

Detection Lyme disease spirochetes, *Borrelia burgdorferi* sensu lato DNA by polymerase chain reaction in ticks in the recreative areas of Szczecin.

18. Tagung der DVG, 24.-28.03.98 in Dresden, Abstr. P 122

SLAJCHERT, T., U.D. KITRON, C.J. JONES, and A. MANNELLI (1997):

Role of eastern chipmunk (*Tamias striatus*) in the epizootiology of Lyme borreliosis in northwestern Illinois, USA.

J. Wildl. Dis. **33**, 40-46

STANEK, G. (1997):

Biology of *Borrelia burgdorferi*, risk of infection of Lyme-borreliosis.

In: Süss, J., O. Kahl (Hrsg.), Tick-borne Encephalitis and Lyme Borreliosis, 4th Int. Potsdam Symp. on Tick-Borne Dis., 21.-22. Februar 1997, Pabst Science Publishers, Lengerich, 230-237

STANEK, G. (1998):

Klinische Falldefinitionen der Lyme-Borreliose für Europa.

In: Talaska, T. (Hrsg.), Für die Praxis: Lyme Borreliose, Chromik Offsetdruck, Brieskow-Finkenheerd, ISBN 3-00-002363-1, 114-117

STRLE, F., Y. CHENG, J.A. NELSON, M.M. PICKEN, J.K. BOUSEMANN, R.N. PICKEN (1995):

Infection rate of *Ixodes ricinus* ticks with *Borrelia afzelii*, *Borrelia garinii*, and *Borrelia burgdorferi* sensu stricto in Slovenia.

Eur. J. Clin. Microbiol. Infect. Dis. **14**, 994-1001

TÄLLEKLINT, L. and T.G.T. JAENSON (1994):

Transmission of *Borrelia burgdorferi* s.l. from mammal reservoirs to the primary vector of Lyme borreliosis, *Ixodes ricinus* (Acari: Ixodidae), in Sweden.

J. Med. Entomol. **31**, 880-886

TALASKA, T. (1998a):

Diagnostische Methoden bei Borrelien-Infektionen - Übersicht -.

In: Talaska, T. (Hrsg.), Für die Praxis: Lyme Borreliose, Chromik Offsetdruck, Brieskow-Finkenheerd, ISBN 3-00-002363-1, 48-59

TALASKA, T. (1998b):

Borreliose-Epidemiologie am Beispiel des Bundeslandes Brandenburg.

In: Talaska, T. (Hrsg.), Für die Praxis: Lyme Borreliose, Chromik Offsetdruck, Brieskow-Finkenheerd, ISBN 3-00-002363-1, 40-47

TELFORD, S.R., T.N. MATHER, S.I. MOORE, M.L. WILSON, A. SPIELMAN (1988):

Incompetence of deer as reservoir of the Lyme disease spirochete.

Am. J. Trop. Med. Hyg. **39**, 105-109

TREVISAN, G., G. STINCO, M. CINCO (1997):

Neonatal skin lesions due to a spirochetal infection: a case of congenital Lyme borreliosis?

Int. J. Dermat. **36**, 677-699

WASMOEN, T.L., R.W. SEBRING, B.M. BLUMER, L.G. CHAVEZ, H.J. CHU, and W.M. ACREE (1992):

Examination of Koch's postulates for *Borrelia burgdorferi* as the causative agent of limb/joint dysfunction in dogs with borreliosis.

JAVMA **201**, 412-418

WEBER, A. und U. HEIM (1989):

Borreliose beim Hund.

Vet. **9**, 14-16

WEGNER, Z., J. STANCZAK, M. RACEWICZ, W. KRUMINIS-LOZOWSKA, B. KUBICA-BIERNAT (1993/94):

Occurrence of *Borrelia* spirochetes in ticks (Acari: Ixodidae) collected in the forest areas in Olsztyn province (north central Poland).

Bull. Inst. Mar. Trop. Med. Gdynia **44/45**, 51-59

WEGNER, Z., J. STANCZAK, M. RACEWICZ, B. KUBICA-BIERNAT, and W. KRUMINIS-LOZOWSKA (1997):

The etiological agent of Lyme disease, *Borrelia burgdorferi*, in ticks (Acari: Ixodidae) from eastern Poland.

Zbl. Bakt. **286**, 93-106

WILLIAMS, W.V., P. CALLEGARI, B. FREUNDLICH, G. KEENAN, D. ELDRIDGE, H. SHIN, M. KREITMAN, D. McCALLUS, and D.B. WEINER (1992):

Molecular diagnosis of *Borrelia burgdorferi* infection (Lyme disease).

DNA Cell Biol. **11**, 207-213

WILSKE, B. and V. PREAC-MURSIC (1993):

Microbiological diagnosis of Lyme Borreliosis.

In: Weber, K., W. Burgdorfer (Hrsg.), Aspects of Lyme Borreliosis, Springer, Berlin, Heidelberg, 267-299

WILSKE, B., V. PREAC-MURSIC and G. SCHIERZ (1985):

Antigenic heterogeneity of European *Borrelia* strains isolated from patients.

Lanc. **1**, 1099

WILSKE, B., U. BUSCH, V. FINGERLE, S. JAURIS-HEIPKE, V. PREAC-MURSIC, D. ROESSLER, and G. WILL (1996):

Immunological and molecular variabilitiy of OspA and OspC. Implications for *Borrelia* vaccine development.

Infect. **24**, 208-212

WILSON, I.G. (1997):

Inhibition and facilitation of nucleic acid amplification.

Appl. Environ. Microbiol. **63**, 3741-3751

WITTENBRINK, M.M., D. THIELE, and H. KRAUSS (1994):

Comparison of dark-field microscopy, culture, and polymerase chain reaction (PCR) for detection of *Borrelia burgdorferi* in field-collected *Ixodes ricinus* ticks.

Zbl. Bakt. **281**, 183-191

WOLF, S. (1996):

Elektronenmikroskopische Untersuchungen zur Ultrastruktur von *Borrelia burgdorferi*.

Vet. med. Diss., FU Berlin

ZHIOUA, E., D. POSTIC, F. RODHAIN, and C. PEREZ-EID (1996):

Infection of *Ixodes ricinus* (Acari: Ixodidae) by *Borrelia burgdorferi* in Ile de France.

J. Med. Entomol. **33**, 694-697