

9. Verzeichnis der eingebundenen Publikationen

- [A] Giessing M, Fischer T, Deger S, Türk I, Schönberger B, Fritzsche L, Neumayer HH, Loening SA, Budde K. Increased frequency of lymphoceles following kidney transplantation under treatment with Sirolimus: a single center experience. *Transplant Proc* 2002; 34:1815-1816.
- [B] Fischer T, Filimonow SI, Taupitz M, Petersein J, Beyersdorff D, Bollow M, Hamm B. Bildqualität und Detektion pathologischer Prozesse im Ultraschallbild: Vergleich von Photopischem Sehen, Tissue Harmonic Imaging und nativem B-Bild. *Fortschr Röntgenstr* 2002; 174:1313-1317.
- [C] Fischer T, Filimonow SI, Petersein J, Beyersdorff D, Mühler M, Bollow M, Badakhshi HR, Hamm B. Sonographie am Krankenbett: Zeitgewinn durch Nutzung eines tragbaren Ultraschallgerätes? *Ultraschall Med* 2002; 23:311-314.
- [D] Fischer T, Mühler M, Kröncke TJ, Lembcke A, Rudolph J, Diekmann F, Ebeling V, Thomas A, Greis C, Hamm B, Filimonow S. Early Postoperative Ultrasound of Kidney Transplants: Evaluation of Contrast Medium Dynamics Using Time-intensity Curves. *Fortschr Röntgenstr* 2004; 176:472-477.

- [E] Fischer T, Ebeling V, Giessing M, Mühler M, Filimonow S, Dieckhöfer J, Lembcke A, Rudolph J, Morgera S, Budde K, Hamm B, Thomas A. A new method for standardized diagnosis following renal transplantation: Ultrasound with contrast enhancement. *Urologe A* 2006; 45: 38-45.
- [F] Fischer T, Filimonow S, Dieckhöfer J, Slowinski T, Mühler M, Lembcke A, Budde K, Neumayer HH, Ebeling V, Giessing M, Thomas A, Morgera S. Diagnosis of Acute Rejection of Kidney Transplants by Ultrasound with Echo Enhancer. *Nephrol Dial Transpl* 2006; 5: (im Druck)
- [G] Fischer T, Filimonow S, Rudolph J, Morgera S, Budde K, Slowinski T, Ebeling V, Taymoorian K, Giessing M, Thomas A. Arrival Time Parametric Imaging: A New Ultrasound Technique for Quantifying Perfusion of Kidney Grafts. *European Journal of Ultrasound* 2006 (im Druck)
- [H] Fischer T, Filimonow S, Mutze S, Morgera S, Thomas A. Die Transplantatniere: Farbkodierte Duplexsonographie und kontrastmittelverstärkte Sonographie in der Beurteilung nephrologischer und chirurgischer Komplikationen. *Fortschr Röntgenstr* 2006; 178:1-10. (im Druck)

10. Allgemeines Literaturverzeichnis

- 1) Murray JE. Renal homotransplantation in identical twins. *Surg Forum* 1955; 6: 432-436.
- 2) Eigler FW. Zur Geschichte der Nierentransplantation in Deutschland. *Zentralbl Chir* 2002; 127: 1001-1008.
- 3) Calne RY, Murray JE. Inhibition of rejection of renal allografts in dogs by Burroughs Wellcome 57-322. *Surg Forum* 1961; 12: 118-120.
- 4) Borel JF, Feurer C, Gubler HU, Stahelin H. Biological effects of cyclosporine A: a new antilymphocytic agent. *Agents Actions* 1976; 6: 468-475.
- 5) Mallek R, Mostbeck G, Kain R, et al. Vaskuläre Nierentransplantatabstoßung – Ist eine duplexsonographische Diagnose möglich? *Fortschr Röntgenstr* 1990; 152: 283.
- 6) Baxter GM, Ireland H, Moss JG, et al. Colour Doppler ultrasound in renal transplant artery stenosis: which Doppler index? *Clin Radiol* 1995; 50: 618-622.
- 7) Mebel M, May G, Althaus P (Hrsg). *Der komplette Nierenersatz! Aufbau und Entwicklung der Nierentransplantation in der DDR*. Pabst Science Publishers, Lengerich, Berlin, Bremen 2003.
- 8) Deutsche Stiftung Organtransplantation, www.dso.de
- 9) Meier-Kriesche HU, Kaplan B. Waiting time on dialysis as the strongest modifiable risk factor for renal transplant outcomes. Paired donor kidney analysis. *Transplantation* 2002; 74: 1377-1381.

- 10) Diethelm AG, Deierhoi MH, Hudson SL, et al. Progress in renal transplantation. A single center study of 3359 patients over 25 years. Ann Surg 1995; 221: 446-458.
- 11) Lee HM, Posner MP, King AL, et al. Status of long-term (25 years) survival of kidney transplant patients. Transplant Proc 1993; 25: 1336-1337.
- 12) Alvarez G, Gonzalez-Molina M, Cabello M, et al. Pulsed and continuous Doppler evaluation of renal dysfunction after kidney transplantation. Eur J Radiol 1991; 12: 108-112.
- 13) Arima M, Ogino T, Hosokawa S, et al. Functional image diagnosis of kidney transplants using ultrasonic Doppler flowmetry and magnetic resonance imaging. Transplant Proc 1989; 21: 1907-1911.
- 14) De-Gaetano AM, Boldrini G, Nanni G, et al. Noninvasive surveillance of allografted kidneys by ultrasonic duplex scanning. Angiology 1989; 40: 705-715.
- 15) Flückner F, Steiner H, Horn S, et al. Farbkodierte Duplexsonographie und Widerstandsindex bei Nierentransplantaten mit Dysfunction. Fortschr Röntgenstr 1990; 153: 692-697.
- 16) Grenier N, Douws C, Morel D, et al. Detection of vascular complications in renal allografts with color Doppler flow imaging. Radiology 1991; 178: 217-223.
- 17) Hollenbeck M. New diagnostic techniques in clinical nephrology. Colour coded duplex sonography of renal transplants – tool or toy for the nephrologist? Nephrol Dial Transplant 1994; 9: 1822-1828.

- 18) Steinberg HV, Nelson RC, Murphy FB, et al. Renal allograft rejection: evaluation by Doppler US and MR Imaging. Radiology 1987; 162: 337-342.
- 19) Radermacher J, Mengel M, Ellis S, et al. The renal arterial resistance index and renal allograft survival. N Engl J Med 2003; 349(2):182-184.
- 20) Restrepo-Schafer IK, Schwerk WB, Muller TF, et al. Intrarenal doppler flow analysis in patients with kidney transplantation and stable transplant function. Ultraschall Med 1999; 20(3): 87-92.
- 21) Letourneau JG, Day DL, Feinberg SB. Ultrasound and computed tomographic evaluation of renal transplantation. Radiol Clin North Am 1987; 25: 267-279.
- 22) Dodd GD, Tublin ME, Shah A, et al. Imaging of vascular complications associated with renal transplants. Am J Roentgenol 1991; 157: 449-459.
- 23) Hollenbeck M, Kutkuhn B, Grabensee B. Colour Doppler ultrasound in the diagnosis of transplant renal artery stenosis. Bildgebung 1994; 61: 248-254.
- 24) Townsend RR, Tomlanowich SJ, Goldstein RB, et al. Combined Doppler and morphologic sonographic evaluation of renal transplant rejection. J Ultrasound Med 1990; 9: 199-206.
- 25) Weskott HP. Power US: Phantom measurement of slow flow. Radiology 1995; 197: 337.
- 26) Waiser J, Schreiber M, Budde K, et al. Prognostic value of the Banff classification. Transpl Int 2000; 13(Suppl 1): 106-111.
- 27) Bude RO, Rubin JM, Adler ED. Power versus conventionel color Doppler sonography: comparison in the depiction of normal intrarenal vasculature. Radiology 1994; 192: 777-780.

- 28) Genkins SM, Sanfilippo FP, Carroll BA. Duplex Doppler sonography of renal transplants: Lack of sensitivity and specificity in establishing pathologic diagnosis. Am J Roentgenol 1989; 152: 535-539.
- 29) Hollenbeck M, Hilbert N, Meusel F, et al. Increasing sensitivity and specificity of Doppler sonographic detection of renal transplant rejection with serial investigation technique. Clin Invest 1994; 72: 609-615.
- 30) Phillips AO, Deane C, O'Donnell P, et al. Evaluation of Doppler ultrasound in primary non-function of renal transplants. Clin Transplant 1994; 8: 83-86.
- 31) Rigsby CM, Burns PM, Weltin GG, et al. Doppler signal quantitation in renal allografts: comparison in normal and rejecting transplants, with pathologic correlation. Radiology 1987; 162: 39-42.
- 32) Mallek R, Mostbeck G, Kain R, et al. Vaskuläre Nierentransplantatabstoßung – Ist eine duplexsonographische Diagnose möglich? Fortschr Röntgenstr 1990; 152: 283.
- 33) Hollenbeck M, Stuhrmann M, Trapp R, et al. Farbkodierte Dopplersonographie zur Früherkennung von Abstoßungsreaktionen nach allogener Nierentransplantation. Dtsch med Wochenschr 1994; 116: 921-927.
- 34) Pozniak MA, Dodd GD, Kelcz F. Ultrasonographic evaluation of renal transplantation. Radiol Clin North Am 1992; 30(5): 1053-1066.
- 35) Fahlenkamp D, Schönberger B, Lindeke A, et al. Laparaskopische Lymphozelendrainage nach Nierentransplantation. Z Urologie 1993; 2: 55-56.
- 36) Martinoli C, Bertolotto M, Crespi G, et al. Duplex Doppler analysis of interlobular arteries in transplanted kidneys. Eur Radiol 1998; 8: 765-769.

- 37) Takano R, Ando Y, Taniguchi N, et al. Power Doppler sonography of the kidney: effect of Valsalva's maneuver. *J Clin Ultrasound* 2001; 29: 384-390.
- 38) Schwerk WB, Restrepo IK, Prinz H. Semiquantitative analysis of intrarenal arterial Doppler flow spectra in healthy adults. *Ultraschall Med* 1993; 14: 117-122.
- 39) Delbeke D, Sacks GA, Sandler MP. Diagnosis of allograft renal vein thrombosis. *Clin Nucl Med* 1989; 14: 415-420.
- 40) Duckett T, Bretan PN, Cochran ST. Noninvasive radiological diagnosis of renal vein thrombosis in renal transplantation. *J Urol* 1991; 46: 403-406.
- 41) Laplante S, Patriquin HB, Robitaille P, et al. Renal vein thrombosis in children: evidence of early flow recovery with Doppler US. *Radiology* 1993; 189: 37-42.
- 42) Birkeland SA, Rohr N, Elbirk A et al. Chronic rejection in kidney transplant patients receiving cytomegalovirus prophylaxis with acyclovir. *Transplant Proc* 1995; 27: 2040-2041.
- 43) Jordan ML, Cook GT, Cardelle CJ. Ten years of experience with vascular complications in renal transplants. *J Urol* 1982; 128: 689-692.
- 44) Baxter GM, Ireland H, Moss JG, et al. Colour Doppler ultrasound in renal transplant artery stenosis: which Doppler index? *Clin Radiol* 1995; 50: 618-622.
- 45) Maia CR, Bittar AE, Goldani JC, et al. Doppler ultrasonography for the detection of renal artery stenosis in transplanted kidneys. *Hypertension* 1992; 19: 207-209.
- 46) Snider JF, Hunter DW, Moradian GP, et al. Transplant renal artery stenosis: evaluation with duplex sonography. *Radiology* 1989; 172: 1027-1030.

- 47) Deane C, Cowan N, Giles J, et al. Arteriovenous fistulas in renal transplants: color Doppler ultrasound observations. *Urol Radiol* 1992; 13: 211-217.
- 48) Fischer T, Mühler M, Kröncke TJ, et al. Early Postoperative Ultrasound of Kidney Transplants: Evaluation of Contrast Medium Dynamics Using Time-intensity Curves. *Fortschr Röntgenstr* 2004; 176(4): 472-477.
- 49) Pallardó LM, Sánchez J, Puig N, et al. Chronic rejection in 500 kidney transplant patients treated with cyclosporine: incidence and risk factors. *Transplant Proc* 1995; 27: 2215-2216.
- 50) Bude RO, Rubin JM, Adler RD. Power versus conventional color Doppler sonography: comparison in the depiction of normal intrarenal vasculature. *Radiology* 1994; 192: 777-780.
- 51) Chiu MI, Katz H, Berlin V. RAPT1, a mammalian homolog of yeast Tor, interacts with the FKBP12/rapamycin complex. *Proc Natl Acad Sci USA* 1994; 91(26): 12574-12578.
- 52) Kahan BD. Efficacy of sirolimus compared with azathioprine for reduction of acute renal allograft rejection: a randomised multicentre study. The Rapamune US Study Group. *Lancet* 2000; 356(9225): 194-202.
- 53) Boedker A, Ooikjar O, Rasmussen F, Loekkegaard H. Lymphocele following renal transplantation: a clinical study. *Transpl Proc* 1990; 22: 154.
- 54) Khatri RB, Stoff JS, Lovewell T, et al. Post-transplant lymphoceles: a critical look into the risk factors, pathophysiology and management. *J Urol* 1993; 150(1): 22-26.
- 55) Nuininga JE, Feitz WF, van Dael KC, et al. Urological complications in pediatric renal transplantation. *Euro Urol* 2001; 39(5): 598-602.

- 56) McCullough CS, Soper NJ, Claymann RV, et al. Laparoscopic drainage of a posttransplant lymphocele. *Transplantation* 1991; 51(3): 725-727.
- 57) Giessing M, Fuller TF, Deger S, et al. 10 Jahre laparaskopische Lebendnierenpende: Von Rarität zu Routine. *Urologe* 2006; 45: 46-52.
- 58) Griffin JF, McNicholas MM. Morphological appearance of renal allografts in transplant failure. *J Clin Ultrasound* 1992; 20: 529-537.
- 59) Jansen O, Rob PM, Schmidtke V, et al. Follow-up study of renal transplants by duplex Doppler and grey scale ultrasound. *Eur J Radiol* 1992; 15: 26-31.
- 60) Renowden SA, Blethyn J, Cochlin DL. Renal transplant sonography: correlation of Doppler and biopsy results in cellular rejection. *Clin Radiol* 1992; 46: 265-269.
- 61) Blane CE, Gagnadoux MF, Brunelle F, et al. Doppler ultrasonography in the early postoperative evaluation of renal transplants in children. *Can Assoc Radiol J* 1993; 44: 176-178.
- 62) Saarinen O. Diagnostic value of resistive index of renal transplants in the early postoperative period. *Acta Radiol* 1991; 32: 166-169.
- 63) Di Palo FQ, Rivolta R, Elli A, et al. Effect of cyclosporin A on renal cortical resistances measured by color doppler flowmetry on renal grafts. *Nephron* 1993; 65: 240-244.
- 64) Stevens PE, Gwyther SJ, Hanson ME, et al. Interpretation of duplex Doppler ultrasound in renal transplants in the early postoperative period. *Nephrol Dial Transplant* 1993; 8: 255-258.

- 65) Hollenbeck M, Hetzel GR, Hilbert N, et al. Dopplersonographische Beurteilung der Effektivität einer Abstoßungstherapie nach Nierentransplantation. Dtsch med Wschr 1995; 120: 277-282.
- 66) Rubin JM, Bude RO, Carson PL, et al. Power Doppler US: potentially useful alternative to mean frequency-based color Doppler US. Radiology 1994; 190: 853-856.
- 67) Delorme S, Krix M, Albrecht T. Ultraschallkontrastmittel - Grundlagen und klinische Anwendung. Fortschr Röntgenstr 2006; 178(2): 155-164.
- 68) Desser TS, Jeffrey Jr RB, Lane MJ, Ralls PW. Tissue harmonic imaging: utility in abdominal and pelvic sonography. J Clin Ultrasound 1999; 27(3): 135-142.
- 69) Stiskal M, Steinbach R, Obholzer G, et al. Tissue Harmonic Imaging Sonographie. Wird die Bildqualität bei Routineultraschalluntersuchungen im Abdomen verbessert? Fortschr Röntgenstr 2000; 172(12): 1006-1010.
- 70) Schoelgens C. Native tissue harmonic imaging. Radiologe 1998; 38(5):420-423.
- 71) Shapiro RS, Wagreich J, Parsons RB, et al. Tissue harmonic imaging sonography: evaluation of image quality compared with conventional sonography. Am J Roentgenol 1998, 171(5): 1203-1206.
- 72) Deetjen P, Speckmann EJ. Physiologie. 2. Aufl. München, Wien, Baltimore: Urban & Schwarzenberg, 1994: 89-100.
- 73) Bleck JS, Gebel M, Manns MP. Quantitative Sonographie. Bedeutung und Perspektive. Internist 2000; 41: 10-16.

- 74) Flinn GS. Grey scale colour coding of abdominal B-scan ultrasonogramms. *J Clin Ultrasound* 1975; 3(3): 179-185.
- 75) Thijssen JM, Oosterveld BJ, Wagner RF. Grey level transforms and lesion detectability in echographic images. *Ultrasound Imaging* 1988; 10(3): 171-195.
- 76) Bleck JS, Gebel M, Hebel R, et al. Tissue characterization using intelligent adaptive filter in the diagnosis of diffuse and focal liver disease. *Ultrasound Med Biol* 1994; 20: 521-528.
- 77) Bleck JS, Gebel M, Witt B, et al. Sonography under Daylight conditions. *Ultraschall Med* 1998; 19: 259-264.
- 78) Gebel M. Neue Modifikationen und Verfahren der Sonographie und ihre praktische Bedeutung. *Internist* 2000; 41: 3-9.
- 79) Bleck JS, Gebel M, Satrapa JD, et al. Approches to objective assessment of differences in tissue textures by ultrasound equipment as a contribution to quality assurance. *Ultraschall Med* 1997; 18(6): 238-243.
- 80) Townsend RR, Tomlanowich SJ, Goldstein RB, et al. Combined Doppler and morphologic sonographic evaluation of renal transplant rejection. *J Ultrasound Med* 1990; 9: 199-206.
- 81) Weskott HP. Power US: Phantom measurement of slow flow. *Radiology* 1995; 197: 337.
- 82) Waiser J, Schreiber M, Budde K, et al. Prognostic value of the Banff classification. *Transpl Int* 2000; 13(1): 106-111.
- 83) Teichgräber UKM, Beuter T, Klühs L, et al. Netzplantchnik zur Zeitplanung bei abdominalen Ultraschalluntersuchungen. *Ultraschall Med* 1999; 20: 224-225.

- 84) Pudszuhn A, Marx C, Malich A, et al. Prospective Analysis of Quantification of Contrast Media Enhanced Power Doppler Sonography of Equivocal Breast Lesions. *Fortschr Röntgenstr* 2003; 175: 495-501.
- 85) Dietrich CF. 3D real time contrast enhanced ultrasonography, a new technique. *Fortschr Röntgenstr* 2002; 174: 160-163.
- 86) Hohmann J, Skrok J, Puls R, et al. Characterization of focal liver lesions with contrast-enhanced low MI real time ultrasound and SonoVue. *Fortschr Röntgenstr* 2003; 175(6): 835-843.
- 87) Quaia E, Bertolotto M, Calderan L, et al. US characterization of focal hepatic lesions with intermittent high-acoustic-power mode and contrast material. *Acad Radiol* 2003 10(7): 739-750.
- 88) Basilico R, Blomley MJ, Harvey CJ, et al. Which continuous US scanning mode is optimal for the detection of vascularity in liver lesions when enhanced with a second generation contrast agent? *Eur J Radiol* 2002; 41(3): 184-191.
- 89) Schlosser T, Pohl C, Veltmann C. Feasibility of the flash-replenishment concept in renal tissue: which parameters affect the assessment of the contrast replenishment? *Ultrasound Med Biol* 2001; 27: 937-944.
- 90) Wei K, Le E, Bin JP, et al. Quantification of renal blood flow with contrast-enhanced ultrasound. *J Am Coll Cardiol* 2001; 37(4):1135-1140.
- 91) Girard MS, Mattrey RF, Baker KG, et al. Comparison of Standard and Second Harmonic B-mode Sonography in the Detection of Segmental Renal Infarction with SonographicContrast in a Rabbit Model. *J Ultrasound Med* 2000; 19:185-192.

- 92) Nilsson A. Contrast-enhanced ultrasound of the kidneys. *Eur Radiol* 2004; 8: 104-109.
- 93) Bokor D, Chambers JB, Rees PJ, et al. Clinical safety of SonoVue, a new contrast agent for ultrasound imaging, in healthy volunteers and in patients with chronic obstructive pulmonary disease. *Invest Radiol* 2001; 36(2): 104-109.
- 94) Correas JM, Claudon M, Tranquart F, et al. Contrast-enhanced ultrasonography: renal applications. *J Radiol* 2003; 84(12): 2041-2054.
- 95) Lucidarme O, Franchi-Abella S, Correas JM, et al. Blood flow quantification with contrast-enhanced US: "entrance in the section" phenomenon-phantom and rabbit study. *Radiology* 2003; 228(2): 298-299.
- 96) Nahar T, Li P, Kuersten B, et al. Detection of resting myocardial perfusion defects by SonoVue myocardial contrast echocardiography. *Echocardiography* 2003; 20(6): 511-517.
- 97) Schlosser T, Veltmann C, Lohmaier S, et al. Determination of the renal blood flow in macro- and microcirculation by means of pulse inversion imaging. *Fortschr Röntgenstr* 2004; 176(5): 724-730.
- 98) Bouakaz A, Krenning BJ, Vletter WB, et al. Contrast superharmonic imaging: a feasibility study. *Ultrasound Med Biol* 2003; 29(4): 547-553.
- 99) Robbin ML, Lockhart ME, Barr RG. Renal imaging with ultrasound contrast: current status. *Radiol Clin North Am* 2003; 41(5): 963-978.
- 100) Lefevre F, Correas JM, Briancon S, et al. Contrast-enhanced sonography of the renal transplant using triggered pulse-inversion imaging: preliminary results. *Ultrasound Med Biol* 2002; 28(3): 303-314.

- 101) Krix M, Kauczor HU, Delorme S. Quantification of tissue perfusion with novel ultrasound methods Radiologe 2003; 43(10): 823-830.
- 102) Pollard RE, Sadlowski AR, Bloch SH, et al. Contrast-assisted destruction-replenishment ultrasound for the assessment of tumor microvasculature in a rat model Technol Cancer Res Treat 2002; 1(6): 459-470.
- 103) Argalia G, Cacciamani L, Fazi R, et al. Contrast-enhanced sonography in the diagnosis of renal artery stenosis: comparison with MR-angiography. Radiol Med 2004; 107(3): 208-217.
- 104) Blebea J, Zickler R, Volteas N, et al. Duplex imaging of the renal arteries with contrast enhancement. Vasc Endovascular Surg 2003; 37(6): 429-436.
- 105) Teixeira OU, Bortolotto LA, Siva HB. The contrast-enhanced Doppler ultrasound with perfluorocarbon exposed sonicated albumin does not improve the diagnosis of renal artery stenosis compared with angiography. J Negat Results Biomed 2004; 3(1): 3.
- 106) Ascenti G, Gaeta M, Magno C, et al. Contrast-enhanced second -harmonic sonography in the detection of pseudocapsule in renal cell carcinoma. AJR Am J Roentgenol 2004; 182(6): 1525-1530.
- 107) Kreft B, Schild HH. Zystische Nierenläsionen. Fortschr Röntgenstr 2003; 175(7): 892-903.
- 108) Veltri A, De Fazio G, Malfitana V, et al. Percutaneous US-guided RF thermal ablation for malignant renal tumors: preliminary results in 13 patients. Eur Radiol 2004; 14(12): 2303-2310.