

RESEARCH ARTICLE

The Slavic *NBN* Founder Mutation: A Role for Reproductive Fitness?

Eva Seemanova^{1*}, Raymonda Varon², Jan Vejvalka³, Petr Jarolim⁴, Pavel Seeman⁵, Krystyna H. Chrzanowska⁶, Martin Digweed², Igor Resnick⁷, Ivo Kremensky⁸, Kathrin Saar⁹, Katrin Hoffmann¹⁰, Véronique Dutrannoy², Mohsen Karbasiyan², Mehdi Ghani¹¹, Ivo Barić¹², Mustafa Tekin¹³, Peter Kovacs¹⁴, Michael Krawczak¹⁵, André Reis¹⁶, Karl Sperling^{2‡}, Michael Nothnagel^{17‡*}

1 Department of Clinical Genetics, Institute of Biology and Medical Genetics, 2nd Medical School, Charles University, Prague, Czech Republic, **2** Institute of Medical and Human Genetics, Charité-Universitätsmedizin Berlin, Germany, **3** Department of Informatics, 2nd Medical School, Charles University, Prague, Czech Republic, **4** Department of Pathology, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, United States of America, **5** DNA Laboratory, Department of Pediatric Neurology, 2nd Medical School, Charles University and University Hospital Motol, Prague, Czech Republic, **6** Department of Medical Genetics, The Children's Memorial Health Institute, Warsaw, Poland, **7** Department of Bone Marrow Transplantation and Cancer Immunotherapy, Hadassah-Hebrew University Medical Center, Jerusalem, Israel, **8** National Genetics Laboratory, University Hospital of Obstetrics and Gynecology, Medical University, Sofia, Bulgaria, **9** Max Delbrück Center for Molecular Medicine (MDC), Berlin-Buch, Germany, **10** Institute of Human Genetics, Martin-Luther University Halle-Wittenberg, Halle, Germany, **11** Tanz Centre for Research in Neurodegenerative Diseases, University of Toronto, Toronto, Canada, **12** Department of Paediatrics, University Hospital Center Zagreb and University of Zagreb, School of Medicine, Zagreb, Croatia, **13** Division of Pediatric Molecular Genetics, Ankara University School of Medicine, Ankara, Turkey, **14** Leipzig University Medical Center, IFB AdiposityDiseases, Leipzig, Germany, **15** Institute of Medical Informatics and Statistics, Christian-Albrechts University, Kiel, Germany, **16** Institute of Human Genetics, Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU), Erlangen, Germany, **17** Cologne Center for Genomics, University of Cologne, Köln, Germany

‡ These authors share senior authorship on this work.

* eva.seemanova@lfmotol.cuni.cz (ES); michael.nothnagel@uni-koeln.de (MN)



OPEN ACCESS

Citation: Seemanova E, Varon R, Vejvalka J, Jarolim P, Seeman P, Chrzanowska KH, et al. (2016) The Slavic *NBN* Founder Mutation: A Role for Reproductive Fitness? PLoS ONE 11(12): e0167984. doi:10.1371/journal.pone.0167984

Editor: Ludmila Prokunina-Olsson, National Cancer Institute, UNITED STATES

Received: June 27, 2016

Accepted: November 23, 2016

Published: December 9, 2016

Copyright: © 2016 Seemanova et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was supported by grants from the Sander Stiftung to KS and ES, by grants from the DFG to AR, KS and MD, by the German Ministry of Education and Research (BMBF) through National Genome Research Network (NGFN) grants to MK and MN, by grant 108-1081870-1885 of the Ministry of Science, Education and Sport of the Republic of Croatia to IB, and by grants AZV 12-31899 A and DRO 00064203 from the Czech Ministry of Health to PS.

Abstract

The vast majority of patients with Nijmegen Breakage Syndrome (NBS) are of Slavic origin and carry a deleterious deletion (c.657del5; rs587776650) in the *NBN* gene on chromosome 8q21. This mutation is essentially confined to Slavic populations and may thus be considered a Slavic founder mutation. Notably, not a single parenthood of a homozygous c.657del5 carrier has been reported to date, while heterozygous carriers do reproduce but have an increased cancer risk. These observations seem to conflict with the considerable carrier frequency of c.657del5 of 0.5% to 1% as observed in different Slavic populations because deleterious mutations would be eliminated quite rapidly by purifying selection. Therefore, we propose that heterozygous c.657del5 carriers have increased reproductive success, i.e., that the mutation confers heterozygote advantage. In fact, in our cohort study of the reproductive history of 24 NBS pedigrees from the Czech Republic, we observed that female carriers gave birth to more children on average than female non-carriers, while no such reproductive differences were observed for males. We also estimate that c.657del5 likely occurred less than 300 generations ago, thus supporting the view that the original mutation predated the historic split and subsequent spread of the 'Slavic people'. We surmise that the higher fertility of female c.657del5 carriers reflects a lower miscarriage rate in

The Sorb study was supported by Federal Ministry of Education and Research (BMBF), Germany, (FKZ: 01EO1501) and by grants from the Collaborative Research Center funded by the German Research Foundation (CRC 1052; C01, B03). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

these women, thereby reflecting the role of the *NBN* gene product, nibrin, in the repair of DNA double strand breaks and their processing in immune gene rearrangements, telomere maintenance, and meiotic recombination, akin to the previously described role of the DNA repair genes *BRCA1* and *BRCA2*.

Introduction

Nijmegen Breakage Syndrome (NBS [MIM: 251260]) is a chromosome instability disorder characterized by microcephaly, growth retardation, immunodeficiency, hypersensitivity to X-irradiation, and an exceptionally high risk for lymphoid malignancy [1–3]. The *NBN* gene (also abbreviated *NBS*, *NBS1*, *ATV*, or *AT-V1*, amongst others) was identified as the cause of NBS in 1998 [4–6] and is located on chromosome 8q21 (GRCh38.p2: 89,933,336–89,984,733). The corresponding gene product, nibrin, is part of the trimeric MRE11/RAD50 complex that is involved in the repair of DNA double strand breaks (DSBs) and in their processing in immune gene rearrangements, telomere maintenance, and meiotic recombination (summarized in [7, 8]). This multi-functionality explains the complex NBS phenotype, particularly the very high incidence of malignancies that form the major cause of early death amongst NBS patients [2]. The majority of NBS patients originates from Eastern Europe and is homozygous for a common founder mutation in the *NBN* gene, namely a deletion of nucleotide positions 657 to 661 of the coding sequence in exon 6 (abbreviated as c.657del5; rs587776650) that leads to two truncated fragments, p26- and p70 nibrin [9].

To the best of our knowledge, *NBN* c.657del5 is the only Slavic founder mutation of high prevalence described so far. It obviously predated the medieval migration of the Slavic tribes. Until recently, our insights into the history of the Slavs, the largest population in Eastern and Southeastern Europe, were mainly based upon linguistic and archaeological evidence. Although a common origin of the Slavs is generally accepted, controversy exists about their original homeland and the timing of their historical diversification. During the early medieval waves of migration into Europe in the 5th and 6th century, the Slavs split into the Eastern Slavs (Belarusians, Russian, Ukrainians), the Western Slavs (e.g. Czechs, Poles, Slovaks, Sorbs), and the Southern Slavs (e.g. Bulgarians, Croats, Serbs, Slovenes) (cited in [10, 11]). A recent study of autosomal, mitochondrial and Y-chromosomal data coined the existence of two sub-groups that presumably existed some 1900 years ago, namely the East-West Slavs (from Poland to the Volga river) and the South Slavs (confined to the Balkan Peninsula) [12]. Moreover, the study revealed pronounced genetic similarities between the Czechs and their immediate Germanic neighbors while, further north, the Sorbs and the surrounding Germans were clearly genetically distinct [13]. However, the time of the initial division of the proto-Slavic tribes was not determined. From the evolutionary history of Y-chromosomal STR haplotype R1a, the Russian Plain branch (including proto-Slavic tribes) living in present-day Russia, Ukraine, Belarus, Poland and the Baltic countries is estimated to have appeared about 4,600 years ago [14].

Interestingly, a high frequency of heterozygous c.657del5 carriers has been reported from the Slavic populations of Poland, the Czech Republic/Slovakia (henceforth abbreviated as CS), the Ukraine, and Germany [15, 16]. Notably, heterozygous carriers were particularly prevalent among cancer patients in these populations (reviewed in [17, 18]), an association that was rigorously confirmed in a family-based study carried out with the Index-Test Method [19].

Whilst heterozygous carriers seemingly reproduce, despite their increased cancer risk, no parenthood of a homozygous individual has been reported so far. The high frequency of

c.657del5 carriers in different Slavic populations therefore seems enigmatic and difficult to explain by genetic drift alone. This can readily be illustrated by a simple selection model with relative fitness values of $1-hs$ and $1-s$ for hetero- and homozygous mutation carriers [20]. Assuming $s > 0.9$ and an initial mutation frequency of 0.1%, the latter is expected to decrease slightly (to $\sim 0.085\%$), rather than increase, within 200 generations if $h = 0$. Moreover, the mutation frequency is bound to fall below 10^{-11} over the same time period if $h = 0.1$, i.e. if heterozygous carriers are of slightly reduced reproductive fitness (data not shown).

In the present study, we provide evidence from a family-based cohort study for a reproductive advantage of female heterozygous *c.657del5* carriers that potentially explains the high frequency of this mutation in Eastern Europe by balancing selection. In addition, we estimated *c.657del5* to be approximately 4000 to 6650 years old, using a large collection from unrelated carriers of Eastern, Western and Southern Slavic origin.

Materials and Methods

Reproductive success of *NBN c.657del5* carriers

We assessed the relative reproductive success of heterozygous *NBN c.657del5* carriers relative to their non-carrier relatives. To this end, we drew upon a large collection of families of NBS patients from the Czech Republic and Slovakia that were originally recruited as part of a study to assess the relative cancer risk of carriers and non-carriers [19]. For more than 20 years, one of us (E.S.) has seen virtually all patients with clinically confirmed NBS in these countries. The present study, more specifically, includes 24 families with 344 members for whom comprehensive medical records were available to us. Seven of these families had been counseled and repeatedly interviewed over 10–20 years whilst another seven were regularly contacted for over 20–25 years. The remaining 10 families were diagnosed and followed between 1994 and 2003. Molecular testing after 1998 confirmed that all NBS cases were homozygous for the *NBN c.657del5* mutation.

Starting from the affected child in each pedigree ('index patient'), as many relatives as possible were contacted through the parents of the index patient. Attempts were made to contact all grandparents, their siblings (if alive), aunts, uncles and cousins who lived in the Czech Republic. Relatives were recruited without prior knowledge of their *c.657del5* carrier status. Of the 490 individuals contacted, 344 agreed to participate in our study.

Upon informed consent, oral interviews were conducted via structured questionnaires administered by one and the same person (E.S.). The questionnaires covered socio-demographics (including reproductive history), general medical history, life style and occupational history. Socioeconomic status was inferred from the educational level attained when leaving school. Family history and gynecological as well as obstetric history were recorded for female participants only. The vast majority of interviews took place at the homes of participants and required around 60 minutes each. Obligate heterozygotes were not included so that the interviewer was consistently 'blinded' with respect to the carrier status of their interviewees. After the interview, a blood sample was taken to assess the *NBN* germline mutation *c.657del5* genotype [21].

We compared the number of offspring of *c.657del5* carriers and non-carriers, separately for males and females. Statistical analyses were carried out with Fisher's exact or an unpaired *t*-test, as appropriate.

Age estimation for *NBN c.657del5*

Sampling of individuals. We used the family data available to us to reconstruct the original *NBN c.657del5* haplotype and to estimate the age of the deletion. In addition, we analyzed

Table 1. Individuals and haplotypes used for the c.657del5 age estimation and ancestral haplotype inference.

Origin	Sampled individuals (heteroz.)	c.657del5 carrying chromos.	Unambiguously inferred c.657del5 carrying haplotypes
Austria	1 (1)	1	0
Belgium	1	2	0
Bulgaria	4 (4)	4	0
Czech Republic/Slovakia (CS)	9 (1)	17	11
Croatia	1	2	2
Germany (Germans)	22 (2)	42	4
Germany (Sorbs from Lusatia)	10 (10)	10	3
Italy	1	2	2
Poland	35	70	64
Russia	10 (1)	19	0
Turkey	3	6	6
Ukraine	6	12	0
<i>Total</i>	<i>103 (19)</i>	<i>187</i>	<i>92</i>

Haplotypes were inferred from both homo- and heterozygous deletion carriers.

doi:10.1371/journal.pone.0167984.t001

Guthrie cards from newborn screening programs, including 1502 samples from Bulgaria, 994 from Croatia, and 1035 from Lusatia (Germany), the latter being part of an extensively phenotyped sample from Eastern Germany [13]. We also asked colleagues for additional samples from NBS patients of Eastern, Western, and Southern Slavic origin (Table 1).

Genotyping and reconstruction of haplotypes. Some 84 homozygous and 19 heterozygous *NBN* c.657del5 carriers were genotyped at markers covering a 5.7 Mb region around the deletion on chromosome 8q21. More specifically, 13 microsatellites located between D8S271 and D8S270 were analyzed (Table 2) in addition to 12 SNPs (Table 3). For 187 chromosomes carrying *NBN* c.657del5 (S1 File), 92 haplotypes could be deduced without ambiguity using the available family data (Table 1). These haplotypes formed the basis of our attempt to estimate the age of the mutation. The unambiguous haplotypes and the most likely albeit ambiguous haplotypes are listed in the Supporting Information (S1 File).

Genetic marker distances. We considered two commonly used genetic maps, namely the Rutgers map [22–25] and the high-resolution recombination map from deCODE Genetics [26]. We abstained from involving the genetic map provided by the Marshfield Clinic [27] because it does not resolve large parts of the investigated chromosomal region. Where possible, we used sex-averaged maps. If the available map information did not include genetic distances for particular markers, as was the case for some microsatellites and all SNPs, these distance were interpolated assuming proportionality between physical and genetic distance, and by drawing upon the closest marker of known genetic location on either side of an uncharted marker. Since each marker had to be assigned a unique location for the age estimation software to function properly (see below), we introduced an artificial genetic distance of 10^{-9} cM between adjacent markers if they had the same genetic location according to the map in question (Table 2).

Age estimation. We used the DMLE+ software [28] for estimating the age of *NBN* c.657del5. This software employs a Bayesian algorithm to infer the location and/or age of a founder mutation based upon patterns of linkage disequilibrium and associated demographic parameters [29]. Numerical integration in DMLE+ is done by Monte Carlo Markov Chain

Table 2. Microsatellite markers and one InDel polymorphism used in this study.

Marker	Available alleles	Sample heterozygosity	Physical distance [kb]	Genetic map [cM]	
				Rutgers	Decode
D8S271	89	0.614	0	0	0
D8S273	85	0.494	393.2	0.0000000001	0.0000000001
rs6150693 [‡]	84	0.023	1,020.1	0.6485	0.453
D8S1800	91	0.423	1,176.7	0.7300	0.5100
AFM289	87	0.130	1,586.4	0.8152	0.6662
H3GT	89	0.189	1,947.2	0.8903	0.8038
H2CA	89	0.209	2,307.9	0.9653	0.9414
D8S88	92	0.257	2,330.6	0.9700	0.9500
<i>NBN</i> c.657del5	92	0.000	2,440.2	1.1128	1.0357
H5CA	82	0.137	2,602.0	1.3235	1.1621
D8S1811	91	0.146	2,714.5	1.4700	1.2500
D8S1724	92	0.181	2,901.6	1.4700000001	1.5700
D8S1146	82	0.181	3,504.9	1.8017	1.7019
D8S1618	85	0.134	3,986.7	2.0665	1.8073
D8S270	86	0.716	4,502.3	2.35	1.92

All markers are located on chromosome 8q21. Physical distances according to human genome assembly hg19 (Genome Reference Consortium GRCh37).

[‡]Note that the InDel polymorphism rs6150693 (position 89607483), which affects 12 base-pairs, has been removed from dbSNP on Oct 17, 2013, due to mapping or clustering errors (http://www.ncbi.nlm.nih.gov/SNP/snp_ref.cgi?type=rs&rs=rs6150693), but is listed in UCSC's Genome Browser (<http://genome.ucsc.edu/>).

doi:10.1371/journal.pone.0167984.t002

(MCMC). Note that this software does not estimate the time to the most recent common ancestor (MRCA), but the actual age of the mutation. We initially used DMLE+ assuming a known location of the mutation (Table 2) together with the following parameters: population growth rate 0.01, mutation population frequency 1.0 (since all haplotypes carried the

Table 3. Additional single-nucleotide polymorphism markers at the 92 unambiguously inferred haplotypes in this study.

Marker	Available alleles	Allele on haplotype	Global MAF	Physical distance [kb]	Genetic map [cM]	
					Rutgers	Decode
rs10111232	90	C	0.0004	2,394.3	1.0529	0.9998
rs1063045	92	T	0.3792	2,429.3	1.0986	1.0272
rs1805794	92	G	0.3570	2,437.4	1.1092	1.0335
<i>NBN</i> c.657del5 (rs587776650)	92	c.657del5	-	2,440.2	1.1128	1.0357
rs2234744	92	T	0.3530	2,449.5	1.1249	1.0430
rs709816	92	T	0.3914	2,452.7	1.1291	1.0455
rs1061302	92	C	0.3528	2,465.3	1.1455	1.0553
rs3736639	92	A	0.3792	2,472.3	1.1546	1.0607
rs1063053	92	T	0.3219	2,476.8	1.1605	1.0643
rs13272541	92	C	<0.0002	2,493.4	1.1821	1.0772
rs1000249	86	A	<0.0002	2,554.3	1.2614	1.1248
rs6994202	86	A	0.2899	2,577.8	1.2920	1.1432
rs11994308	86	C	0.0899	3,142.2	1.6022	1.6226

All markers are located on chromosome 8q21. Physical distances according to human genome assembly hg19 (Genome Reference Consortium GRCh37). All SNPs were found to be monomorphic in our sample. **MAF**: minor allele frequency, obtained from dbSNP (<http://www.ncbi.nlm.nih.gov/snp>).

doi:10.1371/journal.pone.0167984.t003

mutation); burn-in iteration 5,000,000; data iterations 10,000,000; number of histogram bars 200; mutation age limits 0–5000 generations; no star-like genealogy; no sequence weights. Subsequently, we considered population growth rates of 0.005 and 0.001 as well in order to assess the effect of this parameter on the age estimate. The R statistical software [30], version 3.2.2, was used for descriptive statistics and for creating graphs. Perl was used for scripting the analysis with DMLE+.

Our study was approved by the Ethics Committee of the 2nd Medical School of Charles University and University Hospital Motol, Prague, Czech Republic. All participants gave signed informed consent, with the exception of the Sorb probands [13]. The respective study was approved by the Ethics Committee of the Leipzig University, Germany, and all participants had given their signed informed consent.

Results

Reproductive success of *NBN* c.657del5 carriers

The reproductive success of female *NBN* c.657del5 carriers was compared to that of their non-carrier relatives in a cohort of 40 heterozygotes and 208 homozygotes for the wild-type allele. The average age was almost the same in the two groups (51.2 vs. 50.9 years, respectively). However, the average number of offspring differed significantly ($P = 0.02$), with 3.03 children born by carriers and 2.36 children born by non-carriers (Table 4). For an age-stratified analysis, we split the cohort into three age groups: <40 years, 41–60 years, and >60 years at the time of the last interview. In all three groups, carriers had more children on average than non-carriers, and this difference was statistically significant in the 41–60 years age group (Table 4). There was no statistically significant difference with respect to school and university attendance between carriers and non-carriers. Furthermore, we also compared the menstrual history of female carriers and non-carriers, characterized by the age at first menstrual period and the age at onset of menopause (females 30 years and older). We observed no significant difference between the two groups in this regard (data not shown). No significant difference in the number of offspring was observed between male mutation carriers and non-carriers (Table 5). Data on survival until reproductive age were not available to us.

The number of spontaneous or induced abortions could be ascertained for 32 of the 40 female carriers and for 145 of the 208 female non-carriers, with both subgroups being of almost the same average age (54.5 and 53.4 years). The relative number of induced abortions was virtually the same whilst the rate of spontaneous abortions was slightly albeit non-significantly lower among carriers (Table 6). The 48 obligate heterozygous females (not included in the above analysis) had 57 living offspring (30 affected by NBS), three fetuses diagnosed with NBS, and 11 pregnancies that were interrupted for other reasons. Two fetuses were lost spontaneously. Of 73 registered pregnancies, only two thus ended spontaneously.

Table 4. Number of offspring of female *NBN* c.657del5 carriers and non-carriers of different age groups.

Age group	Carriers of c.657del5				Non-carriers of c.657del5				P
	N	Average age	Number of offspring	Ratio	N	Average age	Number of offspring	Ratio	
<41	12	32.3	21	1.75	77	32.2	123	1.60	0.59
41–60	17	51.6	62	3.65	61	51.3	160	2.62	0.01
>60	11	71.2	38	3.45	70	71.0	207	2.96	0.47
All groups	40	51.2	121	3.03	208	50.9	490	2.36	0.02

Average age is given in years. The two-tailed P-values were obtained from unpaired t-test.

doi:10.1371/journal.pone.0167984.t004

Table 5. Number of offspring of male *NBN* c.657del5 carriers and non-carriers of different age groups.

Age group	Carriers of c.657del5				Non-carriers of c.657del5				P
	N	Average age	Number of offspring	Ratio	N	Average age	Number of offspring	Ratio	
<41	20	31.5	34	1.70	62	34.9	109	1.76	0.80
41–60	21	50.7	61	2.90	66	51.3	158	2.39	0.25
>60	18	70.1	51	2.83	40	71.2	99	2.48	0.45
All groups	59	50.1	146	2.47	168	50.0	366	2.18	0.20

Average age is given in years. The two-tailed P-values were obtained from unpaired t-test.

doi:10.1371/journal.pone.0167984.t005

Age of the *NBN* c.657del5 mutation

We applied the DMLE+ software to a set of microsatellite markers to estimate the age of *NBN* c.657del5. In order to assess the sensitivity of these analyses to the underlying assumptions, we assumed two different genetic maps and three different population growth rates. With an average population growth of 1% per generation, the mean posterior age estimate was 265 and 267 generations, respectively, for the Rutgers and the deCODE genetic map. This would roughly correspond to approximately 4000 to 6650 years, depending upon the presumed generation time (15 to 25 years; [Table 7](#)). Larger estimates of 343 to 369 and 509 to 698 generations were obtained for growth rates of 0.5% and 0.1%, respectively. These figures would correspond to an age range of 7000 to 12,500 years, assuming a generation time of 20 years ([Table 7](#) and [Fig 1](#)).

Reconstruction of ancestral mutation haplotype

Of the 92 haplotypes that carried *NBN* c.657del5 and that could be deduced without ambiguity ([S1 File](#)), 22 (23.9%) carried identical alleles at all markers between the two most distant microsatellites D8S271 (allele 257) and D8S270 (allele 200). This haplotype ([Table 8](#)) was found in individuals from Poland, Germany, CS and Lusatia (Sorbs) and most likely represents the ancestral haplotype. The remaining 95 haplotypes from 40 homozygous and 15 heterozygous c.657del5 carriers (out of a total of 187 chromosomes) could not be resolved without ambiguity and are best guesses ([S1 File](#)). Among these haplotypes, we observed the putative ancestral haplotype not only in individuals from the countries mentioned above, but also from Bulgaria, Russia, and the Ukraine. Some 30% of chromosomes carrying the founder mutation showed the deduced founder haplotype ([Table 8](#)). Moreover, all 187 chromosomes carrying c.657del5 showed the same SNP alleles ([Table 3](#)). Therefore, the deletion is not only confined to individuals of Slavic origin but is most likely due to a single mutational event. All differences observed

Table 6. Reproductive histories for the subset of female *NBN* c.657del5 carriers and non-carriers who were older than 30 years, ascertained by the index test method and had information on the number of spontaneous or induced abortions available.

	Carriers of c.657del5	Non-carriers of c.657del5	P
Number of mothers	32	145	
Average age (years)	54.5	53.4	
All registered pregnancies	120	436	
Number of living offspring (proportion of all pregnancies)	102 (85.0%)	351 (80.5%)	0.46
Number of spontaneous abortions (proportion of all pregnancies)	7 (5.8%)	41 (9.4%)	
Number of induced abortions (proportion of all pregnancies)	11 (9.2%)	44 (10.1%)	

The two-tailed P-values were obtained from Fisher’s exact test.

doi:10.1371/journal.pone.0167984.t006

Table 7. Age estimates for the *NBN* c.657del5 deletion.

	Years per generation	Mean (95% credible interval)	
		Rutgers map	Decode map
Assumed population growth of 1.0%			
Generations		264.8 (149.5–512.4)	267.2 (143.9–475.5)
Years	15	3972.0 (2241.8–7686.1)	4008.0 (2158.5–7131.8)
	20	5296.0 (2989.1–10,248.2)	5344.0 (2878.1–9509.1)
	25	6620.0 (3736.4–12,810.2)	6680.0 (3597.6–11,886.4)
Assumed population growth of 0.5%			
Generations		369.1 (186.4–702.0)	343.0 (185.4–759.2)
Years	15	5536.5 (2796.3–10,530.2)	5145.0 (2781.3–11,387.7)
	20	7382.0 (3728.4–14,040)	6860.0 (3708.4–15,183.6)
	25	9227.5 (4660.5–17,550.4)	8575.0 (4635.5–18,979.6)
Assumed population growth of 0.1%			
Generations		508.8 (166.9–1470.8)	698.0 (185.3–2261.6)
Years	15	7632.0 (2504.0–22,062.0)	10,470.0 (2779.3–33,924.3)
	20	10,176.0 (3338.6–29,416.0)	13,960.0 (3705.7–45,232.4)
	25	12,720 (4173.3–36,770.0)	17,450.0 (4632.1–56,540.5)

doi:10.1371/journal.pone.0167984.t007




Fig 1. Posterior distribution of age estimates for *NBN* c.657del5. Box plots of the posterior age distribution refer to a population growth of 1.0%, 0.5% and 0.1%, respectively, and two genetic maps.

doi:10.1371/journal.pone.0167984.g001

Table 8. Illustration of past recombination events in NBN c.657del5 deletion carrying haplotypes.

Origin	D8S271	D8S273	InsDel	D8S1800	AFM289	H3GT	H2CA	D8S88	NBN	H5CA	D8S1811	D8S1724	D8S1146	D8S1618	D8S270
aH	257	158	ins	145	189	216	232	90	c.657del5	189	108	142	258	274	200
Po	261	164		139						185	106	146	262		186
Po	266										82	138	258		196
Po	267	160		141											
Po	255	156		139											
Po										185	106	136	262		196
Po	255	154			191	196	230						262		186
Po	265			139						185		140	262		188
Po	255	160		139	191					185	106	140	264		186
Po					187	210	236	76							
Po	259	154		139											
Po	259	154		139											
Po	259	154		139											
Po	259	154		139											
Po	265	160		139											
Po												140	262		194
Po												140			192
Po	267	160	del												
Po	255	156		139											
Cr	255	156		139										286	186
Ru	255	160		139											
Ru	255	160		139											
Ru	259			139	191						100	136	268		194
Ru	265	160		139	191										
Ru	261	160		139	191										
Ru	261	160		139	187	208									
Ru	255	160		141	191										
CS	267	154		141		210	236								
CS										187	106				194
Uk	265	160		141	191										
Uk	265	160		141	191										
Uk	255	154		139	193	210							262		198
Uk	265			139	187										
Uk	265	154		139											
So															
Bu	265	156		139	191										190
Ge	263	154		141	191	210	236	86							

(Continued)

Table 8. (Continued)

Origin	D8S271	D8S273	InsDel	D8S1800	AFM289	H3GT	H2CA	D8S88	NBN	H5CA	D8S1811	D8S1724	D8S1146	D8S1618	D8S270
Ge	166	166		141	191										
Ge	265	154													
Ge	261	154		141											
Ge	255	154		141	191										
Ge															
Ge	265	154		145	189	196	230	96							
Ge	261	156		139	191										
Ge	255	162		143											
Ge	263	154		141											
Ge	261	154		139											
Ge	255		del	139	191	218									
Au	255	156	del	139											
It		166		139	191	214									
Tu	261														194
Tu	261														194

aH: Reconstructed ancestral haplotype around the NBN c.657del5 mutation. The ancestral haplotype is based on the analysis of 13 microsatellite markers and one ins/del polymorphism (see Table 2). The 12 analyzed SNPs (Table 3) were located between markers D8S88 and D8S1146. Alleles deviating from the ancestral haplotype are specified. **Origin:** Po (Poland), Cr (Croatia), Ru (Russia), CS (Czech Republic/Slovakia), Uk (Ukraine), So (Sorbs), Bu (Bulgaria), Ge (Germany), Au (Austria), I (Italy), and Tu (Turkey). **Italicized origin:** haplotypes that could not be inferred without ambiguity. **Alleles in bold/shaded areas:** likely past recombination events, assuming a single event only (see main text for details).

doi:10.1371/journal.pone.0167984.t008

for individual microsatellite alleles are explicable by past recombination or mutation owing to the non-negligible recombination rate between the STR markers (Table 2) and the high mutation rates of microsatellites of 10^{-4} to 10^{-2} per generation.

We next aimed at inferring likely recombination events using an ad-hoc approach that postulated a recombination if two or more adjacent markers in the flanking regions of the original haplotype differed from the putative ancestral alleles. Using this approach, we identified more than 50 likely recombination events, most of them representing single events (Table 8). A rare deletion polymorphism was documented in one chromosome from Germany (255-158-del-139 at D8S271-D8S273-rs6150693-D8S1800) and one chromosome from Austria (255-156-del-139 at D8S271-D8S273-rs6150693-D8S1800) as well as in one Polish individual (267-160-del-145 at D8S271-D8S273-rs6150693-D8S1800). The flanking microsatellites on two chromosomes from Turkey shared the same alleles (261 at D8S271, 194 at D8S270) that were different from the original alleles (257 at D8S271, 200 at D8S270), rendering a common origin of the respective disease mutations rather likely.

Discussion

We estimated the age of the *NBN* c.657del5 founder mutation and sought for an explanation of its widespread occurrence in Slavic populations. Our calculations place the original mutation at around 266 generations ago assuming a growth rate of 1% per generation, and at around 356 generations ago with a rate of 0.5%. These estimates translate into roughly 5300 to 7100 years, given a generation time of 20 years. These estimates are consistent with a previous assessment [14] of the age of the R1a Y-STR haplotype (4600 years), which is indicative of the Russian Plain branch (including proto-Slavic tribes) and which is present in Russia, Ukraine, Belarus, Poland and the Baltic countries. Furthermore, our estimates are in concordance with the proposed time of the split between the Proto-Balto-Slavic languages and the Indo-European tree around 4500–7000 YBP, but would predate the proposed split between Slavic and Baltic branches between 3500 and 2500 YBP (summarized in [12]). While this does not preclude the presence of *NBN* c.657del5 at low frequency in the steppe Yamnaya people before their admixture with Central and Western Europeans around 4500 YBP [46], our findings render far more likely a mutation origin in the ancestor population of all Slavic peoples, either before or after its admixture with the Yamnaya. Unfortunately, our data do not allow formal distinction between these two scenarios.

It should be noted that DMLE+ does not estimate the time to the most recent common ancestor (MRCA) of extant deletion carriers, but the actual mutational age. Thus, the deletion may have existed in the population at low frequency for some time before the MRCA. As regards the underlying population growth rate, we deem 1% to be a realistic assumption. While it is still possible that the true rate may have been closer to the (commonly used) value of 0.5%, rates of 0.1% or less appear unrealistic in view of the massive population expansion that took place during the late Neolithic and the Bronze Age. That different age estimates resulted from the use of two genetic maps is explicable in terms of slightly larger genetic distances in the Rutgers map, rendering recombination more likely and reducing the time span required to explain an observed haplotype pattern, compared to the deCODE map.

While the *NBN* c.657del5 mutation is found in all Slavic populations, a rather high heterozygote frequency (0.5–1.0%) has been reported for Ukraine (Eastern Slavs), Poland, CS, and Sorbs in Germany (Western Slavs), and for Bulgaria (Southern Slavs) (Table 9). These figures may still represent underestimates, because all of the countries are characterized by admixture between the Slavic and previous indigenous population. They also experienced massive

Table 9. Frequency of heterozygous *NBN* c.657del5 deletion carriers in different populations.

Country	Absolute number	Relative number	Allele frequency	Reference
Poland				
Mazowsze	10/1,620	1/162 ^a	0.31%	Steffen et al. 2004[31]
Gdansk	21/4,000	1/190 ^a	0.26%	Kanka et al. 2007[32]
Szczecin	3/530	1/176 ^b	0.28%	Gorski et al. 2003[33]
Szczecin	9/1,500	1/167 ^{a,b}	0.30%	Cybulski et al. 2004[34]
Malopolska	12/2,274	1/190 ^a	0.26%	Varon et al. 2000[15]
Wielkopolska	16/2,090	1/131 ^a	0.38%	Ziolkowska et al. 2006[35]
var. Voivodships	6/1,000	1/167 ^a	0.30%	Chrzanowska et al. 2006 [36]
Czech Republic				
	8/1,234	1/154 ^a	0.33%	Varon et al. 2000[15]
	4/383 (34 y.)	1/96 ^b	0.52%	Drabek et al. 2002[37]
	5/1,411	1/282 ^b	0.18%	Pardini et al. 2009[38]
	2/915	1/457 ^b	0.11%	Mateju et al. 2012[39]
Ukraine				
	5/908	1/182 ^a	0.28%	Varon et al. 2000[15]
Belarus				
Minsk	1/1,014	1/1,014	0.05%	Bogdanova et al. 2008[40]
Russia				
	0/548	0/548 ^c	0.09%	Resnick et al. 2003[41]
	2/348 (38.6 y.)	1/174 ^b	0.29%	Buslov et al. 2005[42]
	0/344 (80.5 y.)	0/344 ^b	0.15%	Buslov et al. 2005[42]
Bulgaria				
	9/1002	1/111 ^a	0.45%	this report
Croatia				
	2/994	1/497 ^a	0.10%	this report
Slovakia				
	3/2,996	1/999 ^a	0.05%	Seemanova et al. 2004[43]
Germany				
not specified	1/886	1/886 ^b	0.06%	Carlomagno et al. 1999[44]
Lusatia (Sorbs)	5/1,035	1/207 ^b	0.24%	this report
Lusatia (Sorbs)	5/170	1/34 ^b	1.49%	Maurer et al. 2010[16]
Northeast Bavaria	6/10,656	1/176 ^a	0.28%	Maurer et al. 2010[16]
Berlin	1/990	1/990 ^a	0.05%	Maurer et al. 2010[16]
Hannover	0/1,017	0/1,017 ^b	0.05%	Bogdanova et al. 2008[40]
China				
	0/804	0/804 ^b	0.06%	He et al. 2012[45]

Study design:

^a newborn;

^b adults (average age);

^c not specified.

doi:10.1371/journal.pone.0167984.t009

population movement after World War II. Thus, while the heterozygote frequency was estimated to be 1 in 167 (9 of 1502) in Bulgaria when based upon a random collection of Guthrie cards, it increased notably to 1 in 111 (9 of 1002) when the analysis was confined to cards from individuals with Slavic surnames.

The high frequency of c.657del5 carriers observed in Northeast Bavaria has been explained by local immigration of Slavic people between the 6th and 9th century [16]. On the other hand, the rather low heterozygote frequency in Slovakia was completely unexpected because of the high frequency of c.657del5 homozygotes, higher than in the adjacent Czech Republic. This discrepancy has been explained by the presence of traditional population isolates in Slovakia and by difficulties in collecting representative samples [43].

Transmission distortion has been put forward as an explanation of the abundance of c.657del5 (reviewed in [47]). Unfortunately, our data do not allow validation of this idea. Two other explanations for the persistence of a deleterious mutation in a population are chance (genetic drift) and reproductive advantage. Generally speaking, genetic drift explains well the increase in frequency of disease-causing mutations in local populations of recent origin, but does rather less convincingly so for a consistent trend across old and geographically separated populations [48]. On the other hand, balancing selection, involving a heterozygote advantage due to higher reproductive fitness, is a more likely explanation for the persistence of a deleterious founder mutation, thereby complementing the scarce evidence in humans (reviewed in [49]).

Based upon our study of the reproductive history of individuals from the Czech Republic, we found that female heterozygous *NBN* c.657del5 carriers gave birth to more children on average than non-carriers, pointing to a higher fertility of the former. No significant differences were observed between male carriers and non-carriers. Of note, our analyses of the number of offspring and of the reproductive histories partially included individuals from the same families. However, since we conditioned our analyses on mutation carrier status, observations in these analyses were conditionally independent and the applied statistical tests remained valid under the assumption of negligible intra-familial, i.e. block-like, effects other than mutation carrier-ship.

Clearly, many factors influence reproductive behaviour in modern societies. In our study, participants were interviewed before their *NBN* c.657del5 genotypes were known, and therefore the interviewer was blind to their *NBN* status. Moreover, the control homozygotes for the wild-type allele came from the same families. We found no evidence for differences between the two groups with respect to age at first menstrual period or at onset of menopause. However, we observed that the increase in offspring number was paralleled by a slight decrease in the number of spontaneous, not induced, abortions. Interestingly, a higher birth rate and lower spontaneous abortion frequency were also reported for female carriers of deleterious mutations in the DNA repair genes *BRCA1* and *BRCA2* [50–52]. From an evolutionary point of view, this link is explicable in terms of genetic variation in DNA repair genes that predisposes to cancer causing better survival under adverse environmental conditions [53].

The *NBN* gene is expressed in human germ cells. Immunofluorescent localisation showed that nibrin is specifically found at meiotic telomeres [54]. In all organisms tested so far, nibrin is required for meiotic recombination (reviewed in [55]). Moreover, it physically interacts with *BRCA1* at the centrosomes and is obviously involved in mitotic centrosome maintenance [56]. Consequently, NBS patients show an impaired development of gonads and ovarian failure in females [2]. The analysis of a humanized *NBN* mouse model provides direct evidence, that the development of male germ cells is somewhat retarded while most female oocytes are arrested at pachytene [57].

Human female meiosis, but not male meiosis, is a highly error-prone process. Fewer than 30% of naturally fertilised human ova survive to term. The vast majority of losses occur before clinical recognition of pregnancy [58, 59]. The leading cause of these losses is a high rate of aneuploidies due to spontaneous non-disjunction during maternal meiosis [60–62]. Moreover, aneuploidies are also the major cause of spontaneous abortions in humans, affecting 10 to 15%

of all recognized pregnancies. In principle, a slightly lower rate of non-disjunction in female carriers of the founder mutation could explain their higher fertility. Although highly speculative at this point, this assertion can be tested empirically.

The unusually high rate of human maternal meiotic non-disjunction and its further increase with age is explained by an interplay between unique features of oogenesis, i.e. recombination failure, loss of cohesion between sister chromatids, and sister centromeres in combination with a relaxed spindle checkpoint [63, 64]. This anomalous chromosomal behaviour still persists during the first mitotic divisions [65] and is paralleled by diminished mRNA levels of several checkpoint proteins and also of cohesin (reviewed in [66]). Carriers of the *NBN* c.657del5 mutation have a distinct gene expression phenotype with about the same number of genes up- and down regulated [67]. Moreover, based upon SDS-PAGE and mass spectrometry, it was shown that the two truncated proteins, p26 and p70, synthesized by carriers of the mutation, bind to proteins that do not interact with full-length nibrin [68]. It is tempting to speculate that these effects are also relevant in maternal germ cells and could explain the increase in fecundity of carriers. Thus, in future experiments, the interactors of these fragments should be studied in oocytes of females with the carrier mutation. In our view, the humanized *NBN* mouse would be an ideal animal model to undertake such studies.

In summary, we presented evidence that the Slavic founder mutation *NBN* c.657del5, which has been implicated in Nijmegen Breakage Syndrome and cancer, may actually confer a reproductive advantage for female carriers due to higher fertility. Our age estimate for this mutation of less than 300 generations for this mutation implies that such an advantage could well explain the allele frequency of between 0.5% and 1.0% in Slavic populations.

Supporting Information

S1 File. Set of 92 haplotypes that could be inferred without ambiguity (lines 97–188) and that formed the basis for the age estimation of the *NBN* c.657del5 deletion plus 95 haplotypes that are best guesses (lines 2–96).

(TXT)

Acknowledgments

We thank the Czechoslovakian Self Support Group of Chromosomal Instability Syndromes for their collaboration and Mrs. Anna Baumgartnerová, Markéta Skalová and Kateřina Gebertová for technical assistance. We also thank Dr. Anke Tönjes and Prof. Michael Stummvoll (University of Leipzig) for providing the samples of the Sorb study.

Author Contributions

Conceptualization: ES K. Sperling M. Krawczak MN.

Data curation: ES RV MN.

Formal analysis: K. Sperling M. Krawczak MN.

Funding acquisition: ES K. Sperling AR MD M. Krawczak MN IB PS KH.

Investigation: ES RV JV PJ PS KHC MD IR IK K. Saar KH VD M. Karbasiyan MG IB MT PK AR K. Sperling.

Methodology: ES K. Sperling M. Krawczak MN.

Project administration: ES K. Sperling MN.

Resources: ES RV JV PJ PS KHC MD IR IK K. Saar KH VD M. Karbasiyan MG IB MT PK AR K. Sperling M. Krawczak MN.

Supervision: ES K. Sperling.

Visualization: ES K. Sperling M. Krawczak MN.

Writing – original draft: K. Sperling M. Krawczak MN.

References

1. Digweed M, Sperling K. Nijmegen breakage syndrome: clinical manifestation of defective response to DNA double-strand breaks. *DNA Repair (Amst)*. 2004; 3(8–9):1207–17. Epub 2004/07/29.
2. Chrzanowska KH, Gregorek H, Dembowska-Baginska B, Kalina MA, Digweed M. Nijmegen breakage syndrome (NBS). *Orphanet J Rare Dis*. 2012; 7:13. doi: [10.1186/1750-1172-7-13](https://doi.org/10.1186/1750-1172-7-13) PMID: [22373003](https://pubmed.ncbi.nlm.nih.gov/22373003/)
3. Digweed M, Reis A, Sperling K. Nijmegen breakage syndrome: consequences of defective DNA double strand break repair. *Bioessays*. 1999; 21(8):649–56. Epub 1999/08/10. doi: [10.1002/\(SICI\)1521-1878\(199908\)21:8<649::AID-BIES4>3.0.CO;2-O](https://doi.org/10.1002/(SICI)1521-1878(199908)21:8<649::AID-BIES4>3.0.CO;2-O) PMID: [10440861](https://pubmed.ncbi.nlm.nih.gov/10440861/)
4. Varon R, Vissinga C, Platzer M, Cerosaletti KM, Chrzanowska KH, Saar K, et al. Nibrin, a novel DNA double-strand break repair protein, is mutated in Nijmegen breakage syndrome. *Cell*. 1998; 93(3):467–76. Epub 1998/05/20. PMID: [9590180](https://pubmed.ncbi.nlm.nih.gov/9590180/)
5. Carney JP, Maser RS, Olivares H, Davis EM, Le Beau M, Yates JR 3rd, et al. The hMre11/hRad50 protein complex and Nijmegen breakage syndrome: linkage of double-strand break repair to the cellular DNA damage response. *Cell*. 1998; 93(3):477–86. PMID: [9590181](https://pubmed.ncbi.nlm.nih.gov/9590181/)
6. Matsuura S, Tsuchi H, Nakamura A, Kondo N, Sakamoto S, Endo S, et al. Positional cloning of the gene for Nijmegen breakage syndrome. *Nat Genet*. 1998; 19(2):179–81. doi: [10.1038/549](https://doi.org/10.1038/549) PMID: [9620777](https://pubmed.ncbi.nlm.nih.gov/9620777/)
7. Yuan J, Chen J. MRE11-RAD50-NBS1 complex dictates DNA repair independent of H2AX. *J Biol Chem*. 2010; 285(2):1097–104. doi: [10.1074/jbc.M109.078436](https://doi.org/10.1074/jbc.M109.078436) PMID: [19910469](https://pubmed.ncbi.nlm.nih.gov/19910469/)
8. Lamarche BJ, Orazio NI, Weitzman MD. The MRN complex in double-strand break repair and telomere maintenance. *FEBS Lett*. 2010; 584(17):3682–95. doi: [10.1016/j.febslet.2010.07.029](https://doi.org/10.1016/j.febslet.2010.07.029) PMID: [20655309](https://pubmed.ncbi.nlm.nih.gov/20655309/)
9. Maser RS, Zinkel R, Petrini JH. An alternative mode of translation permits production of a variant NBS1 protein from the common Nijmegen breakage syndrome allele. *Nat Genet*. 2001; 27(4):417–21. doi: [10.1038/86920](https://doi.org/10.1038/86920) PMID: [11279524](https://pubmed.ncbi.nlm.nih.gov/11279524/)
10. Rebala K, Mikulich AI, Tsybovsky IS, Sivakova D, Dzapinkova Z, Szczerkowska-Dobosz A, et al. Y-STR variation among Slavs: evidence for the Slavic homeland in the middle Dnieper basin. *J Hum Genet*. 2007; 52(5):406–14. doi: [10.1007/s10038-007-0125-6](https://doi.org/10.1007/s10038-007-0125-6) PMID: [17364156](https://pubmed.ncbi.nlm.nih.gov/17364156/)
11. Mielnik-Sikorska M, Daca P, Malyarchuk B, Derenko M, Skonieczna K, Perkova M, et al. The history of Slavs inferred from complete mitochondrial genome sequences. *PLoS One*. 2013; 8(1):e54360. doi: [10.1371/journal.pone.0054360](https://doi.org/10.1371/journal.pone.0054360) PMID: [23342138](https://pubmed.ncbi.nlm.nih.gov/23342138/)
12. Kushniarevich A, Utevska O, Chuhryaeva M, Agdzhoyan A, Dibirova K, Uktverlyte I, et al. Genetic Heritage of the Balto-Slavic Speaking Populations: A Synthesis of Autosomal, Mitochondrial and Y-Chromosomal Data. *PLoS One*. 2015; 10(9):e0135820. doi: [10.1371/journal.pone.0135820](https://doi.org/10.1371/journal.pone.0135820) PMID: [26332464](https://pubmed.ncbi.nlm.nih.gov/26332464/)
13. Veeramah KR, Tonjes A, Kovacs P, Gross A, Wegmann D, Geary P, et al. Genetic variation in the Sorbs of eastern Germany in the context of broader European genetic diversity. *Eur J Hum Genet*. 2011; 19(9):995–1001. doi: [10.1038/ejhg.2011.65](https://doi.org/10.1038/ejhg.2011.65) PMID: [21559053](https://pubmed.ncbi.nlm.nih.gov/21559053/)
14. Rozhanskii I. L. K AA. Haplogroup R1a, its subclades and branches in Europe during the last 9000 years. *Advances in Anthropology*. 2012; 2(3):139–56.
15. Varon R, Seemanova E, Chrzanowska K, Hnateyko O, Piekutowska-Abramczuk D, Krajewska-Walasek M, et al. Clinical ascertainment of Nijmegen breakage syndrome (NBS) and prevalence of the major mutation, 657del5, in three Slav populations. *Eur J Hum Genet*. 2000; 8(11):900–2. Epub 2000/11/28. doi: [10.1038/sj.ejhg.5200554](https://doi.org/10.1038/sj.ejhg.5200554) PMID: [11093281](https://pubmed.ncbi.nlm.nih.gov/11093281/)
16. Maurer MH, Hoffmann K, Sperling K, Varon R. High prevalence of the NBN gene mutation c.657-661del5 in Southeast Germany. *J Appl Genet*. 2010; 51(2):211–4. PMID: [20453309](https://pubmed.ncbi.nlm.nih.gov/20453309/)
17. Demuth I, Digweed M. The clinical manifestation of a defective response to DNA double-strand breaks as exemplified by Nijmegen breakage syndrome. *Oncogene*. 2007; 26(56):7792–8. doi: [10.1038/sj.onc.1210876](https://doi.org/10.1038/sj.onc.1210876) PMID: [18066092](https://pubmed.ncbi.nlm.nih.gov/18066092/)
18. Gao P, Ma N, Li M, Tian QB, Liu DW. Functional variants in NBS1 and cancer risk: evidence from a meta-analysis of 60 publications with 111 individual studies. *Mutagenesis*. 2013; 28(6):683–97. doi: [10.1093/mutage/get048](https://doi.org/10.1093/mutage/get048) PMID: [24113799](https://pubmed.ncbi.nlm.nih.gov/24113799/)

19. Seemanova E, Jarolim P, Seeman P, Varon R, Digweed M, Swift M, et al. Cancer risk of heterozygotes with the NBN founder mutation. *J Natl Cancer Inst*. 2007; 99(24):1875–80. doi: [10.1093/jnci/djm251](https://doi.org/10.1093/jnci/djm251) PMID: [18073374](https://pubmed.ncbi.nlm.nih.gov/18073374/)
20. Gillespie JH. *Population genetics: a concise guide*. 2nd ed. Baltimore, MD, USA: The Johns Hopkins University Press; 2004.
21. Seeman P, Gebertova K, Paderova K, Sperling K, Seemanova E. Nijmegen breakage syndrome in 13% of age-matched Czech children with primary microcephaly. *Pediatr Neurol*. 2004; 30(3):195–200. Epub 2004/03/23. doi: [10.1016/j.pediatrneurol.2003.07.003](https://doi.org/10.1016/j.pediatrneurol.2003.07.003) PMID: [15033202](https://pubmed.ncbi.nlm.nih.gov/15033202/)
22. Kong X, Matisse TC. MAP-O-MAT: internet-based linkage mapping. *Bioinformatics*. 2005; 21(4):557–9. doi: [10.1093/bioinformatics/bti024](https://doi.org/10.1093/bioinformatics/bti024) PMID: [15374870](https://pubmed.ncbi.nlm.nih.gov/15374870/)
23. Lander ES, Green P. Construction of multilocus genetic linkage maps in humans. *Proc Natl Acad Sci U S A*. 1987; 84(8):2363–7. PMID: [3470801](https://pubmed.ncbi.nlm.nih.gov/3470801/)
24. Matisse TC, Chen F, Chen W, De La Vega FM, Hansen M, He C, et al. A second-generation combined linkage physical map of the human genome. *Genome Res*. 2007; 17(12):1783–6. doi: [10.1101/gr.7156307](https://doi.org/10.1101/gr.7156307) PMID: [17989245](https://pubmed.ncbi.nlm.nih.gov/17989245/)
25. Matisse TC, Gitlin JA. MAP-O-MAT: marker-based linkage mapping on the World Wide Web. *Am J Hum Genet*. 1999; 65(4):A435.
26. Kong A, Gudbjartsson DF, Sainz J, Jonsdottir GM, Gudjonsson SA, Richardsson B, et al. A high-resolution recombination map of the human genome. *Nat Genet*. 2002; 31(3):241–7. Epub 2002/06/08. doi: [10.1038/ng917](https://doi.org/10.1038/ng917) PMID: [12053178](https://pubmed.ncbi.nlm.nih.gov/12053178/)
27. Broman KW, Murray JC, Sheffield VC, White RL, Weber JL. Comprehensive human genetic maps: individual and sex-specific variation in recombination. *Am J Hum Genet*. 1998; 63(3):861–9. Epub 1998/08/27. doi: [10.1086/302011](https://doi.org/10.1086/302011) PMID: [9718341](https://pubmed.ncbi.nlm.nih.gov/9718341/)
28. Reeve JP, Rannala B. DMLE+: Bayesian linkage disequilibrium gene mapping. *Bioinformatics*. 2002; 18(6):894–5. PMID: [12075030](https://pubmed.ncbi.nlm.nih.gov/12075030/)
29. Rannala B, Reeve JP. High-resolution multipoint linkage-disequilibrium mapping in the context of a human genome sequence. *Am J Hum Genet*. 2001; 69(1):159–78. doi: [10.1086/321279](https://doi.org/10.1086/321279) PMID: [11410841](https://pubmed.ncbi.nlm.nih.gov/11410841/)
30. R Development Core Team. *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria 2015.
31. Steffen J, Varon R, Mosor M, Maneva G, Maurer M, Stumm M, et al. Increased cancer risk of heterozygotes with NBS1 germline mutations in Poland. *Int J Cancer*. 2004; 111(1):67–71. Epub 2004/06/09. doi: [10.1002/ijc.20239](https://doi.org/10.1002/ijc.20239) PMID: [15185344](https://pubmed.ncbi.nlm.nih.gov/15185344/)
32. Kanka C, Brozek I, Skalska B, Siemiatkowska A, Limon J. Germline NBS1 mutations in families with aggregation of Breast and/or ovarian cancer from north-east Poland. *Anticancer Res*. 2007; 27(4C):3015–8. PMID: [17695489](https://pubmed.ncbi.nlm.nih.gov/17695489/)
33. Gorski B, Debniak T, Masojc B, Mierzejewski M, Medrek K, Cybulski C, et al. Germline 657del5 mutation in the NBS1 gene in breast cancer patients. *Int J Cancer*. 2003; 106(3):379–81. doi: [10.1002/ijc.11231](https://doi.org/10.1002/ijc.11231) PMID: [12845677](https://pubmed.ncbi.nlm.nih.gov/12845677/)
34. Cybulski C, Gorski B, Debniak T, Gliniewicz B, Mierzejewski M, Masojc B, et al. NBS1 is a prostate cancer susceptibility gene. *Cancer Res*. 2004; 64(4):1215–9. PMID: [14973119](https://pubmed.ncbi.nlm.nih.gov/14973119/)
35. Ziolkowska I, Mosor M, Nowak J. Regional distribution of heterozygous 657del5 mutation carriers of the NBS1 gene in Wielkopolska province (Poland). *J Appl Genet*. 2006; 47(3):269–72. doi: [10.1007/BF03194635](https://doi.org/10.1007/BF03194635) PMID: [16877808](https://pubmed.ncbi.nlm.nih.gov/16877808/)
36. Chrzanowska KH, Piekutowska-Abramczuk D, Popowska E, Gladkowska-Dura M, Maldyk J, Syczewska M, et al. Carrier frequency of mutation 657del5 in the NBS1 gene in a population of Polish pediatric patients with sporadic lymphoid malignancies. *Int J Cancer*. 2006; 118(5):1269–74. doi: [10.1002/ijc.21439](https://doi.org/10.1002/ijc.21439) PMID: [16152606](https://pubmed.ncbi.nlm.nih.gov/16152606/)
37. Drabek J, Hajduch M, Gojova L, Weigl E, Mihal V. Frequency of 657del(5) mutation of the NBS1 gene in the Czech population by polymerase chain reaction with sequence specific primers. *Cancer Genet Cytogenet*. 2002; 138(2):157–9. PMID: [12505263](https://pubmed.ncbi.nlm.nih.gov/12505263/)
38. Pardini B, Naccarati A, Polakova V, Smerhovsky Z, Hlavata I, Soucek P, et al. NBN 657del5 heterozygous mutations and colorectal cancer risk in the Czech Republic. *Mutat Res*. 2009; 666(1–2):64–7. doi: [10.1016/j.mrfmmm.2009.04.004](https://doi.org/10.1016/j.mrfmmm.2009.04.004) PMID: [19393249](https://pubmed.ncbi.nlm.nih.gov/19393249/)
39. Mateju M, Kleiblova P, Kleibl Z, Janatova M, Soukupova J, Ticha I, et al. Germline mutations 657del5 and 643C>T (R215W) in NBN are not likely to be associated with increased risk of breast cancer in Czech women. *Breast Cancer Res Treat*. 2012; 133(2):809–11. doi: [10.1007/s10549-012-2049-x](https://doi.org/10.1007/s10549-012-2049-x) PMID: [22491912](https://pubmed.ncbi.nlm.nih.gov/22491912/)

40. Bogdanova N, Schurmann P, Waltes R, Feshchenko S, Zalutsky IV, Bremer M, et al. NBS1 variant I171V and breast cancer risk. *Breast Cancer Res Treat.* 2008; 112(1):75–9. doi: [10.1007/s10549-007-9820-4](https://doi.org/10.1007/s10549-007-9820-4) PMID: [18049891](https://pubmed.ncbi.nlm.nih.gov/18049891/)
41. Resnick IB, Kondratenko I, Pashanov E, Maschan AA, Karachunsky A, Togoiev O, et al. 657del5 mutation in the gene for Nijmegen breakage syndrome (NBS1) in a cohort of Russian children with lymphoid tissue malignancies and controls. *Am J Med Genet A.* 2003; 120A(2):174–9. doi: [10.1002/ajmg.a.20188](https://doi.org/10.1002/ajmg.a.20188) PMID: [12833396](https://pubmed.ncbi.nlm.nih.gov/12833396/)
42. Buslov KG, Iyevleva AG, Chekmariova EV, Suspitsin EN, Togo AV, Kuligina E, et al. NBS1 657del5 mutation may contribute only to a limited fraction of breast cancer cases in Russia. *Int J Cancer.* 2005; 114(4):585–9. doi: [10.1002/ijc.20765](https://doi.org/10.1002/ijc.20765) PMID: [15578693](https://pubmed.ncbi.nlm.nih.gov/15578693/)
43. Seemanova E, Pohanka V, Seeman P, Misovicova N, Behunova J, Kvasnicova M, et al. [Nijmegen breakage syndrome in Slovakia]. *Cas Lek Cesk.* 2004; 143(8):538–41; discussion 42. Epub 2004/09/28. PMID: [15446459](https://pubmed.ncbi.nlm.nih.gov/15446459/)
44. Carlomagno F, Chang-Claude J, Dunning AM, Ponder BA. Determination of the frequency of the common 657Del5 Nijmegen breakage syndrome mutation in the German population: no association with risk of breast cancer. *Genes Chromosomes Cancer.* 1999; 25(4):393–5. PMID: [10398434](https://pubmed.ncbi.nlm.nih.gov/10398434/)
45. He M, Di GH, Cao AY, Hu Z, Jin W, Shen ZZ, et al. RAD50 and NBS1 are not likely to be susceptibility genes in Chinese non-BRCA1/2 hereditary breast cancer. *Breast Cancer Res Treat.* 2012; 133(1):111–6. doi: [10.1007/s10549-011-1700-2](https://doi.org/10.1007/s10549-011-1700-2) PMID: [21811815](https://pubmed.ncbi.nlm.nih.gov/21811815/)
46. Haak W, Lazaridis I, Patterson N, Rohland N, Mallick S, Llamas B, et al. Massive migration from the steppe was a source for Indo-European languages in Europe. *Nature.* 2015; 522(7555):207–11. doi: [10.1038/nature14317](https://doi.org/10.1038/nature14317) PMID: [25731166](https://pubmed.ncbi.nlm.nih.gov/25731166/)
47. Huang LO, Labbe A, Infante-Rivard C. Transmission ratio distortion: review of concept and implications for genetic association studies. *Hum Genet.* 2013; 132(3):245–63. doi: [10.1007/s00439-012-1257-0](https://doi.org/10.1007/s00439-012-1257-0) PMID: [23242375](https://pubmed.ncbi.nlm.nih.gov/23242375/)
48. Rotter JI, Diamond JM. What maintains the frequencies of human genetic diseases? *Nature.* 1987; 329(6137):289–90. doi: [10.1038/329289a0](https://doi.org/10.1038/329289a0) PMID: [3114647](https://pubmed.ncbi.nlm.nih.gov/3114647/)
49. Gemmell NJ, Slate J. Heterozygote advantage for fecundity. *PLoS One.* 2006; 1:e125. doi: [10.1371/journal.pone.0000125](https://doi.org/10.1371/journal.pone.0000125) PMID: [17205129](https://pubmed.ncbi.nlm.nih.gov/17205129/)
50. Smith KR, Hanson HA, Mineau GP, Buys SS. Effects of BRCA1 and BRCA2 mutations on female fertility. *Proc Biol Sci.* 2012; 279(1732):1389–95. doi: [10.1098/rspb.2011.1697](https://doi.org/10.1098/rspb.2011.1697) PMID: [21993507](https://pubmed.ncbi.nlm.nih.gov/21993507/)
51. Smith KR, Hanson HA, Hollingshaus MS. BRCA1 and BRCA2 mutations and female fertility. *Curr Opin Obstet Gynecol.* 2013; 25(3):207–13. doi: [10.1097/GCO.0b013e32835f1731](https://doi.org/10.1097/GCO.0b013e32835f1731) PMID: [23411475](https://pubmed.ncbi.nlm.nih.gov/23411475/)
52. Kwiatkowski F, Arbre M, Bidet Y, Laquet C, Uhrhammer N, Bignon YJ. BRCA Mutations Increase Fertility in Families at Hereditary Breast/Ovarian Cancer Risk. *PLoS One.* 2015; 10(6):e0127363. doi: [10.1371/journal.pone.0127363](https://doi.org/10.1371/journal.pone.0127363) PMID: [26047126](https://pubmed.ncbi.nlm.nih.gov/26047126/)
53. Voskarides K. Genetic epidemiology of cancer predisposition DNA repair genes is probably related with ancestral surviving under adverse environmental conditions. *Genet Test Mol Biomarkers.* 2014; 18(8):533–7. doi: [10.1089/gtmb.2014.0053](https://doi.org/10.1089/gtmb.2014.0053) PMID: [24933130](https://pubmed.ncbi.nlm.nih.gov/24933130/)
54. Lombard DB, Guarente L. Nijmegen breakage syndrome disease protein and MRE11 at PML nuclear bodies and meiotic telomeres. *Cancer Res.* 2000; 60(9):2331–4. PMID: [10811102](https://pubmed.ncbi.nlm.nih.gov/10811102/)
55. Borde V. The multiple roles of the Mre11 complex for meiotic recombination. *Chromosome Res.* 2007; 15(5):551–63. doi: [10.1007/s10577-007-1147-9](https://doi.org/10.1007/s10577-007-1147-9) PMID: [17674145](https://pubmed.ncbi.nlm.nih.gov/17674145/)
56. Shimada M, Sagae R, Kobayashi J, Habu T, Komatsu K. Inactivation of the Nijmegen breakage syndrome gene leads to excess centrosome duplication via the ATR/BRCA1 pathway. *Cancer Res.* 2009; 69(5):1768–75. doi: [10.1158/0008-5472.CAN-08-3016](https://doi.org/10.1158/0008-5472.CAN-08-3016) PMID: [19244116](https://pubmed.ncbi.nlm.nih.gov/19244116/)
57. Difilippantonio S, Celeste A, Kruhlak MJ, Lee Y, Difilippantonio MJ, Feigenbaum L, et al. Distinct domains in Nbs1 regulate irradiation-induced checkpoints and apoptosis. *J Exp Med.* 2007; 204(5):1003–11. doi: [10.1084/jem.20070319](https://doi.org/10.1084/jem.20070319) PMID: [17485521](https://pubmed.ncbi.nlm.nih.gov/17485521/)
58. Macklon NS, Geraedts JP, Fauser BC. Conception to ongoing pregnancy: the 'black box' of early pregnancy loss. *Hum Reprod Update.* 2002; 8(4):333–43. PMID: [12206468](https://pubmed.ncbi.nlm.nih.gov/12206468/)
59. Santos MA, Kuijk EW, Macklon NS. The impact of ovarian stimulation for IVF on the developing embryo. *Reproduction.* 2010; 139(1):23–34. doi: [10.1530/REP-09-0187](https://doi.org/10.1530/REP-09-0187) PMID: [19710204](https://pubmed.ncbi.nlm.nih.gov/19710204/)
60. Petersen MB, Mikkelsen M. Nondisjunction in trisomy 21: origin and mechanisms. *Cytogenet Cell Genet.* 2000; 91(1–4):199–203. doi: [56844](https://doi.org/10.1155/0000-0877(2000)91%3C199::AID-HUGA101%3E3.0.CO;2) PMID: [11173856](https://pubmed.ncbi.nlm.nih.gov/11173856/)
61. Hassold T, Hunt P. To err (meiotically) is human: the genesis of human aneuploidy. *Nat Rev Genet.* 2001; 2(4):280–91. doi: [10.1038/35066065](https://doi.org/10.1038/35066065) PMID: [11283700](https://pubmed.ncbi.nlm.nih.gov/11283700/)

62. Oliver TR, Feingold E, Yu K, Cheung V, Tinker S, Yadav-Shah M, et al. New insights into human nondisjunction of chromosome 21 in oocytes. *PLoS Genet*. 2008; 4(3):e1000033. doi: [10.1371/journal.pgen.1000033](https://doi.org/10.1371/journal.pgen.1000033) PMID: [18369452](https://pubmed.ncbi.nlm.nih.gov/18369452/)
63. Delhanty JD. Mechanisms of aneuploidy induction in human oogenesis and early embryogenesis. *Cytogenet Genome Res*. 2005; 111(3–4):237–44. doi: [10.1159/000086894](https://doi.org/10.1159/000086894) PMID: [16192699](https://pubmed.ncbi.nlm.nih.gov/16192699/)
64. Nagaoka SI, Hassold TJ, Hunt PA. Human aneuploidy: mechanisms and new insights into an age-old problem. *Nat Rev Genet*. 2012; 13(7):493–504. doi: [10.1038/nrg3245](https://doi.org/10.1038/nrg3245) PMID: [22705668](https://pubmed.ncbi.nlm.nih.gov/22705668/)
65. Vanneste E, Voet T, Le Caignec C, Ampe M, Konings P, Melotte C, et al. Chromosome instability is common in human cleavage-stage embryos. *Nat Med*. 2009; 15(5):577–83. doi: [10.1038/nm.1924](https://doi.org/10.1038/nm.1924) PMID: [19396175](https://pubmed.ncbi.nlm.nih.gov/19396175/)
66. Gorbsky GJ. The spindle checkpoint and chromosome segregation in meiosis. *FEBS Journal*. 2014; 282:2471–87.
67. Cheung VG, Ewens WJ. Heterozygous carriers of Nijmegen Breakage Syndrome have a distinct gene expression phenotype. *Genome Res*. 2006; 16(8):973–9. doi: [10.1101/gr.5320706](https://doi.org/10.1101/gr.5320706) PMID: [16809669](https://pubmed.ncbi.nlm.nih.gov/16809669/)
68. Cilli D, Mirasole C, Pennisi R, Pallotta V, D'Alessandro A, Antoccia A, et al. Identification of the interactors of human nibrin (NBN) and of its 26 kDa and 70 kDa fragments arising from the NBN 657del5 founder mutation. *PLoS One*. 2014; 9(12):e114651. doi: [10.1371/journal.pone.0114651](https://doi.org/10.1371/journal.pone.0114651) PMID: [25485873](https://pubmed.ncbi.nlm.nih.gov/25485873/)