

8 LITERATURVERZEICHNIS

1. Aghabeigi B, Hiranaka D, Keith DA, Kelly JP, Crean SH:
Effect of orthognathic surgery on the temporomandibular joint in patients with anterior open bite.
Int J Adult Orthod Orthognath Surg 2001; 16: 153-160.
2. Al-Din OFS, Coghlan KM, Magennis P:
Sensory nerve disturbance following Le Fort I osteotomy.
Int J Oral Maxillofac Surg 1996; 25:13-19.
3. Ataoglu H, Uckan S, Karaman AI, Uyar Y:
Bimaxillary orthognathic surgery in a patient with long face: A case report.
Int J Adult Orthod Orthognath Surg 1999; 14: 304-309.
4. Ayoub AF, Millett DT, Hasan S:
Evaluation of skeletal stability following surgical correction of mandibular prognathism.
Br J Oral Maxillofac Surg 2000; 38: 305-311.
5. Bailey LJ, Duong HL, Proffit WR:
Surgical Class III treatment: Long-term stability and patient perceptions of treatment outcome.
Int J Adult Orthod Orthognath Surg 1998; 13: 35-44.
6. Berg R:
Post-retention analysis of treatment problems and failures in 264 consecutively treated cases.
Er J Oprthod 1979; 1: 55-68.
7. Bertolini F, Russo V, Sansebastiano S:
Pre- and postsurgical psycho-emotional aspects of the orthognathic surgery patient.
Int J Adult Orthod Orthognath Surg 2000; 15: 16-23.
8. Bothur S, Blomqvist JE, Isaksson S:
Stability of Le Fort I osteotomy with advancement: a comparison of single maxillary surgery and a two-jaw procedure.
J Oral Maxillofac Surg 1998; 56: 1029-1034.
9. Bremerich A:
Die Rehabilitation von Dysgnathiepatienten aus der Sicht der Betroffenen.
(Rehabilitations of dysgnathic patients from the Patients' viewpoint.)
Zahnärztliche Praxis 1991; 42: 55-58.
10. Brix F, Lambrecht JT:
Individuelle Schädelmodellherstellung auf der Grundlage computertomographischer Informationen.
Fortschr Kiefer Gesichtschir 1987; 32: 74-77.

11. Buchanan IB, Shaw WC, Richmond S, O'Brien KD, Andrews M:
A comparison of the reliability and validity of the PAR Index and Summers' Occlusal Index.
Er J Orthod 1993; 15: 27-31.
12. Bumann A, Lotzmann U:
Funktionsdiagnostik und Therapieprinzipien.
In: Rateitschak KH (Hrsg.): Farbatlanten der Zahnmedizin, Band 12
Thieme; Stuttgart, 2000.
13. Bühl A, Zöfel P:
SPSS für Windows Version 10
Addison-Wesley Verlag; München, 2000.
14. Chen N, Neal CE, Lingenbrink P, Bloomquist D:
Neurosensory changes following orthognathic surgery.
Int J Adult Orthod Orthognath Surg 1999; 14: 259-267.
15. Cohn-Stock G:
Die chirurgische Immediatregulierung der Kiefer, speziell die chirurgische Behandlung der Prognathie.
Vjschr Zahnheilk 1921; 37: 320.
16. Costa F, Robiony MD, Politi M:
Stability of Le Fort I osteotomy in maxillary advancement: review of the literature.
Int J Adult Orthod Orthognath Surg 1999; 14: 207-213.
17. Costa F, Robiony M, Politi M:
Stability of Le Fort I osteotomy in maxillary inferior repositioning: Review of the literature.
Int J Adult Orthod Orthognath Surg 2000; 15: 197-204.
18. Csaszar GR, Niederpellmann H:
Reliability of bimaxillary surgical planning with the 3-D orthognathic surgery simulator.
Int J Adult Orthod Orthognath Surg 2000; 15: 51-58.
19. Cutbirth M, Van Sickels JE, Thrash WJ:
Condylar resorption after bicortical screw fixation of mandibular advancement.
J Oral Maxillofac Surg 1998; 56: 178-183.
20. Dal Pont G:
Retromolar osteotomy for the correction of prognathism.
J Oral Surg Anesth Hosp Dent Serv 1961; 19: 42-47.
21. De Clerk CAS, Abeloos JSV, Mommaerts MY, Neyt LF:
Temporomandibular joint symptoms in an orthognathic surgery population.
J Craniomaxillofac Surg 1995; 23: 195-199.

22. De Vries K, Devriese RP, Hovinga J, van den Akker HP:
Facial palsy after sagittal split osteotomies.
J Craniomaxillofac Surg 1993; 21:60-53.
23. Dimitroulis G:
A simple classification of orthognathic surgery complications.
Int J Adult Orthod Orthognath Surg 1998; 13: 79-87.
24. Ellis E:
Bimaxillary surgery using an intermediate splint to position the maxilla.
J Oral Maxillofac Surg 1999; 57: 53-56.
25. Fellner U, Schlömer R:
Der PAR-Index – eine praktikable Methode der Qualitätssicherung.
Kieferorthop 1996; 10: 10-20.
26. Firestone AR, Hasler RU, Ingervall B:
Treatment results in dental school orthodontic patients in 1983 and 1993.
Angle Orthod 1999; 69: 19-26.
27. Fischer K, von Konow L, Brattström V:
Open bite: stability after bimaxillary surgery-2-year treatment outcomes in 58 patients.
Eur J Orthod 2000; 22: 711-718.
28. Flanary CM, Barnwell GM, Van Sickels JE, Littlefield JH, Rugh AL:
Impact of orthognathic surgery on normal and abnormal personality dimensions:
A 2-year follow-up study of 61 patients.
Am J Orthod Dentofac Orthop 1990; 98: 313-322.
29. Fox NA:
The first 100 cases: a personal audit of orthodontic treatment assessed by the PAR
(peer assessment rating) index.
Br Dent J 1993; 174: 290-297.
30. Fridrich KL, Williamson LW:
Modification of the interocclusal splint for orthognathic surgery.
J Oral Maxillofac Surg 1989; 47: 96-97.
31. Gattinger J, Obwegeser J:
Chirurgische Kieferorthopädie und kraniofaziale Fehlbildungschirurgie.
In : Schwenzer N, Ehrenfeld M (Hrsg.): Zahn-Mund-Kiefer-Heilkunde; Spezielle Chirurgie, Band 2; 3. Auflage;
Thieme; Stuttgart; 2002; 235-273.
32. Ghali GE; Sikes JW:
Intraoral vertical ramus osteotomy as the preferred treatment for mandibular prognathism.
J Oral Maxillofac Surg 2000; 58: 313-315.

33. Grewe JM, Hagan D:
Malocclusion indices: A comparative evaluation.
Am J Orthod 1972; 61: 286-294.
34. Handwerker H.O.:
Allgemeine Sinnesphysiologie.
In: Schmidt RF, Thews G (Hrsg.): Physiologie des Menschen; 26. Auflage,
Springer; Berlin; 1995; 195-215.
35. Harzer:
Diagnostik; Fernröntgenanalyse und Kephalometrie
In: Harzer W (Hrsg.): Lehrbuch der Kieferorthopädie,
Hanser Verlag; München; 1999; 101-111
36. Hoffmann GR, Staples G, Moloney FB:
Cephalometric alterations following facial advancement surgery: Statistical
evaluation. A review of aesthetic evaluation of the face and an attempt to assess
treatment validity.
J Cranio Maxillofac Surg 1994; 22: 214-219.
37. Holmann JK, Hans MG, Nelson S, Powers MP:
An assessment of extraction versus nonextraction orthodontic treatment using the
peer assessment rating (PAR) index
Angle Orthod 1998; 68: 527-534.
38. Hoppenreijts TJM, van der Linden FPGM, Freihofer HPM, Stoelinga PJW,
Tuinzing, DB, Jacobs BTH, van't Hof MA:
Stability of transverse maxillary dental arch dimensions following orthodontic-
surgical correction of anterior open bites
Int J Adult Orthod Orthognath Surg 1998; 13: 7-22.
39. Hwang SJ, Haers PE, Zimmermann A, Oechslin C, Seifert B, Sailer HF:
Surgical risk factors for condylar resorption after orthognathic surgery.
Oral Surg, Oral Med, Oral Pathol 2000; 89: 542-552.
40. Hwang SJ, Haers PE, Seifert B, Sailer HF:
Non-surgical risk factors for condylar resorption after orthognathic surgery.
J Cranio Maxillofac Surg 2004; 32: 103-111.
41. Ive J, Mc Neill RW, West RA:
Mandibular advancement: skeletal and dental changes during fixation.
J. Oral Surg 1977; 35: 881-886.
42. Iwase M, Sugimori M, Kurachi Y, Nagumo M:
Changes in bite force and occlusal contacts in patients treated for mandibular
prognathism by orthognathic surgery.
J Oral Maxillofac Surg 1998; 56: 850-856.

43. Jacobson R, Sarver DM:
The predictability of maxillary repositioning in LeFort I orthognathic surgery.
Am J Orthod Dentofac Orthop 2002; 122: 142-154.
44. Jünger TH, Ruf S, Eisfeld J, Howaldt HP:
Cephalometric assessment of sagittal jaw base relationship prior to orthognathic surgery: The role of anterior cranial base inclination.
Int J Adult Orthod Orthognath Surg 2000; 15: 290-298.
45. Kallela I, Laine P, Suuronen R, Iizuka T, Pirinen S, Lindqvist C:
Skeletal stability following mandibular advancement and rigid fixation with polylactide biodegradable screws.
Int J Oral Maxillofac Surg 1998; 27: 3-8.
46. Keeling S, Dolce C, Van Sickels JE, Bays RA, Clark GM, Rugh JD:
A comparative study of skeletal and dental stability between rigid and wire fixation for mandibular advancement.
Am J Orthod Dentofacial Orthop 2000; 117: 638-649.
47. Kerstens HCJ, Tuinzing DB, van der Kwast WAM:
Temporomandibular joint symptoms in orthognathic surgery.
J Craniomaxillofac Surg 1989; 17: 215-218.
48. Kiyak HA:
Psychological aspects of orthognathic surgery.
Psychol Health 1993; 8: 197-212.
49. Koole R, Egyedi P:
The case of postoperative orthodontics in orthognathic surgery.
J Cranio-Max-Fac Surg 1990; 18: 293-296.
50. Krenkel C, Lixl G:
Ein Modelloperationsgerät zur Planung und Simulation von Ok- und Ukoosteotomien.
Zahnärztl Prax 1991; 42: 471-473.
51. Lapp TH:
Bimaxillary surgery without the use of an intermediate splint to position the maxilla.
J Oral Maxillofac Surg 1999; 57: 57-60.
52. Little RM:
The Irregularity Index: A quantitative score of mandibular anterior alignment.
Am J Orthod 1975; 68: 554-563.
53. Lobb WK, Ismail AI, Andrews CL, Spracklin TE:
Evaluation of orthodontic treatment using the Dental Aesthetic Index.
Am J Orthod Dentofac Orthop 1994; 106: 70-75.

54. Luhr H G:
Stabile Fixation von Oberkiefer-Mittelgesichtsfrakturen durch Mini-Kompressionsplatten.
Dtsch Zahnärztl Z 1979; 34: 851.
55. Magnusson T, Alhborg G, Finne K, Nethander G, Svartz K:
Changes in temporomandibular joint pain-dysfunction after surgical correction of dentofacial anomalies.
Int J Oral Maxillofac Surg 1986; 15: 707-714.
56. Major PW, Philippson GE, Glover KE, Grace MG:
Stability of maxilla downgrafting after rigid or wire fixation.
J Oral Maxillofac Surg 1996; 54: 1287-1291.
57. Marchetti C, Gentile L, Bianchi A, Bassi M:
Semirigid fixation of the mandible in bimaxillary orthognathic surgery: stability after 18 months.
Int J Adult Orthod Orthognath Surg 1999; 14: 37-45.
58. Medeiros PJ, Camargo ES, Vitral R, Rocha R:
Orthodontic-surgical approach in a case of severe open-bite associated with functional macroglossia.
Am J Orthod Dentofacial Orthoped 2000; 118: 347-351.
59. Michel C, Reuther J:
Orthopädische Chirurgie des Gesichtsschädels.
In: Hausamen, JE, Machtens E, Reuther J (Hrsg.): Mund-, Kiefer- und Gesichtschirurgie; Kirschner'sche allgemeine und spezielle Operationslehre; Band II; Dritte Auflage; Springer; Berlin; 1995; 360-401.
60. Mihalik CA, Proffit WR, Phillips C:
Long-term follow-up of Class II adults treated with orthodontic camouflage: A comparison with orthognathic surgery outcomes.
Am J Orthod Dentofacial Orthop 2003; 123: 266-278.
61. Mobarak KA, Krogstad O, Espeland L, Lyberg T:
Long-term stability of mandibular setback surgery: A follow-up of 80 bilateral Sagittal split osteotomy patients.
Int J Adult Orthod Orthognath Surg 2000; 15: 83-95.
62. Mobarak KA, Krogstad O, Espeland L, Lyberg T:
Stability of extraoral vertical ramus osteotomy: Plate fixation versus maxillomandibular / skeletal suspension wire fixation.
Int J Adult Orthod Orthognath Surg 2000; 15: 97-113.
63. Moldez MA, Sugawara J, Umemori M, Mitani H, Kawamura H:
Long-term dentofacial stability after bimaxillary surgery in skeletal Class III open bite patients.
Int J Adult Orthod Orthognath Surg 2000; 15: 309-319.

64. Motegi E, Hatch JP, Rugh JD, Yamaguchi H:
Health-related quality of life and psychosocial function 5 years after orthognathic surgery.
Am J Orthod Dentofac Orthop 2003; 124: 138-143.
65. Moyers RE:
Quality assurance in orthodontics.
Am J Orthod Dentofac Orthop 1990; 97: 3-9.
66. Neal CE, Kiyak HA:
Patient perceptions of pain, paresthesia and swelling after orthognathic surgery.
Int J Adult Orthod Orthognath Surg 1991; 6: 169-181.
67. Obwegeser H:
Zur Operationstechnik bei der Progenie und anderen Unterkieferanomalien.
In: Trauner R, Obwegeser H (Hrsg.): Deutsche Zahn-, Mund- und Kieferheilkunde 23; 1955; 1
68. Obwegeser H:
Surgical correction of small or retrodisplaced maxillae.
J Plast Reconstr Surg 1969; 43:351.
69. Obwegeser JA:
Osteosynthesis using biodegradable Poly-p-dioxanon (PDS II) in Le Fort I-osteotomy without postoperative intermaxillary fixation.
J Cranio Maxillofac Surg 1994; 22: 129-137.
70. Panula K, Somppi M, Finne K, Oikarinen K:
Effects of orthognathic surgery on temporomandibular joint dysfunction.
A controlled prospective 4-year follow-up study.
Int J Oral Maxillofac Surg 2000; 29: 183-187.
71. Paulus GW:
Moderne funktionelle und ästhetische Aspekte bei kieferorthopädischen Operationen.
Info aus Ortho und Kfo Jan 1990.
72. Pepersack WJ:
Tooth vitality after alveolar segmental osteotomy.
J. Maxillofac Surg 1979; 1:85-91.
73. Perez MM, Sameshima GT, Sinclair PM:
The long-term stability of Le Fort I maxillary downgrafts with rigid fixation to correct vertical maxillary deficiency.
Am J Orthod Dentofac Orthop. 1997; 112: 104-108.
74. Phillips C, Kiyak HA, Bloomquist D, Turvey TA:
Perceptions of recovery and satisfaction in the short term after orthognathic surgery.
J Oral Maxillofac Surg 2004; 62: 535-544.

75. Proffit WR, Phillips C, Dann C, Turvey TA:
Stability after surgical-orthodontic correction of skeletal Class III malocclusion.
I. Mandibular setback
Int J Adult Orthod Orthognath Surg 1991; 6: 7-18.
76. Rakosi T:
Atlas und Anleitung zur praktischen Fernröntgenanalyse.
2. Aufl. München: Hanser; 1988.
77. Reitzik M:
Skeletal and dental changes after surgical correction of mandibular prognathism.
J. Oral Surg 1980; 38: 109-116.
78. Richmond S, Shaw WC, Roberts CT, Andrews M:
The PAR Index (Peer Assessment Rating): methods to determine outcome of orthodontic treatment in terms of improvement and standards.
Er J Orthod 1992; 14: 180-187.
79. Richmond S, Andrews M:
Orthodontic treatment standards in Norway.
Er J Orthod 1993; 15: 7-15.
80. Richmond S, O'Brien K, Buchanan I, Burden D:
An introduction to occlusal indices.
ORTHO-CARE (UK) Limited;
University of Manchester; Mudent Press; 1994; 15-24.
81. Richter U, Richter F, Klein J, Michel CH:
Das vertikale und sagittale Rezidiv nach kieferorthopädisch-kieferchirurgischer Korrektur distaler Bisslagen.
Inf Orthod Kieferorthop 1990; 4: 461-479.
82. Rosenberg A, Sailer HF:
A prospective study on changes in the sensibility of the oral mucosa and the mucosa of the upper lip after Le Fort I osteotomy.
J Cranio Maxillofac Surg 1994; 22: 286-293.
83. Rotter BE, Zeitler DL:
Stability of the Le Fort I Maxillary osteotomy after rigid internal fixation.
J Oral Maxillofac Surg 1999; 57: 1080-1089.
84. Santler G:
Computergestützter 3-D-Operationssimulator.
Mund Kiefer Gesichts Chir 2000; 4: 39-44.
85. Scheerlinck JPO, Stoelinga PJW, Blijdorp PA, Brouns JJA, Nijs MLL:
Sagittal split advancement osteotomies stabilized with miniplates. A 2-5 year follow-up.
Int J Oral Maxillofac Surg 1994; 23: 127-131.

86. Scheuermann M:
Postoperative Veränderungen nach chirurgischer Kieferorthopädie zur Korrektur der mandibulären Retrognathie mit und ohne maxillärer Prognathie.
Zahnmedizinische Dissertation, Universität Frankfurt am Main; (1993)
87. Schopf P:
Kieferorthopädische Apparatesysteme (Teil 1)
In: Schopf P (Hrsg.): Curriculum Kieferorthopädie; Band I, 3. Auflage;
Quintessenz; Berlin, 2000; 287-291.
88. Schultes G, Gaggl A, Kärcher H:
Periodontal disease associated with interdental osteotomies after orthognathic surgery
J Oral Maxillofac Surg 1998; 56: 414-417.
89. Schwarz AM:
Die Röntgenostatik.
Urban & Schwarzenberg; Wien, 1958.
90. Scott AA, Hatch JP, Rugh JD, Hoffmann TJ, Rivera SM; Dolce C, Bays RA:
Psychosocial predictors of satisfaction among orthognathic surgery patients.
Int J Adult Orthod Orthognath Surg 2000; 15: 7-15.
91. Segner D, Hasund A:
Individualisierte Kephalometrie. 3. Auflage
Segner-Verlag; 1998.
92. Sienholz Dorina:
Skelettale und Weichteilveränderungen nach chirurgischer Dysgnathiebehandlung.
Dissertation 1992 HU-Berlin.
93. Spiessl B:
Osteosynthese bei sagittaler Osteotomie nach Obwegeser-Dal Pont.
In: Schuchardt K (Hrsg.): Fortschritte der Kiefer- und Gesichts-Chirurgie, Bd. 18
Thieme; Stuttgart 1974; 145.
94. Steinhäuser EW:
Bone screws and plates in orthognathic surgery.
Int J Oral Surg 1982; 11: 209.
95. Steinhäuser EW:
Historical development of orthognathic surgery.
J Cranio Maxillofac Surg 1996; 24: 195-204.
96. Summers CJ:
The occlusal index: A system for identifying and scoring occlusal disorders.
Am J Orthod 1971; 59: 552-567.

97. Tessier P:
The definitive plastic surgical treatment of the severe facial deformities of craniofacial dysostosis: Crouzon's and Apert's diseases.
Plast Reconstr Surg 1971; 48: 419.
98. Turbill EA, Richmond S, Wright JL:
A critical assessment of orthodontic standards in England and Wales (1990-1991) in relation to changes in prior approval.
Br J Orthod 1996; 23: 221-228.
99. Van de Perre JPA, Stoelinga PJW, Blijdorp PA, Brouns JJA, Hoppenreijns TJM:
Perioperative morbidity in maxillofacial orthopaedic surgery: a retrospective study.
J Cranio Maxillofac Surg 1996; 24: 263-270.
100. Van den Braber W, Van der Glas H, Van der Bilt A, Bosman F:
Masticatory function in retrognathic patients, before and after mandibular advancement surgery.
J Oral Maxillofac Surg 2004; 62: 549-554.
101. Van Merkesteyn JP, Groot RH, van Leeuwaarden R, Kroon FHM:
Intra-operative complications in sagittal and vertical ramus osteotomies.
Int J Oral Maxillofac Surg 1987; 16: 665-670.
102. Van Sickels JE, Dolce C, Keeling S, Tiner BD, Clark GM, Rugh JD:
Technical factors accounting for stability of a bilateral sagittal split osteotomy advancement: wire osteosynthesis versus rigid fixation.
Oral Surg Oral Med Oral Pathol Oral Radiol Endod 2000; 89: 19-23.
103. Vedofte P, Njattestadt A:
Pulp sensibility and pulp necrosis after LeFort I osteotomy.
J Craniomaxillofac Surg 1989; 17:167-171.
104. Voges H:
Veränderungen der Ästhetik nach Dysgnathie-Operationen und ihre psychische Verarbeitung – eine katamnestische Untersuchung.
Dissertation zur Erlangung eines Doktors der Medizin Universität Düsseldorf 1990.
105. Wagner S, Reyneke JP:
The Le-Fort I downsliding osteotomy: A study of long-term hard tissue stability.
Int J Adult Orthod Orthognath Surg 2000; 15: 37-49.
106. Waite PD, Tejera TJ, Anucul B:
The stability of maxillary advancement using Le Fort I osteotomy with and without genial bone grafting.
Int J Oral Maxillofac Surg 1996; 25: 264-267.

107. Wassmund M:
Frakturen und Luxationen des Gesichtsschädels.
Meusser; Leipzig; 1927.
108. Wassmund M:
Lehrbuch der praktischen Chirurgie des Mundes und der Kiefer. Bd I,
Meusser; Leipzig; 1935.
109. Watted N, Bill JS, Peters C:
Therapiekonzept zur kombinierten kieferorthopädisch-kieferchirurgischen Behandlung von Klasse-II-Dysgnathien mit Short-face-Syndrom.
Mund Kiefer Gesichts Chir 2000; 4: 118-124.
110. Watzke IM, Tucker MR, Turvey TA:
Lag screw versus position screw techniques for rigid internal fixation of sagittal osteotomies: a comparison of stability.
Int J Adult Orthod Orthognath Surg 1991; 6: 19-27.
111. Westermark A, Bystedt H, von Konow L:
Patients' evaluation of the final result of sagittal split osteotomy: Is it influenced by impaired sensitivity of the lower lip and chin?
Int J Adult Orthod Orthognath Surg 1999; 14: 135-139.
112. Westermark A, Englesson L, Bongenhielm U:
Neurosensory function after sagittal split osteotomy of the mandible: A comparison between subjective evaluation and objective assessment.
Int J Adult Orthod Orthognath Surg 1999; 14: 268-275.
113. Westermark A, Shayeghi M, Thor A:
Temporomandibular dysfunction in 1,516 patients before and after orthognathic surgery.
Int J Adult Orthod Orthognath Surg 2001; 16: 145-151.
114. White CS, Dolwick MF:
Prevalence and variance of temporomandibular dysfunction in orthognathic surgery patients.
Int J Adult Orthod Orthognath Surg 1992; 7: 7-14.
115. Wolford LM, Chemello PD, Hilliard F:
Occlusal plane alteration in orthognathic surgery- Part I: Effects on function and esthetics.
Am J Orthod Dentofac Orthop 1994; 106: 304-316.
116. Wright GW, Geggie Aac:
Bilateral temporomandibular joint ankylosis after bimaxillary surgery.
J Oral Maxillofac Surg 1998; 56: 1437-1441.
117. Wunderer S:
Die Prognathieoperation mittels frontal gestieltem Maxillafragment.
Österr Z Stomat 1962; 59: 98.

118. Xia J, Samman N, Yeung RWK, Shen SG, Wang D, Ip HHS, Tideman H: Three-dimensional virtual reality surgical planning and simulation workbench for orthognathic surgery.
Int J Adult Orthod Orthognath Surg 2000; 15: 265-282.
119. Yamaguchi H, Naruse T, Sueishi K, Arakawa Y, Suzuki T, Isshiki Y, Noma H, Uchiyama T, Saito C, Shigematsu T, Nakao M: Long-term stability following surgical orthodontic treatment of mandibular prognathisms: Investigation by means of lateral X-ray cephalogram.
Bull Tokyo dent Coll 1996; 37: 167-175.
120. Zimmermann M.:
Das somatoviszerale sensorische System.
In: Schmidt R.F., Thews G. (Hrsg.): Physiologie des Menschen; 26. Auflage;
Springer; Berlin; 1995; 216-235.