

References

- Agrawal, R.K., Spahn, C.M.T., Penczek, P., Grassucci, R.A., Nierhaus, K.H. and Frank, J. (2000) Visualization of tRNA movements on the *Escherichia coli* 70S ribosome during the elongation cycle. *J. Cell. Biol.*, **150**, 447-459.
- Alakhov, Y.B., Baranov, P.V., Ovodov, S.J., Ryabova, L.A., Spirin, A.S. and Morozov, I.J. (1995) Method of preparing polypeptides in cell-free translation system. USA.
- Baca, A.M. and Hol, W.G. (2000) Overcoming codon bias: a method for high-level overexpression of *Plasmodium* and other AT-rich parasite genes in *Escherichia coli*. *Int. J. Parasitol.*, **30**, 113-118.
- Ban, N., Nissen, P., Hansen, J., Moore, P.B. and Steitz, T.A. (2000) The complete atomic structure of the large ribosomal subunit at 2.4 Å resolution. *Science*, **289**, 905-920.
- Baneyx, F. and Georgiou, G. (1991) Construction and characterization of *Escherichia coli* strains deficient in multiple secreted proteases: protease III degrades high-molecular-weight substrates *in vivo*. *J. Bacteriol.*, **173**, 2696-2703.
- Berger, S.L. (1987) Quantifying ³²P-labeled and unlabeled nucleic acids. *Methods Enzymol.*, **152**, 49-54.
- Björk, G.R. (1995) Genetic dissection of synthesis and function of modified nucleosides in bacterial transfer RNA. In Cohn, W.E. and Moldave, K. (eds.), *Progress in Nucleic Acid Research and Molecular Biology*, Vol 50. Academic Press Inc., 525 B Street, Suite 1900, San Diego, CA 92101-4495, Vol. 50, pp. 263-338.
- Blaha, G. and Nierhaus, K.H. (2001) Features and functions of the ribosomal E site. *Cold Spring Harbor Symposia on Quantitative Biology*, **65**, 135-145.
- Bonner, G., Lafer, E.M. and Sousa, R. (1994) Characterization of a set of T7 RNA polymerase active site mutants. *J. Biol. Chem.*, **269**, 25120-25128.

- Bonner, G., Patra, D., Lafer, E.M. and Sousa, R. (1992) Mutations in T7 RNA polymerase that support the proposal for a common polymerase active site structure. *EMBO J.*, **11**, 3767-3775.
- Borsook, H., Deasy, C.L., Haagensmit, A.J., Keighley, G. and Lowy, P.H. (1950) Incorporation *in vitro* of labeled amino acids into rat diaphragm proteins. *J. Biol. Chem.*, **186**, 309-315.
- Bouadloun, F., Donner, D. and Kurland, C.G. (1983) Codon-specific missense errors *in vivo*. *EMBO J.*, **2**, 1351-1356.
- Bremer, H. and Dennis, P.P. (1996) Modulation of chemical composition and other parameters of the cell by growth rate. In Neidhardt, F.C., III, R.C., Ingraham, J.L., Lin, E.C.C., Low, K.B., Magasanik, B., Reznikow, W.S., Riley, M., Schaechter, M. and Umberger, H.E. (eds.), *Escherichia coli and Salmonella*. ASM Press, Washington D.C., Vol. 2, pp. 1553-1569.
- Bulmer, M. (1988) Are codon usage patterns in unicellular organisms determined by selection-mutation balance? *J. evol. Biol.*, **1**, 15-26.
- Burkhardt, N., Jünemann, R., Spahn, C.M.T. and Nierhaus, K.H. (1998) Ribosomal tRNA binding sites: Three-sites models of translation. *Crit. Rev. Biochem. Mol. Biol.*, **33**, 95-149.
- Caskey, C.T., Forrester, W.C. and Tate, W. (1984) Peptide chain termination. *Gene Expression proceeding of The Alfred Benzon Symposium, Copenhagen Munkgard*, **19**, 457-466.
- Cate, J.H., Yusupov, M.M., Yusupova, G.Z., Earnest, T.N. and Noller, H.F. (1999) X-ray crystal structures of 70S ribosome functional complexes. *Science*, **285**, 2095-2104.
- Chalfie, M., Tu, Y., Euskirchen, G., Ward, W.W. and Prasher, D.C. (1994) Green fluorescent protein as a marker for gene expression. *Science*, **263**, 802-805.

- Chamberlin, M. and Ring, J. (1973) Characterization of T7-specific ribonucleic acid polymerase. 1. General properties of the enzymatic reaction and the template specificity of the enzyme. *J. Biol. Chem.*, **248**, 2235-2244.
- Cheetham, G.M. and Steitz, T.A. (1999) Structure of a transcribing T7 RNA polymerase initiation complex. *Science*, **286**, 2305-2309.
- Chen, H.Y., Bjercknes, M., Kumar, R. and Jay, E. (1994) Determination of the optimal aligned spacing between the Shine-Dalgarno sequence and the translation initiation codon of *Escherichia coli* mRNAs. *Nucleic Acids Res.*, **22**, 4953-4957.
- Craigien, W.J., Cook, R.G., Tate, W.P. and Caskey, C.T. (1985) Bacterial peptide chain release factors: conserved primary structure and possible frameshift regulation of release factor 2. *Proc. Natl. Acad. Sci. USA*, **82**, 3616-3620.
- Cramer, A., Whitehorn, E.A., Tate, E. and Stemmer, W.P. (1996) Improved green fluorescent protein by molecular evolution using DNA shuffling. *Nat. Biotechnol.*, **14**, 315-319.
- de Smit, M.H. and van Duin, J. (1990) Secondary structure of the ribosome binding site determines translational efficiency: A quantitative analysis. *Proc. Natl. Acad. Sci. USA*, **87**, 7668-7672.
- Dekker, N., Cox, R.C., Kramer, R.A. and Egmond, M.R. (2001) Substrate specificity of the integral membrane protease OmpT determined by spatially addressed peptide libraries. *Biochemistry*, **40**, 1694-701.
- Dong, H.J., Nilsson, L. and Kurland, C.G. (1996) Co-variation of tRNA abundance and codon usage in *Escherichia coli* at different growth rates. *J. Mol. Biol.*, **260**, 649-663.
- Frank, J., Zhu, J., Penczek, P., Li, Y.H., Srivastava, S., Verschoor, A., Radermacher, M., Grassucci, R., Lata, R.K. and Agrawal, R.K. (1995) A model of protein synthesis based on cryo-electron microscopy of the *E. coli* ribosome. *Nature*, **376**, 441-444.

- Freistoffer, D.V., Pavlov, M.Y., MacDougall, J., Buckingham, R.H. and Ehrenberg, M. (1997) Release factor RF3 in *E. coli* accelerates the dissociation of release factors RF1 and RF2 from the ribosome in a GTP-dependent manner. *EMBO J.*, **16**, 4126-4133.
- Gabashvili, I.S., Agrawal, R.K., Spahn, C.M.T., Grassucci, R.A., Svergun, D.I., Frank, J. and Penczek, P. (2000) Solution structure of the *E. coli* 70S ribosome at 11.5 angstrom resolution. *Cell*, **100**, 537-549.
- Gale, E.F. and Folkes, J.P. (1954) Effect of nucleic acids on protein synthesis and amino-acid incorporation in disrupted staphylococcal cells. *Nature*, **173**, 1223-1227.
- Gallie, D.R. and Kado, C.I. (1989) A translational enhancer derived from tobacco mosaic virus is functionally equivalent to a Shine-Dalgarno sequence. *Proc. Natl. Acad. Sci. USA*, **86**, 129-132.
- Ganoza, M.C., Aoki, H., Burkhardt, N. and Murphy, B.J. (1996) The ribosome as 'affinity matrix': Efficient purification scheme for translation factors. *Biochimie*, **78**, 51-61.
- Garcia, G.M., Mar, P.K., Mullin, D.A., Walker, J.R. and Prather, N.E. (1986) The *E. coli dnaY* gene encodes an arginine transfer RNA. *Cell*, **45**, 453-459.
- Gold, L. (1988) Posttranscriptional regulatory mechanisms in *Escherichia coli*. *Ann. Rev. Biochem.*, **57**, 199-233.
- Gold, L.M. and Schweiger, M. (1969) Synthesis of phage-specific alpha- and beta-glucosyl transferases directed by T-even DNA *in vitro*. *Proc. Natl. Acad. Sci. USA*, **62**, 892-898.
- Goldman, E., Rosenberg, A.H., Zubay, G. and Studier, F.W. (1995) Consecutive low-usage leucine codons block translation only when near the 5' end of a message in *Escherichia coli*. *J Mol Biol*, **245**, 467-473.
- Grajevskaja, R.A., Ivanov, Y.V. and Saminsky, E.M. (1982) 70-S ribosomes of *Escherichia coli* have an additional site for deacylated tRNA binding. *Eur. J. Biochem.*, **128**, 47-52.

- He, B., Rong, M., Lyakhov, D., Gartenstein, H., Diaz, G., Castagna, R., McAllister, W.T. and Durbin, R.K. (1997) Rapid mutagenesis and purification of phage RNA polymerases. *Protein Express. Purif.*, **9**, 142-151.
- Henaut, A. and Danchin, A. (1996) Analysis and predictions from *Escherichia coli* sequences, or *E. coli* in silico. In Neidhardt, F.C. (ed.) *Escherichia coli and Salmonella typhimurium cellular and molecular biology*, Vol. 2, pp. 2047-2066.
- Huala, E., Moon, A.L. and Ausubel, F.M. (1991) Aerobic inactivation of *Rhizobium meliloti* NifA in *Escherichia coli* is mediated by *lon* and two newly identified genes, *snoB* and *snoC*. *J. Bacteriol.*, **173**, 382-390.
- Ikemura, T. (1985) Codon usage and tRNA content in unicellular and multicellular organisms. *Mol. Biol. Evol.*, **2**, 13-34.
- Ikemura, T. and Ozeki, H. (1983) Codon usage and transfer RNA contents: organism-specific codon-choice patterns in reference to the isoacceptor contents. *Cold Spring Harb. Symp. Quant. Biol.*, **47 Pt 2**, 1087-1097.
- Iost, I. and Dreyfus, M. (1995) The stability of *Escherichia coli lacZ* mRNA depends upon the simultaneity of its synthesis and translation. *EMBO J.*, **14**, 3252-3261.
- Iost, I., Guillerez, J. and Dreyfus, M. (1992) Bacteriophage T7 RNA polymerase travels far ahead of ribosomes *in vivo*. *J. Bacteriol.*, **174**, 619-622.
- Ivanov, I.G., Alexandrova, R., Dragulev, B., Leclerc, D., Saraffova, A., Maximova, V. and Abouhaidar, M.G. (1992) Efficiency of the 5'-terminal sequence (omega) of tobacco mosaic virus RNA for the initiation of eukaryotic gene translation in *Escherichia coli*. *Eur J Biochem*, **209**, 151-156.
- Jewett, M.C. and Swartz, J.R. (2004a) Mimicking the *Escherichia coli* cytoplasmic environment activates long-lived and efficient cell-free protein synthesis. *Biotechnol. Bioeng.*, **86**, 19-26.

- Jewett, M.C. and Swartz, J.R. (2004b) Substrate replenishment extends protein synthesis with an *in vitro* translation system designed to mimic the cytoplasm. *Biotechnol. Bioeng.*, **87**, 465-472.
- Kane, J.F. (1995) Effects of rare codon clusters on high-level expression of heterologous proteins in *Escherichia coli*. *Curr. Opin. Biotechnol.*, **6**, 494-500.
- Keller, E.B. and Littlefield, J.W. (1957) Incorporation of ¹⁴C-amino acids into ribonucleoprotein particles from the Ehrlich mouse ascites tumor. *J. Biol. Chem.*, **224**, 13-30.
- Kim, D.M. and Swartz, J.R. (1999) Prolonging cell-free protein synthesis with a novel ATP regeneration system. *Biotechnol. Bioeng.*, **66**, 180-188.
- Kim, D.M. and Swartz, J.R. (2000) Prolonging cell-free protein synthesis by selective reagent additions. *Biotechnol. Prog.*, **16**, 385-390.
- Kim, D.M., Swartz, J. R. (2000) Oxalate improves protein synthesis by enhancing ATP supply in cell-free system derived from *E. coli*. *Biotechnol. Lett.*, **22**, 1537-1542.
- Komine, Y., Adachi, T., Inokuchi, H. and Ozeki, H. (1990) Genomic organization and physical mapping of the transfer RNA genes in *Escherichia coli* K12. *J. Mol. Biol.*, **212**, 579-598.
- Kozak, M. (1999) Initiation of translation in prokaryotes and eukaryotes. *Gene*, **234**, 187-208.
- Kurland, C. and Gallant, J. (1996) Errors of heterologous protein expression. *Curr. Opin. Biotechnol.*, **7**, 489-493.
- Lederman, M. and Zubay, G. (1967) DNA-directed peptide synthesis. 1. A comparison of T2 and *Escherichia coli* DNA-directed peptide synthesis in two cell-free systems. *Biochim. Biophys. Acta*, **149**, 253-258.
- Lewicki, B.T.U., Margus, T., Remme, J. and Nierhaus, K.H. (1993) Coupling of rRNA transcription and ribosomal assembly *in vivo* - formation of active ribosomal subunits in *Escherichia coli* requires transcription of rRNA

- genes by host RNA polymerase which cannot be replaced by bacteriophage-T7 RNA polymerase. *J. Mol. Biol.*, **231**, 581-593.
- Liiv, A., Tenson, T. and Remme, J. (1996) Analysis of the ribosome large subunit assembly and 23S rRNA stability *in vivo*. *J. Mol. Biol.*, **263**, 396-410.
- Lill, R., Robertson, J.M. and Wintermeyer, W. (1984) tRNA binding sites of ribosomes from *Escherichia coli*. *Biochemistry*, **23**, 6710-6717.
- Lipmann, F. (1963) Messenger ribonucleic acid. *Prog. Natl. Acad. Res.*, **1**, 135-161.
- Littlefield, J.W., Keller, E.B., Gross, J. and Zamecnik, P.C. (1955) Studies on cytoplasmic ribonucleoprotein particles from the liver of the rat. *J. Biol. Chem.*, **217**, 111-123.
- Loechel, S., Inamine, J.M. and Hu, P.C. (1991) A novel translation initiation region from *Mycoplasma genitalium* that functions in *Escherichia coli*. *Nucleic Acids Res.*, **19**, 6905-6911.
- Londei, P. (2001) Translation initiation models in prokaryotes and eukaryotes. *Encyclopedia of Life Sciences*.
- Looman, A.C., Bodlaender, J., Comstock, L.J., Eaton, D., Jhurani, P., de Boer, H.A. and van Knippenberg, P.H. (1987) Influence of the codon following the AUG initiation codon on the expression of a modified lacZ gene in *Escherichia coli*. *EMBO J.*, **6**, 2489-2492.
- Makarova, O.V., Makarov, E.M., Sousa, R. and Dreyfus, M. (1995) Transcribing of *Escherichia coli* genes with mutant T7 RNA polymerases: Stability of lacZ mRNA inversely correlates with polymerase speed. *Proc. Natl. Acad. Sci. USA*, **92**, 12250-12254.
- Mathews, D.H., Sabina, J., Zuker, M. and Turner, D.H. (1999) Expanded sequence dependence of thermodynamic parameters improves prediction of RNA secondary structure. *J. Mol. Biol.*, **288**, 911-940.

- Matthaei, J.H. and Nirenberg, M.W. (1961) Characteristics and stabilization of DNAase-sensitive protein synthesis in *E. coli* extracts. *Proc. Natl. Acad. Sci. USA*, **47**, 1580-1588.
- Mccarthy, J.E.G. and Brimacombe, R. (1994) Prokaryotic translation: The interactive pathway leading to initiation. *Trends Genet.*, **10**, 402-407.
- Miroux, B. and Walker, J.E. (1996) Over-production of proteins in *Escherichia coli*: mutant hosts that allow synthesis of some membrane proteins and globular proteins at high levels. *J. Mol. Biol.*, **260**, 289-298.
- Moazed, D. and Noller, H.F. (1989) Intermediate states in the movement of transfer RNA in the ribosome. *Nature*, **342**, 142-148.
- Mohanty, B.K. and Kushner, S.R. (1999) Analysis of the function of *Escherichia coli* poly(A) polymerase I in RNA metabolism. *Mol. Microbiol.*, **34**, 1094-1108.
- Moore, P.B. and Steitz, T.A. (2002) The involvement of RNA in ribosome function. *Nature*, **418**, 229-235.
- Nakamura, Y., Gojobori, T. and Ikemura, T. (2000) Codon usage tabulated from international DNA sequence databases: status for the year 2000. *Nucleic Acids Res.*, **28**, 292.
- Nierhaus, K.H. (1990) The allosteric three-site model for the ribosomal elongation cycle: features and future. *Biochemistry*, **29**, 4997-5008.
- Nierhaus, K.H., Beyer, D., Dabrowski, M., Schäfer, M.A., Spahn, C.M.T., Wadzack, J., Bittner, K.-U., Burkhardt, N., Diedrich, G., Jünemann, R., Kamp, D., Voss, H. and Stuhmann, H.B. (1995) The elongating ribosome: Structural and functional aspects. *Biochem. Cell Biol.*, **73**, 1011-1021.
- Nierhaus, K.H., Wadzack, J., Burkhardt, N., Jünemann, R., Meerwinck, W., Willumeit, R. and Stuhmann, H.B. (1998) Structure of the elongating ribosome: Arrangement of the two tRNAs before and after translocation. *Proc. Natl. Acad. Sci. USA*, **95**, 945-950.

- Nirenberg, M.W. and Matthaei, J.M. (1961) The dependence of cell-free protein synthesis in *E. coli* upon naturally occurring or synthetic polyribonucleotides. *Proc. Natl. Acad. Sci. USA*, **47**, 1588-1602.
- Olins, P.O., Devine, C.S., Rangwala, S.H. and Kavka, K.S. (1988) The T7 phage *gene 10* leader RNA, a ribosome-binding site that dramatically enhances the expression of foreign genes in *Escherichia coli*. *Gene*, **73**, 227-235.
- Palade, G.E. (1955) A small particulate component of the cytoplasm. *J. Biophys. Biochem. Cytol.*, **1**, 59-68.
- Petersen, H.U., Danchin, A. and Grunberg-Manago, M. (1976) Toward an understanding of the formylation of initiator tRNA methionine in prokaryotic protein synthesis. I. *In vitro* studies of the 30S and 70S ribosomal-tRNA complex. *Biochemistry*, **15**, 1357-1362.
- Prasher, D.C. (1995) Using GFP to see the light. *Trends Genet.*, **11**, 320-323.
- Prasher, D.C., Eckenrode, V.K., Ward, W.W., Prendergast, F.G. and Cormier, M.J. (1992) Primary structure of the *Aequorea victoria* green-fluorescent protein. *Gene*, **111**, 229-233.
- Ramakrishnan, V. (2002) Ribosome structure and the mechanism of translation. *Cell*, **108**, 557-572.
- Ramakrishnan, V. and Moore, P.B. (2001) Atomic structures at last: the ribosome in 2000. *Curr. Opin. Struct. Biol.*, **11**, 144-154.
- Rheinberger, H.-J. (2004) *A history of protein biosynthesis and ribosome research*.
- Rheinberger, H.-J. and Nierhaus, K.H. (1980) Simultaneous binding of the 3 tRNA molecules by the ribosome of *E coli*. *Biochem. Internatl.*, **1**, 297-303.
- Rheinberger, H.-J., Sternbach, H. and Nierhaus, K.H. (1981) Three tRNA binding sites on *Escherichia coli* ribosomes. *Proc. Natl. Acad. Sci. USA*, **78**, 5310-5314.
- Ross, J. (2001) mRNA turnover. *Encyclopedia of Life Sciences*.

- Sardesai, N.Y., Green, R. and Schimmel, P. (1999) Efficient 50S ribosome-catalyzed peptide bond synthesis with an aminoacyl minihelix. *Biochemistry*, **38**, 12080-8.
- Saxena, P. and Walker, J.R. (1992) Expression of *argU*, the *Escherichia coli* gene coding for a rare arginine tRNA. *J. Bacteriol.*, **174**, 1956-1964.
- Selinger, D.W., Saxena, R.M., Cheung, K.J., Church, G.M. and Rosenow, C. (2003) Global RNA half-life analysis in *Escherichia coli* reveals positional patterns of transcript degradation. *Genome Res.*, **13**, 216-223.
- Shine, J. and Dalgarno, L. (1974) The 3'-terminal sequence of *E. coli* 16S rRNA: Complementarity to nonsense triplets and ribosome binding sites. *Proc. Natl Acad. Sci. USA*, **71**, 1342-1346.
- Spahn, C.M.T. and Nierhaus, K.H. (1998) Models of the elongation cycle: An evaluation. *Biol. Chem.*, **379**, 753-772.
- Sprengrart, M.L. and Porter, A.G. (1997) Functional importance of RNA interactions in selection of translation initiation codons. *Mol. Microbiol.*, **24**, 19-28.
- Srivastava, A.K. and Schlessinger, D. (1990) Mechanism and regulation of bacterial ribosomal RNA processing. *Annu. Rev. Microbiol.*, **44**, 105-129.
- Stark, H., Mueller, F., Orlova, E.V., Schatz, M., Dube, P., Erdemir, T., Zemlin, F., Brimacombe, R. and Heel, M.v. (1995) The 70S *Echerichia coli* ribosome at 23 Å resolution: Fitting the ribosomal RNA. *Structure*, **3**, 815-821.
- Studier, F.W. and Moffatt, B.A. (1986) Use of bacteriophage T7 RNA polymerase to direct selective high-level expression of cloned genes. *J. Mol. Biol.*, **189**, 113-130.
- Studier, F.W., Rosenberg, A.H., Dunn, J.J. and Dubendorff, J.W. (1990) Use of T7 RNA polymerase to direct expression of cloned genes. *Methods Enzymol.*, **185**, 60-89.
- Ueda T. and K., W. (2001) tRNA synthesis and regulation. *Enciclopedia of Life Sciences*.

- Ugarov, V.I., Morozov, I.Y., Jung, G.Y., Chetverin, A.B. and Spirin, A.S. (1994) Expression and stability of recombinant RQ-mRNAs in cell-free translation systems. *FEBS Lett.*, **341**, 131-134.
- Voss, S. and Skerra, A. (1997) Mutagenesis of a flexible loop in streptavidin leads to higher affinity for the Strep-tag II peptide and improved performance in recombinant protein purification. *Protein. Eng.*, **10**, 975-982.
- Wadzack, J., Burkhardt, N., Jünemann, R., Diedrich, G., Nierhaus, K.H., Frank, J., Penczek, P., Meerwinck, W., Schmitt, M., Willumeit, R. and Stuhmann, H.B. (1997) Direct localization of the tRNAs within the elongating ribosome by means of neutron scattering (Proton-spin contrast-variation). *J. Mol. Biol.*, **266**, 343-356.
- Walz, A., Pirrotta, V. and Ineichen, K. (1976) Lambda repressor regulates the switch between PR and Prm promoters. *Nature*, **262**, 665-669.
- Watson, J.D. (1963) Involvement of tRNA in the synthesis of proteins. *Science*, **140**, 17-26.
- Watson, J.D. (1964) The synthesis of proteins upon ribosomes. *Bull. Soc. Chim. Biol.*, **46**, 1399-1425.
- Weiss, R.B., Murphy, J.P. and Gallant, J.A. (1984) Genetic screen for cloned release factor genes. *J. Bacteriol.*, **158**, 362-364.
- Williams, K.J. and Piddock, L.J. (1998) Accumulation of rifampicin by *Escherichia coli* and *Staphylococcus aureus*. *J. Antimicrob. Chemother.*, **42**, 597-603.
- Wilson, D.N. and Nierhaus, K.H. (2003) The ribosome through the looking glass. *Angew. Chem. Int. Ed. Engl.*, **42**, 3464-3486.
- Wilson, K.S. and Noller, H.F. (1998) Mapping the position of translational elongation factor EF-G in the ribosome by directed hydroxyl radical probing. *Cell*, **92**, 131-139.

- Winnick, T. (1950) Studies on the mechanism of protein synthesis in embryonic and tumor tissues. II. Inactivation of fetal rat liver homogenates by dialysis, and reactivation by the adenylic acid system. *Arch. Biochem.*, **28**, 338-447.
- Wu, C.J. and Janssen, G.R. (1996) Translation of vph mRNA in *Streptomyces lividans* and *Escherichia coli* after removal of the 5' untranslated leader. *Mol. Microbiol.*, **22**, 339-355.
- Yusupov, M.M., Yusupova, G.Z., Baucom, A., Lieberman, K., Earnest, T.N., Cate, J.H. and Noller, H.F. (2001) Crystal structure of the ribosome at 5.5 Å resolution. *Science*, **292**, 883-896.
- Zamecnik, P.C. (1969) An historical account of protein synthesis, with current overtones - a personalized view. *Cold Spring Harbor Symposia on Quantitative Biology*, **34**, 1-16.
- Zaniewski, R., Petkaites, E. and Deutscher, M.P. (1984) A multiple mutant of *Escherichia coli* lacking the exoribonucleases RNase II, RNase D, and RNase BN. *J. Biol. Chem.*, **259**, 11651-11653.
- Zavialov, A.V., Buckingham, R.H. and Ehrenberg, M. (2001) A posttermination ribosomal complex is the guanine nucleotide exchange factor for peptide release factor RF3. *Cell*, **107**, 115-124.
- Zubay, G. (1973) *In vitro* synthesis of protein in microbial systems. *Annu. Rev. Genet.*, **7**, 267-287.
- Zuker, M. (2003) Mfold web server for nucleic acid folding and hybridisation prediction. *Nucleic Acids Res.*, **31**, 3406-3415.