

GLOSSARY

1. INTRODUCTION	8
2. LITERATURE	10
2.1. Significance of pyrogens	10
2.2. Pyrogens	10
2.3. Mechanisms of pyrogenic effects	13
2.3.1. Binding proteins	13
2.3.2. Receptors	14
2.3.3. Intracellular signalling	16
2.4. Cytokines	17
2.4.1. Interleukin-1 β	17
2.4.2. Interleukin-6	18
2.4.3. Tumor Necrosis Factor- α	19
2.4.4. Effects of cytokines	19
2.5. Interleukin-8	20
2.5.1. Effects of Interleukin-8	21
2.6. Pyrogen measurement	22
2.6.1. Rabbit pyrogen test	22
2.6.2. Limulus Amoebocyte Lysate Test (LAL)	24
2.6.3. Whole blood test	25
2.6.3.1. Incubation	25
2.6.3.2. ELISA	25
2.6.4. WEHI	25

3. MATERIALS AND METHODS	26
3.1. Substances	26
3.1.1. Whole blood incubation	26
3.1.2. ELISA	26
3.1.3. WEHI	28
3.2. Animals	28
3.3. Equipment	28
3.3.1. Whole blood incubation	28
3.3.2. ELISA	29
3.3.3. WEHI	29
3.4. Methods	29
3.4.1. Whole blood incubations	29
3.4.2. ELISAs	30
3.4.2.1. Human ELISAs	30
3.4.2.2. Rabbit ELISAs	30
3.4.3. WEHI assay	30
3.5. Statistics	31
4. RESULTS	32
Part 1: Comparison of the in vivo fever reaction of the rabbit with the in vitro cytokine production measured by ELISA system	32
rabbit IL-1 β	33
rabbit IL-8	33
rabbit IL-6	34
rabbit TNF- α	35

Part 2: Comparison of the in vitro cytokine production of humans and rabbits towards a variety of stimuli	37
4.1. <i>E. coli</i> O113	37
human IL-1 β	
human IL-8	38
human IL-6	38
human TNF- α	39
 4.2. <i>E. coli</i> O111	
human IL-1 β	40
rabbit IL-1 β	41
human IL-8	41
rabbit IL-8	42
human IL-6	43
rabbit IL-6	43
human TNF- α	44
rabbit TNF- α	44
 4.3. <i>E. coli</i> O128	45
 4.4. <i>Salmonella typhosa</i>	46
 4.5. <i>Salmonella typhimurium</i>	47
 4.6. <i>Shigella flexneri</i> 1A	48
 4.7. <i>Serratia marcescens</i>	48
 4.8. <i>Bacillus subtilis</i>	49
human IL-1 β	49
rabbit IL-1 β	50
human IL-8	51
rabbit IL-8	51
human IL-6	52

rabbit IL-6	52
human TNF- α	53
rabbit TNF- α	54
Part 3:	
Less potent Gram-negative stimuli measured in the rabbit IL-1 β and rabbit IL-8 assay	
4.9. <i>Bordetella pertussis</i>	55
4.10. <i>Pseudomonas aeruginosa</i>	57
Part 4: Establishment of a bioassay (WEHI) in order to compare TNF- α production in human and canine whole blood	
	58
5. DISCUSSION	64
6. SUMMARY	73
7. REFERENCES	75
8. APPENDIX	
8.1. Abbreviations	100
8.2. Tables	102