

7. References

- Akashi, S., Shimazu, R., Ogata, H., Nagai, Y., Takeda, K., et al. (2000)
Cutting edge: cell surface expression and lipopolysaccharide signalling via the Toll-like receptor 4-MD-2 complex on mouse peritoneal macrophages
J Immunol 164: 3471-75
- Akira, S., Takeda, K., and Kaisho, T. (2001)
TOLL-like receptors: critical proteins linking innate and acquired immunity
Nat Immunol 2: 675-680
- Alheim, K., Chai, Z., and Fantuzzi, G. (1997)
Hyperresponsive febrile reactions to interleukin (IL) 1 alpha and IL-1 beta, and altered brain cytokine mRNA and serum cytokine levels, in IL-1 beta deficient mice
Proc Natl Acad Sci USA 94; 2681-2686
- Atkins, E., Bodel, P., and Francis L. (1967)
Release of an endogenous pyrogen in vitro from rabbit mononuclear cells
J Exp Med 126: 357-86
- Banks, W. A., Ortiz, L., Plotkin, S. R., and Kastin, A. J. (1991)
Human interleukin (IL) 1 α , murine IL-1 α and murine IL-1 β are transported from blood to brain in the mouse by a shared saturable mechanism
J Pharmacol Exp Ther 259: 988-996
- Banks, W. A., Kastin, A. J., and Gutierrez, E. G. (1994)
Penetration of interleukin-6 across the murine blood-brain barrier
Neurosci Lett 26; 179: 53-56
- Blatteis, C. M. (1992)
Role of the OVLT in the febrile response to circulating pyrogens
Prog Brain Res 91; 409-412.

Blease, K., Chen, Y., Hellewell, P. G., and Burke-Gaffney, A. (1999)
Lipoteichoic acid inhibits lipopolysaccharide-induced adhesion molecule expression and
IL-8 release in human lung microvascular endothelial cells
J Immunol 163: 6139-47

Bristow, A. F., Mosley, K., and Poole, S. (1991)
Interleukin-1 beta production in vivo and in vitro in rats and mice measured using
specific immunoradiometric assays
J Mol Endocrinol 7: 1-7

Burrows, S. D., Doyle, M. L., Murphy, K. P., Franklin, S. G., White, J. R., Brooks, I.,
McNulty, D. E., Scott, M. O., Knutson, J. R., Porter, D., Young, P. R., and Hensley, P.
(1994)
Determination of the monomer-dimer equilibrium of interleukin-8 reveals it is a
monomer at physiological concentrations
Biochemistry 33: 12741-45

Cannon, J. G., Tompkins, R. G., Gelfand, J. A., Michie, H. R., Stanford, G. G., van der
Meer, H. W. M., Endres, S., Lonnemann, G., Corsetti, J., Chrenow, B., Wilmore, D. W.,
Wolff, S. M., Burke, J. F., and Dinarello, C. A. (1990)
Circulating interleukin-1 and tumor necrosis factor in septic shock and experimental
endotoxin fever
J Infect Dis 161: 79-84

Cao, Z., Henzel, J., and Gao, X. (1996)
IRAK: A kinase associated with the interleukin-1 receptor
Science 271: 1128-1131

Cao, Z., Xiong, J., Takeuchi, M., Kurama, T., and Goeddel, D. V. (1996)
Traf 6 is a signal transducer for interleukin-1
Nature 383: 443-446

Cartmell T. , Mitchell D., and Lamond F. J. (2002)
Route of administration differentially affects fevers induced by Gram-negative and
Gram-positive pyrogens in rabbits
Exp Physiol (England) 87: 391-9

Chai, Z., Gatti, S., Toniatti, C., Poli, V., and Bartfai, T. (1996)
Interleukin (IL)-6 gene expression in the central nervous system is necessary for fever
response to lipopolysaccharide or IL-1 β : a study on IL-6 deficient mice
J Exp Med 183 : 311-316

Co Tui and Schrift, M. H. (1942)
A tentative test for pyrogen in infusion fluids
Proc Soc Exp Biol Med 49: p 320

Cooper, A. I., Kirkman, E., Little, R. A., Hopkins, S. J., and Rothwell, N. J. (1991)
Mechanisms involved in the fever and hypermetabolism induced by turpentine injection
in the rat
Proc Nutr Soc 50; 149A

Couturier, C., Jahns, G., Kazatchkine, M. D., and Haeffner-Cavaillon, N. (1992)
Membrane molecules which trigger the production of interleukin-1 and tumor necrosis
factor- α by lipopolysaccharide-stimulated human monocytes
Eur J Immunol 22: 1461-1466

Deeter, L. B., Martin L. W., and Lipton, J. M. (1989)
Age- and sex-related differences in febrile response to peripheral pyrogens in the rabbit
Gerontology 35; 297-304

Deiters, U., Gumenscheimer, M., Galanos, C., and Muhlradt, P. F. (2003)
Toll-like receptor 2- and 6-mediated stimulation by macrophage-activating lipopeptide 2
induces lipopolysaccharide (LPS) cross tolerance in mice, which results in protection
from tumor necrosis factor alpha but in partial protection from lethal LPS doses
Infect Immun 71: 4456-62

Dinarello, C. A., Bodel, P. T., and Atkins, E. (1968)
The role of the liver in the production of fever and in pyrogenic tolerance
Trans Assoc Am Physicians 81: 334-343

Dinarello, C. A., Shparber, M., Kent, E. F. Jr., and Wolff, S. M. ((1981))
Production of leukocytic pyrogen from phagocytes of neonates
J Infect Dis 144: 337-43

Dinarello, C. A. (1984)
Interleukin-1
Rev Infect Dis 6: 51-95

Dinarello, C. A., Cannon, J. G., Wolff, S. M., Bernheim, H. A., Beutler, B., Cerami, A., Figari, I. S., Palladino, M. A., and O'Connor J. V. (1986)
Tumor necrosis factor (cachectin) is an endogenous pyrogen and induces production of interleukin-1
J Exp Med 163: 1133-1150

Dinarello, C. A., Cannon, J. G., Mancilla, J., Bishai, I., Lees, J., and Coceani, F. (1991)
Interleukin-6 as an endogenous pyrogen: induction of prostaglandin E2 in brain but not in peripheral blood mononuclear cells
Brain Res 562: 199-206

Dinarello C. A. (1998)
Interleukin-1 beta, interleukin-18, and the interleukin-1 beta converting enzyme
Ann N Y Acad Sci 856: 1-11

De Winter, R. J., Manten, A., de Jong Y. P., Adams, R., van Deventer, S. J., and Lie K. I. (1997)
Interleukin 8 released after acute myocardial infarction is mainly bound to erythrocytes
Heart 78: 598-602

Duan, H. and Dixit, V. M. (1997)
RAIDD is a new `death` adapter molecule
Nature 385: 86-89

Duncombe, A. S., and Brenner, M. K. (1988)
Is circulating tumor necrosis factor bioactive?
N Engl J Med 319: 1227-1228

Elin, R. J., Sandberg, A. L., and Rosenstreich, D. (1976)
Comparison of the pyrogenicity, limulus activity, mitogenicity and complement reactivity
of several bacterial endotoxins and related compounds
J Immunol 117: 1238-1242

Elsbach, P. and Weiss, J. (1992)
Phagocytic cells: Oxygen-independent antimicrobial systems.
In: Inflammation: Basic principles and clinical correlates. 2nd edn. Edited by Gallstein,
Goldstein, Snyderman. New York: Raven Press Ltd.: 603-636

Engel, A., Kern, W. V., Murdter, G., and Kern, P. (1994)
Kinetics and correlation with body temperature of circulating interleukin-6, interleukin-8,
tumor necrosis factor alpha and interleukin-1 beta in patients with fever and neutropenia
Infection 22: 160-164

Erickson, S. L., de Sauvage, F. J., Kikly, K., Carver Moore, K., Pitts Meek, S., Gillett, N.,
Sheehan, K. C., Schreiber, R. D., Goeddel, D. V. and Moore, M. W. (1994)
Decreased sensitivity to tumour necrosis factor but normal T-cell development in TNF
receptor-2-deficient mice
Nature 372: 560-63

Fitzgerald, K. A., Palsson-McDermot, E. M., Bowie, A. G., Jefferies, C. A., Mansell A. S.,
Brady, G., Dunne, A., Gray, P., Harte, M. T., McMurray, D., Smith, D. E., Sims, J. E.,
Bird, T. A., and O'Neill, L. A (2001)
MAL (MyD88-adaptor-like) is required for Toll like receptor 4 signal transduction
Nature 413: 78-83

Flynn, A., Finke, J. H., and Hilfiker, M. L. (1982)
Placental mononuclear phagocytes as a source of interleukin-1
Science 218: 475-7

Fomsgaard, A., Worsaae, H., and Bendtzen, K. (1988)
Detection of tumor necrosis factor from lipopolysaccharide-stimulated human
mononuclear cells by enzyme-linked immunosorbent assay and cytotoxicity bioassay
Scand J Immunol 27: 143-147

Frey, E. A., Miller, D. S., Jahr, T. G., Sundan, A., Bazil, V., Espenvik, T., Finlay, B. B.,
and Wright, S. D. (1992)
Soluble CD14 participates in the response of cell to lipopolysaccharide
J Exp Med 176: 1665-1671

Fukui, A., Inoue, N., Matsumoto, M., Nomura, M., Yamada, K., Matsuda, Y.,
Toyoshima, K., and Seya, T. (2001)
Molecular cloning and functional characterization of chicken TOLL-like receptors. A
single chicken TOLL-like receptor covers multiple molecular patterns
J Biol Chem 276: 47143-47149

Gazzano-Santorio, H., Parent, B., Grinna, L., Horwitz, A., Parsons, T., Theofan, G.,
Elsbach, P., Weiss, J., and Conlon, P. J. (1992)
High affinity binding between Bactericidal/Permeability Increasing protein and the
Lipid A region of LPS
Infect Immun 60: 4754-4761

Gearing, A. J. H., Cartwright, J. E., and Wadhwa, M. (1994)
25. Biological and immunological assays for cytokines. In ed. Thomson A.W. The
Cytokine Handbood, 2nd edn. Academic Press Ltd., London, 507-524

Grant, R. (1950)
Emotional hypothermia in rabbits
Am J Physiol 160: 285

Greenfeder, S. A., Nunes, P., Kwee, L., Labow, M., Chizzonite, R. A., and Ju, G. (1995)
Molecular cloning and characterization of a second subunit of the interleukin 1 receptor
complex
J Biol Chem 270: 13757-13765

Greisman, S. E. and Hornick, R. B. (1969)
Comparative pyrogenic reactivity of rabbit and man to bacterial endotoxin
Proc Soc Exp Biol Med 131: 1154

Guschin, D., Rogers, N., Briscoe, J., Witthuhn, B., Watling, D., Horn, F., Pellegrini, S.,
Yasukawa, K., Heinrich, P., Stark, G. R., et al. (1995)
A major role for the protein tyrosine kinase JAK1 in the JAK/STAT signal transduction
pathway in response to Interleukin-6
EMBO J (England) 14: 1421-29

Gutierrez, E. G., Banks, W. A., and Kastin, A. J. (1994)
Blood-borne interleukin-1 receptor antagonist crosses the blood-brain barrier
J Neuroimmunol 55: 153-160

Hailman, E., Lichenstein H. S., Wurfel, M. M., Miller, D. S., Johnson, D. A., Kelley, M.,
Busse, L. A., Zukowski, M. M., and Wright, S. D. (1994)
Lipopolysaccharide (LPS)-binding protein accelerates the binding of LPS to CD14
J Exp Med 179: 269-277

Hartung, T. and Wendel, A. (1996)
Detection of pyrogens using human whole blood
In Vitro Toxicology (9) 4: 353-59

Hartung, T., Fennrich, S., Fischer, M., Montag-Lessing, T. and Wendel, A. (2000)
Prevalidation of an alternative to the rabbit pyrogen test based on human whole blood
Balls, M., van Zeller, A. M., Halder, M. E. (eds.): 991-999
2000 Elsevier Science B. V.

Hayashi, F., Smith, K. D., Ozinsky, A., Hown, T. R., Yi, E. C., Goodlett, E. R., Eng, E. K., Akira, S., Underhill, D. M., and Aderem, A. (2001)

The innate immune response to bacterial flagellin is mediated by Toll-like receptor 5
Nature 410: 1099-1103

Hemmi, H., Takeuchi, O., Kawai, T., Kaisho, T., Sato, S., Sanjo, H., Matsumoto, M., Hoshino, K., Wagner, H., Takeda, K., and Akira, S. (2000)

A Toll-like receptor recognizes bacterial DNA

Nature 408: 740-45

Hirata, H., Takahashi, A., Kobayashi, S., Yonehara, S., Sawai, H., Okazaki, T., Yamamoto, K., and Sasada, M. (1998)

Caspases are activated in a branched protease cascade and control distinct downstream processes in Fas-induced apoptosis

J Exp Med 187: 587-600

Hirschfeld, M., Weis J. H., Vogel, S. N., and Weis J. J. (2000)

Cutting Edge: repurification of lipopolysaccharide eliminates signalling through both human and murine toll-like receptor 2

J Immunol 165: p 618-622

Hoffmann, E., Dittrich-Breiholz, O., Holtmann, H., et al.(2002)

Multiple control of interleukin-8 gene expression

J Leukoc Biol 72: 847-55

Hoffmann, S., Peterbauer, A., Schindler, S., Fennrich, S., Poole, S., Mistry, Y., Montag-Lessing, T., Spreitzer, I., Löschner, B., et al., (2005)

International validation of novel pyrogen tests based on human moncytoid cells

J Immunol Methods 298; 161-73

Hoffmann, S., Luederitz-Puechel, U., Montag-Lessing T., and Hartung, T. (2005)

Optimisation of pyrogen testing in parenterals according to different pharmacopoeias by probabilistic modelling

J Endotoxin Res 11; 25-31

Holtmann, H., Winzen, R., Holland, P., et al. (1999)
Induction of interleukin-8 synthesis integrates effects on transcription and mRNA
degradation from at least three cytokine- or stress-activated signal transduction
pathways
Mol Cell Biol 19: 6742-53

Horng, T., Barton, G. M., and Medzhitov, R. (2001)
TIRAP: an adapter molecule in the TOLL signalling pathway
Nat Immunol 2: 835-841

Horng, T. and Medzhitov, R. (2001)
Drosophila MyD88 is an adapter in the TOLL signalling pathway
Proc Natl Acad Sci USA 98: 12654-12658

Hort, E. and Penfold, W. (1912)
Microorganisms and their relation to fever
J Hyg 12: 361

Hort, E. and Penfold, W. (1912)
A critical study of experimental fever
Proc R Soc Lond (Biol) 85: 174

Hoshino, K., Takeuchi, T., Sanjo H., Ogawa T., Takeda Y., and Akira S. (1999)
Cutting edge: Toll-like receptor 4 (TLR-4) deficient mice are hyporesponsive to
lipopolysaccharide: evidence for tlr-4 as the LPS gene product
J Immunol 162: 3749-52.

Hsu, H., Xiong, J., and Goeddel, D. V. (1995)
The TNF receptor 1- associated protein TRADD signals cell death and NF-κB activation
Cell 81: 505-12

Inohara, N., Ogura, Y., Chen, F. F., Muto, A., and Nunez, G. (2001)
Human Nod 1 confers responsiveness to bacterial lipopolysaccharides
J Biol Chem 276: 2551-54

Jiang, Q., Akashi, S., Miyake, K., and Pett, H. R. (2000)
Cutting edge: Lipopolysaccharide induces physical proximity between CD14 and Toll-like receptor 4 (tlr4) prior to nuclear translocation of NF-kB
J Immunol 165: 3541-44

Jones, S. A., Wolf, M., Qin, S., Mackay, C. R. and Baggolini, M. (1996)
Different functions for the interleukin 8 receptors of human neutrophil leukocytes
NADPH oxidase and phospholipase D are activated through IL-8R1 but not IL-8R2
Proc Natl Acad Sci USA 93: 6682-86

Jones, B. W., Heldwein, K. A., Means, T. K., Saukkonen, J. J., Fenton, M. J. (2001)
Differential roles of TOLL-like receptors in the elicitation of proinflammatory responses by macrophages
Ann Rheum Dis 60 (Suppl 3): p iii6-12

Kabir, S., Rosenstreich, D. L., and Mergenhagen, S. E. (1978)
Bacterial endotoxin and cell membranes
In: Jeljaszewic, J., Wadstrom, T. (eds) Bacteria toxins and cell membranes. Academic, London: 59-87

Kampschmidt, R. F., Upchurch, H. F., and Worthington, M. L. (1983)
Further comparisons of endogenous pyrogens and leukocytic endogenous mediators
Infect Immun 126: 2474-9

Kawai, T., Adachi, O., Ogawa, T., Takeda, K., and Akira, S. (1999)
Unresponsiveness of MyD88-deficient mice to endotoxin
Immunity 11: 115-122

Kawai, T., Takeuchi, O., Fujita, T., Inoue, J., Muhlradt, P. F., Sato, S., Hoshino, K., and Akira, S. (2001)
Lipopolysaccharide stimulates the MyD88-independent pathway and results in activation of IFN-regulatory factor 3 and the expression of a subset of lipopolysaccharide-inducible genes
J Immunol 167: 5887-5894

- Kawasaki, K., Akashi, S., Shimazu, R., Yoshida, T., Miyake, K., and Nishijima, M. (2000) Mouse TOLL-like receptor4/MD-2 complex mediates lipopolysaccharide-mimetic signal transduction by Taxol
J Biol Chem 275: 2251-2254
- Kettelhut, I. C., and Goldberg, A. L. (1988) Tumor necrosis factor can induce fever in rats without activating protein breakdown in muscle or lipolysis in adipose tissue
J Clin Invest 81: 1384-1389
- Kirkland, T. N., Finley, F., Leturcq, D., Moriarty, A., Lee, J.-D., Ulevitch, R. D., and Tobias, P. S. (1993) Analysis of lipopolysaccharide binding by CD14
J Biol Chem 268: 24818-24823
- Klir, J. J., McClellan, J. L., and Kluger, M. J. (1994) Interleukin-1 beta causes the increase in anterior hypothalamic IL-6 during LPS-induced fever in rats
Am J Physiol 266: R1845-R1848
- Kobayashi, K., Inohara, N., Hernandez, L. D., Galan, J. E., Nunez, G., Janeway, C. A., Medzhitov, R., and Flavell, R. A. (2002) RICK/Rip2/CARDIAK mediates signalling for receptors of the innate and adaptive immune systems
Nature 416 (6877): 194-199
- Kozak, W., Kluger M. J., and Soszynski D. (1998) IL-6 and IL-1 β in fever. Studies using cytokine-deficient (knockout) mice
Ann NY Acad Sci 856: 33-47
- Laburn, H. P., Rosendorff, C., and Willies, G. (1974) A role for noradrenaline and cyclic AMP in prostaglandin E1 fever
J Physiol 240: 49P-50P

Lee, S. and Rivier, C. (1994)

Hypophysiotropic role and hypothalamic gene expression of corticotropin-releasing factor and vasopressin in rats injected with interleukin-1 β systemically or into the brain ventricles

J Neuroendocrinol 6: 217-224

Leonard, W. J. (1996)

STATS and cytokine specificity

Nature Med 2: 968-69

Levin, J. and Bang, F. B. (1964)

The role of endotoxin in the extravascular coagulation of Limulus blood

Bull Johns Hopkins Hosp: 265-274

Lien, E., Means, T. K., Heine, H., Yoshimura, A., Kusumoto, S., Fukase, K., Fenton, M.

J., Oikawa, M., Qureshi, N., Monks, B., Finberg, R. W., Ingalls, R. R., and Golenbock,

D. T. (2000)

TOLL-like receptor 4 imparts ligand-specific recognition of bacterial lipopolysaccharide

J Clin Invest 105: 497-504

Luheshi, G., Miller, A. J., Brouwer, S., Dascombe, M. J., Rothwell, N. J., and Hopkins,

S. J. (1996)

Interleukin-1 receptor antagonist inhibits endotoxin fever and systemic interleukin-6 induction in the rat

Am J Physiol 270 (1 Pt 1): E91-5

Luheshi, G., Stefferl, A., Turnbull, A., Dascombe, M. J., Brouwer, S., Hopkins, S. J., and

Rothwell, N. J. (1997)

Febrile response to tissue inflammation involves both peripheral and brain IL-1 and TNF- α in the rat

Am J Physiol 272: p R862-R868

Mannion, B. A., Weiss, J., and Elsbach, P. (1990)
Separation of lethal and sublethal effect of polymorphonuclear leucocytes on
Escherichia coli
J Clin Invest 86: 631-641

Medzhitov, R. and Janeway Jr., C. A. (1997)
Innate immunity: the virtues of a non-clonal system of recognition
Cell 91: 295-298

Medzhitov, R., Preston-Hurlburt, P., Janeway Jr., C. A. (1997)
A human homologue of the Drosophila TOLL protein signals activation of adaptive
immunity
Nature 388: 394-397

Medzhitov, R., Preston-Hurlburt, G., Timans, J. C., Kastelein, RA and Bazan, J. F.
(1998)
A family of human receptors structurally related to Drosophila Toll
Proc Natl Acad Sci USA 95: 588-93

Michie, H. R., Manogue, K. R., Spriggs, D. R., Revhaug, A., O'Dwyer, S., Dinarello,
C. A., Cerami, A., Wolf, S. M., and Wilmore, D. W. (1988)
Detection of circulating tumor necrosis factor after endotoxin administration
N Engl J Med 318: 1481- 1486

Morath, S., Geyer A. and Hartung, T. (2001)
Structure-function relationship of cytokine induction by Lipoteichoic Acid from
Staphylococcus aureus (Brief Definite Report)
J Exp Med 193: 393-397

Morath, S., Stadelmaier, A., Geyer, A., Schmidt, R. R., and Hartung, T. (2002)
Synthetic lipoteichoic acid from *Staphylococcus aureus* is a potent stimulus of cytokine release
J Exp Med 195: 1635-40

Morimoto, A., Murakami, N., Sakata, Y., Watanabe, T., and Yamaguchi, K. (1990)
Functional and structural differences in febrile mechanisms between rabbits and rats
J Physiol 427: 227-239

Morrison, D. C. (1989)
The case for specific lipopolysaccharide receptors expressed on mammalian cells
Microb. Pathogen 7: 389-398

Moser, B., Barella, L., Mattei, S., Schumacher, C., Boulay, F., Colombo, M. P., and Baggolini, M. (1993)
Expression of transcripts for two interleukin 8 receptors in human phagocytes, lymphocytes, and melanoma cells
Biochem J 294: 285-92

Mott, J.C. (1967)
The relation of blood volume to body weight and arterial haemoglobin levels in rabbits
J Physiol 191: 131-40

Mukaida, N., Shiroo, M., and Matsushima, K. (1989)
Genomic structure of the human monocyte-derived neutrophil chemotactic factor IL-8
J Immunol 143; 1366-71

Murakami, N., Sakata, Y., and Watanabe T. (1990)
Central action sites of Interleukin-1 β for inducing fever in rabbits
J Physiol 428: 299-312

Muzio, M., Ni, J., Feng, P., and Dixit, V. M. (1997)
IRAK (Pelle) family member IRAK-2 and MyD88 as proximal mediators of IL-1 signalling
Science 278: 1612-15

Nakamura, H., Seto, Y., Motoyoshi, S., Kadokawa, T., and Sunahara, N. (1988)
Recombinant human tumor necrosis factor causes long-lasting and prostaglandin-mediated fever, with little tolerance in rabbits
J Pharmacol Exp Ther 245: 336-341

Nijsten, M. W., de Groot, E. R., Ten Duis, H. J., Klasen, H. J., and Aaden, L. A. (1987)
Serum levels of interleukin-6 and acute phase response
Lancet 2: 921

Ooi, C. E., Weiss, J., Doerfler, M. E., and Elsbach, P. (1991)
Endotoxin-neutralizing properties of the 25 kD N-terminal fragment and the newly isolated 30 kD C-terminal fragment of the 55-60 kD Bactericidal/Permeability Increasing protein of human neutrophils
J Exp Med 174: 649-655

Ozinsky, A., Underhill, D. M., Fontenot, J. D., Hajjar, A. M., Smith, K. D., Wilson, C. B., Schroeder, L., and Aderem, A. (2000)
The repertoire for pattern recognition of pathogens by the innate immune system is defined by cooperation between TOLL-like receptors
Proc Natl Acad Sci USA 97: 13766-1377

Petersen, C. M., Moller, B. K. (1988)
Immunological reactivity and bioactivity of tumour necrosis factor
Lancet 23: 934-935

Poltorak, A., He, X., Smirnova, I., Liu, M. Y., Huffel, C. V., Du, X., Birdwell, D., Alejos, E., Silva, M., Galanos, C., Freudenberg, M., Ricciardi-Castagnoli, P., Layton, B., and Beutler, B. (1998)
Defective LPS-signaling in C3H/HeJ and C57BL/10ScCr mice: mutations in the tlr4 gene
Science 282: 2085-2088

Poltorak, A., Ricciardi-Castagnoli, P., Citterio, S., and Beutler, B. (2000)
Physical contact between lipopolysaccharide and TOLL-like receptor 4 revealed by
genetic complementation
Proc Natl Acad Sci USA 97: 2163-2167

Probey, T. F., and Pittman, M. (1945)
The pyrogenicity of bacterial contaminants found in biologic products
J Bacteriol 50: 397-411

Qureshi, ST., Lariviere, L., Leveque, G., Clermont, S., Moore, KJ., Gros, P., and
Malo, D. (1999)
Endotoxin-tolerant mice have mutations in toll-like receptor 4 (Tlr4)
J Exp Med 189: 615-25.

Rajarathnam, K., Sykes, BD., Kay, CM., Dewald, B., Geiser, T., Baggolini, M., and
Clark-Lewis, I. (1994)
Neutrophil activation by monomeric interleukin-8
Science 264: 90-92

Ray, A., Redhead, K., Selkirk, S., Poole, S.. (1991)
Variability in LPS composition, antigenicity and reactogenicity of phase variants of
Bordetella pertussis.
FEMS Microbiol Lett 63: 211-7

Rietschel, E. T., Brade, H., Brade, L., Brandenburg, K., Schade F. U., Seydel U.,
Zahringer U., Galanos, C., Luderitz, O., Westphal O., Labischinski, H., Kusumoto S.,
and Shiba T. (1987)
Lipid A, the endotoxic centre of bacterial lipopolysaccharides: relation of chemical
structure to biological activity
Prog Clin Biol Res 231: 25-34

Rietschel, E. T., Kirikae, T., Schade, F. U., Ulmer, A. J., Holst, O., Brade, H., Schmidt, G., Mamat, U., Grimmecke H.D., Kusumoto, S., and Zahringer, U. (1993)
The chemical structure of bacterial endotoxin in relation to bioactivity
Immunobiology 187 (3-5): 169-90

Rollinghoff, M., and Warner, N. L. (1973)
Specificity of in vivo tumor rejection assessed by mixing immune spleen cells with target and unrelated tumor cells
Proc Soc Exp Biol Med 144: 813-8

Rothe, J., Lesslauer, W., Lotscher, H., Lang, Y., Koebel, P., Kontgen, F., Althage, A., Zinkernagel, R., Steinmetz, M., and Bluethmann, H. (1993)
Mice lacking the tumour necrosis factor receptor 1 are resistant to TNF-mediated toxicity but highly susceptible to infection by *Listeria monocytogenes*
Nature 364: 798-802

Rothwell, N. J., Busbridge, N. J., Humphray, H., and Hissey, P. (1990)
Central actions of interleukin-1 β on fever and thermogenesis
The Physiological and Pathological Effects of Cytokines: 307-311. Wiley-Liss, Inc., New York

Rothwell, N. J. and Hopkins, S. J. (1995)
Cytokines and the nervous system II: Actions and mechanisms of action
TiNS 18: 130-136

Rosendorff, C. (1976)
The pathogenesis of fever: neurochemical aspects
Isr J Med Sci 12: 960-973

Rosendorff, C. (1976)
Neurochemistry of fever
S Afr J Med Sci 41: 23-48

Rossi, V., Breviario, F., and Ghezzi, P. (1985)
Prostacyclin synthesis induced in vascular cells by interleukin-1
Science 229: 174-176

Schindler S, Bristow A, Cartmell T., Hartung, T., and Fennrich, S.(2003)
Comparison of the reactivity of human and rabbit blood towards pyrogenic stimuli
ALTEX 20: 59-63

Schumann, R. R. (1992)
Function of lipopolysaccharide binding protein and CD14, the receptor for LPS/LBP
complexes: a short review
Res Immunol 143: 11-15

Schutt, C., Schilling, T., Grunwald, U., Schonfeld, W., and Kruger, C. (1992)
Endotoxin-neutralizing capacity of soluble CD14
Res Immunol 143: 71-8

Seibert, F. B. (1925)
The cause of many febrile reactions following intravenous injections
Am J Phys 71: 621

Sheagren, J. N., and Wolff, S. M. (1966)
Demonstration of endogenous pyrogen in afebrile rabbits
Nature 210(35): 539-40

Shimazu, R., Akashi, S., Ogata, H., Nagai, Y., Fukudome, K., et al. (1999)
MD-2, a molecule that confers lipopolysaccharide responsiveness on toll-like receptor 4
J Exp Med 189: 1777-82

Simon, P. L. and Willoughby, T. F. (1981)
The role of subcellular factors in pulmonary immune function: physicochemical
characterization of two distinct species of lymphocyte-activating factor produced by
rabbit alveolar macrophages
J Immunol 126: 1534-41

Smith, B., Kluger, K., and Kluger, M. J. (1992)
Human IL-1 receptor antagonist partially suppresses fever but not plasma levels of IL-6
in Fischer rats
Am J Physiol 263: R653-655

Stahl, N., Boulton, T. G., Farruggela, T., Ip, N. Y., Davis, S., Witthuhn, B. A., Quelle, F. W., Silvennoinen, O., Barbieri, G., Pellegrini, S., et al. (1994)
Association and activation of Jak-Tyk Kinases by CNTF-LIF-OSM-IL-6 beta receptor components
Science 263: 92-95

Standiford, T. J., Arenberg, A., Danforth, J. M., Kunkel, S. L., van Otteren, G. M., and Strieter, R. M. (1994)
Lipoteichoic acid induces secretion of interleukin-8 from human blood monocytes:
a cellular and molecular analysis
Infect Immun: 62: 119-25

Stanger, B. Z., Leder, P., Lee, T.-H., Kim, E., and Seed, B. (1995)
RIP: a novel protein containing a death domain that interacts with Fas-Apo-1 (CD95) in yeast and causes cell death
Cell 81: 513-23

Stein, D., Roth, S., Vogelsang, E., and Nusslein-Volhard, C. (1991)
The polarity of the dorsoventral axis in the Drosophila embryo is defined by an extracellular signal
Cell 65: 725-735

Stitt, J. T. (1985)
A study of the variability in the febrile responses of rabbits to endogenous pyrogen
J Appl Physiol 59: 1254-7

Suda, T., Tozawa, F., Ushiyama, T., Sumimoto, T., Yamada, M., and Demura, H. (1990)
Interleukin-1 stimulates corticotropin-releasing factor gene expression in rat
hypothalamus
Endocrinology 126: 1223-1228

Takeda, H. and Kotani, S. (1992)
in: D. C. Morrison, J. L. Ryans (Eds), Bacterial Endotoxic Lipopolysaccharides, Vol. 1
Molecular Biochemistry and Cellular Biology, CRC Press, Boca Raton, USA: 107-131

Takeda, K., and Akira, S. (2001)
Regulation of innate immune responses by Toll-like receptors
Jpn J Infect Dis 54: 209-19

Takeuchi, O., Hoshino, K., Kawai, T., Sanjo, H., Takanda, H., Takeda, K., and Akira, S. (1999)
Differential roles of TLR2 and TLR4 in recognition of gram-negative and gram-positive bacterial cell wall components
Immunity 11: 443-51

Takeuchi, O., Kaufmann, A., Grote, K., Kawai, T., Hoshino, K., Morr, M., Muhlradt, P. F., and Akira, S. (2000)
Cutting edge: preferentially the R- stereoisomer of the mycoplasmal lipopeptide macrophage-activating lipopeptide-2 activates immune cells through a TOLL-like receptor 2- and MyD88-dependent signaling pathway
J Immunol 164: 554-557

Takeuchi, O., Takeda, K., Hoshino, K., Adachi, O., Ogawa, T., et al (2000)
Cellular responses to bacterial cell wall components are mediated through MyD88-dependent signaling cascades
Int Immunol 12: 113-17

Takeuchi, O., Kawai, T., Muhlradt, P. E., Radolf, J. D., Zychlinsky, A., et al. (2001)
Discrimination of bacterial lipopeptides by toll-like receptor 6
Int Immunol 13: 933-40

Takeuchi, O., and Akira, S. (2001)
TOLL-like receptors: their physiological role and signal transduction system
Int Immunopharmacol 1: 625-635

Tanamoto, K., Zahringer, U., Mckenzie, G. R., Galanos, C., Rietschel, E.T.,
Luderitz, O., Kusumoto, S., and Shiba T. (1984)
Biological activity of synthetic lipid A analogs: pyrogenicity, lethal toxicity,
anticomplement activity and induction of gelation of Limulus amoebocyte lysate
Infect Immun 44: 421-426

Tennant, D. M., and Ott, W. H. (1953)
Tolerance to bacterial pyrogens in the rabbit
J Am Pharm Assoc 42: 614

Thornberry, N. A., Bull, H. G., Calaycay, J. R., Chapman, K. T., Howard, A. D.,
Kostura, M. J., Miller, D. K., Molineaux, S. M., Weidner, J. R., et al. (1992)
A novel heterodimeric cysteine protease is required for interleukin-1 β processing in
monocytes
Nature 356: 768-774

Tobias, P. S., Mathison, J. C., and Ulevitch, R. J. (1988)
A family of lipopolysaccharide binding proteins involved in response to
Gram-negative sepsis
J Biol Chem 263: 13479-13781

Tobias, P. S., Mathison, J., Mintz, D., Lee, J.-D., Kravchenko, V., Kato, K., Pugin, J., and
Ulevitch, R. J. (1992)
Participation of Lipopolysaccharide dependent macrophage activation.
Am J Respir Cell Mol Biol 7: 239-245

Tobias, P. S. and Ulevitch, R. D. (1993)
Lipopolysaccharide binding protein and CD14 in LPS-dependent macrophage activation
Immunobiol 187: 227-232

Tobias, P. S., Soldau, K., Gegner, J. A., Mintz, D., and Ulevitch, R. J. (1995)
Lipopolysaccharide binding protein-mediated complexation of lipopolysaccharide with
soluble CD14
J Biol Chem 270: 10482-10488

Tsagarakis, S., Gillies, G., Rees, L. H., Besser, M., and Grossmann, A. (1989)
Interleukin-1 directly stimulates the release of corticotropin releasing factor from rat
hypothalamus
Neuroendocrinology 49: 98-101

Ulevitch, R. J. (1993)
Recognition of bacterial endotoxin by receptor dependent mechanisms
Adv Immunol 53: 267-288

Underhill, D. M. and Ozinsky, A. (2002)
Toll-like receptors: key mediators of microbe detection
Curr Opin Immunol 14: 103-110

Van Dam, A. M., De Vries, H. E., Kuiper J., Zulstra, F. J., DeBoer, A. G., Tilders, F. J.,
and Berkenbosch, F. (1996)
Interleukin-1 receptors on rat brain endothelial cells: A role in neuroimmune interaction?
FASEB J. 10: 351-356

van Dijck, P., and van de Voorde, H. (1977)
Factors affecting pyrogen testing in rabbits
Dev Biol Stand 34: 57-63

van Zee, K. J., Coyle, S. M., Calvano, S. E., Oldenburg, H. S. A., Stiles, D. M., Pribble,
J., Catalano, M., Moldawer, L. L., and Lowry, S. F. (1995)
Influence of IL-1 receptor blockade on the human response to endotoxemia
J Immunol 154; 1499-1507

Von Aulock, S., Morath, S., Hareng, L., et al. (2003)
Lipoteichoic acid from *Staphylococcus aureus* is a potent stimulus for neutrophil recruitment (In Process Citation)
Immunobiology 208: 413-22

von Haller A. (1757-1766)
Elements physiologiae corporis humanae, Vol 3 (1757-1766)

Wachtel, R. E., and Tsuji, K. (1977)
Comparison of the limulus amoebocyte lysates and correlation with the USP pyrogen test
Appl and Environment Microbiology 33: 1265-1269

Weary, M., Pearson, F. C., Bohan, J., and Donahue, G. (1982)
Endotoxins and their detection in the LAL test
Alan R. Liss, Inc., 150 5th Avenue New York NY 10011, 365-379

Weiss, J. P., Elsbach, P., Shu, C., Castillo, J., Grinna, I., Horwitz, A., and Theofan, G. (1992)
Human Bactericidal/permeability-increasing protein and a recombinant N-terminal fragment cause killing of serum-resistant Gram-negative bacteria in whole blood and inhibit TNF release induced by the bacteria
J Clin Invest 90: 1122-1130

Werts, C., Tapping, R. I., Mathison, J. C., Chuang, T. H., Kravchenko, V., Saint Girons, I., Haake, D. A., Godowski, P. J., Hayashi, F., Ozinsky, A., Underhill, D. M., Kirschning, C. J., Wagner, H., Adererm, A., Tobias, P. S., and Ulevitch, R. J. (2001)
Leptospiral lipopolysaccharide activates cells through a TLR2-dependent mechanism
Nat Immunol 2: 346-352

Wesche, H., Henzel, WJ., Shillinglaw, W., Li, S., and Cao, Z. (1997)
MyD88: an adaptor protein that recruits IRAK to the IL-1 receptor complex
Immunity 7: 837-47

Westphal, O., Luderitz, O., and Biester F. (1952)
Extraction of bacteria with phenol/water
Naturforsch B.: Anorg Chem Org Chem Biochem Biophys Biol 7B: 148

Westphal, O., Westphal, U., and Sommer, T. (1977)
The history of pyrogen research
In Microbiology-1977, D. Schlesinger, ed. American Society for Microbiology;
Washington, D. C., 221.

Wright, S. D., Ramos, R. A., Tobias, P. S., Ulevitch, R. J. and Mathison, J. C. (1990)
CD14, a receptor for complexes of lipopolysaccharide (LPS) and LPS binding protein
Science 249: p 1431-1433

Wright, G. W., Ooi, C. E., Weiss, J., and Elsbach, P. (1990)
Purification of a cellular (granulocyte) and an extracellular (serum) Phospholipase A2 that
participate in the destruction of *Escherichia coli* in a rabbit inflammatory exsudate
J Biol Chem 265: 6675-6681

Wu, D., LaRosa, G. J., and Simon, M. I. (1993)
G protein-coupled signal transduction pathways for interleukin-8
Science 261: 101-3

Wurfel, M. M., Kunitake, S. T., Lichenstein, H., Vane, J. P., and Wright, S. D. (1994)
Lipopolsaccharide (LPS)-binding protein is carried on lipoproteins and acts as a
cofactor for the ventralization of LPS
J Exp Med 180: 1025-1035

Wurfel, M. W. and Wright, S. (1997)
Lipopolsaccharide-binding proteins and soluble CD14 transfer lipopolysaccharide to
phospholipid bilayers
J Immunol 158: 3925-3934

Yamamoto, M., Sato, S., Hemmi, H., Sanjo, H., Uematsu, S., Kaisho T., et al. (2002)
Essential role for TIRAP in activation of the signalling cascade shared by TLR2 and
TLR4

Nature 420: 324-329

Yang, R. B., Mark, M. R., Gray, A., Huang, A., Xie, M. H., Zhang, M., Goddard, A.,
Wood, W. I., Gurney, A. L., and Godowski, PJ (1998)

Toll like receptor 2 mediates lipopolysaccharide induced cellular signalling

Nature 395: 284-88

Yin, E. T., Galanos, C., Kinsky, S., Bradshaw, R. A., Wessler, S., Luderitz, O., and
Sarmiento, M. E. (1972)

lysate induced by purified lipopolysaccharides and lipid A from gram-negative bacteria
Biochem Biophys Acta 26: 284

Zhong, W. W., Burke, P.A., Drotar, M.E., et et al. (1995)

Effects of Prostaglandin E2, cholera toxin and 8-bromocyclic AMP on
lipopolysaccharide-induced gene expression of cytokines in human macrophages
Immunology 84: 446-52