

10. Literaturverzeichnis

- Acri, J.B., Brown, K.J., Saah, M.I., Grunberg, N.E. (1995):
Strain and age differences in acoustic startle responses and effects of nicotine in rats.
Pharmacol. Biochem. Behav. 50: 191-198
- Aghajanian, G.K., Andrade, R., Azmitia, E.C., Baumgarten, H.G. (1997):
Handbook of Experimental Pharmacology.
Serotonergic Neurons and 5-HT Receptors in the CNS.
Berlin, Heidelberg. New York, Springer-Verlag.
- Alekseeva, M.S., Elkin, V.I.; Fedorov, V.K. (1964):
Comparative genetic studies on the lability of the nervous system in rats with a high degree of sensitivity to sound stimuli and wistar rats.
Zh. Vyssh. Nerv. Deiat. Im. I. P. Pavlova. 14:110-5. Russian.
- American Psychiatric Association“ (1994):
Diagnostic and Statistical Manual of Mental Disorders, 4. Auflage,
American Psychiatric Association“ (Hrsg.), Washington, D.C.
- Andrews, N., File, S.E. (1993):
Handling history of rats modifies behavioural effects of drugs in the elevated plus-maze test of anxiety.
Eur. J. Pharmacol. 235: 109-112
- Archer, J. (1973):
Tests for emotionality in rats and mice: A review
Animal Behaviour 21: 205-235
- Askew, H., (1997):
Behandlung von Verhaltensproblemen bei Hund und Katze.
Blackwell Wissenschafts-Verlag, Berlin. Wien.
- Azmitia, E.C. (1978):
The serotonin-producing neurons of the midbrain median and dorsal raphe nuclei.
In: Iversen LL, Iversen SD und Snyder SH, (eds.).
Handbook of Psychopharmacology, Vol. 9.
New York, Plenum Press; pp. 233-314
- Azmitia, E.C. (2001):
Modern views on an ancient chemical: serotonin effects on cell proliferation, maturation, and apoptosis.
Brain Res. Bull. 56: 413-24

- Barnes, N.M., Sharp, T. (1999):
A review of central 5-HT receptors and their function.
Neuropharmacology 38 (8): 1083-152
- Baskin, L.M. (1976):
Verhalten der Klauentiere.
Moskau; Nauka; pp. 296
- Baumgarten, H.G. (1991):
Neuroanatomie und Neurophysiologie des zentralen 5-HT-Systems.
In: Heinrich, K., Hippus, H., Pödingner, W. (eds.). *Serotonin - ein funktioneller Ansatz für die psychiatrische Diagnose und Therapie?*;
Springer-Verlag, Berlin, Heidelberg; pp. 17-44
- Baumgarten, H.G., Grozdanovic, Z. (1997):
Anatomy of central serotonergic projection systems.
In: Baumgarten, H.G., Göthert, M. (eds.). *Serotonergic neurons and 5-HT receptors in the CNS*.
Springer-Verlag, Berlin, Heidelberg; pp. 41-89
- Barnett, S.A. (1975):
The rat. A study in behavior.
Chicago; London: The University of Chicago Press.
- Beaver, B.V. (1989):
Environmental enrichment for laboratory animals.
Ilar News 31 (2): 5-11
- Bennet, J. L., Aghajanian, G. K. (1974):
J. Life Science 15: 1935-1944
- Birmelin, I. (1990):
Behaviour of pet animals.
Dtsch. Tierärztl. Wochenschr. 97: 243-247
- Blakely, R, De Felice, L.J., Hartzell, H.C. (1994):
Molecular physiology of norepinephrine and serotonin transporters.
J. Exp. Biol. 196: 263-281
- Blanchard, R.J., Yudko, E.B., Rodgers, R.J., Blanchard, D.C. (1993):
Defense system psychopharmacology: an ethological approach to the pharmacology of fear and anxiety.
Behav. Brain Res. 58: 155-165

- Blanchard, R.J., Rodgers, R.J., Blanchard, D.C. (1991):
Risk assessment and animal models of anxiety.
In: B. Olivier, J. Mos, Slangen, J.L. (Hrsg.). *Animal models in Psycholopharmakology*.
Birkhäuser, Basel; pp. 117-134
- Boadle-Biber, M.C. (1993):
Regulation of serotonin synthesis.
Prog. Biophys. molec. Biol. 60: 1-15
- Bodnoff, S.R., Suranyi-Cadotte, B., Quirion, R., Meany, M.J. (1989):
A comparison of the effects of diazepam versus several typical and atypical
antidepressant drugs in an animal model of anxiety.
Psychopharmacol. 97: 277-279
- Borbely, A.A., Huston, J.P., Waser, P.G. (1975):
Control of sleep states in the rat by short light-dark cycles.
Brain Res. 95: 89-101
- Bouchon, R., Will, B. (1982):
Effects of early enriched and restricted environments on the exploratory and locomotor
activity of dwarf mice.
Behavioral and Neural Biology 35 (2): 174-186
- Brain, P.F., Büttner, D., Cost, A. P., Gregory, J.A., Koolhaas, J., Militzer, K., Ödberg, F.O.,
Scharmann, W., Stauffacher, M. (1993):
In: *International Workshop on the Accomodation of Laboratory Animals in Accordance
with Animal Welfare*.
Requirements, Berlin; pp. 1-14
- Britton, D.R., Britton, K.T. (1981):
A sensitive open field measure of anxiolytic drug activity.
Pharmacol. Biochem. Behav. 15: 577-582
- Broadhurst, P.L. (1958):
Determinants of emotionality in the rat: III. Strain differences.
J. Comp. Physiol. Psychol. 51: 55-59
- Broadhurst, P.L. (1960):
Experiments in psychogenetics.
In: H.J. Eysenck (eds.): *Experiments in personality, psychopharmacology*. Vol. I.
Routledge and Kegan Paul, London; pp. 729-763
- Bronstein, P.M., Hirsch, S.M. (1976):
Ontogeny of defensive reactions in Norway rats.
J. Comp. Physiol. Psychol. 90: 620 – 629

- Bunin, M.A., Wightman, R.M. (1999):
Paracrine neurotransmission in the CNS: involvement of 5-HT.
Trends Neurosci. 22 (9): 377-82
- Cadogan, A.K., Kendall, D.A., Fink, H., Mmarsden, C.A. (1994):
Social interaction increases 5-HT release and cAMP efflux in the rat ventral hippocampus
in vivo.
Behav. Pharmacol. 5: 299-305
- Calhoun, J.B. (1962):
The ecology and sociology of the Norway rat.
U.S. Government Printing Office, Washington
- Carli, M., Balducci, C., Samanin, R. (2000):
Low doses of 8-OH-DPAT prevent the impairment of spatial learning caused by
introhypocampal scopolamine through 5-HT(1A) receptors in the dorsal raphe.
Br. J. Pharmacol. 131: 375-81
- Castle, W.E. (1947):
Proc. Natl. Acad. Sci. USA 33: 109-117
- Chaouloff, F., Kulikov, A., Saieau, A., Castanon, N., Mormede, P. (1995):
Male Fischer 344 and Lewis rats display differences in locomotor reactivity, but not in
anxiety-related behaviors: relationship with the hippocampal serotonergic system.
Brain Res. 693: 169-178
- Chojnacka-Wojcik, E. (1995):
5-Hydroxytryptamine in the central nervous system.
Pharmakol. 47: 219-235
- Christmas, A.J., Maxwell, D.R. (1970):
A comparison of the effects of some benzodiazepines and other drugs on aggressive and
exploratory behaviour in mice and rats.
Neuropharmacology 9: 17-29
- Commissaris, R.L., Lyness, W.H., Rech, R.H. (1981):
The effects of *d*-lysergic acid diethylamide (LSD), 2,5-dimethoxy-4-methylamphetamine
(DOM), pentobarbital and methaqualone on punished responding in control and 5,7-
dihydroxytryptaminetreated rats.
Pharmacol. Biochem. Behav. 14: 617-623
- Cole, J.C., Rodgers, R.J. (1993):
An ethological analysis of the effects of chlordiazepoxide and bretazenil in the murine
elevated plus-maze.
Behav. Pharmacol. 4: 573-580

- Costall, B., Hendrie, C.A., Kelly, M.E., Naylor, R.J. (1987):
Actions of sulphiride and tiapride in a simple model of anxiety in mice.
Neuropharmacology 26: 195-200
- Costall, B., Jones, B.J., Kelly, M.E., Naylor, R.J., Tomkins, D.M. (1989):
Exploration of mice in a black and white test box: validation as a model of anxiety.
Pharmacol. Biochem. Behav. 32: 777-785
- Crabbe, J.C., Wahlsten, D., Dudek, B.C. (1999):
Genetics of mouse behaviour: interactions with laboratory environment.
Science 284: 1670-1672
- Crampe, H. (1877):
Landwirtsch. Jahrb. 14: 539-619
- Cummings, J.P., Felten, D.L. (1979):
A raphe dendrite bundle in the rabbit medulla.
J. Comp. Neurol. 183(1): 1-23
- Dahlström, A., Fuxe, K. (1964):
Evidence for the existence of monoamine-containing neurons in the central nervous system.
I. Demonstration of monoamines in the cell bodies of brain stem neurons.
Acta Physiol. Scand. 62 [Suppl 232]: 1-55
- Diaz J-L. (1988):
Brain weights correlate with behavioral parameters in individual inbred mice housed in a common and enriched environment.
Behavioral and Neural Biology 50 (2): 164-183
- Drawer, K., Ennulat, K.J. (Hrsg.) (1977):
Tierschutzgerechtes Töten von Wirbeltieren. *Tierschutzpraxis.*
Gustav Fischer Verlag Stuttgart, New York; pp. 293-300
- Duisberi, D. (1981):
Das Verhalten der Tiere.
Vergleichsaspekten. *Die Welt.* Moskau; 479
- Durcan, M.J., Lister, R.G. (1989):
Does directed exploration influence locomotion activity in a holeboard test?
Behav. Neural. Biol. 51: 121-125
- Edwards, J.G. (1991):
Clinical anxiety and its treatment.
Neuropeptides 19 (Suppl.): 1-10

- Erpsamer V., Asero B. (1952):
Identification of enteramine, specific hormone of enterochromaffin system, as 5-hydroxytryptamine.
Nature 168: 800-801
- Estler, C. (1994):
Pharmakologie und Toxokologie. Lehrbuch für Mediziner, Veterinärmediziner, Pharmazeuten und Naturwissenschaftler.
Schattauer, Erlangen.: 4. Auflage
- Fernandez-Guasti, A., Hong, E., Lopez-Rubalcava, C. (1992):
Species differences in the mechanism through which the serotonergic agonists indorenate and ipsapirone produce their anxiolytic action.
Psychopharmacol. 107: 61-68
- File, S. E. (1977):
Effects of parachlorophenylalanine and amphetamine on habituation of exploration.
Pharmac. Biochem. Behav. 6: 151-156
- File, S.E. (1978):
Can social interaction be used to measure anxiety?
Brain J. Pharmacology 62: 19-24
- File, S.E. (1980):
The use of social interaction as a method for detecting anxiolytic activity of chlordiazepoxide-like drugs.
Neurosci. Methods 2: 219-238
- File, S.E. (1985a):
What can be learned from the effects of benzodiazepines on exploratory behavior?
Neurosci. Biobehav. Rev. 9: 45-54
- File, S.E. (1985b):
Models of anxiety.
Br. J. Clin. Pract. Suppl. 39: 15-19
- File, S.E. (1987):
The contribution of behavioural studies to the neuropharmacology of anxiety.
Neuropharmacology 26: 877-886
- File, S.E. (1990):
Age and anxiety: increased anxiety, decreased anxiolytic, but enhanced sedative, response to chlordiazepoxide in old rats.
Hum. Psychopharmacol. 5: 169-173

- File, S.E. (1992):
Behavioral detection of anxiolytic action.
In: Elliot J.H., Heal D.J., Marsden, C.A. (eds.). *Experimental Approaches to Anxiety and Depression*.
Wiley, Chichester; Chapter III: 25-44
- File, S.E., Wardill, A.G. (1975a):
The reliability of the hole-board apparatus.
Psychopharmacologia 44: 47-51
- File, S.E., Wardill, A.G. (1975b):
Validity of head-dipping as a measure of exploration in a modified hole-board.
Psychopharmacology 44: 53-59
- File, S.E., Hyde, J.R.G. (1977):
The effects of *p*-chlorophenylalanine and ethanolamine-*O*-sulphate in an animal test of anxiety.
Br. J. Pharm. Pharmac. 20: 735-738
- File, S.E., Hyde, J.R.G. (1978):
Can social interaction be used to measure anxiety?
Br. J. Pharmacol. 62: 19-24
- File, S.E., Vellucci, S.V. (1979):
Behavioural and biochemical measures of stress in hooded rats from different sources.
Physiol. Behav. 22: 31-35
- File, S.E., Pellow, S. (1985):
No cross-tolerance between the stimulatory and depressant actions of benzodiazepines in mice.
Behav. Brain Res. 17: 1-7
- Fischer, H. (1971):
Vergleichende Pharmakologie von Überträgersubstanzen in tiersystematischer Darstellung.
Berlin, Heidelberg, New York: Springer Verlag.
- Flossendorf, B. (1988):
Angst.
In: R. Asanger u.G. Wenninger (Hrsg.): *Handwörterbuch der Psychologie*.
4. Aufl. Verlagsunion Psychologie, München; pp. 34 - 36
- Franck, S. (2001):
Untersuchungen des Angstverhaltens der Transgenen Ratte TGR(mRen2)²⁷.
Dissertation des Institutes für Pharmakologie und Toxikologie der FU- Berlin.

- Freud, S., Breuer, J. (1993):
Studien über Hysterie.
Fischer Taschenbuch-Verlag, Frankfurt a.M., 1. Auflage
- Frussa-Filho, R., Otoboni, J.R., Uema, F.T., Sa-Rocha, L.C. (1991):
Evaluation of memory and anxiety in rats observed in the elevated plus-maze: effects of age and isolation.
Braz. J. Med. Biol. Res. 24: 725-728
- Gaddum, J.H., Picarelli, Z.P. (1957):
Two kinds of tryptamine receptors.
Br. J. Pharmacol. 12: 323-328
- Garattini, S., Valzelli, L. (1965):
Serotonin.
Amsterdam. Elsevier.
- Gardner, E.B., Boitano, J.J., Mancino, N.S., D'Amico, D.P., Gardner, E.L. (1975):
Environmental enrichment and deprivation: Effects on learning, memory and exploration.
J. Physiology and Behavior 14 (3): 321-327
- Geller, I., Seifter, J. (1960):
The effects of meprobamate, barbiturates, d-amphetamine and promazine on experimentally induced conflict in the rat.
Psychopharmacologia 1: 482-495
- Gentsch, C., Lichtsteiner, M., Feer, H. (1988):
Genetic and environmental influences on behavioral and neurochemical aspects of emotionality in rats.
In: *Experientia* (44): Reviews, Birkhäuser Verlag, Basel; pp. 482-490
- Gerlai, R. (1996):
Gene-targeting studies of mammalian behavior: Is it the mutation or the background genotype?
Trends Neurosci. 19: 177-181
- Gesner, C. (1669):
Thier-Buch (partial edition of *Historiae animalium* in German), reprinted 1980 by Schlütersche Verlagsanstalt, Hannover.
- Gonyou, H.W. (1994):
Why the study of Animal Behaviour is associated with the Animal Welfare Issue.
J. Animal Sci. 72: 2171-2177

- Griebel, G., Belzung, C., Misslin, R., Vogel, E. (1993):
The free-exploratory paradigm: An effective method for measuring neophobic behaviour in mice and testing potential neophobia-reducing drugs.
Behav. Pharmacol. 4: 637-644
- Griebel, G. (1995):
5-Hydroxytryptamine-interacting drugs in animal models of anxiety disorders: more than 30 years of research.
Pharmacol. Biochem. Behav. 65: 319-395
- Green, S., Hodges, H. (1991):
Animal models of anxiety.
In: Willner P. (ed.). Behavioral Models in Psychopharmacology.
Cambridge University Press.
- Grzimek, B. (1968):
Enzyklopädie des Tierreiches.
Vol. 11. Zürich: Kindler Verlag.
- Guitart, X., Kogan, J.H., Berhof, M., Terwillinger, R.Z., Aghajanian, G.K., Nestler, E.J. (1993):
Lewis and Fischer rat strains display differences in biochemical, electrophysiological and behavioral parameters: Studies in the nucleus accumbens and locus coeruleus of drug naive and morphine-treated animals.
Brain Res. 611: 7-17
- Handley, S.L., Mithani, S. (1984):
Effects of alpha-adrenoceptor agonists and antagonists in a maze-exploration model of „fear“-motivated behaviour.
Nanyn Schmiedberg's Arch. Pharmacol. 327: 1-5
- Handley, S.L., M^cBlane, J.W. (1993):
5-HT drugs in animal models of anxiety.
Psychopharmacol. 112: 13-20
- Hall, C.S. (1934):
Emotional behaviour in the rat. I. Defecation and urination as measures of individual differences in emotionality.
J. Comp.Psychol. 18: 385-403
- Haemisch, A. (1994):
Auswirkungen einer räumlich strukturierten Käfighaltung auf soziale Prozesse in Gruppen männlicher Mäuse.
Tierlaboratorium 17: 148-157

- Heinz, A., Jones, D.W., Mazzanti, C., Goldman, D., Ragan, P., Hommer, D., Linnoila, M., Weinberger, D.R. (2000):
A relationship between serotonin transporter genotype and in vivo protein expression and alcohol neurotoxicity.
Biol. Psychiatry 47: 643-649
- Hendrie, C.A., Weiss, S.M., Eilam, D. (1996):
Exploration and predation models of anxiety: Evidence from laboratory and wild species.
Pharmacol. Biochem. Behav. 54/1: 13-20
- Hedrich, H.J. (2000):
History, strains and models
in: G.J. Krinke (Hrsg.): The laboratory rat.
Academic Press, London, San Diego. Pp. 3 - 7
- Hlinak, Z., Rozmarova, E. (1986):
The locomotor-exploratory behaviour of laboratory male rats tested under the „red“ and „white“ light condition.
Activ. Nerv. Sup. 3: 28
- Horter, M. (1986):
Verhaltensweisen der Ratte als Ausdruck von Wohlbefinden und Unwohlsein unter besonderer Berücksichtigung der Wildform.
In: Militzer K. (Hrsg.). Wege zur Beurteilung tiergerechter Haltung von Labor-, Zoo- und Haustieren.
Parey, Berlin und Hamburg; pp. 33-34
- Hoyer, D., Clarke, D.E., Fozard, J.R., Hartig, P.R., Martin, G.R., Mylecharane, E.J., Saxena, P.R., Humphrey, P.P. (1994):
International Union of Pharmacology classification of receptors for 5-hydroxytryptamine (Serotonin).
Pharmacol. Rev. 46 (2): 157-203
- Hoyer, D., Martin, G. (1997):
5-HT receptor classification and nomenclature: towards a harmonization with the human genome.
Pharmacology 36: 419-428
- Hsia, L.C., Wood-Gush, G.M. (1984):
Social facilitation in the feeding behaviour of pigs and the effect of rank.
Appl. Anim. Ethol. 11: 265-270
- Hughes, R.N. (1968):
Behaviour of male and female rats with a free choice of two environments differing in novelty.
Animal Behav. 16: 92-96

- Imhof, J.T., Coelho, Z.M., Schmitt, M.L., Morato, G.S., Carobrez, A.P. (1993):
Influence of gender and age on performance of rats in the elevated plus maze apparatus.
Behav. Brain Res. 56: 177-180
- Iversen, S.D. (1984):
5-HT and anxiety.
Neuropharmacol. 23: 1553-1560
- Jacobs, B.L., Gannon, P.J., Azmitia, E.C. (1984):
Atlas of serotonergic cell bodies in the cat brainstem: an immunocytochemical analysis.
Brain Res. Bull. 13: 1-31
- Janowska, E., Pucilowski, O., Kostowski, W. (1991):
Chronic oral treatment with diltiazem or verapamil decreases isolation-induced activity impairment in elevated plus-maze.
Behav. Brain Res. 43: 155-158
- Johnston, A.L., File, S.E. (1991):
Sex differences in animal tests of anxiety.
Physiol. Behav. 49: 245-250
- Juhr, N.C. (1977):
Ultraschall-Laute von Ratten - Indices für eine Verhaltensbeurteilung.
Tierlaboratorium 4: 240 – 252
- Juhr, N.C. (1990):
Verhaltensindikatoren für Wohlbefinden.
Tierlaboratorium 13: 192 - 220
- King, H.D. (1919):
J. Exp. Zool. 29, 135-175.
- Konusova, A.V., Popova, N.K. (1982):
Role of central and peripheral serotonergic mechanisms.
Fiziol. Zh. SSSR Im I M. Sechenova. 68(10):1416-20. Russian
- Koriakina, L.A., Lobacheva, I.I., Popova, N.K. (1979):
Catecholamine sensitivity of the hypophyseal-adrenal system in the early ontogeny of rats.
Zh. Ontogenez. 10 (3): 270-5. Russian
- Kosten, T.A., Miserendino, M.J., Chi, S., Nestler, E.J. (1994):
Fischer and Lewis rat strains show differential cocaine effects in conditioned place preference and behavioral sensitization but not in locomotor activity or conditioned taste aversion.
J. Pharmacol. Exp. Ther. 269: 137-144

- Kovalevskij, K.L. (1963):
Auto-Infection (Intestinal) in radiation sickness and its prevention in Wistar white Rats.
Akad. Nauk. SSSR Inst. Genet. 30:315-20. Russian
- Kruschinskij, L.B. (1960):
Verhaltensbildung des Tieres bei normalen und pathologischen Bedingungen.
Akademischer Verlag., MGU., Moskau; pp. 260
- Kulikov, A., Auguerre, S., Berton, O., Ramos, A., Mormede, P. (1997):
Central serotonergic systems in the spontaneously hypertensive and Lewis rat strains that differ in the elevated plus-maze test of anxiety.
J. Pharmacol. Experiment. Behav. 281: 775-784
- Laininger, M. (1989):
Kann mit einfachen Verhaltenstests das Wohlbefinden von Labortieren beurteilt werden? Verhaltensuntersuchungen am Beispiel der Ratte.
Dissertation des Institutes für Pharmakologie und Toxikologie der FU- Berlin.
- Lal, H., Emmett-Oglesby, M.W. (1983):
Behavioral analogues of anxiety animal models.
Neuropharmacology 22 (12B): 1423-1441
- Lantz, D.E. (1909):
US Dept. of Agriculture, Biol. Survey Bull. 33: 9-54.
- Larsson, K. (1956):
Conditioning and sexual behavior in the male albino rat.
Almquist & Wiksell, Stockholm
- Leach, M.C., Ambrose N., Bowell V.J., Morton, D.B. (2000):
The development of a novel form of mouse cage enrichment.
Journal of Applied Animal Welfare Science 3 (2): 81-91
- Lesch, K.P., Balling, U., Gross, J., Strauss, K., Wolozin, B.L., Murphy, D.L., Riederer, P. (1994):
Organization of the human serotonin transporter gene.
J. Neural Transm. Gen. Sect. 95: 157-162
- Lesch, K.P., Bengel, D., Heils, A., Sabol, S.Z., Greenberg, B.D., Petri, S., Benjamin J., Müller C.R., Hamer D.H., Murphy D.L. (1996):
Association of anxietyrelated traits with a polymorphism in the serotonin transporter gene regulatory region.
Science 274: 1527-1531
- Lesch, K.P., Meyer, J., Flügge, G., Hinney, A., Hebebrand, J., Klauck, S.M. (1997):
The 5-HT transporter gene-linked polymorphic region (5-HTTLPR) in evolutionary perspective: Alternative biallelic variation in rhesus monkeys.
J. Neural Transm. Gen. Sect. 104: 1259-1266

- Lesch, K.P., Mössner, R. (1998):
Genetically driven variation in serotonin uptake: is there a link to affective spectrum, neurodevelopmental, and neurodegenerative disorders?
Biol. Psychiatry 44: 179-192
- Lister, R.G. (1987a):
The use of a plus-maze to measure anxiety in the mouse.
Psychopharmacology 92: 180-185
- Lister, R.G. (1987b):
The effects of ethanol on exploration in DBA/2 and C57BL/6 mice.
Alcohol 4: 17-19
- Lister, R.G. (1990):
Ethologically-based animal models of anxiety disorders.
Pharmacol. Ther. 46: 321-340
- Little, K.Y., McLaughlin, D.P., Zhang, L., Livermore, C.S., Dalack, G.W., McFinton, P.R., Del Proposto, Z.S., Hill, E., Cassin, B.J., Watson, S.J., Cook, E.H. (1998):
Cocaine, ethanol, and genotype effects on human midbrain serotonin transporter binding sites and mRNA levels.
Am. J. Psychiatry 155: 207- 213
- Lorens, S.A. (1978):
Some behavioral effects of serotonin depletion depend on method: A comparison of 5,7-dihydroxytryptamine, p-chlorophenylalanine, p-chloroamphetamine and electrolytic raphe lesions.
Ann. NY Acad. Sci. 305: 532-555
- Lorenz, K. (1971):
Studies in Animal and Human Behavior.
Harvard University Press, Cambridge
- Lorenz, K. (1994):
Aggression.
Moskau; Progress; pp. 272
- Lorz, A. (1987):
Tierschutzgesetz mit Rechtsverordnungen und europäischen Übereinkommen.
Kommentar, 3. völlig neubearbeitete und erweiterte Auflage.
C.H. Beck'sche Verlagsbuchhandlung, München.
- Manosewitz, M. (1970):
Early environmental enrichment and mouse behavior.
Journal of Comparative and Physiological Psychology 71 (3): 459-466

- Manosewitz, M., Montmayor, R.J. (1972):
Interaction of environmental enrichment and genotype.
Journal of Comparative and Physiological Psychology 79 (1): 67-76
- Manosewitz, M., Joel, U. (1973):
Behavioral effects of environmental enrichment in randomly bred mice.
Journal of Comparative and Physiological Psychology 85 (2): 373-382
- Marcucci, F., Mussini, E., Fanelli, R., Garattini, S. (1970):
Species differences in diazepam metabolism – I. Metabolism of diazepam metabolites.
Biochem. Pharmacol. 19: 1847-1851
- Martin, J.E., Edwards, S.A. (1994):
Feeding behaviour of outdoor sows: the effects of diet quantity and type.
Applied Animal Science 41.N1-2: 63-74
- Marsden, C.A., Bickerdike, M., Cadogan, A., Wrigth, I., Rex, A., Fink, H. (1993):
Serotonergic mechanisms and animal models of anxiety.
In: M. Hamon, H. Ollat, M.-H. Thibbot, (eds.). *Anxiety: Neurobiology, Clinic and Therapeutic Perspectives Colloque Inserm*; pp. 75-82
- Marsden, C. A., Beckett, S. R. G., Wilson, W., Bickerdike, M., Fink, H., Rex, A., Cadogan, A.-K. (1995):
Serotonin involvement in animal models of anxiety and panic.
In: A. Takade, G. Curzon (eds.). *Serotonin in the central nervous system and Periphery*. Elsevier, Amsterdam; pp. 135-143
- Marsden, C.A. (1996):
The neuropharmacology of serotonin in the central nervous system.
In: J.P. Feighner, Boyer (eds.). *Selective serotonin re-uptake inhibitors*. Wiley, Chichester; pp. 1-33
- McBlane, J.W, Critchley, M.A.E., Handley, S.L. (1992):
Light intensity influences response to 8-OH-DPAT in the elevated X-maze model of anxiety.
Br. J. Pharmacol. 105 (Suppl.): 221
- McCay, C.M. (1973):
Studies of animals in the fasting.
In: Verzar, F. (ed.) *Notes on the History of Nutrition Reserch*. Berne: Hans Huber; pp. 109
- Menard, J., Treit, D. (1999):
Effects of centrally administered anxiolytic compounds in animal models of anxiety.
Neurosc. Biobehav. Rev. 23: 591-613

- Miczek, K.A., Weertz, E.M., Vivian, J.A., Barros, H.M. (1995):
Aggression, anxiety and vocalization in animals: GABA-A und 5-HT anxiolytics.
Psychopharmacol. 121: 38-56
- Miczek, K.A., Maxson, S.C., Fish, E.W., Faccidomo, S. (2001):
Aggressive behavioural phenotypes in mice.
Behav. Brain. Res. 125: 167-81
- Misslin, R., Ropartz, P. (1981):
Effects of methamphetamine on novelty-seeking behaviour by mice.
Psychopharmacology 75: 39-43
- Mitchell, P.J. (1993):
Ethological studies of the social behaviour of the rat.
Anim. Technol. 44 (2): 109-120
- Molewijk, H.E., van der Poel, A.M., Olivier, B. (1995):
The ambivalent behaviour „stretched approach posture“ in the rat as a paradigm to
characterize anxiolytic drugs.
Psychopharmacol. 121: 81-90
- Molliver, M.E. (1987):
Serotonergic neuronal systems: what their anatomic organization tells us
about function.
J. Clin. Psychopharmacol. 7 (6 Suppl): 3-23
- Montgomery, S.A. (1955):
The relation between fear induced by novel stimulation and exploratory behaviour.
J. Comp. Physiol. Psychol. 48: 254-260
- Naumenko, E.V., Popova, N.K., Starygin, A.G. (1971):
Pituitary-adrenal system of animals in groups and in isolation.
Zh. Obshch. Biol. 32 (6): 731-9. Russian.
- Nevison, C.M., Hurst, J.L., Barnard, C.J. (1999):
Strain-specific effects of cage enrichment in male laboratory
mice (*Mus musculus*).
Animal Welfare 8 (4): 361-379
- Nikulina, E.M., Popova, N.K. (1983):
Genetische Analyse der Angreiferaggressivität bei Mäusen.
Genetik. 19: 1105-1110. Russian
- Nikulina, E.M., Borodin, P.M., Popova, N.K. (1985):
Veränderung einiger Formen des Agressivverhaltens und Monoaminmenge im Gehirn bei
Wildratten unter Selektionsprozessen in Richtung Domestikation.
Journal zentrales Nervensystem 35: 703-708. Russian

- Noirot, E. (1972):
Ultrasounds and maternal behaviour in small rodents.
Dev. Psychobiol. 5: 371-387
- Oliverio, A., Eleftheriou, B.E., Bailey, D.W. (1973):
Exploratory activity: Genetic analysis of its modification by scopolamine and amphetamine.
Physiol. Behav. 10: 893-899
- Onaivi, E.S., Maguire, P.A., Tsai, N.F., Davies, M.F., Loew, G.H. (1992):
Comparison of behavioral and central BDZ binding profile in three rat lines.
Pharmacol. Biochem. Behav. 43: 825-831
- Overstreet, D.H., Rezvani, A.H. (1996):
Behavioral differences between two inbred strains of Fawn-Hooded rat: a model of serotonin dysfunction.
Psychopharmacol. 128: 328-330
- Owens, M.J., Nemeroff C.B. (1994):
Role of serotonin in the pathophysiology of depression: focus on the serotonin transporter.
Clin. Chem. 40: 288-295
- Palm, M. (1998):
The incidence of chronic progressive nephrosis in young Sprague-Dawley rats from two different breeders.
Lab. Anim. 32: 477-482
- Paxinos, G., Watson. C. (1997):
The rat brainstereotaxic coordinates.
Academic Press, Orlando.
- Pellow, S., Chopin, P., File, S. E., Briley, M. (1985):
Validation of open/closed arm entries in an elevated plus-maze as a measure of anxiety in the rat.
J. Neurosci. Methods 14: 149-167
- Pellow, S., File, S.E. (1986):
Anxiolytic and anxiogenic drug effects on exploratory activity in an elevated-plus-maze: a novel test of anxiety in the rat.
Pharmacol. Biochem. Behav. 43: 471-477
- Peripato, A.C., De Brito, R.A., Vaughn, T.T., Pletscher, L.S., Marioli, S.R.,
Cheverud, J.M. (2002):
Quantitative trait loci for maternal performance for offspring survival in mice.
Genetics 162: 1341-53

- Persch, A. (1994):
Die Bedeutung des Explorationsverhaltens bei Versuchstieren
Tierärztliche Umschau 49: 539-545
- Pfeuffer, C. (1996):
Wahlversuche zur Haltung von Laborratten
Vet. Med. Diss. FU- Berlin.
- Philipeaux, C.R. (1856):
Studies on the importance of the adrenal glands.
Hebd. Seances Acad. Sci. 43: 904-906
- Poole, T.B. (1992):
The nature and evolution of behavioural needs of mammals
Animal Welfare 1: 203-220
- Popova, N.K. (1971):
Intragroup behavior and the pituitary-adrenal system in albino rats.
Fiziol. Zh. SSSR Im I M. Sechenova. 57(11):1723-7. Russian.
- Popova, N.K., Naumenko, E.V. (1972):
Dominance relations and the pituitary-adrenal system in rats.
J. Anim. Behav. 20(1):108-11. Russian.
- Popova, N.K., Naumenko, E.V., Kolpakov, V.G. (1978):
Serotonin and Behavior.
Nauka, Nivisibirsk; pp. 304. Russian
- Popova, N.K., Avgustinovitch, D.F., Kolpakov, V.G., Plyusnina, I. (1998):
Specific (3H) 8-OH DPAT binding in brain regions of rats genetically predisposed to various defense behaviour strategy.
Pharmacology Biochemistry and Behaviour 59: 793-797
- Porsolt, R.D., Bertin, A., Jalfre, M. (1978):
„Behavioral despair“ in rats and mice: strain differences and the effects of imipramine.
Eur. J. Pharmacol. 51: 291-294
- Postnov, Iu.V., Orlov, S.N., Shevchenko, A.S. (1975a):
Disorder of erythrocyte membrane permeability in spontaneous genetic hypertension in rats.
J. Kardiologija 15 (10): 88-92. Russian
- Postnov, Iu.V., Orlov, S.N., Shevchenko, A.S. (1975b):
Characteristics of erythrocyte membrane permeability for potassium ions in rats with familial spontaneous hypertension.
J. Kardiologija 15 (11): 61-4. Russian

- Ramos, A., Berton, O., Mormede, P., Chaouloff, F. (1997):
A multiple-test study of anxiety-related behaviours in six inbred rat strains.
Behav. Brain Res. 85: 57-69
- Rapport, M.M., Green, A.A., Page, I.H. (1948a):
Partial purification of the vasoconstrictor in beef serum.
J. Biol. Chem. 174: 735-738
- Rapport, M.M., Green, A.A., Page, I.H. (1948b):
Crystalline Serotonin.
Science 108: 329-330
- Renner, M.J., Rosenzweig, M.R. (1987):
Enriched and impoverished environments: Effects on Brain and Behavior.
Springer Verlag, New York.
- Renner, M.J. (1987):
Experience-dependent changes in exploratory behavior in the adult rat
(*Rattus norvegicus*): Overall activity level and interactions with objects.
Journal of Comparative Psychology 101 (1): 94-100
- Rex, A., Marsden, C.A., Fink, H. (1993):
Effect of diazepam on cortical 5-HT release and behavior in the guinea-pig on exposure
the elevated plus maze.
Psychopharmacol. 110: 490-496
- Rex, A., Sondern, U., Voigt, M., Franck, S., Fink, H. (1996):
Strain differences in fear-motivated behavior of rats.
Pharmacol. Biochem. Behav. 54: 107-111
- Rex, A., Fink, H. (1997):
Behavioural and neurochemical differences in Fischer 344 and Harlan-Wistar rats under
aversive conditions.
Soc. Neurosci. Abstr. 23 (929.4): 2384
- Rex, A., Voigt, J.P., Voits, M., Fink, H. (1998):
Pharmacological evaluation of a modified open-field test sensitive
to anxiolytic drugs.
Pharmacol. Biochem. Behav. 59: 677-683
- Rex, A., Voigt, J.P., Fink, H. (1999):
Behavioral and neurochemical differences between Fischer 344 and Harlan-Wistar rats
raised identically.
Behav. Genet. 29: 187-192

- Rex, A., Thomas, H., Hörtnagl, H., Voits, M., Fink, H. (2002):
Behavioural and microdialysis study after neurotoxic lesion of the dorsal raphe nucleus in rats.
Pharmacol., Biochem. And Behav. 74: 587-593
- Ridet, I., Privat, A. (2000):
Volume transmission.
Trends Neurosci. 23 (2): 58-9
- Rilov, A.M. (1985):
Aggressionsänderungen bei männlichen Ratten unter Einwirkung von Nahrungsdeprivation.
Journal zentrales Nervensystem. 35: 875-878, Russian
- Rodgers, R.J., Cole, J.C. (1994):
The elevated-plus-maze: pharmacology, methodology and ethology.
In: S.J. Cooper, C.A. Hendrie (eds.). Ethology and Psychopharmacology.
John Wiley, Chichester Ltd; pp. 9-44
- Rodgers, R.J., Cao, B.J., Dalvi, A., Holmes, A. (1997):
Animal models of anxiety: an ethological perspective.
Brazilian Journal of Medical and Biological Research 30: 289-304
- Roeder, J.J., Chetecuti, Y., Will, B. (1980):
Behavior and length of survival of populations of enriched and impoverished rats in the presence of a predator.
Biology of Behavior 5: 361-369
- Sambraus, H.H. (1997):
Grundbegriffe im Tierschutz.
In: H.H. Sambraus u. A. Steiger (Hrsg.): Das Buch vom Tierschutz.
Verlag Enke, Stuttgart; pp. 30-39
- Savory, W.S. (1863):
Measurement of the quality of proteins.
Lancet 2: 381-383
- Scharmann, W. (1991):
Improved housing of mice, rats and guinea-pigs:
A contribution to the refinement of animal experiments
Atlas 19: 108-114
- Scheibel, M.E., Tomiyasu, U., und Scheibel, A.B. (1975):
Do raphe nuclei of the reticular formation have a neurosecretory or vascular sensor function?
Exp. Neurol. 47(2): 316-29

- Schildlein, S., Huston, J.P., Schwarting, R.K. (2000):
Injections of tacrine and scopolamine into the nucleus accumbens: opposing effects of immediate vs delayed posttrial treatment on memory of an open field.
Neurobiol. Learn. Mem. 73: 21-30
- Shanks, N., Griffiths, J., Amisman, H. (1994):
Norepinephrine and serotonin alterations following chronic stressor exposure: Mouse strain differences.
Pharmacol. Biochem. Behav. 65: 57-65
- Smith, T.A. (1971):
The occurrence, metabolism and functions of amines in plants.
Biol. Rev. Camb. Philos. Soc. 46(2): 201-41
- Smidt, D., Andreae U., Unselm J. (1980):
Ist "Wohlbefinden" messbar?
Tierzüchter 32 (8): 338-340
- Snyder, S.H. (1978):
Neurotransmitter Receptor Binding.
In: H. I. Yamamura, S. J. Enna, M. J. Kuhar.
Raven Press, New York; pp. 1-12
- Schmitt, U., Hiemke, Ch. (1998):
Strain differences in open-field and elevated plus-maze behaviour of rats without and with pretest handling.
Pharmacol. Biochem. Behav. 59: 807-811
- Sheldon, M.H. (1968):
Exploratory behaviour: the inadequacy of activity measures.
Psycho. Sci. 11: 38
- Shepherd, J.K., Grewal, S.S., Fletcher, A., Bill, D.J., Dourish, C.T. (1994):
Behavioural and pharmacological characterisation of the elevated "zero-maze" as an animal model of anxiety.
Psychopharmacol. 116: 56-64
- Seiferle, E. (1960):
Schmerz und Angst bei Tier und Mensch.
Deutsche Tierärztliche Wochenschrift 67: 332-334
- Soghomonian, J.J., Descarries, L., Watkins, K.C. (1989):
Serotonin innervation in adult rat neostriatum. II. Ultrastructural features: a radioautographic and immunocytochemical study.
Brain Research 481 (1): 67-86

- Stefanick, M.L. (1983):
The circadian patterns of spontaneous seminal emission, sexual activity and penile reflexes in the rat.
Physiol. Behav. 31: 737-743
- Steinbusch, H.W.M., Verhofstad, A.A.J., Joosten, H.W.J. (1978):
Localization of serotonin in the central nervous system by immunohistochemistry: description of a specific and sensitive technique and some applications.
Neuroscience 3: 811-819
- Steinbusch, H.W.M. (1981):
Distribution of serotonin-immunoreactivity in the central nervous system of the rat-cell bodies and terminals.
Neuroscience 6: 557-618
- Stöhr, T., Schulte-Wermeling, D., Weiner, I., Feldon, J. (1998):
Rat strain differences in open-field behavior and the locomotor stimulating and rewarding effects of amphetamine.
Pharmacol. Biochem. Behav. 59: 813-818
- Stricklin, W.R., Mench, J.A. (1987):
Social organization.
Veter. Clin. N. America: Food Anim. Pract. 3: 307-322
- Sudakov, S.K., Medvedeva, O.F., Rusakova, I.V., Terebilina, N.N., Goldberg, S.R. (2001):
Differences in genetic predisposition to high anxiety in two inbred rat strains: role of substance P, diazepam binding inhibitor fragment and neuropeptide Y.
Psychopharmacol. 154: 327-35
- Takahashi, L.K., Grossfeld, S., Lore, R.K. (1980):
Attack and escape in the laboratory rat: A modification of the colony-intruder procedure.
Behav. Neural. Biol. 29: 512 - 517
- Thomas, H., Fink, H., Sohr, T.R., Voits, M. (2000):
Lesion of the median raphe nucleus: a combined behavioral and microdialysis study in rats.
Pharmacol. Biochem. Behav. 65: 15-21
- Townsend, P. (1997):
Use of in-cage shelters by laboratory rats.
Animal Welfare 6 (2): 95-103
- Treit, D. (1985):
Animal models for the study of anti-anxiety agents: a review.
Neurosci. Biobehav. Reviews 9: 203-222

- Treit, D., Menard, J. (1998):
Animal models of anxiety and depression.
In: A.A. Boulton, G.B. Baker, A.N. Bateson (eds.). *Neuromethods* 34. In Vivo
Neuromethods.
Humana Press Inc., Totowa, New Jersey, pp. 89-148
- Trullas, R., Skolnick, P.H. (1993):
Differences in fear motivated behaviors among inbred mouse strains.
Psychopharmacol. 111: 323-331
- Twarog, B.M., Page, I.H. (1953):
Serotonin content of some mammalian tissues and urine and a method for its
determination.
J. Physiol. 175: 157-161
- Uhl, G.R., Johnson, P.S. (1994):
Neurotransmitter transporters: Three important gene families for neuronal function.
J. Exp. Biol. 196: 229-236
- Van der Staay, F.J., Blokland, A. (1996):
Behavioral differences between outbred Wistar, inbred Fischer 344, Brown Norway, and
hybrid Fischer 344 X Brown Norway rats.
Physiol. and Behav. 60: 97-109
- Van Hartesveldt, C., Moore, B., Hartman, B.K. (1986):
Transient midline raphe glial structure in the developing rat.
J. Comp. Neurol. 253 (2): 174-84
- Vanhoutte, P.M., Saxena, P.R., Paoletti, R., Brunello, N., Jackson, N., Jackson, A.S. (1993):
Serotonin: From cell biology to pharmacology and therapeutics.
Kluwer Academic, Dordrecht, The Netherlands.
- Varley, M.A. (1995):
Behavioral patterns of the weaned piglet.
The Pig Journal 34: 71-97
- Vogel, J.R., Beer, B., Clody, D.E. (1971):
A simple and reliable conflict procedure for testing anti-anxiety agents.
Psychopharmacologia 21: 1-7
- Walsh, R.N., Cummins, R.A. (1976):
The open-field test: A critical review.
Psychol. Bull. 83: 482-504
- Weiss, S.M., Lightowler, S., Stanhope, K.J., Kennett, G.A., Dourish, C. (2000):
Measurement of anxiety in transgenic mice.
Rev. Neurosci. 11: 59-74

- Whitaker-Azmitia, P.M., Lauder, J.M., Shemmer, A., Azmitia, E.C. (1987):
Postnatal changes in serotonin receptors following prenatal alterations in serotonin levels: further evidence for functional fetal serotonin receptors.
Brain Res. 430: 285-289
- Wise, C.D., Berger, B.D., Stein, L. (1972):
Benzodiazepines: anxiety-reducing activity by reduction of serotonin turnover in the rat brain.
Science 177: 180-183
- World Health Organisation (1992):
The ICD-10 Classification of Mental and Behavioural Disorders
World Health Organisation (Hrsg.), Genf.