

## INVITED COMMENTARY

# How can we model subsurface stormflow at the catchment scale if we cannot measure it?

Peter Chiffard<sup>1</sup>  | Theresa Blume<sup>2</sup> | Katja Maerker<sup>3</sup> | Luisa Hopp<sup>4</sup> | Ilja van Meerveld<sup>5</sup> | Thomas Graef<sup>6</sup> | Oliver Gronz<sup>7</sup> | Andreas Hartmann<sup>8</sup> | Bernhard Kohl<sup>9</sup> | Edoardo Martini<sup>10</sup> | Christian Reinhardt-Imjela<sup>11</sup> | Martin Reiss<sup>1</sup> | Michael Rinderer<sup>8</sup> | Stefan Achleitner<sup>12</sup>

<sup>1</sup>Department of Geography, Philipps-University Marburg, Marburg, Germany

<sup>2</sup>Department of Hydrology, German Research Centre for Geosciences Potsdam, Potsdam, Germany

<sup>3</sup>Department of Physical Geography, (formerly Katja Heller) University of Technology Dresden, Dresden, Germany

<sup>4</sup>Department of Hydrology, University of Bayreuth, Bayreuth, Germany

<sup>5</sup>Department of Geography – Hydrology and Climate, University of Zürich, Zurich, Switzerland

<sup>6</sup>German Environment Agency, Dessau-Roßlau, Germany

<sup>7</sup>Department of Physical Geography, University of Trier, Trier, Germany

<sup>8</sup>Department of Hydrology, Albert-Ludwigs-University of Freiburg, Freiburg im Breisgau, Germany

<sup>9</sup>Department of Natural Hazards, Austrian Research Centre for Forests Innsbruck, Innsbruck, Austria

<sup>10</sup>Institute of Environmental Physics, Ruprecht-Karls-University of Heidelberg, Heidelberg, Germany

<sup>11</sup>Department of Applied Physical Geography, Freie Universität Berlin, Berlin, Germany

<sup>12</sup>Unit for Hydraulic Engineering, University of Innsbruck, Innsbruck, Austria

## Correspondence

Peter Chiffard, Department of Geography, Philipps-University Marburg, Marburg, Germany.

Email: peter.chiffard@geo.uni-marburg.de

## Funding information

Deutsche Forschungsgemeinschaft (German Research Foundation; DFG), Grant/Award Number: CH 870/5-1


## 1 | INTRODUCTION

Subsurface stormflow (SSF) can be a dominant run-off generation process in humid mountainous catchments (e.g., Bachmair & Weiler, 2011; Blume & van Meerveld, 2015; Chiffard, Didszun, & Zepp, 2008). Generally, SSF develops in structured soils where bedrock or a less permeable soil layer is overlaid by a more permeable soil layer and vertically percolating water is deflected, at least partially, in a lateral downslope direction due to the slope inclination. SSF can also occur when groundwater levels rise into more permeable soil layers and water flows laterally through the more permeable layers to the stream (“transmissivity feedback mechanism”; Bishop, Grip, & O'Neill, 1990). The different existing terms for SSF in the hydrological literature such as shallow subsurface run-off, interflow, lateral flow, or soil water flow reflects

the different underlying process concepts developed in various experimental studies in different environments by using different experimental approaches at different spatial and temporal scales (Weiler, McDonnell, Tromp-van Meerveld, & Uchida, 2005). Intersite comparisons and the extraction of general rules for SSF generation and its controlling factors are still lacking, which hampers the development of appropriate approaches for modelling SSF. But appropriate prediction of SSF is essential due to its clear influence on run-off generation at the catchment scale (e.g., Chiffard et al., 2010; Zillgens, Merz, Kirnbauer, & Tilch, 2005), on the formation of floods (e.g., Markart et al., 2013, 2015) and on the transport of nutrients or pollutants from the hillslopes into surface water bodies (Zhao, Tang, Zhao, Wang, & Tang, 2013). However, a precise simulation of SSF in models requires an accurate process understanding including, knowledge about water

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2019 The Authors Hydrological Processes Published by John Wiley & Sons Ltd



**FIGURE 1** Caricature of the investigation and simulation of subsurface stormflow (SSF) (Illustration: Edoardo Martini)

pathways, residence times, magnitude of water fluxes, or the spatial origin of SSF within a given catchment because such factors determine the transport of subsurface water and solutes to the stream. But due to its occurrence in the subsurface and its spatial and temporal variability, determining and quantifying the processes generating SSF is a challenging task as they cannot be observed directly. Therefore, it is logical to ask whether we can really model SSF correctly if we cannot measure it well enough on the scale of interest (Figure 1).

This commentary reflects critically on whether current experimental concepts and modelling approaches are sufficient to predict the contribution of SSF to the run-off at the catchment scale. This applies in particular to the underlying processes, controlling factors, modelling approaches, research gaps, and innovative strategies to trace SSF across different scales.

## 2 | WHAT HAVE WE LEARNED FROM ALL THE EXPERIMENTAL STUDIES ABOUT SSF CARRIED OUT IN VARIOUS CATCHMENTS AT DIFFERENT SPATIAL AND TEMPORAL SCALES?

Experimental studies on SSF generation have been carried out in different mountainous catchments with steep, well drained soils (e.g., Maimai, New Zealand [e.g., McGlynn & McDonnell, 2003]), shallow soils with bedrock outcrops (Panola, United States [e.g., van Meerveld, Seibert, & Peters, 2015]), and Fudoji, Japan [e.g., Uchida, Asano, Mizuyama, & McDonnell, 2004]), or catchments with periglacial drift deposits (Bohlmicke, Germany [e.g., Chiffard et al., 2008]), and Ore Mountains, Germany [e.g., Heller & Kleber, 2016]). These studies have resulted in comprehensive process knowledge about SSF, which was synthesized in several reviews (e.g., Ghasemizade & Schirmer, 2013)

and have led to the identification of controlling factors of SSF like initial soil moisture content (e.g., Blume, Zehe, & Bronstert, 2009; Chiffard & Zepp, 2008; Martini et al., 2015), water table development at the soil-bedrock interface (e.g., Anderson, Weiler, Alila, & Hudson, 2010; Jost, Schume, Hager, Markart, & Kohl, 2012), preferential flow paths (e.g., Laine-Kaulio, Backnäs, Karvonen, Koivusalo, & McDonnell, 2014; Sidle, Noguchi, Tsuboyama, & Laursen, 2001; Uchida, Kosugi, & Mizuyama, 2001), hillslope characteristics (e.g., Bachmair & Weiler, 2012), drainable porosity (e.g., Weiler & McDonnell, 2006), precipitation thresholds (e.g., Hopp, McDonnell, & Condon, 2011; Kienzler & Naef, 2008a, 2008b; Peralta-Tapia, Sponseller, Tetzlaff, Soulsby, & Laudon, 2014), soil properties (e.g., Bachmair, Weiler, & Nützmann, 2009; Hopp & McDonnell, 2009), soil depth (e.g., Tromp-Van Meerveld & McDonnell, 2006a, 2006b), or bedrock topography (e.g., Freer et al., 2002). Nevertheless, little of this understanding has been incorporated into current hydrological models. As catchment hydrologists, we are particularly interested in run-off generation at the catchment scale that seems more controlled by the interplay of processes than the details of individual ones. In other words, landscape heterogeneity and process complexity at the small scale can lead to typical emergent response behaviour at the catchment scale (McDonnell et al., 2007). From this, it follows that landscape structure can inform us about the dominant run-off generation mechanisms that are most often hidden in the subsurface and therefore so difficult to observe across the entire catchment. For instance, generation of SSF on low mountain ranges in middle Europe is strongly influenced by the widespread periglacial cover beds, which are a typical example for stratified soils (Hübner, Günther, Heller, Noell, & Kleber, 2016; Hübner, Heller, Günther, & Kleber, 2015; Kleber & Terhorst, 2013; Moldenhauer, Heller, Chiffard, Hübner, & Kleber, 2013). Although in soil science the Substrate-Oriented-Soil-Evolution-Model (Lorz, Heller, & Kleber, 2011) underlines the importance of stratified soils and lithological

discontinuities as a key element controlling ecological processes, in hydrologic research, less attention has been paid to the stratification of soils as a major trigger of lateral water paths (e.g., Reinhardt-Imjela, Maerker, Schulte, & Kleber, 2018; Reiss & Chiffard, 2015; Zhang, Lin, & Doolittle, 2014). The existence of a non-linear and threshold-type response of SSF to precipitation (e.g., Ali et al., 2015; Graham, Woods, & McDonnell, 2010) adds to the challenge of both measuring and modelling this process. Indeed, the detection of these thresholds helps to classify behaviours of different hillslopes; however, the controlling factors and processes responsible for these thresholds are not yet fully clear. Therefore, it is also still unclear if these thresholds are transferable to other sites (Zhao et al., 2013).

One possible way forward would be a comprehensive site inter-comparison and an innovative strategy to compare and combine all the obtained first-order controls to assess the SSF generation. Previous attempts have provided very informative results (Bachmair, Weiler, & Troch, 2012; Uchida, Tromp-van Meerveld, & McDonnell, 2005). But on the other hand, we should ask ourselves whether it is expedient to compare all these various experimental studies investigating SSF generation carried out in different hydroclimatic regimes, at different scales or by using different experimental approaches. A better way forward might be the development of a systematic method-orientated measurement program, which combines a mixture of appropriate methods specifically targeted to the identification and characterization of SSF generation and which will be applied across well-instrumented catchments covering different spatial scales within similar environments. The experimental investigation can then focus on understanding process heterogeneity and complexity in connection with controlling factors and landscape structure. Thus, it is indispensable to develop a method-orientated research approach, which is specifically targeted to SSF generation, covers standard as well as innovative methods, and can be applied across different catchments within similar environments. This kind of standardized and systematic protocol to capture and characterize SSF will help to improve the representation of subsurface processes in spatially distributed hydrological models.

### 3 | ARE THE EXISTING HYDROLOGICAL WATERSHED MODEL CONCEPTS REFLECTING SSF ADEQUATELY?

The simulation of catchment-scale run-off generation and the associated water balance in the unsaturated zone, including SSF, strongly varies with respect to the model concept and spatial scale of prediction. Methods range from detailed physically based approaches such as the Richards' equation (Beven & German, 2013) or the kinematic wave method (e.g. Flügel & Smith, 1999) to less complex conceptual models such as the Soil Conservation Service Curve Number methods. The conceptual models do not require detailed process knowledge, but we make the claim that for the development of process-based hydrological models, a good understanding of the generation of SSF and the incorporation of process knowledge is essential (e.g.,

Bachmair, Weiler, & Nützmann, 2010; McGuire, Weiler, & McDonnell, 2007; Zhu & Lin, 2009). However, parametrizing the experimentally identified and quantified SSF knowledge at the catchment scale is a problem, as at this scale, the spatial heterogeneity of soil properties and the spatial organization of the specific pathways in the subsurface are largely undeterminable (Lin & Zhou, 2008). And even if information on all model parameters was available at the catchment scale, the question posed by Tromp-van Meerveld and Weiler (2008, p. 25) "How much model complexity is needed to explain the observed subsurface flow response [...]?" still remains unsolved. Indeed, it could be that the process complexity (e.g., matrix and preferential flow) and the natural variability of environmental properties (e.g., soil properties) collapse to a relatively simple functional relationship between a functional trait (e.g., soil moisture patterns) and catchment-scale run-off response (McDonnell et al., 2007).

Hydrogeophysical methods may have the potential to identify subsurface flow paths (e.g., Angermann et al., 2017; Binley et al., 2015) or, at least, soil heterogeneity (Martini et al., 2017), but their use is mostly limited to the hillslope scale (Vereecken et al., 2015). Here, subsurface flow paths are more likely to be connected over shorter rather than longer distances, which leads to higher effective flow velocities (e.g., Anderson, Weiler, Alila, & Hudson, 2009; Wienhöfer & Zehe, 2014). Preferential flow processes should be taken into account when calculating SSF in rainfall-run-off models at the catchment scale, but the representation of preferential flow is a particular challenge for all model concepts (e.g., Gerke, Germann, & Nieber, 2010; Hartmann, 2016). It would require model parameters (such as macropore density) at high temporal and spatial resolution, something that we can only determine at the scale of a soil column or a plot (e.g. Rinderer & Seibert, 2012). In addition, the spatial discretization (e.g., control volume or pixel) required if these approaches are used at the catchment scale is often in the order of tens to hundreds of metres and thus one or two orders of magnitude larger than the scale at which these physical relations (e.g., Richards' equation) originally have been developed. So it is doubtful that these models can still be referred to as "physically based" (Köhne, Köhne, & Šimůnek, 2009). In order to incorporate hydrological heterogeneity at scales larger than the plot or hillslope, hydrological models have used certain simplifications and assumptions. For instance, TOPMODEL (Beven et al., 1979) is based on the assumption that under steady-state flow conditions, the slope of the groundwater table is parallel to the slope of the surface topography. Only then is the topographic index derived from a digital elevation model such as the Topographic Wetness Index (Beven & Kirkby, 1979) a good proxy to estimate the groundwater table across a catchment. Other models, such as PDM (Probability Distributed Model, Moore, 2007) or VarKarst-R (Hartmann et al., 2015; Hartmann, Gleeson, Wada, & Wagener, 2017), implement subsurface heterogeneity by using Pareto functions.

Although these approaches are useful modelling concepts for capturing flow in the saturated zone, they do not explicitly incorporate SSF processes (Rinderer, van Meerveld, & Seibert, 2014; Seibert, Bishop, Rodhe, & McDonnell, 2003). Therefore, new modelling concepts are necessary, which explicitly incorporate the process

knowledge and first-order controls of SSF that were obtained in many experimental studies at various spatial scales. In addition, new ways have to be found to parameterize rainfall-run-off models adequately to calculate SSF at scales larger than a soil column or an experimental plot. Instead of continuing to investigate the process complexity of SSF with more and more experimental studies at different sites, we call for a concerted, method-orientated experimental approach carried out in accordance with the new approaches to include SSF explicitly in catchment-scale rainfall-run-off models. We thus generate process knowledge based on a systematic measurement program, which in turn allows us to parameterize and calibrate SSF modules in rainfall-run-off models at the catchment scale.

#### 4 | CAN WE REALLY VERIFY SIMULATED SSF?

Rainfall-run-off models are used for studies that are either investigative or predictive (Blöschl & Sivapalan, 1995). In both cases, the simulated SSF is mainly calibrated and validated based on single rainfall-run-off events (e.g., artificial sprinkling experiments) for which tracer hydrological data and information on specific run-off components are available (e.g., Markart et al., 2015; Uhlenbrook, Roser, & Tilch, 2004). However, it is obvious that these single events with steady-state conditions are not sufficient to capture the whole range of SSF response that depends on factors such as initial conditions and rainfall intensities and is often threshold dependent. Furthermore, the quality of a run-off model is still assessed by comparing modelled and observed total run-off measured at a gauge often situated at the catchment outlet. This is not expedient for studying the generation of SSF. However, currently, SSF in both types of rainfall-run-off models (investigative and predictive) is still an unvalidated parameter, which is adjusted (calibrated) to fit the model output against available discharge observations. The assumption is that if the model discharge fits the discharge observations satisfyingly, SSF is also simulated correctly. This is not necessarily true, particularly when considering changes in SSF contributions over the course of an event. High frequency measurements of chemical tracers and stable water isotopes (e.g.,  $^{18}\text{O}$  and  $^2\text{H}$ ) in streams and soils have the potential to gain better insights into SSF (e.g., Mueller et al., 2014; Sprenger et al., 2018). However, this comes with additional challenges: for soil water isotope data, the choice of the sampling method (e.g., wick sampler and suction cups) predetermines whether the more tightly bound or the more mobile soil water is extracted (Landon, Delin, Komor, & Regan, 1999). Even the choice of laboratory has been shown to influence the results of isotope analysis (Orlowski et al., 2018). In addition, hydrochemical signatures that can give insights into the biogeochemical-hydrological process links at different spatial scales (e.g., McKnight, Burns, Barnard, & Gabor, 2015; Ponton, West, Feakins, & Galy, 2014) are promising approaches to identify subsurface flow networks. The use of such "tracers" (e.g., N, DOC,  $^{13}\text{C}$ ,  $^{15}\text{N}$ , and microbial communities; Blume & van Meerveld, 2015; Sanderman, Lohse, Baldock, & Amundson, 2009; Sebestyen et al., 2008), their chemical characteristics (e.g., biodegradable organic carbon and excitation

emission matrix; Barnard, Burns, McKnight, Gabor, & Brooks, 2014; Burns, 2014; Hood, Williams, & McKnight, 2005), and their depth distribution in soils (Gabor, Eilers, McKnight, Fierer, & Anderson, 2014; Hassouna, Massian, Dudal, Pech, & Theraulaz, 2010; Wynn, Harden, & Fries, 2006) in combination with traditional tracers (e.g.,  $^{18}\text{O}$ ,  $^2\text{H}$ , and  $\text{SiO}_2$ ) may offer new opportunities for testing hydrological models. Nevertheless, limited consideration has been given to assimilating these approaches into rainfall-run-off models (Ebert, McKnight, Lajtha, Hartnett, & Jaffe, 2013).

#### 5 | CONCLUSIONS

Existing empirical studies have revealed different facets of SSF across catchments in different environments. Nevertheless, deficits still exist in the capability to use this knowledge to generalize our process understanding on subsurface flow dynamics at the catchment scale. This might be due to the fact that a generally accepted organizational framework for site intercomparison is not yet available. We have to consider whether such a framework is absolutely necessary or approaches based on a few representative monitoring sites and an upscaling approach based on landscape structure seem a promising way forward.

Nevertheless, many catchment hydrological models do not even consider SSF, whereas the more detailed, physically based models that do so are difficult to parameterize or validate without spatial information on catchment states (e.g., soil properties) and dynamics (e.g., antecedent soil moisture). The run-off response at the catchment scale is also most likely not dominated by the same detailed processes that we observe at the point or plot scale but instead by an emergent behaviour that results from an interplay between small-scale and large-scale processes. So how can we simulate subsurface flow at the catchment scale if we cannot even measure it?

We advocate for a more systematic design of future empirical studies on SSF across a range of representative landscapes which are concerned with the need not only to improve process understanding but also to develop new modelling approaches. Instead of model validation on the basis of single rainfall-run-off events for which tracer hydrological data and hydrometric measurements of selected run-off components are available at a single gauged trench, it is instead necessary to continuously monitor subsurface run-off components at the catchment outlet as well as on its hillslopes with high temporal resolution over longer time spans. We see great potential in exploiting the potential of distributed sensor networks and new tracers or combination of tracers that can give direct or indirect information on SSF. To address this concern, we need to collaborate with colleagues from neighbouring disciplines that have developed biogeochemical methods that can also be highly informative for SSF (e.g., microbial communities and nanoparticles as tracers). These new types of datasets will bring us one step further towards knowing what, where, and when to measure and how complex our models need to be in order to make our SSF simulations more realistic than they are today.

## ACKNOWLEDGMENTS

The authors gratefully acknowledge the financial support provided by the Deutsche Forschungsgemeinschaft (German Research Foundation; DFG) supporting the scientific network: "Subsurface stormflow—A well-recognized but still challenging process in catchment hydrology research" (CH 870/5-1). Moreover, the authors would like to extend their special thanks to the scientific guests Doerthe Tetzlaff, Nicola Fohrer, Yuko Asano, Axel Bronstert, Gunnar Lischeid, Jan Seibert, Taro Uchida, Markus Weiler, and Harald Zepp for their participation in our discussion on open questions related to SSF.

## ORCID

Peter Chiffard  <https://orcid.org/0000-0001-5325-3819>

## REFERENCES

- Ali, G., Tetzlaff, D., McDonnell, J. J., Soulsby, C., Carey, S., Laudon, H., ... Shanley, J. (2015). Comparison of threshold hydrologic response across northern catchments. *Hydrological Processes*, 29(16), 3575–3591. <https://doi.org/10.1002/hyp.10527>
- Anderson, A. E., Weiler, M., Alila, Y., & Hudson, R. O. (2009). Subsurface flow velocities in a hillslope with lateral preferential flow. *Water Resources Research*, 45, W11407. <https://doi.org/10.1029/2008WR007121>
- Anderson, A. E., Weiler, M., Alila, Y., & Hudson, R. O. (2010). Piezometric response in zones of a watershed with lateral preferential flow as a first-order control on subsurface flow. *Hydrological Processes*, 24, 2237–2247. <https://doi.org/10.1002/hyp.7662>
- Angermann, L., Jackisch, C., Allroggen, N., Sprenger, M., Zehe, E., Troncke, J., ... Blume, T. (2017). Form and function in hillslope hydrology: Characterization of subsurface flow based on response observations. *Hydrology and Earth System Sciences*, 21, 3727–3748. <https://doi.org/10.5194/hess-21-3727-2017>
- Bachmair, S., & Weiler, M. (2011). New dimensions of hillslope hydrology. In D. F. Levia, D. Carlyle-Moses, & T. Tanaka (Eds.), *Forest hydrology and biogeochemistry: Synthesis of past research and future directions—Ecological studies*, 216 (pp. 455–481). Dordrecht: Springer.
- Bachmair, S., & Weiler, M. (2012). Hillslope characteristics as controls of subsurface flow variability. *Hydrology and Earth System Sciences*, 16, 3699–3715. <https://doi.org/10.5194/hess-16-3699-2012>
- Bachmair, S., Weiler, M., & Nützman, G. (2009). Controls of land use and soil structure on water movement: Lessons for pollutant transfer through the unsaturated zone. *Journal of Hydrology*, 369, 241–252. <https://doi.org/10.1016/j.jhydrol.2009.02.031>
- Bachmair, S., Weiler, M., & Nützman, G. (2010). Benchmarking of two dual-permeability models under different land use and land cover. *Vadose Zone Journal*, 9(2), 226–237. <https://doi.org/10.2136/vzj2009.0089>
- Bachmair, S., Weiler, M., & Troch, P. A. (2012). Intercomparing hillslope hydrological dynamics: Spatio-temporal variability and vegetation cover effects. *Water Resources Research*, 48. <https://doi.org/10.1029/2011WR011196>
- Barnard, H. R., Burns, M. A., McKnight, D. M., Gabor, R. S. & Brooks, P. D. (2014). Reconciling stream dissolved organic matter with snowmelt-driven subsurface flowpaths in a montane, headwater catchment. – American Geophysical Union, Fall Meeting 2014, abstract #H53K-03, <http://adsabs.harvard.edu/abs/2014AGUFM.H53K.03B>; 09.06.2015; 14:00 MEZ).
- Beven, K., & Germann, P. (2013). Macropores And Water Flow In Soils Revisited. *Water Resources Research* 49, 3071–3092.
- Beven, K. J., Kirby, M. J., Kirkby, M. J., Kirby, M. J., Kirkby, M. J., & Hydrologiques, H. S. S. (1979). A physically based, variable contributing area model of basin hydrology. *Hydrological Sciences Bulletin*, 24(1), 43–69. <https://doi.org/10.1080/02626667909491834>
- Beven, K. J., & Kirkby, M. J. (1979). A physically based, variable contributing area model of basin hydrology. *Hydrological Sciences Bulletin*, 24, 43–69. <https://doi.org/10.1080/02626667909491834>
- Binley, A., Hubbard, S. S., Huisman, J. A., Revil, A., Robinson, D. A., Singha, K., & Slater, L. D. (2015). The emergence of hydrogeophysics for improved understanding of subsurface processes over multiple scales. *Water Resources Research*, 51, 3837–3866. <https://doi.org/10.1002/2015WR017016>
- Bishop, K. H., Grip, H., & O'Neill, A. (1990). The origins of acid runoff in a hillslope during storm events. *Journal of Hydrology*, 116, 35–61. [https://doi.org/10.1016/0022-1694\(90\)90114-D](https://doi.org/10.1016/0022-1694(90)90114-D)
- Blöschl, G., & Sivapalan, M. (1995). Scale issues in hydrological modelling: A review. *Hydrological Processes*, 9, 251–290. <https://doi.org/10.1002/hyp.3360090305>
- Blume, T., & van Meerveld, H. J. I. (2015). From hillslope to stream: Methods to investigate subsurface connectivity. *Wiley Interdisciplinary Reviews Water*, 2(3), 177–198. <https://doi.org/10.1002/wat2.1071>
- Blume, T., Zehe, E., & Bronstert, A. (2009). Use of soil moisture dynamics and patterns at different spatio-temporal scales for the investigation of subsurface flow processes. *Hydrology and Earth System Sciences*, 13, 1215–1234. <https://doi.org/10.5194/hess-13-1215-2009>
- Burns, M. A. (2014). Hillslope dissolved organic matter transport and transformation in a semi-arid headwater catchment. – Geography theses & dissertations. Paper 3. ([http://scholar.colorado.edu/cgi/viewcontent.cgi?article=1002&context=geog\\_gradetds](http://scholar.colorado.edu/cgi/viewcontent.cgi?article=1002&context=geog_gradetds); 09.06.2015, 14:00 MEZ).
- Chiffard, P., Didszun, J., & Zepp, H. (2008). Analysis of runoff generation at different scales in areas characterized by periglacial cover beds (Sauerland, Germany) (in German). *Grundwasser*, 13, 27–41. <https://doi.org/10.1007/s00767-007-0058-1>
- Chiffard, P., Kirnbauer, R., Zepp, H., Tilch, N., Didszun, J., Zillgens, B., ... Uhlenbrook, S. (2010). Tracing runoff generation processes through different spatial scales in low and high mountain ranges. *IAHS Publication*, 336, 90–95.
- Chiffard, P., & Zepp, H. (2008). Erfassung der zeitlichen Variabilität der Abflussbereitschaft eines Einzugsgebietes auf Grundlage von Bodenfeuchtemessungen. *Hydrologie Und Wasserwirtschaft*, 52(3), 98–109.
- Ebert, J., McKnight, D., Lajtha, K., Hartnett, H., & Jaffe, R. (2013). Carbon science investigations. *Network News*, 26(3). <http://news.itsernet.edu/print/2880> (05.09.2015, 12:00 MEZ)
- Flügel, W.-a., & Smith, R. E. (1999). Integrated process studies and modeling simulations of hillslope hydrology and interflow dynamics using the HILLS model. *Environmental Modeling Software*, 14, 153–160.
- Freer, J., McDonnell, J. J., Beven, K. J., Peters, N. E., Burns, D. A., Hooper, R. P., & Aulenbach, B. (2002). The role of bedrock topography on subsurface storm flow. *Water Resources Research*, 38(12), 51–516. <https://doi.org/10.1029/2001WR000872>
- Gabor, R. S., Eilers, K., McKnight, D. M., Fierer, N., & Anderson, S. P. (2014). From the litter layer to the saprolite: Chemical changes in water-soluble soil organic matter and their correlation to microbial community composition. *Soil Biology and Biochemistry*, 68, 166–176. <https://doi.org/10.1016/j.soilbio.2013.09.029>

- Gerke, H., Germann, P., & Nieber, J. (2010). Preferential and unstable flow: From the pore to the catchment scale. *Vadose Zone Journal*, 9, 207–212. <https://doi.org/10.2136/vzj2010.0059>
- Ghasemzade, M., & Schirmer, M. (2013). Subsurface flow contribution in the hydrological cycle: Lessons learned and challenges ahead—A review. *Environment and Earth Science*, 69, 707–718. <https://doi.org/10.1007/s12665-013-2329-8>
- Graham, C. B., Woods, R., & McDonnell, J. J. (2010). Hillslope threshold response to rainfall: (1) A field based forensic approach. *Journal of Hydrology*, 393, 65–76. <https://doi.org/10.1016/j.jhydrol.2009.12.015>
- Hartmann, A. (2016). Putting the cat in the box: Why our models should consider subsurface heterogeneity at all scales. *WIREs Water*, 3, 478–486. <https://doi.org/10.1002/wat2.1146>
- Hartmann, A., Gleeson, T., Rosolem, R., Pianosi, F., Wada, Y., & Wagener, T. (2015). A large-scale simulation model to assess karstic groundwater recharge over Europe and the Mediterranean. *Geoscientific Model Development*, 8(6), 1729–1746. <https://doi.org/10.5194/gmd-8-1729-2015>
- Hartmann, A., Gleeson, T., Wada, Y., & Wagener, T. (2017). Enhanced groundwater recharge rates and altered recharge sensitivity to climate variability through subsurface heterogeneity. *Proceedings of the National Academy of Sciences*, 114(11), 2842–2847. <https://doi.org/10.1073/pnas.1614941114>
- Hassouna, M., Massian, C., Dudal, Y., Pech, N., & Theraulaz, F. (2010). Changes in water extractable organic matter (WEOM) in a calcareous soil under field conditions with time and soil depth. *Geoderma*, 155, 75–85. <https://doi.org/10.1016/j.geoderma.2009.11.026>
- Heller, K., & Kleber, A. (2016). Hillslope runoff generation influenced by layered subsurface in a headwater catchment in Ore Mountains, Germany. *Environment and Earth Science*, 75(11), 1–15. <https://doi.org/10.1007/s12665-016-5750-y>
- Hood, E. W., Williams, M. W., & McKnight, D. M. (2005). Sources of dissolved organic matter (DOM) in a Rocky Mountain stream using chemical fractionation and stable isotopes. *Biogeochemistry*, 74(2), 231–255. <https://doi.org/10.1007/s10533-004-4322-5>
- Hopp, L., & McDonnell, J. J. (2009). Connectivity at the hillslope scale: Identifying interactions between storm size, bedrock permeability, slope angle and soil depth. *Journal of Hydrology*, 376, 378–391. <https://doi.org/10.1016/j.jhydrol.2009.07.047>
- Hopp, L., McDonnell, J. J., & Condon, P. (2011). Lateral subsurface flow in a soil cover over waste rock in a humid temperate environment. *Vadose Zone Journal*, 10, 1–13.
- Hübner, R., Günther, T., Heller, K., Noell, U., & Kleber, A. (2016). Monitoring infiltration and subsurface stormflow in layered slope deposits with ERT and hydrometric measurements—The capillary barrier effect as crucial factor. *Hydrology and Earth System Sciences*, 21, 5181–5199.
- Hübner, R., Heller, K., Günther, T., & Kleber, A. (2015). Monitoring hillslope moisture dynamics with surface ERT for enhancing spatial significance of hydrometric point measurements. *Hydrology and Earth System Sciences*, 19, 225–240. <https://doi.org/10.5194/hess-19-225-2015>
- Jost, G., Schume, H., Hager, H., Markart, G., & Kohl, B. (2012). A hillslope scale comparison of tree species influence on soil moisture dynamics and runoff processes during intense rainfall. *Journal of Hydrology*, 420, 112–124.
- Kienzler, P. M., & Naef, F. (2008a). Temporal variability of subsurface stormflow formation. *Hydrology and Earth System Sciences*, 12, 257–265. <https://doi.org/10.5194/hess-12-257-2008>
- Kienzler, P. M., & Naef, F. (2008b). Subsurface storm flow formation at different hillslopes and implications for the “old water paradox”. *Hydrological Processes*, 22(1), 104–116. <https://doi.org/10.1002/hyp.6687>
- Kleber, A., & Terhorst, B. (2013). Mid-latitude slope deposits (cover beds). *Developments in Sedimentology*, 66, 302.
- Köhne, J. M., Köhne, S., & Šimůnek, J. (2009). A review of model applications for structured soils: A water flow and tracer transport. *Journal of Contaminant Hydrology*, 104(1–4), 4–35. <https://doi.org/10.1016/j.jconhyd.2008.10.002>
- Laine-Kaulio, H., Backnäs, S., Karvonen, T., Koivusalo, H., & McDonnell, J. J. (2014). Lateral subsurface stormflow and solute transport in a forested hillslope: A combined measurement and modeling approach. *Water Resources Research*, 50, 8159–8178. <https://doi.org/10.1002/2014WR015381>
- Landon, M. K., Delin, G. N., Komor, S. C., & Regan, C. P. (1999). Comparison of the stable-isotopic composition of soil water collected from suction lysimeters, wick samplers, and cores in a sandy unsaturated zone. *Journal of Hydrology*, 224, 45–54. [https://doi.org/10.1016/S0022-1694\(99\)00120-1](https://doi.org/10.1016/S0022-1694(99)00120-1)
- Lin, H., & Zhou, X. (2008). Evidence of subsurface preferential flow using soil hydrologic monitoring in the Shale Hills catchment. *European Journal of Soil Science*, 59, 34–49.
- Lorz, C., Heller, K., & Kleber, A. (2011). Stratification of the regolith continuum—A key property for processes and functions of landscapes. *Zeitschrift für Geomorphologie*, 55, 277–292. <https://doi.org/10.1127/0372-8854/2011/0055S3-0062>
- Markart G., Römer, A., Bieber, G., Pirkl, H., Klebinder, K., Hörfarer, C., Ahl, A., Ita, A., Jochum, B., Kohl, B., Meissl, G., Motschka, K., Ottowitz, D., Schattauer, I., Sotier, B., Strasser, M., Suntinger, K., & Winkler, E. (2013). Abschätzung der Bandbreiten von Fließgeschwindigkeiten des oberflächennahen Zwischenabflusses in alpinen Einzugsgebieten. Endbericht - 3. Projektjahr an die Österreichische Akademie der Wissenschaften (ÖAW), Nationalkomitee Hydrologie Österreichs (IHP), ISBN-online: 978-3-7001-7392-2; <https://doi.org/10.1553/Shallow>
- Markart, G., Römer, A., Bieber, G., Pirkl, H., Klebinder, K., Hörfarer, C., ... Motschka, K. (2015). Assessment of shallow interflow velocities in alpine catchments for the improvement of hydrological modelling. In G. Lollino, M. Arattano, M. Rinaldi, O. Giustolisi, J. C. Marechal, & G. E. Grant (Eds.), *Engineering geology for society and territory - Volume 3* (pp. 611–615). River Basins: Reservoir Sedimentation and Water Resources.
- Martini, E., Werban, U., Zacharias, S., Pohle, M., Dietrich, P., & Wollschläger, U. (2017). Repeated electromagnetic induction measurements for mapping soil moisture at the field scale: Validation with data from a wireless soil moisture monitoring network. *Hydrology and Earth System Sciences*, 21, 495–513. <https://doi.org/10.5194/hess-21-495-2017>
- Martini, E., Wollschläger, U., Kögler, S., Behrens, T., Dietrich, P., Reinstorf, F., ... Zacharias, S. (2015). Spatial and temporal dynamics of hillslope-scale soil moisture patterns: Characteristic states and transition mechanisms. *Vadose Zone Journal*, 14. <https://doi.org/10.2136/vzj2014.10.0150>
- McDonnell, J. J., Sivapalan, M., Vaché, K., Dunn, S., Grant, G., Haggerty, R., ... Weiler, M. (2007). Moving beyond heterogeneity and process complexity: A new vision for watershed hydrology. *Water Resources Research*, 43, W07301.
- McGlynn, B. L., & McDonnell, J. J. (2003). Role of discrete landscape units in controlling catchment dissolved organic carbon dynamics. *Water Resources Research*, 9(4), 1090. <https://doi.org/10.1029/2002WR001525>
- McGuire, K. J., Weiler, M., & McDonnell, J. J. (2007). Integrating tracer experiments with modeling to assess runoff processes and water

- transit times. *Advances in Water Resources*, 30(4), 824–837. <https://doi.org/10.1016/j.advwatres.2006.07.004>
- McKnight, D. M., Burns, M. A., Barnard, H., & Gabor, R. S. (2015). Transport of dissolved organic matter from the hillslope to the stream: Using fluorescence spectroscopy to identify changing sources during snowmelt. *Geophysical Research Abstracts*, 17, EGU2015–EGU5219.
- Moldenhauer, K.-M., Heller, K., Chiffard, P., Hübner, R., & Kleber, A. (2013). Influence of cover beds on slope hydrology. In A. Kleber, & B. Terhorst (Eds.), *Mid-latitude slope deposits (cover beds): Developments in sedimentology*, 66 (pp. 127–152). Amsterdam: Elsevier. <https://doi.org/10.1016/B978-0-444-53118-6.00004-0>
- Moore, R. J. (2007). The PDM rainfall–runoff model. *Hydrology and Earth System Sciences*, 11(1), 483–499. <https://doi.org/10.5194/hess-11-483-2007>
- Mueller, M. H., Alaoui, A., Kuells, C., Leistert, H., Meusburger, K., Stumpp, C., ... Alewell, C. (2014). Tracking water pathways in steep hillslopes by  $d^{18}O$  depth profiles of soil water. *Journal of Hydrology*, 519, 340–352. <https://doi.org/10.1016/j.jhydrol.2014.07.031>
- Orlowski, N., Breuer, L., Angeli, N., Boeckx, P., Brumbt, C., Cook, C. S., ... McDonnell, J. J. (2018). Inter-laboratory comparison of cryogenic water extraction systems for stable isotope analysis of soil water. *Hydrology and Earth System Sciences*, 22, 3619–3637. <https://doi.org/10.5194/hess-22-3619-2018>
- Peralta-Tapia, A., Sponseller, R. A., Tetzlaff, D., Soulsby, C., & Laudon, H. (2014). Connecting precipitation inputs and soil flow pathways to stream water in contrasting boreal catchments. *Hydrological Processes*, 29(16), 3546–3555.
- Ponton, C., West, A. J., Feakins, S. J., & Galy, V. (2014). Leaf wax biomarkers in transit record river catchment composition. *Geophysical Research Letters*, 41, 6420–6427. <https://doi.org/10.1002/2014GL061328>
- Reinhardt-Imjela, C., Maerker, K., Schulte, A., & Kleber, A. (2018). Implications of hydraulic anisotropy in periglacial cover beds for flood simulation in low mountain ranges (Ore Mountains, Germany). *Die Erde*, 149(2–3), 86–101.
- Reiss, M., & Chiffard, P. (2015). Depth function of manganese (Mn) concentration in soil solutions: Hydropedological translocation of trace elements in stratified soils. *European Journal of Soil Science*, 4(3), 169–177.
- Rinderer, M., & Seibert, J. (2012). Soil information in hydrologic models: Hard data, soft data, and the dialog between experimentalists and modelers. In H. Lin (Ed.), *Hydropedology: Synergistic integration of soil science and hydrology* (pp. 515–536). USA: Elsevier.
- Rinderer, M., van Meerveld, H. J., & Seibert, J. (2014). Topographic controls on shallow groundwater levels in a steep, prealpine catchment: When are the TWI assumptions valid? *Water Resources Research*, 50, 6067–6080. <https://doi.org/10.1002/2013WR015009>
- Sanderman, J., Lohse, K. A., Baldock, J. A., & Amundson, R. (2009). Linking soils and streams: Sources and chemistry of dissolved organic matter in a small coastal watershed. *Water Resources Research*, 45(3), W03418. <https://doi.org/10.1029/2008WR006977>
- Sebestyen, S. D., Boyer, E. W., Shanley, J. B., Kendall, C., Doctor, D. H., Aiken, G. R., & Ohte, N. (2008). Sources, transformations, and hydrological processes that control stream nitrate and dissolved organic matter concentrations during snowmelt in an upland forest. *Water Resources Research*, 44, W12410. <https://doi.org/10.1029/2008WR006983>
- Seibert, J., Bishop, K., Rodhe, A., & McDonnell, J. J. (2003). Groundwater dynamics along a hillslope: A test of the steady state hypothesis. *Water Resources Research*, 39(1), 1014–1023.
- Side, R. C., Noguchi, S., Tsuboyama, Y., & Laursen, K. (2001). A conceptual model of preferential flow systems in forested hillslopes: Evidence of self-organization. *Hydrological Processes*, 15, 1675–1692. <https://doi.org/10.1002/hyp.233>
- Sprenger, M., Tetzlaff, D., Buttle, J., Carey, S. K., McNamara, J. P., Laudon, H., ... Soulsby, C. (2018). Storage, mixing, and fluxes of water in the critical zone across northern environments inferred by stable isotopes of soil water. *Hydrological Processes*, 32, 1720–1737. <https://doi.org/10.1002/hyp.13135>
- Tromp-Van Meerveld, I., & McDonnell, J. J. (2006a). Threshold relations in subsurface stormflow: 1. A 147-storm analysis of the Panola hillslope. *Water Resources Research*, 42, W02410. <https://doi.org/10.1029/2004WR003778>
- Tromp-Van Meerveld, I., & McDonnell, J. J. (2006b). Threshold relations in subsurface stormflow: 2. The fill and spill hypothesis. *Water Resources Research*, 42, W02411. <https://doi.org/10.1029/2004WR003800>
- Tromp-Van Meerveld, I., & Weiler, M. (2008). Hillslope dynamics modeled with increasing complexity. *Journal of Hydrology*, 361(1–2), 24–40. <https://doi.org/10.1016/j.jhydrol.2008.07.019>
- Uchida, T., Asano, Y., Mizuyama, T., & McDonnell, J. J. (2004). Role of upslope soil pore pressure on lateral subsurface storm flow dynamics. *Water Resources Research*, 40, W12401. <https://doi.org/10.1029/2003WR002139>
- Uchida, T., Kosugi, K., & Mizuyama, T. (2001). Effects of pipeflow on hydrological process and its relation to landslide: A review of pipeflow studies in forested headwater catchments. *Hydrological Processes*, 15, 2151–2174. <https://doi.org/10.1002/hyp.281>
- Uchida, T., Tromp-van Meerveld, I., & McDonnell, J. J. (2005). The role of lateral pipe flow in hillslope runoff response: An intercomparison of non-linear hillslope response. *Journal of Hydrology*, 311, 117–133. <https://doi.org/10.1016/j.jhydrol.2005.01.012>
- Uhlenbrook, S., Roser, S., & Tilch, N. (2004). Hydrological process representation at the meso-scale: The potential of a distributed, conceptual catchment model. *Journal of Hydrology*, 291(3–4), 278–296. <https://doi.org/10.1016/j.jhydrol.2003.12.038>
- van Meerveld, H. J., Seibert, J., & Peters, N. E. (2015). Hillslope-riparian-stream connectivity and flow directions at the Panola Mountain Research Watershed. *Hydrological Processes*, 29(16), 3556–3574. <https://doi.org/10.1002/hyp.10508>
- Vereecken, H., Huisman, J. A., Hendricks Franssen, H. J., Brüggemann, N., Bogena, H. R., Kollet, S., ... Vanderborght, J. (2015). Soil hydrology: Recent methodological advances, challenges, and perspectives. *Water Resources Research*, 51(4), 2616–2633. <https://doi.org/10.1002/2014WR016852>
- Weiler, M., & McDonnell, J. J. (2006). Testing nutrient flushing hypotheses at the hillslope scale: A virtual experiment approach. *Journal of Hydrology*, 319, 339–356. <https://doi.org/10.1016/j.jhydrol.2005.06.040>
- Weiler, M., McDonnell, J. J., Tromp-van Meerveld, I., & Uchida, T. (2005). Subsurface stormflow. In M. G. Anderson, & J. J. McDonnell (Eds.), *Encyclopedia of hydrological sciences*, Volume 3, Part 10 (p. 14). USA: Wiley.
- Wienhöfer, J., & Zehe, E. (2014). Predicting subsurface stormflow response of a forested hillslope—The role of connected flow paths. *Hydrology and Earth System Sciences*, 18, 121–138. <https://doi.org/10.5194/hess-18-121-2014>
- Wynn, J. G., Harden, J. W., & Fries, T. L. (2006). Stable carbon isotope depth profiles and soil organic carbon dynamics in the lower Mississippi Basin. *Geoderma*, 131, 89–109. <https://doi.org/10.1016/j.geoderma.2005.03.005>
- Zhang, J., Lin, H., & Doolittle, J. (2014). Soil layering and preferential flow impacts on seasonal changes of GPR signals in two contrasting

- soils. *Geoderma*, 213, 560–569. <https://doi.org/10.1016/j.geoderma.2013.08.035>
- Zhao, P., Tang, X., Zhao, P., Wang, C., & Tang, J. (2013). Identifying the water source for subsurface flow with deuterium and oxygen-18 isotopes of soil water collected from tension lysimeters and cores. *Journal of Hydrology*, 503, 1–10. <https://doi.org/10.1016/j.jhydrol.2013.08.033>
- Zhu, Q., & Lin, H. S. (2009). Simulation and validation of concentrated subsurface lateral flow paths in an agricultural landscape. *Hydrology and Earth System Sciences*, 13, 1503–1518. <https://doi.org/10.5194/hess-13-1503-2009>
- Zillgens, B., Merz, B., Kirnbauer, R., & Tilch, N. (2005). Analysis of the runoff response of an alpine catchment at different scales. *Hydrology and Earth System Sciences*, 2, 1923–1960. <https://doi.org/10.5194/hessd-2-1923-2005>

**How to cite this article:** Chiffard P, Blume T, Maerker K, et al. How can we model subsurface stormflow at the catchment scale if we cannot measure it? *Hydrological Processes*. 2019;33: 1378–1385. <https://doi.org/10.1002/hyp.13407>