

Aus den NeuroCure Exzellenzcluster und Institut der
Neurophysiologie
der Medizinischen Fakultät Charité – Universitätsmedizin Berlin

DISSERTATION

**Distinct Functions of Syntaxin-1 in Neuronal Maintenance,
Synaptic Vesicle Docking, and Fusion in Mouse Neurons**

zur Erlangung des akademischen Grades
Doctor of Philosophy (PhD)

vorgelegt der Medizinischen Fakultät
Charité – Universitätsmedizin Berlin

von

Gülçin Vardar

aus Bakirköy, der Türkei

Datum der Promotion: 8. Dezember 2017

Table of Contents

Abstract.....	3
Zusammenfassung.....	5
Affidavit.....	7
Excerpt from the Journal Summary List (ISI Web of Knowledge).....	9
Publication: <i>Distinct Functions of Syntaxin-1 in Neuronal Maintenance, Synaptic Vesicle Docking, and Fusion in Mouse Neurons</i>	11
Curriculum Vitae.....	25
List of publications.....	29
Acknowledgements.....	31

Abstract

Neurons relay information via neurotransmitter release through the fusion of the neurotransmitter filled presynaptic vesicles with the plasma membrane mediated by the sophisticated interplays of presynaptic proteins. Syntaxin-1 (Stx1) is one of the core elements of this vesicular release machinery. As a member of the soluble N-ethylmaleimide-sensitive factor attachment protein receptor (SNARE) protein family, it essentially functions in vesicle fusion, and in the prior preparatory steps, the vesicle docking, and priming.

Knock-out studies in mammalian systems addressing Stx1's functions in neurotransmission has been complicated by the functional redundancy between the two isoforms Stx1A, and Stx1B, and by the embryonic lethality in the absence of both isoforms. In this project, we aimed to circumvent these complications to investigate the roles of Stx1 in neurotransmitter release, and in neuronal maintenance, and generated two Stx1A/1B double knock-out (DKO) mouse models:

1) The constitutive Stx1A/1B DKO mice exhibited a widespread neuronal loss throughout the whole brain at embryonic day (E) 16 progressing towards E18. The cultured cortical neurons from E16 Stx1A/1B DKO mice were also severely lethal as almost no neurons remained alive at day *in vitro* 4. Comparison with the lethality of Munc18-1 null neurons in the same culture conditions manifested a higher dependence of neuronal maintenance on Stx1 than on Munc18-1.

2) The conditional deletion of Stx1B via Cre/LoxP system on an Stx1A null background (Stx1A/1B cDKO) revealed a perpetual requirement of Stx1 for neuronal survival as the deletion of Stx1 in mature neurons ultimately caused neuronal death. Additionally, neurotransmitter release measurements by electrophysiological recordings and presynaptic ultrastructural analysis by electron microscopy showed a complete inhibition of vesicle priming and fusion, and a severe impairment in vesicle docking in the dissociated hippocampal Stx1A/1B cDKO neurons. Lentiviral expression of an Stx1A mutant (A240V, V244A) in Stx1A/1B cDKO neurons rescued the neuronal survival but not the vesicle fusion and priming implying distinct functions of Stx1 in neuronal maintenance and neurotransmitter release. Strikingly, this mutant restored vesicle docking, which indicates a further distinction between the functions of Stx1 in vesicle docking and fusion.

Future studies will focus on structure-function analysis of Stx1 in neuronal maintenance and neurotransmitter release by utilizing lentiviral expression of Stx1 mutants in Stx1A/1B cDKO mouse neurons. Such a detailed analysis of the functions of Stx1's different domains in vesicle docking, priming, and fusion would enhance our understanding of the intricate mechanisms underlying neurotransmission. Moreover, discovery of the pathways involving Stx1 with a function in neuronal maintenance would shed light onto the interdependence of neurodegeneration and synaptic functions.

Zusammenfassung

Neurone übertragen Informationen untereinander durch die Freisetzung von Neurotransmittern, wobei die mit Neurotransmittern gefüllten präsynaptischen Vesikel mit der Plasmamembran fusionieren. Dieser elementare Prozess wird durch ein komplexes Zusammenspiel präsynaptischer Proteine reguliert. Eines der Kernbestandteile dieses vesikulären Proteinapparates ist Syntaxin-1 (Stx1). Als ein Teil der soluble N-ethylmaleimide-sensitive factor attachment protein receptor (SNARE) Proteinfamilie wirkt es bei der Vesikelfusion, sowie in den vorangehenden Vorbereitungsschritten, bei dem Andocken und Priming mit.

Knock-out Analysen der Funktionen von Stx1 in der Neurotransmitterfreisetzung wurden einerseits durch die kompensatorischen Effekte der zwei neuronalen Isoformen Stx1A und Stx1B und andererseits durch die embryonale Letalität in Abwesenheit beider Isoformen erschwert. Das Ziel dieses Projektes war es daher, diese Komplikationen zu umgehen und die Funktionen von Stx1 bei der Neurotransmitterfreisetzung und auch bei der Überlebensfähigkeit der Neurone zu untersuchen. Zu diesem Zweck haben wir zwei Doppelknockout (DKO) Mausmodelle generiert.

1) Konstitutive Stx1A/1B DKO Mäuse zeigten ein gehirnweites Absterben der Neurone vom embryonalen Tag (E) 16 an, welches bis E18 fortschritt. Das Absterben kam auch bei kultivierten kortikalen Neuronen von E16 Stx1A/1B DKO Mäusen vor: nur sehr wenige Neurone überlebten bis Tag 4 *in vitro*. Ein direkter Vergleich des Verlusts von Munc18-1 und Stx1 unter den gleichen Zellkulturbedingungen zeigte eine größere Abhängigkeit von Stx1 als von Munc18-1 beim Überleben der Neurone.

2) Die konditionale Deletion von Stx1B mit Hilfe des Cre/LoxP-Systems in Stx1A null Neurone (Stx1A/1B cDKO) zeigte eine permanente Notwendigkeit von Stx1 für das neuronale Überleben, da die Deletion von Stx1 in reifen Neuronen zum Zelltod führte. Außerdem konnten mit elektrophysiologischen Messungen der Neurotransmitterfreisetzung und mit Ultrastrukturanalyse der synaptischen Kompartimente durch Elektronenmikroskopie eine komplette Hemmung der Vesikelfusion und des Vesikelprimings, sowie schwere Beeinträchtigungen des Andockens von Vesikeln in dissoziierten hippokampalen Neuronen gezeigt werden. Die lentivirale Expression eines Stx1A-Mutanten (A240V, V244A) konnte das neuronale Überleben, jedoch nicht die Fusion oder das Priming von Vesikeln retten. Dies deutet auf verschiedene Funktionen von Stx1 bei der Überlebensfähigkeit der Neurone und in der Neurotransmitterfreisetzung hin. Erstaunlicherweise stellte dieser Mutant auch die angedockten Vesikel wieder her, was einen weiteren Unterschied in den Funktionen von Stx1 im Andocken und in der Fusion von Vesikeln offenbart.

Zukünftige Studien werden sich auf die Struktur-Funktionsanalyse von Stx1 in Bezug auf das Überleben von Neuronen und die Neurotransmitterfreisetzung durch Nutzung lentiviraler Expression von Stx1-Mutanten in Stx1A/1B cDKO Mausneuronen konzentrieren. Eine solch detaillierte Analyse der Funktionen der verschiedenen Domänen von Stx1 im Andocken, Priming und Fusion von Vesikeln würde unser Verständnis der komplexen Mechanismen, die der Neurotransmitterfreisetzung zugrunde liegen, fördern. Darüberhinaus würden Erkenntnisse über die Funktionen von Stx1 im neuronalen Überleben Aufschluss über die wechselseitige Abhängigkeit von Neurodegeneration und synaptischen Funktionen geben.

Affidavit

I, Gülçin Vardar, certify under penalty of perjury by my own signature that I have submitted the thesis on the topic “Distinct Functions of Syntaxin-1 in Neuronal Maintenance, Synaptic Vesicle Docking, and Fusion in Mouse Neurons” I wrote this thesis independently and without assistance from third parties, I used no other aids than the listed sources and resources.

All points based literally or in spirit on publications or presentations of other authors are, as such, in proper citations (see "uniform requirements for manuscripts (URM)" the ICMJE www.icmje.org) indicated. The section on methodology (in particular practical work, laboratory requirements, statistical processing) and results (in particular images, graphics and tables) corresponds to the URM (s.o) and are answered by me. My contribution in the selected publication for this dissertation corresponds to those that are specified in the following joint declaration with the responsible person and supervisor.

The importance of this affidavit and the criminal consequences of a false affidavit (section 156,161 of the Criminal Code) are known to me and I understand the rights and responsibilities stated therein.

Date

Signature

Detailed Declaration of Contribution

Gülçin Vardar had the following share in the following publication:

Publication : Vardar G, Chang S, Arancillo M, Wu YJ, Trimbuch T, Rosenmund C. Distinct Functions of Syntaxin-1 in Neuronal Maintenance, Synaptic Vesicle Docking, and Fusion in Mouse Neurons. J Neurosci. 2016

Contribution in detail:

Design of the experiments:

The experiments were designed by Vardar G and Rosenmund C

Performance of the experiments:

The generation of the mouse lines by Arancillo M.

Cesarean section of pregnant females from timed breedings by Vardar G together with Wu YJ (Figure 1).

Histology and Western Blots by Vardar G (Figures 1 and 3).

Culture preparations by Vardar G (Figures 2-7).

Lentiviral production by Trimbuch T together with Vardar G (Figures 2-7).

Immunocytochemistry by Vardar G (Figures 2, 5, and 6).

Electrophysiological recordings by Vardar G (Figures 5 and 6).

High pressure freezing of cultured neurons and acquisition of electron microscopy images by Chang S together with Vardar G (Figure 7).

Analysis of the experiments:

The analysis of the data was performed by Vardar G (Figures 1-7).

Writing and editing the publication:

The publication was written and edited by Vardar G together with Rosenmund C.

Signature, date and stamp of the supervising University teacher

Signature of the doctoral candidate

ISI Web of KnowledgeSM

Journal Citation Reports[®]

[WELCOME](#) [HELP](#)

2015 JCR Science Edition

Journal Summary List

[Journal Title Changes](#)

Journals from: **subject categories NEUROSCIENCES** [VIEW CATEGORY SUMMARY LIST](#)

Sorted by: [SORT AGAIN](#)

Journals 1 - 20 (of 256)

Navigation: [1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10]

Page 1 of 13

[MARK ALL](#) [UPDATE MARKED LIST](#)

Ranking is based on your journal and sort selections.

Mark	Rank	Abbreviated Journal Title <i>(linked to journal information)</i>	ISSN	JCR Data ⁱ						Eigenfactor [®] Metrics ⁱ	
				Total Cites	Impact Factor	5-Year Impact Factor	Immediacy Index	Articles	Cited Half-life	Eigenfactor [®] Score	Article Influence [®] Score
<input type="checkbox"/>	1	NAT REV NEUROSCI	1471-003X	33792	29.298	35.142	5.603	58	8.0	0.07244	16.183
<input type="checkbox"/>	2	BEHAV BRAIN SCI	0140-525X	7873	20.415	23.842	1.700	10	>10.0	0.01286	11.186
<input type="checkbox"/>	3	TRENDS COGN SCI	1364-6613	21382	17.850	23.872	2.444	72	8.6	0.04719	10.507
<input type="checkbox"/>	4	NAT NEUROSCI	1097-6256	51112	16.724	16.874	4.100	221	7.8	0.15468	9.141
<input type="checkbox"/>	5	ANNU REV NEUROSCI	0147-006X	13125	14.265	22.563	3.000	21	>10.0	0.02068	11.493
<input type="checkbox"/>	6	NEURON	0896-6273	78507	13.974	16.092	2.927	385	8.3	0.22223	8.421
<input type="checkbox"/>	7	MOL PSYCHIATR	1359-4184	15797	13.314	13.274	2.892	167	5.6	0.04440	4.877
<input type="checkbox"/>	8	PROG NEUROBIOL	0301-0082	11877	13.177	11.181	1.318	44	9.7	0.02031	3.730
<input type="checkbox"/>	9	TRENDS NEUROSCI	0166-2236	18656	12.504	13.470	2.464	69	>10.0	0.03012	5.928
<input type="checkbox"/>	10	ACTA NEUROPATHOL	0001-6322	14701	11.360	10.616	2.743	109	6.7	0.03692	4.062
<input type="checkbox"/>	11	BIOL PSYCHIAT	0006-3223	42289	11.212	10.799	3.653	190	8.1	0.07406	3.724
<input type="checkbox"/>	12	BRAIN	0006-8950	46207	10.103	10.545	2.320	259	8.9	0.08378	4.110
<input type="checkbox"/>	13	ANN NEUROL	0364-5134	32995	9.638	10.880	2.277	177	>10.0	0.05944	4.357
<input type="checkbox"/>	14	J PINEAL RES	0742-3098	6914	9.314	7.140	2.391	87	6.1	0.00779	1.142
<input type="checkbox"/>	15	FRONT NEUROENDOCRIN	0091-3022	3407	8.852	9.045	1.297	37	6.5	0.00725	3.054
<input type="checkbox"/>	16	NEUROSCI BIOBEHAV R	0149-7634	18522	8.580	10.504	1.498	233	6.2	0.04521	3.615
<input type="checkbox"/>	17	CEREB CORTEX	1047-3211	26911	8.285	7.881	1.423	409	7.1	0.06806	3.376
<input type="checkbox"/>	18	SLEEP MED REV	1087-0792	4392	7.341	9.636	2.375	48	6.6	0.01033	3.450
<input type="checkbox"/>	19	NEUROSCIENTIST	1073-8584	4096	7.295	7.609	1.977	44	6.9	0.01008	2.899
<input type="checkbox"/>	20	MOL NEURODEGENER	1750-1326	2312	6.510	5.599	0.523	65	4.1	0.00933	1.966

[MARK ALL](#) [UPDATE MARKED LIST](#)

Journals 1 - 20 (of 256)

Navigation: [1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | 10]

Page 1 of 13

[Acceptable Use Policy](#)
Copyright © 2017 [Thomson Reuters](#).





Journal Citation Reports®

WELCOME ? HELP

2015 JCR Science Edition

Journal Summary List

[Journal Title Changes](#)

Journals from: **subject categories NEUROSCIENCES** [VIEW CATEGORY SUMMARY LIST](#)

Sorted by: [SORT AGAIN](#)

Journals 21 - 40 (of 256)

Navigation icons: back, forward, search, etc.

Page 2 of 13

MARK ALL UPDATE MARKED LIST

Ranking is based on your journal and sort selections.

Mark	Rank	Abbreviated Journal Title <i>(linked to journal information)</i>	ISSN	JCR Data ⁱ						Eigenfactor® Metrics ⁱ	
				Total Cites	Impact Factor	5-Year Impact Factor	Immediacy Index	Articles	Cited Half-life	Eigenfactor® Score	Article Influence® Score
<input type="checkbox"/>	21	NEUROPSYCHOPHARMACOL	0893-133X	22869	6.399	7.825	1.794	291	6.7	0.04998	2.648
<input type="checkbox"/>	22	CURR OPIN NEUROBIOL	0959-4388	13090	6.373	6.958	1.671	152	8.5	0.03549	3.503
<input type="checkbox"/>	23	NEUROPSYCHOL REV	1040-7308	2314	6.061	7.594	0.440	25	6.6	0.00476	2.461
<input type="checkbox"/>	24	CEPHALALGIA	0333-1024	7906	6.052	4.363	1.333	132	7.8	0.01187	1.140
<input type="checkbox"/>	25	GLIA	0894-1491	11548	5.997	5.411	1.285	158	7.7	0.02050	1.688
<input checked="" type="checkbox"/>	26	J NEUROSCI	0270-6474	172201	5.924	6.780	1.148	1342	8.5	0.35220	2.779
<input type="checkbox"/>	27	BRAIN BEHAV IMMUN	0889-1591	9464	5.874	6.020	1.462	238	4.9	0.02397	1.791
<input type="checkbox"/>	28	BRAIN STRUCT FUNCT	1863-2653	3534	5.811	6.803	1.307	244	3.0	0.01503	2.510
<input type="checkbox"/>	29	J PSYCHIATR NEUROSCI	1180-4882	2690	5.570	6.207	1.136	44	6.5	0.00551	2.034
<input type="checkbox"/>	30	PAIN	0304-3959	32930	5.557	6.244	1.119	269	>10.0	0.04647	2.053
<input type="checkbox"/>	31	NEUROIMAGE	1053-8119	79475	5.463	6.797	1.160	770	6.7	0.17977	2.321
<input type="checkbox"/>	32	MOL NEUROBIOL	0893-7648	5142	5.397	5.392	1.155	283	3.4	0.01331	1.480
<input type="checkbox"/>	33	BRAIN PATHOL	1015-6305	4403	5.256	4.485	1.026	77	7.5	0.00877	1.547
<input type="checkbox"/>	34	ALZHEIMERS RES THER	1758-9193	1164	5.197	5.126	0.921	76	2.5	0.00537	1.725
<input type="checkbox"/>	35	FRONT MOL NEUROSCI	1662-5099	1592	5.154		0.329	79	3.2	0.00782	
<input type="checkbox"/>	36	NEUROBIOL AGING	0197-4580	19205	5.153	5.193	1.261	429	5.7	0.04573	1.662
<input type="checkbox"/>	37	SOC COGN AFFECT NEUR	1749-5016	4695	5.101	5.532	0.848	210	3.4	0.01834	2.069
<input type="checkbox"/>	38	HUM BRAIN MAPP	1065-9471	17184	4.962	5.638	0.931	378	6.5	0.04035	2.084
<input type="checkbox"/>	39	MOL AUTISM	2040-2392	847	4.961	5.184	0.921	63	2.5	0.00450	1.956
<input type="checkbox"/>	40	NEUROPHARMACOLOGY	0028-3908	17341	4.936	4.709	1.288	365	5.8	0.03930	1.457

MARK ALL UPDATE MARKED LIST

Journals 21 - 40 (of 256)

Navigation icons: back, forward, search, etc.

Page 2 of 13

[Acceptable Use Policy](#)
Copyright © 2017 Thomson Reuters.



Published by Thomson Reuters

Distinct Functions of Syntaxin-1 in Neuronal Maintenance, Synaptic Vesicle Docking, and Fusion in Mouse Neurons

Gülçin Vardar,^{1,2}  Shuwen Chang,^{1,2} Marife Arancillo,²  Yuan-Ju Wu,²  Thorsten Trimbuch,^{1,2} and  Christian Rosenmund^{1,2}

¹Department of Neurophysiology and ²NeuroCure Cluster of Excellence, Charité Universitätsmedizin Berlin, 10117 Berlin, Germany

Neurotransmitter release requires the formation of soluble *N*-ethylmaleimide-sensitive factor attachment protein receptor (SNARE) complexes by SNARE proteins syntaxin-1 (Stx1), synaptosomal-associated protein 25 (SNAP-25), and synaptobrevin-2 (Syb2). In mammalian systems, loss of SNAP-25 or Syb2 severely impairs neurotransmitter release; however, complete loss of function studies for Stx1 have been elusive due to the functional redundancy between Stx1 isoforms Stx1A and Stx1B and the embryonic lethality of Stx1A/1B double knock-out (DKO) mice. Here, we studied the roles of Stx1 in neuronal maintenance and neurotransmitter release in mice with constitutive or conditional deletion of Stx1B on an Stx1A-null background. Both constitutive and postnatal loss of Stx1 severely compromised neuronal viability *in vivo* and *in vitro*, indicating an obligatory role of Stx1 for maintenance of developing and mature neurons. Loss of Munc18-1, a high-affinity binding partner of Stx1, also showed severely impaired neuronal viability, but with a slower time course compared with Stx1A/1B DKO neurons, and exogenous Stx1A or Stx1B expression significantly delayed Munc18-1-dependent lethality. In addition, loss of Stx1 completely abolished fusion-competent vesicles and severely impaired vesicle docking, demonstrating its essential roles in neurotransmission. Putative partial SNARE complex assembly with the SNARE motif mutant Stx1A^{AV} (A240V, V244A) was not sufficient to rescue neurotransmission despite full recovery of vesicle docking and neuronal survival. Together, these data suggest that Stx1 has independent functions in neuronal maintenance and neurotransmitter release and complete SNARE complex formation is required for vesicle fusion and priming, whereas partial SNARE complex formation is sufficient for vesicle docking and neuronal maintenance.

Key words: autapse; cell culture; Munc18; SNARE; vesicle priming

Significance Statement

Syntaxin-1 (Stx1) is a component of the synaptic vesicle soluble *N*-ethylmaleimide-sensitive factor attachment protein receptor (SNARE) complex and is essential for neurotransmission. We present the first detailed loss-of-function characterization of the two Stx1 isoforms in central mammalian neurons. We show that Stx1 is fundamental for maintenance of developing and mature neurons and also for vesicle docking and neurotransmission. We also demonstrate that neuronal maintenance and neurotransmitter release are regulated by Stx1 through independent functions. Furthermore, we show that SNARE complex formation is required for vesicle fusion, whereas partial SNARE complex formation is sufficient for vesicle docking and neuronal maintenance. Therefore, our work provides insights into differential functions of Stx1 in neuronal maintenance and neurotransmission, with the latter explored further into its functions in vesicle docking and fusion.

Introduction

Syntaxin-1 (Stx1) is a neuronal soluble *N*-ethylmaleimide-sensitive factor attachment protein receptor (SNARE) protein located at the plasma membrane (t-SNARE) and involved in

synaptic transmission by interactions with another neuronal t-SNARE, synaptosomal-associated protein 25 (SNAP-25), and the vesicle-SNARE (v-SNARE), synaptobrevin-2 (Syb2). The

Received April 21, 2016; revised May 20, 2016; accepted June 9, 2016.

Author contributions: G.V. and C.R. designed research; G.V., S.-W.C., M.A., Y.-J.W., and T.T. performed research; G.V. analyzed data; G.V. and C.R. wrote the paper.

This work was supported by the NeuroCure Cluster of Excellence ExC257 and the Medical Neuroscience PhD training program (G.V.), the European Research Council (Advanced Grant SynVGlut), and the German Research Council (Collaborative Research Grant SFB958 to C.R.). We thank Mathijs Verhage for the Munc18-1 KO mouse line; 11

the Charité Viral Core facility and Katja Pötschke, Bettina Brokowski, Berit Söhl-Kielczynski, Rike Dannenberg, Sabine Lenz, and Pascal Fenske for technical assistance; Eric Reynolds for comments on the manuscript; and Melissa Herman for discussions and critical feedback on the manuscript.

The authors declare no competing financial interests.

Correspondence should be addressed to Christian Rosenmund, NeuroCure Cluster of Excellence, Charité Universitätsmedizin Berlin, Charitéplatz 1, Charité-Cross-Over, 10117 Berlin, Germany. E-mail: christian.rosenmund@charite.de.

DOI:10.1523/JNEUROSCI.1314-16.2016

Copyright © 2016 the authors 0270-6474/16/367911-14\$15.00/0

α -helical SNARE motifs of Stx1, SNAP-25, and Syb2 form the ternary SNARE complex, which is thought to catalyze vesicle fusion by bringing vesicular and plasma membranes in close proximity (Rizo and Rosenmund, 2008; Rizo and Südhof, 2012).

Gene deletion studies of the neuronal SNARE proteins in vertebrate and invertebrate model organisms illustrate the crucial roles of the SNAREs in vesicle fusion and in the preceding preparatory steps, vesicle docking and priming. Both in Syb2 knock-out (KO) mice and SNAP-25 KO mice, Ca^{2+} -evoked neurotransmitter release is abrogated, but a low level of spontaneous neurotransmitter release is sustained (Schoch et al., 2001; Washbourne et al., 2002; Delgado-Martínez et al., 2007). However, loss of the neuronal Sec1/Munc18-like protein Munc18-1, the tight binding partner of Stx1 (Hata et al., 1993), results in a remarkably more severe phenotype in mammals with a complete inhibition of neurotransmitter release accompanied by neuronal degeneration (Verhage et al., 2000), with similar results obtained in flies (Harrison et al., 1994).

The widespread neuronal degeneration in Munc18-1 KO mice was first proposed as a direct consequence of lack of synaptic input (Verhage et al., 2000). However, this proposal was challenged by the findings from a Munc13-1/2 double KO (DKO) mouse model, which is also completely synaptically silent but without any cytotoxic effects (Varoqueaux et al., 2002). Ultimately, it was proposed that the underlying mechanism for the cell-autonomous degeneration of Munc18-1 KO neurons is related to another function of Munc18-1 independent of synaptic transmission (Heeroma et al., 2004). So far, only deletion of Stx1 in flies has manifested an equally severe phenotype among SNAREs, with a complete inhibition of neurotransmission accompanied by cell lethality (Schulze et al., 1995; Schulze and Bellen, 1996; Burgess et al., 1997). The tight interactions between Stx1 and Munc18-1 and the similarities between their loss-of-function phenotypes strongly suggest the involvement of Stx1 and Munc18-1 as a complex in neuronal maintenance.

In mammals, Stx1 has two ubiquitously expressed neuronal isoforms, Stx1A and Stx1B, which share 84% homology (Bennett et al., 1992). Investigation of Stx1 function in neurotransmitter release has been rendered impracticable in mammals for two main reasons: mutual compensatory effects of Stx1A and Stx1B (Fujiwara et al., 2006; Gerber et al., 2008; Arancillo et al., 2013; Zhou et al., 2013; Wu et al., 2015) and embryonic lethality due to constitutive deletion of Stx1A/1B (Mishima et al., 2014). Here, we present a comprehensive study of Stx1's roles in neuronal maintenance and neurotransmitter release by bypassing these obstacles through the utilization of two Stx1A/1B DKO mouse models. The first, a constitutive deletion of Stx1A/1B, led to severe neuronal lethality *in vitro* and *in vivo*, indicating the absolute requirement for Stx1 in neuronal maintenance. By using the second mouse model, a Stx1A/1B conditional DKO (cDKO), we showed that deletion of Stx1 in mature neurons also caused severe impairments in neuronal viability, demonstrating Stx1 dependency for maintenance throughout the neuronal life span. In addition, loss of Stx1 completely inhibited vesicle fusion and priming and resulted in significant reduction in docked vesicles, showing the essential roles of Stx1 for each step of neurotransmitter release: vesicle docking, priming, and fusion. Finally, exogenous expression of a mutant isoform Stx1A^{AV} (A240V, V244A), which forms SNARE complexes with reduced stability (Kee et al., 1995; Fergestad et al., 2001), rescued neuronal survival and vesicle docking in Stx1-deficient neurons, but could not rescue any form of vesicle fusion. We conclude that the func-

tions of Stx1 in neuronal maintenance and neurotransmitter release are independent from one another, that full SNARE complex assembly is crucial for vesicle fusion and priming, and that partial SNARE complex assembly is sufficient for vesicle docking.

Materials and Methods

Animal maintenance and generation of mouse lines. All procedures for animal maintenance and experiments were in accordance with the regulations of and approved by the animal welfare committee of Charité Medical University and the Berlin state Government Agency for Health and Social Services under license number T0220/09. The generation of Stx1A KO (−/−), Stx1B Het (+/−), Stx1B^{FL/FL}, and Munc18-1 KO mouse lines were described previously (Verhage et al., 2000; Gerber et al., 2008; Arancillo et al., 2013; Wu et al., 2015). Stx1A KO mice were intercrossed with Stx1B Het or Stx1B^{FL/FL} to generate Stx1A/1B DKO or Stx1A/1B cDKO mouse lines, respectively.

Histology and Western blot. Mouse embryos of either sex were obtained by cesarean section of pregnant females from timed heterozygous breedings. For histological analysis of the brain tissues, whole heads of mouse embryos at embryonic day (E) 16 and E18 were immersion fixed in 4% paraformaldehyde (PFA) in 0.1 M sodium phosphate buffer (PB) for two nights at 4°C. The tissues were incubated in 0.4 M and then 0.8 M sucrose in 0.1 M PB, overlaid with Tissue-Tek, and snap-frozen in n-hexane at −70°C for cryoprotection. The tissues were then cryosectioned sagittally using an HM 500 OM cryostat (MICROM International) with a thickness setting of 20 μ m. The sections were mounted directly on gelatin-coated glass slides. Every sixth section was subjected to Nissl staining with 0.2% cresyl violet acetate in 20 mM acetic acid solution and then examined under an Olympus SZX16 microscope with an Olympus DP70 digital camera. For Western blot, whole-brain lysates were obtained from E16 or E18 embryos depending on the experiment. The samples were dissolved in lysis buffer containing 50 mM Tris/HCl, pH 7.9, 150 mM NaCl, 5 mM EDTA, 1% Triton X-100, 0.5% sodium deoxycholate, 1% Nonidet P-40, and 1 tablet of Complete Protease Inhibitor (Roche) for 10 ml of solution. Equal amounts of solubilized proteins were loaded in 8–12% SDS-PAGE and subsequently transferred to nitrocellulose membranes. The membranes were subjected to the following primary antibodies overnight at 4°C: mouse monoclonal anti- β -tubulin III (1:2000) as an internal control, rabbit polyclonal Munc18-1 (1:2000; both purchased from Sigma-Aldrich), mouse monoclonal anti-Stx1A (1:1000), rabbit polyclonal anti-Stx1B (1:1000), mouse monoclonal anti-SNAP-25 (1:2000), mouse monoclonal anti-Syb-2 (1:2000), rabbit polyclonal anti-Munc13-1 (1:1000), mouse monoclonal anti-synaptotagmin-1 (1:1000), rabbit polyclonal anti-complexin-1/2 (1:500), and mouse monoclonal anti-Rab-5 (1:500; all purchased from Synaptic Systems). Horseradish peroxidase-conjugated goat secondary antibodies (Jackson ImmunoResearch) were applied for 1 h at room temperature and detected with ECL Plus Western Blotting Detection Reagents (GE Healthcare Biosciences) in Fusion FX7 image and analytics system (Vilber Lourmat). Ratiometric quantification of protein expression levels was performed using ImageJ software. Data from Stx1A KO and Stx1B Het/Stx1A KO animals were pooled together after no significant differences were observed between these genotypes.

Neuronal cultures and neuronal count. Cortical cultures were prepared from E16 mice of either sex, which were obtained by cesarean section of pregnant females from timed heterozygous breedings. Hippocampal cultures were prepared from mice of either sex at postnatal day (P) 0–2 as described previously (Xue et al., 2007; Arancillo et al., 2013). Briefly, the neurons were seeded with densities as follows: 50 \times 10³/well (22 mm diameter) for neuronal counting, 30 \times 10³/well (22 mm diameter) for immunocytochemistry on continental astrocyte feeder layers as high-density cultures, and 3 \times 10³ neurons/well (35 mm diameter) for electrophysiology on micro-island astrocyte feeder layers. The astrocyte feeder layers were prepared 1–2 weeks before the neuronal culture preparation as described previously (Arancillo et al., 2013). The cortical cultures were transduced with lentiviruses immediately after plating at day *in vitro* (DIV) 0 and the hippocampal cultures were transduced at DIV

1–7 by lentiviral infection according to the experiment. The cultures were incubated in Neurobasal-A medium supplemented with B-27 (Invitrogen), 50 IU/ml penicillin, and 50 μ g/ml streptomycin at 37°C between the days of neuronal counting or until DIV 13–17 before immunocytochemical or electrophysiological analysis.

The viability of the neurons *in vitro* was defined as the number of surviving neurons at different time points between DIV 1 and DIV 42 according to the experiment. Phase-contrast bright-field images of 10 randomly selected fields of 1.23 mm² per well and of 2 wells per group in each culture were acquired with a DMI4000 microscope, DFC 345 FX camera, HCX PL FLUOTAR 10 \times objectives, and LAS-AF software (all from Leica). The neurons between DIV 1 and DIV 7 were counted manually with ImageJ software. After DIV 7, when the nuclear localization signal (NLS)-green fluorescent protein (GFP) and/or NLS-red fluorescent protein (RFP) were at adequate levels for detection, additional fluorescent images were acquired with excitation wavelengths of 488 and 555 nm from the same selected fields and the neurons were counted with the 3D Objects Counter function in Fiji software. To determine the survival of the Stx1A/1B DKO and Munc18-1 KO neurons, the number of neurons in each group was normalized to the number of seeded neurons as a common reference, whereas the rescue efficiencies of lentiviral reintroduction of Stx1A, Stx1B, or Munc18-1 were calculated as the normalized ratio of the number of surviving neurons to the number of Stx1A KO or wild-type (WT) neurons, respectively. To evaluate the rates of cell death of Stx1A/1B cDKO neurons expressing Cre recombinase, the number of neurons was normalized to the number of neurons in control groups at DIV 7 for the cultures infected at DIV 1. For the cultures infected at DIV 3, 5, or 7, the number of neurons within each group at DIV 7 was used as reference for normalization. The ability of WT or mutant Stx1A to rescue the neurons against cell death was examined by coinfection with viruses to express Cre recombinase and determined as the ratio of the number of neurons at DIV 24–25 to the number of neurons at DIV 11.

Lentiviral constructs. All lentiviral constructs were based on the vector FUGW (Lois et al., 2002). The cDNA for mouse Stx1A (NM_016801), mouse Stx1B (NM_024414), or rat Munc18-1 (NM_013038) were cloned after a NLS-GFP-P2A expression cassette controlled by human synapsin-1 promoter. The P2A sequence allows for the bicistronic expression of nuclear-targeted GFP and the corresponding cDNA (Kim et al., 2011). Empty vectors, including only an NLS-GFP-P2A or NLS-RFP-P2A expression cassette, served as controls. A lentivirus construct, which induces expression of improved Cre recombinase (iCre) fused to NLS-RFP-P2A under control of human synapsin-1 promoter, was used to permit excision of the LoxP flanked Stx1B DNA. The point mutations A240V and V244A were introduced into Stx1A cDNA (Stx1A^{AV}) with the QuikChange Site-Directed Mutagenesis Kit (Stratagene). All lentiviral particles were provided by the Viral Core Facility of the Charité Berlin and were prepared as described previously (Lois et al., 2002). Briefly, HEK293T cells were cotransfected with 10 μ g of shuttle vector, 5 μ g of helper plasmids pCMVdr8.9, and 5 μ g of pVSV.G with X-tremeGENE 9 DNA transfection reagent (Roche Diagnostic). Virus containing cell culture supernatant was collected after 72 h and filtered for purification. Aliquots were flash-frozen in liquid nitrogen and stored at -80°C . Viruses were titrated visually with mass-cultured WT hippocampal mouse neurons using the fluorescence reporter. For infection, $\sim 1 \times 10^6$ infectious virus units per 22- or 35-mm-diameter well were added onto the cultured hippocampal or cortical mouse neurons between DIV 0 and 7 depending on the experiment.

Immunofluorescence labeling. High-density cortical cultures were fixed at DIV 2, 4, or 22, whereas high-density hippocampal cultures were fixed at DIV 12, 15, or 21. Both cultures were fixed in 4% PFA for 20 min, permeabilized with 1 \times PBS containing 0.1% Tween 20 (PBS-T) for 20 min, and blocked with PBS-T containing 2.5% normal goat serum (NGS) and 2.5% normal donkey serum (NDS) for 1 h. The cortical cultures fixed at DIV 2 and 4 were incubated overnight at 4°C with chicken polyclonal anti-MAP2 (Merck) primary antibody (1:2000) in PBS-T containing NGS and NDS. The cortical cultures fixed at DIV 22 and the hippocampal cultures were subjected to guinea pig polyclonal anti-Bassoon (Synaptic Systems) primary antibody (1:2500), which was used

as a synaptic marker, in addition to the primary antibodies against Stx1A (1:1000) and Stx1B (1:1000) mentioned above, overnight at 4°C. Subsequently, the cells were incubated in the following fluorescently labeled secondary antibodies (1:500) according to the experiments: Alexa Fluor 647 (A647) donkey anti-chicken IgG or A647 donkey anti-mouse IgG, rhodamine red donkey anti-guinea pig IgG (all purchased from Jackson ImmunoResearch), and Pacific blue goat anti-rabbit IgG (Invitrogen). The coverslips were mounted on glass slides with MOWIOL mounting agent (Sigma-Aldrich). Immunofluorescence labeling was visualized under an Olympus IX81 epifluorescent microscope with 60 \times optical magnification, MicroMax:1300YHS camera (Princeton Instruments) and MetaMorph software (Molecular Devices) using Z-stacks. The images were analyzed offline with ImageJ as described previously (Arancillo et al., 2013). Briefly, the Z-stacks were overlapped with maximum intensity and the background was subtracted with a rolling ball of a radius of 30 pixels. Two to four regions exhibiting only neuronal extensions were cropped per image and analyzed as independent measurements. Bassoon-positive spots in these crops were determined using the threshold function with MaxEntropy and the created areas were saved as ROIs. The intensities of Bassoon, Stx1A, and Stx1B signals were measured using the corresponding ROIs. The average ratio of the Stx1A and Stx1B signals to the Bassoon signal was calculated for each crop and the ratios were normalized to the control group.

Electrophysiology. Whole-cell patch-clamp recordings were performed on autaptic hippocampal neurons at DIV 13–17 at room temperature with a Multiclamp 700B amplifier controlled by Clampex 9.2 software (both from Molecular Devices). Data were digitally sampled at 10 kHz and low-pass filtered at 3 kHz with an Axon Digidata 1322A digitizer (Molecular Devices). The standard extracellular solution was applied with a fast perfusion system (1–2 ml/min) and contained the following (in mM): 140 NaCl, 2.4 KCl, 10 HEPES, 10 glucose, 2 CaCl₂, and 4 MgCl₂ (~ 300 mOsm; pH 7.4). Borosilicate glass patch pipettes were pulled with a multistep puller, yielding a final tip resistance of 2–5 M Ω when filled with KCl based intracellular solution containing the following (in mM): 136 KCl, 17.8 HEPES, 1 EGTA, 4.6 MgCl₂, 4 ATP-Na₂, 0.3 GTP-Na₂, 12 creatine phosphate, and 50 U/ml phosphocreatine kinase (~ 300 mOsm; pH 7.4). Neurons transduced with viruses were visually identified by RFP and GFP expression before recordings. Series resistance was compensated by 70% and monitored over the course of each recording. The recordings were analyzed offline using Axograph X Version 1.5.4 (Axograph Scientific).

Glutamatergic hippocampal neurons were clamped at -70 mV at steady state and EPSCs were evoked by a 2 ms depolarizing step to 0 mV. EPSCs were then recorded in the presence of 3 μM NBQX (Tocris Bioscience), which is an AMPA receptor antagonist. The averages of the traces recorded in the presence of NBQX were subtracted by using subtraction function in AxographX from the average of the traces recorded in the external solution without NBQX to cancel stimulus artifacts. Spontaneous release was determined by monitoring miniature EPSCs (mEPSCs) for 30–60 s at -70 mV. To correct false-positive events, mEPSCs were again recorded in the presence of 3 μM NBQX. The recorded traces were filtered at 1 kHz and mEPSCs were detected with a template function in Axograph X. The readily releasable pool (RRP) size was determined by a 5 s application of extracellular solution that additionally contained 500 mM sucrose (Rosenmund and Stevens, 1996). The solutions were exchanged with a fast microperfusion device (SF-77B; Warner Instruments).

High-pressure freezing and electron microscopy. Hippocampal neurons were plated onto 6 mm sapphire disks with a density of 50×10^3 to 100×10^3 cells. The neurons were coinfecting at DIV 2 with viruses to express Cre recombinase in combination with control viruses or with viruses to express Stx1A^{WT} or Stx1A^{AV}. At DIV 12–14, the samples were mounted into a sandwich support with extracellular solution containing 15% Ficoll (Sigma-Aldrich) to avoid ice crystal damage and frozen with an HPM 100 Leica high-pressure freezer under a pressure of 2100 bar. Samples were then transferred into cryovials containing 1% glutaraldehyde, 1% osmium tetroxide, 1% Milli-Q water (Millipore) in anhydrous acetone and processed in an AFS2 automated freeze-substitution device (Leica) followed by a temperature ramp from -90°C (8°C/h) to -50°C , from

–50°C (6°C/h) to –20°C, constant temperature for 12 h at –20°C, and a ramp from –20°C (5°C/h) to 20°C. After the freeze-substitution step, samples were contrasted with 0.1% uranyl acetate for 1 h and embedded in pure epoxy epon 812. Finally, samples were placed into capsules filled with pure epoxy epon 812 and further polymerized for 48 h at 60°C. Areas of ~200 μm^2 containing cells were randomly selected and 40-nm-thick sections were obtained with a Leica Ultracut and collected on Formvar-coated (0.5%) 200-mesh copper grids. Those sections were further examined on a Tecnai G20 electron microscope operated at 80–200 keV (FEI). Electron micrographs (2048 × 2048 pixels) were collected blindly using a Velata camera (FEI) with an 89,000× magnification (pixel size 0.57 nm). Data were analyzed blindly using an analysis program developed for ImageJ and MATLAB (Watanabe et al., 2013). Active zones (AZs) were defined by a corresponding postsynaptic density attached to the presynaptic bouton and docked vesicles were defined as those in direct contact with the plasma membrane. Synaptic vesicle distribution was analyzed by calculating the shortest distance from each vesicle to the presynaptic active zone.

Statistical analysis. Data are presented as means \pm SEM. For statistical analysis, data from two groups were subjected to Student's two-tailed *t* test and data from more than two groups were subjected to one-way ANOVA followed by Tukey's *post hoc* test with GraphPad Prism 5.0b. *p*-values >0.05 were considered insignificant.

Results

Deletion of both Stx1 isoforms Stx1A and Stx1B is prenatally lethal

The neuronal t-SNARE protein Stx1 is essential for neuronal maintenance and neurotransmitter release in worms and flies (Schulze et al., 1995; Schulze and Bellen, 1996; Burgess et al., 1997); however, dissecting these functions in mice has been unfeasible due to the functional redundancy of Stx1 isoforms Stx1A and Stx1B (Fujiwara et al., 2006; Gerber et al., 2008; Wu et al., 2015). To eliminate the reciprocal compensatory effects of Stx1A and Stx1B and to thereby study the essential functions of Stx1, we established an Stx1A/1B DKO mouse line by interbreeding heterozygous Stx1B Het (+/–) mice with homozygous Stx1A KO (–/–) mice, which were generated as described previously (Gerber et al., 2008; Wu et al., 2015). Because Stx1A KO mice reportedly display normal development, life span, and neurotransmitter release (Fujiwara et al., 2006; Gerber et al., 2008; Arancillo et al., 2013), all the mice used in this study were held on Stx1A-null background.

Similar to a previous report (Mishima et al., 2014), deletion of both Stx1 isoforms resulted in embryonic lethality. In addition, we observed gross morphological abnormalities at E16–E18 (Fig. 1A), at which state the Stx1A/1B DKO embryos were already distinguishable by a hunched back and distorted body posture visible to the naked eye (Fig. 1A), a phenotype shared by the other SNARE KO and Munc13-1/2 DKO mutants (Schoch et al., 2001; Washbourne et al., 2002; Varoqueaux et al., 2005). Loss of Stx1B was confirmed by Western blot analysis of whole-brain lysates derived from E18 mice (Fig. 1B). To assess the impact of Stx1 deletion on brain assembly, we examined major brain structures by Nissl staining of the brain tissues of Stx1A KO, Stx1B Het/Stx1A KO, and Stx1A/1B DKO embryos dissected at E16 and E18 (Fig. 1C). Based on the observation of no significant differences between Nissl-stained brain tissues from Stx1A KO and Stx1B Het/Stx1A KO, data from these groups were pooled as control. Complete deletion of Stx1 led to partial cell loss in the cerebellum and brainstem at E16 (Fig. 1C). At E18, cell loss in the brainstem and the cerebellum was even more substantial and became visible in the cerebral cortex and hypothalamus (Fig. 1C), indicating that the regular brain structures are first formed but lost at later embryonic stages: a similar phenotype to Munc18-1 KO mice (Ver-

hage et al., 2000). These findings indicate that Stx1 function is essential during embryonic brain development.

Being high-affinity interaction partners (Hata et al., 1993), Munc18-1 and Stx1 regulate one another's expression levels either by serving as chaperones (Rowe et al., 1999, 2001), by assisting the proper localization, or by enhancing the stability of one another (Toonen et al., 2005). It has been shown that Munc18-1 deletion results in an ~70% reduction in both Stx1A and Stx1B levels (Toonen et al., 2005), whereas a ~25% reduction in Munc18-1 protein level is observed either when Stx1B is deleted (Mishima et al., 2014; Wu et al., 2015) or expressed at severely low levels (Gerber et al., 2008; Arancillo et al., 2013; Zhou et al., 2013). Conversely, deletion of Stx1A alone has no effect on Munc18-1 expression, most likely due to the compensatory action of Stx1B (Fujiwara et al., 2006; Mishima et al., 2014). To address whether the deletion of both isoforms further enhances the decrease in Munc18-1 levels, we performed Western blot analysis using whole-brain lysates derived from E18 mice. Munc18-1 expression was ~40% reduced in the Stx1A/1B DKO embryos compared with the Stx1A KO (Fig. 1D). Moreover, unlike in Stx1A, Stx1B, or Munc18-1 KO mice (Verhage et al., 2000; Toonen et al., 2005; Fujiwara et al., 2006; Wu et al., 2015), we observed significant reductions from ~20% to 60% in expression levels of other tested exocytotic proteins SNAP-25, Syb2, synaptotagmin-1 (Syt1), Munc13-1, and complexin-1 (Cpx1), as well as the early endosomal protein Rab5, in Stx1A/1B DKO embryos (Fig. 1D), suggesting that widespread cell loss and/or synaptic loss in the embryonic mouse brains results in reduced levels of synaptic proteins or that the removal of Stx1 downregulates the expression of other proteins involved in secretion.

Stx1 is required for neuronal maintenance *in vitro*

To investigate whether Stx1A/1B DKO neurons can survive *in vitro*, dissociated cortical neurons from E16 mice were cultured at high density (see Materials and Methods). The neurons were cocultured with astrocyte feeder layers. The astrocytes were all derived from WT newborn mice to provide comparable trophic support to both Stx1A/1B DKO and control neurons. At DIV 1, ~60% of control Stx1A KO neurons initially plated were present in the culture, the cell density remained largely unchanged for 3 weeks (Fig. 2A), and the neurons appeared healthy overall (Fig. 2C). On the contrary, the number of surviving Stx1A/1B DKO neurons dropped to 17% within 1 d and reached ~1% at DIV 4, suggesting that Stx1 is also required for neuronal maintenance *in vitro* (Fig. 2A). No Stx1A/1B DKO neurons survived at DIV 22 (Fig. 2A). Even though deletion of Stx1 resulted in severe cell loss, the remaining neurons at DIV 2 and 4 could form dendritic processes as reflected by positive immunofluorescence labeling for MAP2 (Fig. 2B), which indicates that Stx1A/1B DKO neurons could still differentiate. However, the intensity of MAP2 signal and the complexity of dendritic branching appeared reduced in Stx1A/1B DKO neurons compared with the Stx1A KO neurons (Fig. 2B).

To restore the neuronal viability *in vitro*, we reintroduced either Stx1A or Stx1B expression in Stx1A/1B DKO neurons by infecting with lentiviruses immediately after culturing (DIV 0). The control Stx1A KO group was infected with lentiviruses to induce expression of only NLS-GFP to monitor the effect of virus load on the viability of neurons. Exogenous expression of Stx1A or Stx1B rescued the Stx1A/1B DKO neurons against cell death by 17% and 11%, respectively, compared with the Stx1A KO neurons (Fig. 2C), with the low rescue efficiency reflecting the short time window of the survival of Stx1A/1B DKO neurons. Immu-




Figure 1. Deletion of both Stx1 isoforms Stx1A and Stx1B is prenatally lethal. **A**, Stx1A/1B DKO mice (right) show gross morphological abnormalities at E18 compared with Stx1A KO (left) and Stx1B Het/Stx1A KO (middle). Scale bar, 1 cm. **B**, Example immunoblot showing Stx1A and Stx1B protein levels (left) and summary graph of quantification of Stx1B protein levels (right) in the whole-brain lysates obtained from E18 control and Stx1A/1B DKO mice. $n = 6$ animals for each genotype. **C**, Example Nissl-stained sagittal brain sections from control (left) and Stx1A/1B DKO (right) mice at E16 (top) and E18 (bottom). Arrows show the brain structures with excessive cell loss. Bs, Brainstem; C, cortex; Ce, cerebellum; HT, hypothalamus; M, midbrain; T, thalamus. Scale bar, 1 mm. $n = 3$ animals for each genotype. **D**, Example immunoblot (left) and summary graph of quantification (right) of Munc18-1, SNAP-25, Syb-2, Munc13, synaptotagmin-1 (Syt-1), complexin-2 (Cpx2), Cpx1, and Rab5 protein levels in the whole-brain lysates obtained from E18 mice. $n = 5$ animals for each genotype. Bar graphs show means \pm SEM. Statistical analysis was applied by one-way ANOVA followed by Tukey's *post hoc* test in **B** and Student's *t* test in **D**; * $p \leq 0.05$, ** $p \leq 0.01$; *** $p \leq 0.001$

nofluorescence analysis of the exogenously expressed Stx1A or Stx1B protein levels, as normalized to the immunofluorescence signal of presynaptic active zone protein Bassoon, revealed similar or slightly elevated expression levels of Stx1A or Stx1B compared with controls (Fig. 2D). The rapid death caused by loss of Stx1 and the revival of a fraction of Stx1A/1B DKO neurons by exogenous expression of Stx1A or Stx1B signify the absolute requirement of Stx1, whereas the similar rescue efficiencies of Stx1A and Stx1B manifest the functional redundancy between Stx1A and Stx1B in terms of neuronal maintenance.

Loss of Stx1 leads to an earlier and more severe neuronal lethality than loss of Munc18-1

The severe cytotoxicity caused by loss of Stx1 is reminiscent of Munc18-1 deficiency (Verhage et al., 2000; Heeroma et al., 2004). However, loss of Munc18-1 KO neurons could be delayed up to DIV 10, when trophic support was provided (Heeroma et al., 2004), suggesting that loss of Stx1 might be more deleterious than loss of Munc18-1. To compare the severity of cell loss in Stx1A/1B DKO and Munc18-1 KO neurons directly in the same culture conditions, we obtained cortical cultures from E16 Munc18-1 WT and KO mouse brains. Similar to the Stx1A KO neurons (Fig. 2A), ~55% of WT neurons initially plated were present in the culture at DIV 1 and the number remained unchanged for the next 3 weeks (Fig. 3A). Conversely, at DIV 1, ~40% of Munc18-1 KO neurons were still alive and the remaining Munc18-1 KO neurons were gradually lost within 10 d (Fig. 3A). In addition, exogenous expression of Munc18-1 in Munc18-1 KO neurons showed an ~70% rescue efficiency in neuronal survival (Fig. 3B). The slower rate of cell loss and the higher rescue capacity of lentiviral expression of Munc18-1 suggest that loss of Munc18-1 is less toxic than loss of Stx1 in neurons.

The similarity between the Stx1A/1B DKO and Munc18-1 KO phenotypes demonstrates the importance of these proteins for neuronal maintenance, most likely as a complex. However, severe reduction in Stx1 expression and/or its destabilization in the absence of Munc18-1 (Toonen et al., 2005) raises the possibility that the lethality of Munc18-1 KO neurons might be a direct consequence of low Stx1 levels. Similarly, impairment in Munc18-1 expression might be the cause of lethality of the Stx1A/1B DKO neurons.




Figure 2. Stx1 is required for neuronal maintenance *in vitro*. **A**, Example images of cultured Stx1A KO and Stx1A/1B DKO cortical neurons at DIV 1 (left) and summary graph of percentage of surviving neurons in the cultures at DIV 1, 2, 4, and 22 (right). $n = 58 - 60$ images at each time point from three independent cultures. Scale bar, $50 \mu\text{m}$. **B**, Example images of immunofluorescence labeling for MAP2 in cultured Stx1A KO and Stx1A/1B DKO cortical neurons at DIV 2 (top) and DIV 4 (bottom). Scale bar, $20 \mu\text{m}$. **C**, Example images (left) and summary graph of percentage (right) of surviving neurons at DIV 22 in the Stx1A KO cultures infected with viruses expressing only NLS-GFP and in the Stx1A/1B DKO cultures infected with viruses expressing either Stx1A or Stx1B. $n = 58 - 60$ images from three independent cultures. **D**, Example images (left) and summary graphs (right) of immunofluorescence labeling for Stx1A (blue) and Stx1B (green) in control and rescued neurons at DIV 22. The average intensities of Stx1A or Stx1B immunofluorescence signals were analyzed at compartments positive for Bassoon immunofluorescence signal (red). Stx1A KO neurons infected with viruses to express only NLS-GFP or Stx1B were used as controls (dashed lines). $n = 42 - 49$ images from three independent cultures. Data points in **A** and bar graphs in **C** and **D** show means \pm SEM. Statistical analysis was applied by Student's *t* test in **A** and by one-way ANOVA followed by Tukey's *post hoc* test in **C** and **D**, $***p \leq 0.001$.

To test whether the decreased Stx1 expression level is the primary factor for the death of Munc18-1 KO neurons and vice versa, we used a cross-rescue approach. For this purpose, we overexpressed either Stx1A or Stx1B in Munc18-1 KO neurons and Munc18-1 in Stx1A/1B DKO neurons *in vitro*. Exogenous expression of Munc18-1 displayed no significant effects on the survival of Stx1A/1B DKO neurons at DIV 4 (Fig. 3C), indicating that increasing Munc18-1 protein levels is not sufficient to rescue Stx1A/1B DKO neurons against cell death. Conversely, expression of either Stx1A or Stx1B in Munc18-1 KO neurons resulted in prolonged survival (Fig. 3D,E). The number of surviving Munc18-1 KO neurons increased from $\sim 5\%$ to $\sim 15\%$ at DIV 6 and from $\sim 0.5\%$ to $\sim 3\%$ at DIV 10, when Stx1A was overexpressed. Interestingly, the rescue efficiency of Stx1B in Munc18-1 KO neurons was significantly higher than the rescue

efficiency of Stx1A. At DIV 6, $\sim 25\%$ of Munc18-1 KO neurons were still present in the culture and this number was reduced only to $\sim 10\%$ at DIV 10, when Stx1B was overexpressed (Fig. 3D,E). We also noticed that loss of Munc18-1 led to a more severe reduction in Stx1B signal than in Stx1A signal as revealed by Western blot analysis of whole-brain lysates derived from E16 Munc18-1 WT and KO animals (Fig. 3F). This result, together with the observation of a better rescue efficiency of Stx1B in Munc18-1 KO neurons, may suggest that Munc18-1 interacts more strongly with Stx1B than with Stx1A. In addition, although it is clear that both Munc18-1 and Stx1 are absolutely required for cell survival, the more cytotoxic effect of loss of Stx1 than loss of Munc18-1 indicates that Stx1, but not Munc18-1, is absolutely essential for neuronal survival. Alternatively, both Stx1 and Munc18-1 have equally important functions in neuronal sur-




Figure 3. Loss of Stx1 leads to an earlier and more severe neuronal lethality than loss of Munc18-1. **A**, Example images of cultured WT and Munc18-1 KO cortical neurons at DIV 1 (left) and summary graph of percentage of surviving neurons in the cultures at DIV 1–10 and at DIV 22 (right). $n = 58–60$ images from three independent cultures. **B**, Example images (left) and summary graph of percentage (right) of surviving neurons at DIV 22 in the WT culture infected with viruses to express only NLS-GFP and in the Munc18-1 KO cultures infected with viruses to express Munc18-1. $n = 60$ images from three independent cultures. **C**, Summary graph of percentage of surviving Stx1A/1B DKO neurons at DIV 4 infected with viruses to express only NLS-GFP or Munc18-1. **D**, Example images of surviving Munc18-1 KO neurons at DIV 6 infected with viruses to express only NLS-GFP, Stx1A, or Stx1B. **E**, Summary graphs of percentage of Munc18-1 KO neurons expressing only NLS-GFP, Stx1A, or Stx1B at DIV 6 and 10. $n = 60$ images from three independent cultures. **F**, Example immunoblot (left) and summary graph (right) showing Stx1A and Stx1B protein levels in whole-brain lysates obtained from E16 WT and Munc18-1 KO animals. $n = 5$ animals for each genotype. Scale bars in **A**, **B**, and **D**, 50 μm . Data points in **A** and bar graphs in **B**, **C**, **E**, and **F** show means \pm SEM. Statistical analysis was applied by Student's *t* test in **A–C** and **F** and by one-way ANOVA followed by Tukey's *post hoc* test in **E**; * $p \leq 0.05$; ** $p \leq 0.01$; *** $p \leq 0.001$.

vival, but another Munc18 isoform, such as Munc18–2, which has been shown to bind to Stx1 (Hata and Südhof, 1995; Rowe et al., 2001; Gulyás-Kovács et al., 2007), partially supports cell viability.

Stx1 is essential for neuronal maintenance, not only at early developmental stages, but also after maturation *in vitro*

The immature death of Stx1A/1B DKO neurons not only restrains the investigation of the Stx1 requirement for neuronal maintenance to the developing neurons, but also renders the studies of the impact of Stx1 loss on synaptic transmission 17

impracticable. Is the Stx1 requirement for neuronal maintenance confined to development or do neurons perpetually need Stx1 function for viability also after maturation? By using a Cre/LoxP recombination system, we generated an Stx1A/1B conditional DKO (cDKO) mouse line in which the Stx1B gene was flanked by two LoxP sites on an Stx1A-null background (Stx1B^{FL/FL}/Stx1A KO) and thereby bypassed the premature death of Stx1-deficient neurons. High-density hippocampal cultures were prepared from newborn homozygous Stx1A/1B cDKO mice and Stx1B was removed *in vitro* by Cre recombinase




Figure 4. Stx1 is essential for neuronal maintenance after maturation *in vitro*. **A**, Example images (left) and summary graph of percentage (right) of surviving cultured hippocampal neurons derived from P0–P2 Stx1B^{FL/FL}/Stx1A KO animals. Stx1A/1B cDKO neurons were generated by infection at DIV 1 with viruses to express Cre recombinase. Cultures infected with viruses to express only NLS-RFP were used as a control. $n = 40$ images from two independent cultures for DIV 11, 15, 17, 19, 21, and 25. $n = 59–62$ from three independent cultures for DIV 7, 9, 13, and 23. **B**, Example images of Stx1A/1B cDKO cultures generated by infection at DIV 3 with viruses to express Cre recombinase (left) and summary graph of percentage of surviving control and Stx1A/1B cDKO neurons infected at DIV 3, 5, and 7 with viruses to express Cre recombinase (right). $n = 39–40$ images from two independent cultures for DIV 11, 15, 19, 27, 33, 39, and 42 and $n = 58–60$ from three independent cultures for DIV 7 and 23. Scale bar, 50 μ m. **C**, Example images of control and Stx1A/1B cDKO cultures at DIV 42. Scale bars, 50 μ m. Data points show means \pm SEM. Statistical analysis was applied by Student's *t* test; * $p \leq 0.05$; ** $p \leq 0.01$; *** $p \leq 0.001$.

expression induced by lentiviral infection at DIV 1. The control neurons were of the same genotype (Stx1B^{FL/FL}/Stx1A KO), but were transduced with viruses expressing only the reporter construct NLS-RFP. The number of surviving neurons was determined between DIV 7 and 25 and normalized to the number of neurons in the control group at DIV 7. The number of the control neurons and the number of the Stx1A/1B cDKO neurons infected with viruses at DIV 1 to express Cre recombinase were comparable between DIV 7 and 11 (Fig. 4A). However, we observed a gradual reduction of viable Stx1A/1B cDKO neurons starting at DIV 13 and the neuronal viability reached 50% at DIV 15, whereas the number of control neurons did not show any significant differences between DIV 7 and 25 (Fig. 4A), showing Stx1 dependency for neuronal maintenance after development.

To further assess the impact of Stx1 loss on neuronal maintenance at later stages after neuronal maturation is accomplished, Cre recombinase expression was induced by infection at DIV 3, 5, and 7, and subsequently the neuronal viability was monitored up to DIV 42. The delayed induction of Cre recombinase expression and thus

the delayed Stx1B turnover ultimately led again to neuronal lethality but with a delayed onset. The viability declined to 50% at DIV 23, 33, and 39 for neurons that were transduced with Cre recombinase at DIV 3, 5, and 7, respectively (Fig. 4B). The time between genetic removal of Stx1 and cell loss was in the range of 2–3 weeks (Fig. 4A, B), suggestive of a slow Stx1 turnover and/or that only a severe drop in Stx1 levels leads to cell death, which is consistent with the findings from previous Stx1 hypomorphic models (Arancillo et al., 2013). The severity of the structural defects at DIV 42 appeared more substantial in the groups infected with Cre recombinase at earlier time points (Fig. 4C). Altogether, these data demonstrate that Stx1 is essential for neuronal maintenance, not only at early developmental stages, but throughout the lifespan of a neuron.

Stx1 is required for vesicle priming and fusion

The *in vivo* and *in vitro* lethality observed in Stx1-deficient neurons demonstrate that Stx1 is essential for neuronal maintenance during development and after maturation; however, it also obscures the investigation of Stx1's functions in neurotransmitter




Figure 5. Stx1 is required for vesicle priming and fusion. **A**, Example images (top left and bottom) and summary graph (top right) of immunofluorescence labeling for Stx1B (green) in control and in Stx1A/1B cDKO neurons infected at DIV 3 or 5 with viruses to express Cre recombinase. The average intensities of Stx1B immunofluorescence signals were measured at compartments positive for Bassoon immunofluorescence signal (red) and the values were normalized to the values obtained from control neurons (dashed lines). $n = 24–41$ images from two independent cultures. **B**, Example traces (left) and summary graph (right) of EPSCs evoked by a 2 ms depolarization in autaptic neurons derived from P0–P2 Stx1A KO and Stx1A/1B cDKO mice. The neurons were infected at DIV 3 with viruses to express Cre recombinase. **C**, Example traces (left) and summary graph (right) of readily releasable pool in Stx1A KO and Stx1A/1B cDKO neurons determined by 500 mM sucrose application for 5 s. **D**, Example traces (left) and summary graph (right) of mEPSCs in Stx1A KO and Stx1A/1B cDKO neurons recorded at holding potential -70 mV. Bar graphs show means \pm SEM. Statistical analysis was applied by one-way ANOVA followed by Tukey's post hoc test; $*p \leq 0.05$, $**p \leq 0.01$, $***p \leq 0.001$. Cell numbers and the number of independent cultures are presented in the graphs in **B–D**. Artifacts and/or action potentials are blanked in **B** and **C**. The traces were filtered at 1 kHz in **D**.

release such that the impacts of Stx1 deficiency on neurotransmission in mammalian neurons could be determined only by severe reduction of Stx1B expression on a Stx1A-null background (Arancillo et al., 2013; Zhou et al., 2013). To find a time window for electrophysiological examination at which the neurons are mature, yet intact and completely devoid of Stx1, we monitored Stx1B protein levels in Stx1A/1B cDKO neurons, which were infected at DIV 3 or 5 with viruses to express Cre recombinase and fixed at DIV 12 and 21 (Fig. 5A). Immunofluorescence labeling in both Stx1A/1B cDKO groups expressing Cre recombinase revealed Stx1B signals at a level of 15–30% compared with the control neurons at DIV 12, whereas at DIV 21, no further reduction in Stx1B signal was observed in Stx1A/1B cDKO neurons infected at DIV 3 (Fig. 5A). This suggests that the values obtained in this group for Stx1B signal at DIV 12 represent background (Fig. 5A). We therefore chose infection at DIV 3 to express Cre recombinase in Stx1A/1B cDKO neurons to eliminate Stx1B because, at DIV 12, these neurons were still alive, appeared intact, and yet Stx1 was completely absent (Figs. 4B, 5A).

To evaluate the impact of loss of Stx1 on neurotransmitter release, we recorded Ca^{2+} -evoked release, spontaneous release, and hypertonic sucrose solution-evoked release from autaptic hippocampal cultures of Stx1A/1B cDKO neurons compared with WT-like Stx1A KO neurons. The control neurons, namely the Stx1A KO neurons expressing either the reporter NLS-RFP or Cre recombinase, as well as Stx1A/1B cDKO neurons expressing only NLS-RFP, produced robust and comparable EPSCs with an average peak amplitude of ~ 4 nA (Fig. 5B). Conversely, loss of both Stx1 isoforms by expression of Cre recombinase in Stx1A/1B cDKO neurons essentially abolished EPSCs (Fig. 5B). 19

Among 27 Stx1A/1B cDKO neurons, 22 of them exhibited no evoked transmitter release and the remaining five had an average EPSC of 0.21 ± 0.06 nA. Furthermore, measurements of the RRP of vesicles evoked by 5 s application of 500 mM hypertonic solution revealed a complete absence of fusion-competent vesicles in Stx1A/1B cDKO neurons expressing Cre recombinase, whereas the control groups exhibited RRP of an average size of ~ 0.4 nC (Fig. 5C). Similarly, spontaneous release was also comparable among the control groups, as revealed by comparable mEPSC frequencies, whereas no mEPSC events were detected in Stx1A/1B cDKO neurons expressing Cre recombinase (Fig. 5D). These data indicate that the neuronal t-SNARE protein Stx1 is absolutely required for vesicle priming and fusion.

Stx1's function in neuronal survival is independent of its function in neurotransmission

Having completely eliminated Stx1 expression, we wanted to test whether the function of Stx1 in neuronal survival and in synaptic transmission are coupled. Whereas the observation of the degeneration of synaptically silent mutant Munc18-1 KO neurons led to the hypothesis that synaptic activity is required for neuronal maintenance (Verhage et al., 2000), unimpaired survival of Munc13-1/2 DKO provided counterevidence for this hypothesis (Varoqueaux et al., 2002). Furthermore, a Stx1 mutant with 2 point mutations interchanging the +4 and +5 hydrophobic layers of the SNARE motif (A240V, V244A; Stx1A^{AV}) is embryonically lethal in flies despite maintaining neurotransmitter release albeit at severely reduced levels (Fergestad et al., 2001). The degeneration phenotype of this mutant has been also shown by its inability to rescue survival of rat neurons subjected to botulinum




Figure 6. Stx1 function in neuronal survival is independent from its function in neurotransmission. **A**, Example images (left) and summary graph (right) of immunofluorescence labeling for Stx1A (blue) in Stx1A/1B cDKO neurons coinfecting at DIV 3 with viruses to express only the reporter constructs or Cre recombinase and Stx1A^{WT} or Stx1A^{AV} and fixed at DIV 14. The average intensities of Stx1A immunofluorescence signals (blue) were measured at compartments positive for Bassoon immunofluorescence signal (red). $n = 52$ – 55 images from three independent cultures. Scale bar, $10 \mu\text{m}$. **B**, Example traces (left) and summary graph (right) of EPSCs. **C**, Example traces (left) and summary graph (right) of RRP. **D**, Example traces (left) and summary graph (right) of mEPSCs. **E**, Example images of Stx1A/1B cDKO cultures at DIV 24 (left) and summary graph of neuronal survival determined as the percentage of cell number at DIV 24/DIV 11 (right). Scale bar, $50 \mu\text{m}$. $n = 60$ images from three independent cultures. Bar graphs show means \pm SEM; $*p \leq 0.05$, $***p \leq 0.001$. Cell numbers and the number of independent cultures are presented in the graphs in **B–D**. Artifacts and/or action potentials are blanked in **B** and **C**. The traces were filtered at 1 kHz in **D**.

neurotoxin C (Peng et al., 2013). With the purpose of finding a Stx1 mutant, which maintains neurotransmitter release, albeit severely impaired, but which is ineffective for neuronal maintenance and thereby to split the functions of Stx1 in neurotransmitter release and neuronal maintenance, we coexpressed Stx1A^{AV} with Cre recombinase in Stx1A/1B cDKO neurons. For that purpose, we first examined the exogenous expression efficiencies of Stx1A^{AV} and WT Stx1A (Stx1A^{WT}) in Stx1A/1B cDKO neurons expressing Cre recombinase. Immunofluorescence labeling of Stx1A^{AV} revealed an $\sim 20\%$ higher signal on average compared with the Stx1A^{WT} at compartments positive for Bassoon signal, indicating that the exogenously expressed Stx1A^{AV} can localize at synapses (Fig. 6A).

We next tested whether neurotransmitter release can be rescued by Stx1A^{AV} compared with Stx1A^{WT}. Expression of Stx1A^{WT} in Stx1A/1B cDKO neurons successfully restored evoked and spontaneous neurotransmitter release, as well as 20

the RRP, back to the levels of control neurons, which includes only endogenous Stx1B, demonstrating again a functional redundancy between Stx1A and Stx1B in terms of neurotransmitter release (Fig. 6B–D). Conversely, Stx1A^{AV} completely failed to rescue evoked and spontaneous release, nor could it restore the RRP in Stx1A/1B cDKO neurons (Fig. 6B–D). Given that Stx1A^{AV} assembles into SNARE complexes, but with reduced stability (Kee et al., 1995; Fergestad et al., 2001), and that, in Stx1A^{AV}, the mutated hydrophobic layers are closer to the C terminus, these data suggest that incomplete SNARE complex formation is not sufficient for vesicle fusion or for vesicle priming.

We therefore wanted to address whether Stx1A^{AV} and, indirectly, whether partial SNARE complex assembly could rescue neuronal survival despite being release incompetent. We induced coexpression of Cre recombinase with Stx1A^{AV} or with Stx1A^{WT} by infecting Stx1A/1B cDKO neurons with viruses at DIV 1. The neurons were then counted at DIV 11 and DIV 24 and the ratio of




Figure 7. Stx1 functions in vesicle docking and fusion can be uncoupled. **A**, Example electron microscopy images of presynaptic nerve terminals from Stx1A/1B cDKO neurons coinfecting at DIV2 with viruses to express the reporter constructs, Cre recombinase and only a reporter construct, and Cre recombinase and Stx1A^{WT} or Stx1A^{AV}. Arrows show the docked vesicles in the zoomed images. Scale bars, 100 nm (top) and 50 nm (bottom). **B**, Summary graphs of AZ length, docked synaptic vesicles (SVs) per section, and SVs within 30 nm of AZs. Bar graphs show means \pm SEM. Statistical analysis was applied by one-way ANOVA followed by Tukey's *post hoc* test; * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$. The number of sections analyzed and the number of independent cultures are presented in the graphs.

the cell numbers at DIV 24/DIV 11 was used as a read-out for neuronal cell survival. Stx1A^{WT} successfully rescued the Stx1A/1B cDKO neurons against cell death, showing again functional redundancy with Stx1B. Surprisingly, Stx1A^{AV} also rescued neuronal survival at levels comparable to the control and Stx1A^{WT} rescue (Fig. 6E). The inability of Stx1A^{AV} to rescue botulinum neurotoxin-mediated neuronal death might be due to the additional mutation, which was included to render Stx1A cleavage resistant (Peng et al., 2013). Conversely, the efficiency of Stx1A^{AV} to rescue neuronal survival in our assay, but not neurotransmitter release, argues that the lack of synaptic activity is not a cause of neuronal degeneration and that Stx1 has independent functions in neuronal maintenance and neurotransmission. It may also suggest that partial SNARE complex assembly is sufficient for neuronal maintenance, but not for vesicle fusion and priming.

Stx1 function in vesicle docking and fusion can be uncoupled

In addition to its functions in neuronal maintenance, vesicle fusion, and vesicle priming, Stx1 plays a role in vesicle docking in mammalian neurons (Gerber et al., 2008; Arancillo et al., 2013; Imig et al., 2014). However, a reduction of more than $\sim 70\%$ in Stx1 expression is required for observable impairments in vesicle docking (Arancillo et al., 2013; Imig et al., 2014), whereas unfeasibility of complete elimination of Stx1 has obscured the analysis of vesicle docking in the absence of Stx1. To evaluate the impact of complete loss of Stx1 in vesicle docking, we performed high-pressure freezing experiments of the cultured Stx1A/1B cDKO hippocampal neurons infected with viruses at DIV 2 and fixed at 21

DIV 12–14. We then examined the ultrastructure of the synapses by using electron microscopy. Stx1A/1B cDKO neurons expressing Cre recombinase revealed a $\sim 50\%$ reduction in the number of docked vesicles compared with the control neurons, whereas no significant differences were observed in AZ length and total vesicle number within 30 nm AZ (Fig. 7). Even though our results point toward a further reduction in vesicle docking compared with previous studies, which have analyzed the effects of severely reduced levels of Stx1B on Stx1A-null background in mice (Arancillo et al., 2013; Imig et al., 2014), the impairment in vesicle docking is not as severe as observed in Munc13-1/2 DKO neurons (Imig et al., 2014). However, our analysis in a conditional KO system is hampered by cytotoxic effects of loss of Stx1 and thus biased toward cells resembling severe hypomorphic expression of Stx1. Although it is evident that Stx1 is essential for vesicle docking, we cannot rule out the involvement of residual Stx1B in the manifestation of a remaining vesicle-docking function.

Although our data demonstrate that full SNARE complex assembly is absolutely essential for vesicle fusion and vesicle priming, as manifested by complete inhibition of neurotransmission by Stx1A^{AV} (Fig. 6), the role of SNARE complex assembly in vesicle docking is still not well defined. To test whether vesicle docking can be recovered in Stx1A/1B cDKO neurons by Stx1A^{AV} and thus by putative partial SNARE complex assembly, we coexpressed Stx1A^{WT} or Stx1A^{AV} with Cre recombinase. Expression of Stx1A^{WT} fully restored vesicle docking back to the control levels, providing evidence for functional redundancy between Stx1A and Stx1B also in terms of vesicle docking (Fig. 7). Remarkably, expression of Stx1A^{AV} completely restored docked vesicles to levels comparable to the control neurons, demonstrating a clear functional difference between vesicle docking and priming. More importantly, the ability of Stx1A^{AV} to dock the vesicles successfully to the plasma membrane is suggestive of the sufficiency of partially assembled SNARE complexes for vesicle docking.

Discussion

The fundamental roles of Stx1 in neuronal maintenance and neurotransmission have been well established by genetic manipulation studies in invertebrates (Schulze et al., 1995; Burgess et al., 1997; Saifee et al., 1998). The situation in mammalian neurons is less clear. The redundancy between the two isoforms Stx1A and Stx1B (Fujiwara et al., 2006; Gerber et al., 2008; Arancillo et al., 2013; Zhou et al., 2013; Wu et al., 2015) and the embryonic lethality in Stx1A/1B DKO mice (Mishima et al., 2014) limited the functional studies of Stx1 only to the conditions of severely reduced expression of Stx1 (Arancillo et al., 2013; Zhou et al., 2013). With our Stx1A/1B DKO mouse models, we bypassed these limitations and investigated the impacts of loss of Stx1 on neurotransmission and neuronal maintenance in a mammalian system. We draw several major conclusions. Stx1 has an abso-

lutely essential function in neuronal maintenance *in vivo* and *in vitro*, not only during development, but throughout the neuronal life span. This function is independent of its function in neurotransmission and the particular role of Stx1 in neurotransmission can be broken down into its functions in vesicle docking and vesicle fusion.

Stx1 is absolutely required for neuronal maintenance

Complete loss of Stx1 by deletion of both Stx1A and Stx1B led to embryonic lethality, which is consistent with a recent study (Mishima et al., 2014) and caused widespread cell loss in embryonic mouse brains as well as premature neuronal lethality *in vitro* (Figs. 1, 2). In addition, *in vitro* genetic removal of Stx1 subsequently caused death of differentiated neurons, demonstrating the importance of Stx1 also for the neuronal maintenance after maturation.

An equal severity of neuronal lethality was not observed so far in other vesicular SNARE KO mutants because loss of Syb2 or SNAP-25 shows no signs of neuronal loss *in vivo* (Schoch et al., 2001; Washbourne et al., 2002), with the latter leading to neuronal death only *in vitro* (Delgado-Martínez et al., 2007). Conversely, loss of Munc18-1, a well defined interaction partner of Stx1, produces a phenocopy of loss of Stx1 with severe neuronal lethality (Verhage et al., 2000), suggesting a functional cooperation between Stx1 and Munc18-1 in neuronal maintenance.

What is the function of Stx1 in neuronal maintenance and is it coupled to its function in neurotransmitter release? Previous studies suggest a primary dysfunction in trophic support to be the underlying mechanism of Stx1-mediated cell death (Kofuji et al., 2014); however, a putative glial dysfunction is incompatible with our experiments because we cocultured the Stx1-deficient neurons with WT astrocytes. Furthermore, selective death of Munc18-1-deficient Purkinje cells *in vivo* showed the cell-autonomous nature of the observed cell loss (Heeroma et al., 2004). Nevertheless, we cannot rule out the involvement of dysfunctions in autocrine signaling pathways.

Lack of synaptic input has also been proposed as a possible mechanism leading to neuronal death (Verhage et al., 2000), yet the unimpaired viability of Syb1/2-deficient and Munc13-1/-2-deficient neurons, despite being release incompetent (Varoqueaux et al., 2002; Zimmermann et al., 2014), shows that neuronal maintenance is independent of neurotransmission. Interestingly, in our conditional Stx1 deletion model, neuronal survival was rescued not only by WT Stx1, but also by a Stx1 mutant (A240V, V244A; Stx1A^{AV}) that is fully deficient in neurotransmitter release (Fig. 6). In view of these findings, we suggest that Stx1, in conjunction with Munc18-1, is required for survival, but in a function that is not related to neurotransmitter release, such as in the ER, Golgi, and posttranslational trafficking processes (Rowe et al., 1999, 2001; Toonen et al., 2005; Arunachalam et al., 2008), in nonvesicular lipid trafficking, and/or in the ER positioning at the plasma membrane contact sites (Petkovic et al., 2014). Defects in these systems would in turn lead to neuronal death. Moreover, Stx1 interacts with glycine and GABA transporters (Blakely et al., 1998; López-Corcuera et al., 2001) and modulates N- and P/Q-type Ca²⁺-channels (Lévêque et al., 1994; Sheng et al., 1994) and M-type K⁺-channels (Regev et al., 2009; Etzioni et al., 2011). A shift in the ionic homeostasis, such as excess amounts of intracellular Ca²⁺, could also produce cytotoxic effects (Valentino et al., 1993; Cataldi, 2013).

A direct cooperation of Stx1 and Munc18-1 in neuronal maintenance is not proven, but likely. As costabilizing chaperones

(Rowe et al., 1999, 2001; Toonen et al., 2005), the putative cooperation of Munc18-1 and Stx1 may in principle be based on an upstream–downstream relationship as implied by our cross-rescue experiments (Fig. 3). Differential efficiencies of Stx1A and Stx1B to rescue Munc18-1 KO neurons (Fig. 3D,E) also points toward a correlation between Munc18-1-dependent Stx1 isoform stability and the severity of phenotype. In fact, loss of Stx1B isoforms is more deleterious than loss of Stx1A (Kofuji et al., 2014; Mishima et al., 2014; Wu et al., 2015) and a decrease in Munc18-1 protein levels is observed only as a result of loss or severe reduction of Stx1B (Gerber et al., 2008; Arancillo et al., 2013; Wu et al., 2015), not as a result of loss of Stx1A (Fujiwara et al., 2006; Gerber et al., 2008). However, the ultimate death of either Stx1- or Munc18-1 deficient neurons is incompatible with such an upstream–downstream relationship. Alternatively, a partial redundancy of Munc18-1 with another isoform, such as Munc18–2, could explain the differential cross rescue activities of Stx1 and Munc18-1. Even though Munc18–2 is not expressed at detectable levels in neurons (Hata and Südhof, 1995), it has been shown to be able to bind to Stx1 (Hata and Südhof, 1995; Rowe et al., 2001; Gulyás-Kovács et al., 2007) and to be expressed at elevated levels in Munc18-1 knock-down PC12 cells (Arunachalam et al., 2008), yet the expression pattern of Munc18 isoforms in the developing brain remains to be elucidated.

Differential functions of Stx1 in vesicle docking and fusion

With our Stx1A/1B cDKO mouse model, we identified a time window when endogenous Stx1 was completely lost with no apparent signs of degenerated neuronal structures, providing the suitable conditions for studies of Stx1's functions in neurotransmission. Electrophysiological recordings from Stx1A/1B cDKO hippocampal neurons revealed complete elimination of neurotransmitter release, including spontaneous and synchronous or asynchronous evoked release (Fig. 5). In addition, vesicle priming was completely inhibited as shown by lack of fusion competent vesicles (Fig. 5). However, a recent study with a similar Stx1A/1B DKO mouse model showed residual asynchronous synaptic activity (Mishima et al., 2014), supporting a model in which release can occur in the absence of Stx1. The asynchronous release observed by Mishima et al. (2014) might be due to up-regulation of, or compensation by, another Stx isoform in a subpopulation of embryonic cortical neurons. Indeed, similar issues of redundancies have also been observed for the other two mammalian synaptic SNAREs. Remaining release activity in SNAP-25-deficient neurons (Washbourne et al., 2002) may well be mediated by SNAP-23 (Delgado-Martínez et al., 2007). In addition, Syb2 is partially redundant to Syb1 and the latter is responsible for remaining release activity (Schoch et al., 2001; Zimmermann et al., 2014) and vesicle-docking activity (Imig et al., 2014) in Syb2 KO neurons. Together, our results fully support the canonical role of SNAREs for any form of neurotransmission.

We also used the Stx1A/1B cDKO models for functional reconstitution experiments. Although exogenous expression of WT Stx1A completely restored vesicle fusion and priming in Stx1A/1B cDKO neurons, the Stx1A^{AV} mutant failed to do so. By the position of the mutations A240V, V244A, which lay at the +4 and +5 layers of SNARE motif, it is plausible that Stx1A^{AV} would assemble with the other SNARE proteins into SNARE complexes. In fact, it has been demonstrated by *in vitro* experiments that Stx1A^{AV} forms ternary SNARE complexes with SNAP-25 and Syb-2 (Kee et al., 1995; Fergestad et al., 2001). However, the reduced stability of the SNARE complexes containing Stx1A^{AV} (Kee et al., 1995; Fergestad et al., 2001) would suggest impair-

ment in full assembly, which in turn may explain the inhibition of neurotransmitter release in our study.

Remarkably, Stx1A^{AV} was able to fully rescue vesicle docking to WT levels in Stx1A/1B cDKO neurons, yet the vesicles were fusion incompetent. This differential effect strongly supports two hypotheses: (1) vesicle docking and vesicle priming are two distinct functional steps that can be differentially assessed by morphological and electrophysiological analysis and (2) vesicle docking is less affected by the stability of SNARE complexes than neurotransmitter release. These phenomena can be explained by the sequential zipper mechanism: SNARE complex assembly is mediated by zippering from the N to the C terminus and partially assembled SNARE complexes are arrested before Ca²⁺ entry. In fact, structural analyses using nuclear magnetic resonance or atomic force microscopies depicted a sequential SNARE complex formation proceeding in two or three steps (Ellena et al., 2009; Gao et al., 2012; Liang et al., 2013). In addition, mutational analysis of SNAP-25 and Syb2 in neurons and chromaffin cells revealed the critical role of the layers from +4 to +6 of the SNARE complex in neurotransmitter release, supporting the hypothesis of sequential assembly of the SNARE complex (Sørensen et al., 2006; Walter et al., 2010). Consistent with these observations, we show here that docking and priming/fusion are associated with different SNARE assembly states, in which the docking state does not require an intact layer +4/+5 structure. Overall, this suggests that these residues might reside between two separate functional domains in the SNARE complex, which are separately responsible for vesicle docking and priming.

References

- Arancillo M, Min SW, Gerber S, Münster-Wandowski A, Wu YJ, Herman M, Trimbuch T, Rah JC, Ahnert-Hilger G, Riedel D, Südhof TC, Rosenmund C (2013) Titration of Syntaxin1 in mammalian synapses reveals multiple roles in vesicle docking, priming, and release probability. *J Neurosci* 33:16698–16714. [CrossRef Medline](#)
- Arunachalam L, Han L, Tassew NG, He Y, Wang L, Xie L, Fujita Y, Kwan E, Davletov B, Monnier PP, Gaisano HY, Sugita S (2008) Munc18-1 is critical for plasma membrane localization of syntaxin1 but not of SNAP-25 in PC12 cells. *Mol Biol Cell* 19:722–734. [Medline](#)
- Bennett MK, Calakos N, Scheller RH (1992) Syntaxin: a synaptic protein implicated in docking of synaptic vesicles at presynaptic active zones. *Science* 257:255–259. [CrossRef Medline](#)
- Blakely RD, Ramamoorthy S, Schroeter S, Qian Y, Apparsundaram S, Galli A, DeFelice LJ (1998) Regulated phosphorylation and trafficking of antidepressant-sensitive serotonin transporter proteins. *Biol Psychiatry* 44:169–178. [CrossRef Medline](#)
- Burgess RW, Deitcher DL, Schwarz TL (1997) The synaptic protein syntaxin1 is required for cellularization of *Drosophila* embryos. *J Cell Biol* 138:861–875. [CrossRef Medline](#)
- Cataldi M (2013) The changing landscape of voltage-gated calcium channels in neurovascular disorders and in neurodegenerative diseases. *Curr Neuropharmacol* 11:276–297. [CrossRef Medline](#)
- Delgado-Martínez I, Nehring RB, Sørensen JB (2007) Differential abilities of SNAP-25 homologs to support neuronal function. *J Neurosci* 27:9380–9391. [CrossRef Medline](#)
- Ellena JF, Liang B, Wiktor M, Stein A, Cafiso DS, Jahn R, Tamm LK (2009) Dynamic structure of lipid-bound synaptobrevin suggests a nucleation-propagation mechanism for trans-SNARE complex formation. *Proc Natl Acad Sci U S A* 106:20306–20311. [CrossRef Medline](#)
- Etzioni A, Siloni S, Chikvashvili D, Strulovich R, Satchani D, Regev N, Greitzer-Antes D, Hirsch JA, Lotan I (2011) Regulation of neuronal M-channel gating in an isoform-specific manner: functional interplay between calmodulin and syntaxin 1A. *J Neurosci* 31:14158–14171. [CrossRef Medline](#)
- Fergestad T, Wu MN, Schulze KL, Lloyd TE, Bellen HJ, Broadie K (2001) Targeted mutations in the syntaxin H3 domain specifically disrupt SNARE complex function in synaptic transmission. *J Neurosci* 21:9142–9150. [Medline](#)
- Fujiwara T, Mishima T, Kofuji T, Chiba T, Tanaka K, Yamamoto A, Akagawa K (2006) Analysis of knock-out mice to determine the role of HPC-1/syntaxin 1A in expressing synaptic plasticity. *J Neurosci* 26:5767–5776. [CrossRef Medline](#)
- Gao Y, Zorman S, Gundersen G, Xi Z, Ma L, Sirinakis G, Rothman JE, Zhang Y (2012) Single reconstituted neuronal SNARE complexes zipper in three distinct stages. *Science* 337:1340–1343. [CrossRef Medline](#)
- Gerber SH, Rah JC, Min SW, Liu X, de Wit H, Dulubova I, Meyer AC, Rizo J, Arancillo M, Hammer RE, Verhage M, Rosenmund C, Südhof TC (2008) Conformational switch of syntaxin-1 controls synaptic vesicle fusion. *Science* 321:1507–1510. [CrossRef Medline](#)
- Gulyás-Kovács A, de Wit H, Milosevic I, Kochubey O, Toonen R, Klingauf J, Verhage M, Sørensen JB (2007) Munc18-1: sequential interactions with the fusion machinery stimulate vesicle docking and priming. *J Neurosci* 27:8676–8686. [CrossRef Medline](#)
- Harrison SD, Broadie K, van de Goor J, Rubin GM (1994) Mutations in the *Drosophila* Rop gene suggest a function in general secretion and synaptic transmission. *Neuron* 13:555–566. [CrossRef Medline](#)
- Hata Y, Südhof TC (1995) A novel ubiquitous form of Munc-18 interacts with multiple syntaxins. Use of the yeast two-hybrid system to study interactions between proteins involved in membrane traffic. *J Biol Chem* 270:13022–13028. [CrossRef Medline](#)
- Hata Y, Slaughter CA, Südhof TC (1993) Synaptic vesicle fusion complex contains unc-18 homologue bound to syntaxin. *Nature* 366:347–351. [CrossRef Medline](#)
- Heeroma JH, Roelandse M, Wierda K, van Aerde KI, Toonen RF, Hensbroek RA, Brussaard A, Matus A, Verhage M (2004) Trophic support delays but does not prevent cell-intrinsic degeneration of neurons deficient for Munc18-1. *Eur J Neurosci* 20:623–634. [CrossRef Medline](#)
- Imig C, Min SW, Krinner S, Arancillo M, Rosenmund C, Südhof TC, Rhee J, Brose N, Cooper BH (2014) The morphological and molecular nature of synaptic vesicle priming at presynaptic active zones. *Neuron* 84:416–431. [CrossRef Medline](#)
- Kee Y, Lin RC, Hsu SC, Scheller RH (1995) Distinct domains of syntaxin are required for synaptic vesicle fusion complex formation and dissociation. *Neuron* 14:991–998. [CrossRef Medline](#)
- Kim JH, Lee SR, Li LH, Park HJ, Park JH, Lee KY, Kim MK, Shin BA, Choi SY (2011) High cleavage efficiency of a 2A peptide derived from porcine teschovirus-1 in human cell lines, zebrafish and mice. *PLoS One* 6:e18556. [CrossRef Medline](#)
- Kofuji T, Fujiwara T, Sanada M, Mishima T, Akagawa K (2014) HPC-1/syntaxin 1A and syntaxin 1B play distinct roles in neuronal survival. *J Neurochem* 130:514–525. [CrossRef Medline](#)
- Lévêque C, el Far O, Martin-Moutot N, Sato K, Kato R, Takahashi M, Seagar MJ (1994) Purification of the N-type calcium channel associated with syntaxin and synaptotagmin. A complex implicated in synaptic vesicle exocytosis. *J Biol Chem* 269:6306–6312. [Medline](#)
- Liang B, Kiessling V, Tamm LK (2013) Prefusion structure of syntaxin-1A suggests pathway for folding into neuronal trans-SNARE complex fusion intermediate. *Proc Natl Acad Sci U S A* 110:19384–19389. [CrossRef Medline](#)
- Lois C, Hong EJ, Pease S, Brown EJ, Baltimore D (2002) Germline transmission and tissue-specific expression of transgenes delivered by lentiviral vectors. *Science* 295:868–872. [CrossRef Medline](#)
- López-Corcuera B, Aragón C, Geerlings A (2001) Regulation of glycine transporters. *Biochem Soc Trans* 29:742–745. [CrossRef Medline](#)
- Mishima T, Fujiwara T, Sanada M, Kofuji T, Kanai-Azuma M, Akagawa K (2014) Syntaxin 1B, but not syntaxin 1A, is necessary for the regulation of synaptic vesicle exocytosis and of the readily releasable pool at central synapses. *PLoS One* 9:e90004. [CrossRef Medline](#)
- Peng L, Liu H, Ruan H, Tepp WH, Stoothoff WH, Brown RH, Johnson EA, Yao WD, Zhang SC, Dong M (2013) Cytotoxicity of botulinum neurotoxins reveals a direct role of syntaxin 1 and SNAP-25 in neuron survival. *Nat Commun* 4:1472. [CrossRef Medline](#)
- Petkovic M, Jemaiel A, Daste F, Specht CG, Izeddin I, Vorkel D, Verbavatz JM, Darzacq X, Triller A, Pfenninger KH, Tareste D, Jackson CL, Galli T (2014) The SNARE Sec22b has a non-fusogenic function in plasma membrane expansion. *Nat Cell Biol* 16:434–444. [CrossRef Medline](#)
- Regev N, Degani-Katzav N, Korngreen A, Etzioni A, Siloni S, Alaimo A, Chikvashvili D, Villarreal A, Attali B, Lotan I (2009) Selective interaction of syntaxin 1A with KCNQ2: possible implications for specific modulation of presynaptic activity. *PLoS One* 4:e6586. [CrossRef Medline](#)

- Rizo J, Rosenmund C (2008) Synaptic vesicle fusion. *Nat Struct Mol Biol* 15:665–674. [CrossRef Medline](#)
- Rizo J, Südhof TC (2012) The membrane fusion enigma: SNAREs, Sec1/Munc18 proteins, and their accomplices—guilty as charged? *Annu Rev Cell Dev Biol* 28:279–308. [CrossRef Medline](#)
- Rosenmund C, Stevens CF (1996) Definition of the readily releasable pool of vesicles at hippocampal synapses. *Neuron* 16:1197–1207. [CrossRef Medline](#)
- Rowe J, Corradi N, Malosio ML, Taverna E, Halban P, Meldolesi J, Rosa P (1999) Blockade of membrane transport and disassembly of the Golgi complex by expression of syntaxin 1A in neurosecretion-incompetent cells: prevention by rbSEC1. *J Cell Sci* 112:1865–1877. [Medline](#)
- Rowe J, Calegari F, Taverna E, Longhi R, Rosa P (2001) Syntaxin 1A is delivered to the apical and basolateral domains of epithelial cells: the role of munc-18 proteins. *J Cell Sci* 114:3323–3332. [Medline](#)
- Saifee O, Wei L, Nonet ML (1998) The *Caenorhabditis elegans* unc-64 locus encodes a syntaxin that interacts genetically with synaptobrevin. *Mol Biol Cell* 9:1235–1252. [CrossRef Medline](#)
- Schoch S, Deák F, Königstorfer A, Mozhayeva M, Sara Y, Südhof TC, Kavalali ET (2001) SNARE function analyzed in synaptobrevin/VAMP knockout mice. *Science* 294:1117–1122. [CrossRef Medline](#)
- Schulze KL, Bellen HJ (1996) *Drosophila* syntaxin is required for cell viability and may function in membrane formation and stabilization. *Genetics* 144:1713–1724. [Medline](#)
- Schulze KL, Broadie K, Perin MS, Bellen HJ (1995) Genetic and electrophysiological studies of *Drosophila* syntaxin-1A demonstrate its role in non-neuronal secretion and neurotransmission. *Cell* 80:311–320. [CrossRef Medline](#)
- Sheng ZH, Rettig J, Takahashi M, Catterall WA (1994) Identification of a syntaxin-binding site on N-type calcium channels. *Neuron* 13:1303–1313. [CrossRef Medline](#)
- Sørensen JB, Wiederhold K, Müller EM, Milosevic I, Nagy G, de Groot BL, Grubmüller H, Fasshauer D (2006) Sequential N- to C-terminal SNARE complex assembly drives priming and fusion of secretory vesicles. *EMBO J* 25:955–966. [CrossRef Medline](#)
- Toonen RF, de Vries KJ, Zalm R, Südhof TC, Verhage M (2005) Munc18-1 stabilizes syntaxin 1, but is not essential for syntaxin 1 targeting and SNARE complex formation. *J Neurochem* 93:1393–1400. [CrossRef Medline](#)
- Valentino K, Newcomb R, Gadbois T, Singh T, Bowersox S, Bitner S, Justice A, Yamashiro D, Hoffman BB, Ciaranello R (1993) A selective N-type calcium channel antagonist protects against neuronal loss after global cerebral ischemia. *Proc Natl Acad Sci U S A* 90:7894–7897. [CrossRef Medline](#)
- Varoqueaux F, Sigler A, Rhee JS, Brose N, Enk C, Reim K, Rosenmund C (2002) Total arrest of spontaneous and evoked synaptic transmission but normal synaptogenesis in the absence of Munc13-mediated vesicle priming. *Proc Natl Acad Sci U S A* 99:9037–9042. [CrossRef Medline](#)
- Varoqueaux F, Sons MS, Plomp JJ, Brose N (2005) Aberrant morphology and residual transmitter release at the Munc13-deficient mouse neuromuscular synapse. *Mol Cell Biol* 25:5973–5984. [CrossRef Medline](#)
- Verhage M, Maia AS, Plomp JJ, Brussaard AB, Heeroma JH, Vermeer H, Toonen RF, Hammer RE, van den Berg TK, Missler M, Geuze HJ, Südhof TC (2000) Synaptic assembly of the brain in the absence of neurotransmitter secretion. *Science* 287:864–869. [CrossRef Medline](#)
- Walter AM, Wiederhold K, Bruns D, Fasshauer D, Sørensen JB (2010) Synaptobrevin N-terminally bound to syntaxin-SNAP-25 defines the primed vesicle state in regulated exocytosis. *J Cell Biol* 188:401–413. [CrossRef Medline](#)
- Washbourne P, Thompson PM, Carta M, Costa ET, Mathews JR, López-Bendito G, Molnár Z, Becher MW, Valenzuela CF, Partridge LD, Wilson MC (2002) Genetic ablation of the t-SNARE SNAP-25 distinguishes mechanisms of neuroexocytosis. *Nat Neurosci* 5:19–26. [Medline](#)
- Watanabe S, Liu Q, Davis MW, Hollopeter G, Thomas N, Jørgensen NB, Jørgensen EM (2013) Ultrafast endocytosis at *Caenorhabditis elegans* neuromuscular junctions. *Elife* 2:e00723. [CrossRef Medline](#)
- Wu YJ, Tejero R, Arancillo M, Vardar G, Korotkova T, Kintscher M, Schmitz D, Ponomarenko A, Tabares L, Rosenmund C (2015) Syntaxin 1B is important for mouse postnatal survival and proper synaptic function at the mouse neuromuscular junctions. *J Neurophysiol* 114:2404–2417. [CrossRef Medline](#)
- Xue M, Reim K, Chen X, Chao HT, Deng H, Rizo J, Brose N, Rosenmund C (2007) Distinct domains of complexin I differentially regulate neurotransmitter release. *Nat Struct Mol Biol* 14:949–958. [CrossRef Medline](#)
- Zhou P, Pang ZP, Yang X, Zhang Y, Rosenmund C, Bacaj T, Südhof TC (2013) Syntaxin-1 N-peptide and Habc-domain perform distinct essential functions in synaptic vesicle fusion. *EMBO J* 32:159–171. [Medline](#)
- Zimmermann J, Trimbuch T, Rosenmund C (2014) Synaptobrevin 1 mediates vesicle priming and evoked release in a subpopulation of hippocampal neurons. *J Neurophysiol* 112:1559–1565. [CrossRef Medline](#)

Mein Lebenslauf wird aus datenschutzrechtlichen Gründen in der elektronischen Version meiner Arbeit nicht veröffentlicht.

List of publications

Publication	Impact factor
Wu YJ, Tejero R, Arancillo M, Vardar G , Korotkova T, Kintscher M, Schmitz D, Ponomarenko A, Tabares L, Rosenmund C. “Syntaxin 1B is important for mouse postnatal survival and proper synaptic function at the mouse neuromuscular junctions.” <i>J Neurophysiol.</i> 2015	2.653 (year 2015/2016)
Vardar G , Chang S, Arancillo M, Wu YJ, Trimbuch T, Rosenmund C. “Distinct Functions of Syntaxin-1 in Neuronal Maintenance, Synaptic Vesicle Docking, and Fusion in Mouse Neurons.” <i>J Neurosci.</i> 2016	5.92 (year 2015/2016)
Grauel MK, Maglione M, Reddy-Alla S, Willmes CG, Brockmann MM, Trimbuch T, Rosenmund T, Pangalos M, Vardar G , Stumpf A, Walter AM, Rost BR, Eickholt BJ, Haucke V, Schmitz D, Sigrist SJ, Rosenmund C. “RIM-binding protein 2 regulates release probability by fine-tuning calcium channel localization at murine hippocampal synapses.” <i>Proc Natl Acad Sci U S A.</i> 2016	9.67 (year 2014)

Acknowledgments

First and foremost, I would like to express my sincere gratitude to my doctoral supervisor Prof. Dr. Christian Rosenmund for his invaluable guidance during my studies. Without his patient support, not only at good times but also at difficult times, I couldn't have become the researcher I am now. Working with him, being able to share his knowledge and experience, having hours-long scientific conversations and discussions were a great pleasure for me. Thank you for letting me be a part of your lab and for encouraging me to pursue a career in science.

Second, I would like to thank to Dr. Yuan-Ju Wu and Dr. Marife Arancillo for sharing their knowledge and for having critical and productive discussions about Syntaxin-1 with me. I also would like to thank to Dr. Shuwen Chang for her collaboration for electron microscopy data and for those beautiful pictures. I want to thank to Dr. Thorsten Trimbuch for virus production, and to Katja Pötschke, Bettina Brokowski, Berit Söhl-Kielczynski, Rike Dannenberg and Sabine Lenz for their technical assistance. Besides, I would like to thank all the members of Rosenmund Lab for such a friendly lab environment, for their constantly available help, for their feedback, for their patience for listening my endless talks about Syntaxin-1 and everything else, and for lunch breaks at Mensa full of fun. Working with you was a very joyful experience. I am also thankful to my former supervisor Dr. Michael Willem for believing in me and to Dr. Caroline Pirkevi, my first-time supervisor who gave me my first project, for inviting me to the life of science and being the role model for me for how to be a scientist.

I am very very thankful to my friend Herr Neumann for his never-ending belief in the scientist in me and for never giving up on being my closest friend, who always understands me. We will always be flat-mates, no matter where we are. Also, thank you Lars for your curiosity about my project even in the middle of a party. A big thank is also to Mareike for those unforgettable and hours long Mensa meetings, which always had such a good effect on my mood, for our beautiful road trip, for your patience for my unexplainable grumpiness, and for being such a good friend. Yes, we have been friends since the beginning of our studies, such a long time. Mr. D'arcy, how can I forget you? Thank you so much for making me realize how much I like science and my job. I would like to thank to Dave for making my way to Berlin and to Daniel for being such a beautiful person. Ezgi, Selim, and Kerem, your presences were so important for me whenever I needed a breath from the stresses of life. And I know you will always be there for me as I will always be there for you (let's not take Mahmut with us). I am also very thankful to Kalan Sağlar for his support throughout my studies. Gülfem, Betül, and Merve, my life-long friends, thank you just for being.

Last but not least, I am especially grateful to my mother for her patience and sacrifices to let her daughter live abroad. Her braveness and strength to follow her own path and to be always open to new things in life always has been and always will be an inspiration for me. I also want to thank to my sisters, who are also very close friends to me. I know coping with me was not always easy, but here I am with your support. And Eylül, my little niece, even though I live far from you, you brought one of the biggest happiness in my life, in our lives. I love you all.