

6. Literaturverzeichnis

1. Adli M, Baethge C, Heinz A, Langlitz N, Bauer M. Is dose escalation of antidepressants a rational strategy after a medium-dose treatment has failed? *Eur Arch Psychiatry Clin Neurosci* 2005; 255: 387-400.
2. APA Commission of Psychotherapies: Process research. In: *Psychotherapy Research. Methodological and Efficacy Issues*. APA Press Washington, DC 1982.
3. Arolt V. Die Häufigkeit psychischer Störungen bei körperlich Kranken. In: Arolt V, Diefenbacher A. (Hrsg.) *Psychiatrie in der klinischen Medizin*. Steinkopff Darmstadt 2004: 19-53.
4. Arolt V, Driessen M, Schürmann A. Indikation zur psychotherapeutischen Intervention bei psychisch Kranken- Ergebnisse der Lübecker Allgemeinkrankenhausstudie. In: Mundt C, Linden M, Barnett W. (Hrsg.) *Psychotherapie in der Psychiatrie*. Springer, Wien, New York 1997: 269-27.
5. Arolt V, Gehrman A, John U, Dilling H. Psychiatrischer Konsiliardienst an einem Universitätsklinikum- Eine empirische Untersuchung zur Leistungscharakteristik. *Nervenarzt* 1995; 66: 347-354.
6. Arolt V, Rothermundt M. Depression bei körperlichen Erkrankungen. In: Arolt V, Diefenbacher A. (Hrsg.) *Psychiatrie in der klinischen Medizin*. Steinkopff Darmstadt 2004: 349- 388.
7. Bankier B, Littman AB. Psychiatric disorders and coronary heart disease in women- a still neglected topic: review of the literature from 1971- 2000. *Psychother Psychosom* 2002; 71: 133- 140.
8. Baldwin RC, Anderson D, Black S, Evans S, Jones R, Wilson K, Iliffe S, Faculty of Old Age Psychiatry Working Group, Royal College of Psychiatrists. Guideline for the management of late-life depression in primary care. *Int J Geriatr Psychiatry* 2003; 18 (9): 829-838.
9. Billowitz A, Friedson W. Are psychiatric consultants' recommendations followed? *Int J Psychiatry Med* 1978-1979; 9: 179-189.
10. Bodlund O, Andersson SO, Mallon L. Effects of consulting psychiatrist in primary care. *Scand Prim Health Care* 1999. 17: 153- 157.
11. Brunner, E, Langer F. *Nichtparametrische Analyse longitudinaler Daten*. R. Oldenbourg Verlag München Wien 1999.

12. Brieger P, Marneros A. Angststörungen bei unipolaren Depressionen. Klinische Diagnosen vs. Standardisiertes Interview. *Nervenarzt* 2005; 76: 586-591.
13. Bryant RA, Harvey AG. Acute stress disorder. American Psychological Association, Washington DC 2000.
14. Carr VJ, Faehrmann C, Lewin TJ, Walton JM, Reid AA. Determining the effect that consultation-liaison psychiatry in primary care has on family physicians psychiatric knowledge and practice. *Psychosomatics* 1997; 38: 217- 229.
15. Creed F. Liaison psychiatry for the 21st century: a review. *J R Soc Med* 1991; 84: 414- 417.
16. Dawson R, Lavori PW, Coryell WH, Endicott J, Keller MB. Maintenance strategies for unipolar depression: an observational study of levels of treatment and recurrence. *J. Affect. Disord.* 1998. 49: 31-44.
17. De Araujo Andreoli PB, De Albuquerque Citero V, De Jesus Mari J. A systematic review of studies of the cost effectiveness of Mental Health consultaion- liaison interventions in General Hospitals. *Psychosomatics* 2003; 44: 499-507.
18. Deckert J. Angsterkrankungen. In: Arolt V, Diefenbacher A. (Hrsg.) *Psychiatrie in der klinischen Medizin.* Steinkopff Darmstadt 2004: 410- 437.
19. De Cruppe W, Hennch C, Buchholz C, Müller A, Eich W, Herzog W. Communication between psychosomatic C-L consultants and general practitioners in a German health care system. *General Hospital Psychiatry* 2005. 27: 63-72.
20. De Jonge P, Huyse FJ, Maarten-Friso Ruinemans G et al.. Timing of Psychiatric Consultations- The Impact of Social Vulnerability and Level of Psychiatric Dysfunction. *Psychosomatics* 2000; 41: 505-511.
21. Deter HC, Weber C, Adolph D et al. Von der angewandten Psychosomatik zur integrierten Medizin- Erfahrungen mit dem Steglitzer Belegbetten-Modell. *Psychosom Med Psychol.* 2004; 54: 161-164.
22. Deutscher Bundestag. Bericht über die Lage der Psychiatrie in der Bundesrepublik Deutschland- Zur psychiatrischen und psychotherapeutisch/psychosomatischen Versorgung der Bevölkerung. Drucksache 7/4200. Heger, Bonn 1975.
23. Diefenbacher A. Erkrankungen im Grenzbereich von Psyche und Soma aus konsiliar-liaisonpsychiatrischer Sicht. *Die Psychiatrie* 2006; 3: 24-9.

24. Diefenbacher A. Consultation- Liaison Psychiatry in Germany. In : Diefenbacher A. (Hrsg.) Consultation- Liaison Psychiatry in Germany, Austria and Switzerland. Karger Basel 2004 (2004a).
25. Diefenbacher A. Konsiliar- und Liaisonpsychiatrie. In: Helmchen H, Henn F, Lauter H, Sartorius N. (Hrsg). Psychiatrie der Gegenwart. Springer-Verlag, Berlin Heidelberg 1999: 434-456.
26. Diefenbacher A, Rothermundt M, Arolt V. Aufgaben der Konsiliarpsychiatrie und –psychotherapie. In: Arolt V, Diefenbacher A. (Hrsg.) Psychiatrie in der klinischen Medizin. Steinkopff Darmstadt 2004: 3-18. (2004b).
27. Diefenbacher A, Strain JJ. Consultation-liaison psychiatry: stability and change over a 10-year period. Gen Hosp Psychiatry 2002; 24: 249-256.
28. DiMatteo MR, Lepper HS, Croghan TW. Depression is a risk factor for non-compliance with medical treatment. Arch Intern Med 2000; 160: 2101- 2107.
29. Druss BG, The mental health/primary care interface in the United States: history, structure, and context. Gen Hosp Psychiatry 2002; 24: 197-202.
30. Druss BG, Hoff RA, Rosenheck RA. Under use of antidepressants in major depression: prevalence and correlates in a national sample of young adults. Journal of Clinical Psychiatry 2000; 61(3): 234-239.
31. Eikelmann B, Harter C. Entwicklung und Nutzung der psychiatrische-psychotherapeutischen Versorgung in Deutschland. Nervenheilkunde 2006; 25: 278- 285.
32. Fleet RP, Lavoie K, Breitman BD. Is panic disorder associated with coronary heart disease? A critical review of the literature. J Psychosom Res 2000; 48: 347- 356.
33. Fritze J, Barth- Stoppik A, Berger M et al. Systematische Behandlung der Depression: Versuch eines Konsenses. Psycho 1998; 24:204- 213.
34. Galan I, Rodriguez-Artalejo F, Zorilla B. Telephone versus face-to-face household interviews in the assessment of health behaviours and preventive practices. Gac Sanit 2004 Nov; 18: 440-450.
35. Gater R, De Almeida B, Sousa E et al. The pathways to psychiatric care: a cross-cultural study. Psychological Med 1991; 21: 761-774.
36. Glassman, AH, Shapiro PA. Depression and the course of artery disease. Am J Psychiatry 1998; 155: 4- 11.
37. Gulbrandsen P, Hjortdahl P, Fugelli P. General practitioners´ knowledge of their patients´ psychosocial problems: multipractice questionnaire survey. BMJ 1997; Apr 5; 314: 1014-1018.

38. Hammer JS, Strain JJ, Lyons JS. Health Service Delivery and Research Outcomes of a Common Departmental Structure for Psychiatric Consultation-Liaison and Social Work Services at a University Hospital. *Gen Hosp Psychiatry* 1991; 13: 95- 105.
39. Harpole LH, Willams JW Jr., Olsen MK et al. Improving depression outcomes in older adults with comorbid medical illness. *Gen Hosp Psychiatry* 2005; 27: 4-12.
40. Härter M, Bermejo I, Schneider F et al. Versorgungsleitlinien zur Diagnostik und Therapie depressiver Störungen in der hausärztlichen Praxis. *Z. ärztl. Fortbild. Qual. sich.* 2003; 97 Suppl. IV:16-35.
41. Haynes RB, Taylor DW, Sackett DL. Compliance in Health Care. John Hopkins University Press Baltimore 1979.
42. Henry JA. Debits and credits in the management of depression. *Br. J. Psychiatry* 1993; 163: 33-39.
43. Herzog T, Stein B. Qualitätsmanagement in der psychotherapeutischen und psychiatrischen CL- Versorgung am Allgemeinkrankenhaus. In: Rudolf G. Eich W. (Hrsg.) *Konsiliar- und Liaisonpsychosomatik und –psychiatrie.* Schattauer Stuttgart New York 2003a; 181-232.
44. Herzog T, Stein B. Basisdokumentation für den Konsiliar- und Liaisondienst: EuroQA-CL (Deutsche Fassung V8, 6/02). In: Rudolf G. Eich W. (Hrsg.) *Konsiliar- und Liaisonpsychosomatik und –psychiatrie.* Schattauer Stuttgart New York 2003b; 227-253.
45. Herzog T, Stein B. Versorgungspraxis durch interne Anbieter (Deutsche ECLW- Studie). In: Rudolf G. Eich W. (Hrsg.) *Konsiliar- und Liaisonpsychosomatik und –psychiatrie.* Schattauer Stuttgart New York 2003c; 88-94.
46. Herzog W, Hartmann A. Psychiatrische, psychosomatische und medizinpsychologische Konsiliar- und Liaisonstätigkeit in der Bundesrepublik Deutschland- Ergebnisse einer Umfrage. *Nervenarzt* 1990; 61: 281- 293.
47. Hull SA, Jones C, Tissier JM et al. Relationship style between GPs and community mental health teams affects referral rates. *Brit J Gen Pract* 2002; 52: 101- 107.
48. Huyse FJ, Herzog T, Lobo A et al. Consultation-liaison psychiatric service delivery . Results from a European study. *Gen Hosp Psychiatry* 2001; 23:124-132.
49. Huyse FJ, Strain JJ, Hammer JS: Interventions in consultation/liaison psychiatry, part I: patterns of recommendations. *Gen Hosp Psychiatry* 1990a; 12: 213-220.

50. Huyse FJ, Strain JJ, Hammer JS: Interventions in consultation/liason psychiatry, part II: concordance. *Gen Hosp Psychiatry* 1990b; 12:221-231.
51. Huyse FJ, Lyons JS, Strain JJ. Evaluation of psychiatric consultations in the general hospital. Multivariate prediction of concordance. *Gen Hosp Psychiatry* 1992; 14: 363-369.
52. Judd LL. Mood disorders in the general population represent an important and worldwide public health problem. *Int Clin Psychopharmacol* 1995; 10: 5-10.
53. Kapfhammer HP. Psychopharmakotherapie depressiver Störungen bei somatischen Erkrankungen: SSRI und andere neue Antidepressiva. In: Diefenbacher A. (Hrsg.) Aktuelle Konsiliarpsychiatrie und Psychotherapie. Thieme Stuttgart 1999; 131-160.
54. Kathol RG, Wenzel RP. Natural History of Symptoms of Depression and Anxiety during Inpatient Treatment on General Medicine Wards. *J Gen Intern Med* 1992; 7: 287-293.
55. Katon W. Panic disorder: relationship to high medical utilization, unexplained physical symptoms, and medical costs. *J Clin Psychiatry* 1996; 57 (suppl 10): 11- 18.
56. Katon W, Gonzales J. A Review of Randomized Trials of Psychiatric Consultation- Liason Studies in Primary Care. *Psychosomatics* 1994; 35: 268-278.
57. Katon W, Roy- Byrne P, Russo J, Cowley D. Cost-effectiveness and cost offset of a collaborative care intervention for primary care patients with panic disorder. *Arch Gen Psychiatry* 2002; 59: 1098-1104.
58. Katon W, Von Korff M, Lin E et al. Stepped collaborative care for primary care patients with persistent symptoms of depression: a randomized trial. *Arch Gen Psychiatry* 1999; 56: 1109-1115.
59. Kitchell MA, Barnes HF, Veith RC et al.. Screening for depression in hospitalized geriatric medical patients. *J Am Geriatr Soc* 1982; 30:174-182.
60. Koenig HG, George LK, Meador KG. Use of antidepressants by nonpsychiatrists in the treatment of medically ill hospitalized depressed elderly patients. *Am J Psych* 1997; 154: 1369- 1375.
61. Koenig HG, Meador KG, Cohen HJ, Blazer DG. Depression in elderly hospitalized patients with medical illness. *Arch Intern Med* 1988; 148:1929-1936.
62. Lang KS. Psychiatrischer Konsil- und Liaisondienst am Allgemeinkrankenhaus. Eine empirische Untersuchung. Dissertationsschrift; Charité-Universitätsmedizin Berlin. Campus Benjamin Franklin 2006.

63. Lepine JP, Gastpar M, Mendlewicz J, Tylee A. Depression in Community: The first pan- European study DEPRES (Depression Research in European Society). *Int. Clin. Psychopharmacology* 1997; 12: 19-29.
64. Levenson JL, Hamer RM, Rossiter LF. Relation of psychopathology in general medical inpatients to use and cost of services. *Am J Psychiatry* 1990; 147: 1498- 1503.
65. Lejoyeux M, Leon E, Rouillon F et al. Epidemiology of suicide and parasuicide. *Encephale* 1994; 20: 215- 225.
66. Linden M, Bohlken J. Compliance und Psychopharmakotherapie. In: Riederer P, Laux G, Pöldinger W (Hrsg.). *Neuro- Psychopharmaka. Ein Therapie- Handbuch. Band 1: Allgemeine Grundlagen der Pharmakopsychiatrie.* Springer: Wien, New York 1992: 201-209.
67. Linden M, Mair W, Achberger M et al. Psychological disorders and their treatment in general practice in Germany. Results of a World Health Organization (WHO) study. *Nervenarzt* 1996; 67: 205-215.
68. Linden M, Gothe H, Ormel J. Der Hausarzt als Gatekeeper. *Dtsch Arztebl* 2004; 101: 2189-2192.
69. Löwe B, Gräfe K, Zipfel S et al. Detecting panic disorder in medical and psychosomatic outpatients. Comparative validation of the Hospital Anxiety and Depression Scale, the Patient Health Questionnaire, a screening question and physicians' diagnosis. *J Psychosom Res* 2003; 55: 515-519.
70. Lobo A, Huyse FJ, Herzog T et al. The European Consultation Liaison Workgroup (ECLW) Collaborative Study. II: Patient Registration Form (PRF) instrument, training and reliability. *J Psychosom Res* 1996; 40: 143-156.
71. Louis AA, Manousos IR, Coletta AP et al. Clinical trials update : The Heart Protection Study, IONA, CARISA, ENRICHED, ACUTE, ALIVE, MADIT II, and REMATCH. *Eur J Heart Fail* 2002; 4: 111- 116.
72. Lyness JM, Caine ED, King DA et al. Psychiatric disorders in older primary care patients. *J Gen Intern Med* 1999; 14: 249-254.
73. Maguire GP, Julier DRL, Hawton KE, Bancroft JH. Psychiatric morbidity and referral on two general medical wards. *Br Med J* 1974; 1: 268- 270.
74. Malt UF, Huyse FJ, Herzog T et al. The European Consultation Liaison Workgroup (ECLW) Collaborative Study. III: Training and reliability of ICD-10 psychiatric diagnoses in the general hospital setting-an investigation of 220 consultants from 14 European countries. *J Psychosomatic Res* 1996; 41: 451- 463.

75. Mayou R. The history of general hospital psychiatry. *Br J Psychiatry*. 1998; 155: 764- 776.
76. Meyer T, Klemme H, Hermann- Lingen C. Persistenz und Effekte von Depressivität und Angstsymptomatik bei internistischen Patienten im ersten Jahr nach stationärer Behandlung. *Z Psychosom Med Psychother* 2002; 48: 174- 191.
77. Midanik LT, Greenfield TK. Telephone versus in-person interviews for alcohol use: results of the 2000 National Alcohol Survey. *Drug Alcohol Depend* 2003 Dec 11; 72 (3): 209-214.
78. Mintz J, Mintz LI, Arruda MJ, Hwang SS. Treatment of depression and the functional capacity to work. *Arch. Gen. Psychiatry* 1992; 49: 761-768.
79. Niklewski G, Lehfeld H. Anpassungsstörungen im Konsiliar-/Liaisondienst- Konzepte und empirische Befunde. In: Arolt V, Diefenbacher A. (Hrsg.). *Psychiatrie in der klinischen Medizin*. Steinkopff Verlag Darmstadt 2004: 389- 394.
80. Popkin MK, Mackenzie TB, Callies AL: Consultation- liaison outcome evaluation system. *Arch Gen Psychiatry* 1983; 40: 215-219.
81. Popkin MK, Mackenzie TB, Hall RCW, Garrard J. Physicians' concordance with consultants' recommendations for psychotropic medication. *Arch Gen Psychiatry* 1979; 36: 386-389.
82. Querido A. Forecast and follow-up: an investigation to the clinical social and mental factors determining the results of hospital treatment. *Br J Prev Soc Med* 1959; 13: 33-49.
83. Rasmussen A, Lunde M, Loldrup Poulsen D et al. A double blind, placebo kontrollen study of sertraline in the prevention of depression in stroke patients. *Psychosomatics* 2003; 44: 216- 221.
84. Roy-Byrne P, Russo J, Dugdale DC et al. Undertreatment of panic disorder in primary care: role of patient and physician characteristics. *J Am Board Fam Pract* 2002;15: 443-450.
85. Roy-Byrne P, Stein M, Bystrisky A et al. Pharmacotherapy of panic disorder: proposed guidelines for family physicians. *J Am Board Fam Pract* 1998; 11:282-290.
86. Rosemann T, Rüter G, Wensing M, Szecsenyi J. Überweisungen vom Hausarzt zum Facharzt: Naht- oder Bruchstelle? *Dtsch Arztebl* 2006; 103 (37): 2387-2392.
87. Saupe R, Diefenbacher A. Konsilpsychiatrie: Sozial- und angewandte Neuropsychiatrie. In: Kipp J (Hrsg.). *Handeln und Atmosphäre im*

- therapeutischen Raum psychiatrischer Abteilungen. S. Roderer Verlag Regensburg 1995: 263-269.
88. Schneider F, Härter M, Kratz S. Unzureichender subjektiver Behandlungsverlauf bei depressiven Patienten in der haus- und nervenärztlichen Praxis. Z. ärztl. Fortbild. Qual.sich.(ZaeFQ) 2003; 97 Suppl. IV: 57-66.
 89. Schnyder U, Feld C, Leuthold A, Buddeberg C. Reference to psychiatric consultation in the discharge letter of general hospital inpatients. Int'l. J. Psychiatry In Medicine 1997; 4: 391-402.
 90. Schulberg HC, Block MR, Madonia MJ et al. The "usual care" of major depression in primary care practice. Arch Fam Med. 1997; 6(4): 340-341.
 91. Schulberg HC, Block MR, Madonia MJ et al. Treating major depression in primary care practice. Eight-month clinical outcomes. Arch Gen Psychiatry 1996; 53 (10): 913-919.
 92. Schwenk TL, Evans DL, Laden SK, Lewis L. Treatment outcome and physician-patient communication in primary care patients with chronic, recurrent depression. Am J Psychiatry. 2004 Oct; 161 (10): 1892-1901.
 93. Simon GE. Evidence review: efficacy and effectiveness of antidepressant treatment in primary care. Gen Hosp Psychiatry 2002; 24: 213- 124.
 94. Smith GC, Clarke DM, Handrinis D, Dunsis A. Consultation-liaison psychiatrists' management of depression. Psychosomatics 1998 ; 39 : 244-252.
 95. Smit F, Willemse G, Koopmanschap M et al. Cost- effectiveness of preventing depression in primary care patients. Br J Psychiatry 2006; 188: 330- 336.
 96. Snaith P, Herrmann C, Buss U. Hospital anxiety and depression scale: HADS-D; deutsche Version; ein Fragebogen zur Erfassung von Angst und Depressivität in der somatischen Medizin; Testdokumentation und Handanweisung. Huber Bern. 1995.
 97. Snyder SL, Strain JJ. Change of diagnosis at termination of psychiatric consultation. Hosp Community Psychiatry 1990; 41: 286-289.
 98. Soellner W, Creed F. European guidelines for training in consultation-liaison psychiatry and psychosomatics: Report of the EACLPP Workgroup on Training in Consultation-Liaison Psychiatry and Psychosomatics. J. Psychosom Res. 2007; 62: 501-509
 99. Spitzer RL, Williams JB, Kroenke K et al. Utility of a new procedure for diagnosing mental disorders in primary care: the PRIME-MD 1000 study. JAMA 1994; 272: 1749-1756.

100. Starfield B. Primary care: balancing health needs, services, and technology. Oxford University Press. New York, 1992.
101. Starfield B, Steinwachs D, Morris I et al. Patient-provider agreement about problems. Influence on outcome care. JAMA 1979a ; 242: 344- 346.
102. Starfield B, Steinwachs D, Morris I et al. Presence of observers at patient-practitioners interactions: Impact on coordination of care and methodologic implications. Am J Public Health 1979b; 69: 1021-1025.
103. Stoppe G. Interaktion von somatischer Erkrankung und Depressionen in der ambulanten Versorgung. In: Gastpar M, Müller WE. Depressionen. Versorgungsstrukturen und Behandlungsperspektiven. Springer- Verlag Berlin Heidelberg 2002: 57- 62.
104. Strebel B, Kuhnigk O, Jüptner M et al. Einstellungen von Studierenden der Medizin zur Psychiatrie. Nervenarzt 2000; 71(3): 2005-2012.
105. Strain JJ, Smith GC, Hammer JS et al.. Adjustment disorder : a multisite study of its utilisation and interventions in the consultation-liaison psychiatry setting. Gen Hosp Psychiatry 1998; 20: 139-149.
106. Sweet L. Telephone interviewing: is it compatible with interpretive phenomenological research? Contemp Nurse 2002 Feb; 12(1): 58-63.
107. The WHO World Mental Health Survey Consortium. Prevalence, severity und unmet need for treatment of mental disorders in the World Health Organization World Mental Health Surveys. JAMA 2004; 291: 2581- 2590.
108. Üstün TB, Sartorius N. Mental illness in general health care. An international study. John Wiley & Sons, Chichester 1995.
109. Weber A, Hörmann G, Köllner V. Psychische und Verhaltensstörungen- Die Epidemie des 21. Jahrhunderts? Dtsch Arztebl 2006; 13: 712- 715.
110. Windgassen K, Weißen PH, Schmidt K. Vorurteile und Urteile: Die psychiatrische Konsiliaruntersuchung aus der Sicht der Patienten. Psychiat Prax. 1997; 24: 134- 137.
111. Wittchen HU, Pittrow D. Prevalence, recognition and management of depression in primary care in Germany: the Depression 2000 study. Hum Psychopharmacol 2002; 17 Suppl. 1: 1-11.
112. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. Acta Psychiatr Scand 1983; 67: 361-370.