

CORRECTION

Correction: Mental disorders and mental health symptoms during imprisonment: A three-year follow-up study


Caroline Gabrysch, Rosemarie Fritsch, Stefan Priebe, Adrian P. Mundt

In the Results section of the Abstract, the third sentence is incorrect. The correct sentence is: The mean GSI improved from 1.6 (SD 0.84) at intake to 1.2 (SD 0.82) at follow-up ($p < 0.001$).

In the Mental disorders at admission and at follow-up section of the Results, the second sentence is incorrect. The correct sentence is: Out of 47 (64%) individuals with major depression at baseline, 19 (26%) still met the diagnostic criteria at follow-up, and 28 (38%) had been remitted at follow-up and there were four new cases.

In the Mental disorders at admission and at follow-up section of the Results, the fourth sentence is incorrect. The correct sentence is: The prevalence of current psychotic disorders decreased from 22 (30%) at baseline to 10 (14%) at follow-up ($p = 0.008$).

There are errors in [Table 2](#). Please see the correct [Table 2](#) here.



OPEN ACCESS

Citation: Gabrysch C, Fritsch R, Priebe S, Mundt AP (2020) Correction: Mental disorders and mental health symptoms during imprisonment: A three-year follow-up study. PLoS ONE 15(4): e0231593. <https://doi.org/10.1371/journal.pone.0231593>

Published: April 3, 2020

Copyright: © 2020 Gabrysch et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Table 2. The prevalence of psychiatric disorders (diagnostic groups) among prisoners at baseline and at three-year follow-up.

	Baseline		Follow-up		McNemar test				p-Value
	N = 73	Prevalence in % (CI)	N = 73	Prevalence in % (CI)	Cases at baseline and follow-up (n)	New cases at follow-up (n)	Remitted cases at follow-up (n)		
Depression ^a	47	64 (53.4–75.3)	23	32 (20.5–42.5)	19	4	28	<0.001	
Anxiety disorders ^b	32	44 (31.5–56.2)	27	37 (26–47.9)	18	9	14	n.s.	
Psychotic disorders ^c	22	30 (19.2–39.7)	10	14 (6.8–23.3)	7	3	15	0.008	
PD	52	71 (60.3–82.2)	51	70 (58.9–80.8)	42	9	10	n.s.	
IDUD	47	64 (53.4–75.3)	36	50 (38.4–61.6)	26	10	21	n.s.	
AUD	22	30 (19.2–41.1)	13	18 (9.6–27.4)	6	7	16	n.s.	

^aIncluding major depression, recurrent major depression

^bincluding current panic disorder, agoraphobia, social anxiety disorder, generalized anxiety disorder, obsessive-compulsive disorder, posttraumatic stress disorder

^cincluding current psychotic disorder, current psychotic mood disorder

PD = personality disorder (including borderline personality disorder and antisocial personality disorder)

IDUD = illicit drug use disorder (including drug dependence and drug abuse)

AUD = alcohol use disorder (including alcohol dependence and alcohol abuse)

n.s. = non-significant.

<https://doi.org/10.1371/journal.pone.0231593.t001>

Reference

1. Gabrysch C, Fritsch R, Priebe S, Mundt AP (2019) Mental disorders and mental health symptoms during imprisonment: A three-year follow-up study. PLoS ONE 14(3): e0213711. <https://doi.org/10.1371/journal.pone.0213711> PMID: 30870479