

6. References

- Albesiano, E., Messmer, B. T., Damle, R. N., Allen, S. L., Rai, K. R., and Chiorazzi, N. (2003). Activation-induced cytidine deaminase in chronic lymphocytic leukemia B cells: expression as multiple forms in a dynamic, variably sized fraction of the clone. *Blood* 102, 3333-3339.
- Antonchuk, J., Sauvageau, G., and Humphries, R. K. (2001). HOXB4 overexpression mediates very rapid stem cell regeneration and competitive hematopoietic repopulation. *Exp Hematol* 29, 1125-1134.
- Antonchuk, J., Sauvageau, G., and Humphries, R. K. (2002). HOXB4-induced expansion of adult hematopoietic stem cells ex vivo. *Cell* 109, 39-45.
- Arakawa, H., Hauschild, J., and Buerstedde, J. M. (2002). Requirement of the activation-induced deaminase (AID) gene for immunoglobulin gene conversion. *Science* 295, 1301-1306.
- Bachl, J., Carlson, C., Gray-Schopfer, V., Dessing, M., and Olsson, C. (2001). Increased transcription levels induce higher mutation rates in a hypermutating cell line. *J Immunol* 166, 5051-5057.
- Bachl, J., Steinberg, C., and Wabl, M. (1997). Critical test of hot spot motifs for immunoglobulin hypermutation. *Eur J Immunol* 27, 3398-3403.
- Bardwell, P. D., Woo, C. J., Wei, K., Li, Z., Martin, A., Sack, S. Z., Parris, T., Edelmann, W., and Scharff, M. D. (2004). Altered somatic hypermutation and reduced class-switch recombination in exonuclease 1-mutant mice. *Nat Immunol* 5, 224-229.
- Barreto, V., Reina-San-Martin, B., Ramiro, A. R., McBride, K. M., and Nussenzweig, M. C. (2003). C-terminal deletion of AID uncouples class switch recombination from somatic hypermutation and gene conversion. *Mol Cell* 12, 501-508.
- Berek, C., Berger, A., and Apel, M. (1991). Maturation of the immune response in germinal centers. *Cell* 67, 1121-1129.
- Blattner, F. R., and Tucker, P. W. (1984). The molecular biology of immunoglobulin D. *Nature* 307, 417-422.
- Bornemann, K. D., Brewer, J. W., Beck-Engeser, G. B., Corley, R. B., Haas, I. G., and Jack, H. M. (1995). Roles of heavy and light chains in IgM polymerization. *Proc Natl Acad Sci U S A* 92, 4912-4916.
- Bosma, G. C., Kim, J., Urich, T., Fath, D. M., Cotticelli, M. G., Ruetsch, N. R., Radic, M. Z., and Bosma, M. J. (2002). DNA-dependent protein kinase activity is not required for immunoglobulin class switching. *J Exp Med* 196, 1483-1495.

- Boyd, A. W., and Schrader, J. W. (1981). The regulation of growth and differentiation of a murine B cell lymphoma. II. The inhibition of WEHI 231 by anti-immunoglobulin antibodies. *J Immunol* 126, 2466-2469.
- Bransteitter, R., Pham, P., Scharff, M. D., and Goodman, M. F. (2003). Activation-induced cytidine deaminase deaminates deoxycytidine on single-stranded DNA but requires the action of RNase. *Proc Natl Acad Sci U S A* 100, 4102-4107.
- Brenner, S., and Milstein, C. (1966). Origin of antibody variation. *Nature* 211, 242-243.
- Buchhagen, D. L., Pedersen, F. S., Crowther, R. L., and Haseltine, W. A. (1980). Most sequence differences between the genomes of the Akv virus and a leukemogenic Gross A virus passaged in vitro are located near the 3' terminus. *Proc Natl Acad Sci U S A* 77, 4359-4363.
- Buermeyer, A. B., Deschenes, S. M., Baker, S. M., and Liskay, R. M. (1999). Mammalian DNA mismatch repair. *Annu Rev Genet* 33, 533-564.
- Burrows, P. D., Beck, G. B., and Wabl, M. R. (1981). Expression of mu and gamma immunoglobulin heavy chains in different cells of a cloned mouse lymphoid line. *Proc Natl Acad Sci U S A* 78, 564-568.
- Butler, J. E. (1998). Immunoglobulin diversity, B-cell and antibody repertoire development in large farm animals. *Rev Sci Tech* 17, 43-70.
- Camacho, S. A., Kosco-Vilbois, M. H., and Berek, C. (1998). The dynamic structure of the germinal center. *Immunol Today* 19, 511-514.
- Carlson, L. M., McCormack, W. T., Postema, C. E., Humphries, E. H., and Thompson, C. B. (1990). Templated insertions in the rearranged chicken IgL V gene segment arise by intrachromosomal gene conversion. *Genes Dev* 4, 536-547.
- Cascalho, M., Wong, J., Steinberg, C., and Wabl, M. (1998). Mismatch repair co-opted by hypermutation. *Science* 279, 1207-1210.
- Casellas, R., Nussenzweig, A., Wuerffel, R., Pelanda, R., Reichlin, A., Suh, H., Qin, X. F., Besmer, E., Kenter, A., Rajewsky, K., and Nussenzweig, M. C. (1998). Ku80 is required for immunoglobulin isotype switching. *Embo J* 17, 2404-2411.
- Celeste, A., Petersen, S., Romanienko, P. J., Fernandez-Capetillo, O., Chen, H. T., Sedelnikova, O. A., Reina-San-Martin, B., Coppola, V., Meffre, E., Difilippantonio, M. J., *et al.* (2002). Genomic instability in mice lacking histone H2AX. *Science* 296, 922-927.
- Chaudhuri, J., Khuong, C., and Alt, F. W. (2004). Replication protein A interacts with AID to promote deamination of somatic hypermutation targets. *Nature* 430, 992-998.

- Chaudhuri, J., Tian, M., Khuong, C., Chua, K., Pinaud, E., and Alt, F. W. (2003). Transcription-targeted DNA deamination by the AID antibody diversification enzyme. *Nature* 422, 726-730.
- Chester, A., Somasekaram, A., Tzimina, M., Jarmuz, A., Gisbourne, J., O'Keefe, R., Scott, J., and Navaratnam, N. (2003). The apolipoprotein B mRNA editing complex performs a multifunctional cycle and suppresses nonsense-mediated decay. *Embo J* 22, 3971-3982.
- Chiang, P. W., Wang, S., Smithivas, P., Song, W. J., Ramamoorthy, S., Hillman, J., Puett, S., Van Keuren, M. L., Crombez, E., Kumar, A., *et al.* (1996). Identification and analysis of the human and murine putative chromatin structure regulator SUPT6H and Supt6h. *Genomics* 34, 328-333.
- Cyr, D. M., Lu, X., and Douglas, M. G. (1992). Regulation of Hsp70 function by a eukaryotic DnaJ homolog. *J Biol Chem* 267, 20927-20931.
- Di Noia, J., and Neuberger, M. S. (2002). Altering the pathway of immunoglobulin hypermutation by inhibiting uracil-DNA glycosylase. *Nature* 419, 43-48.
- Diaz, M., Verkoczy, L. K., Flajnik, M. F., and Klinman, N. R. (2001). Decreased frequency of somatic hypermutation and impaired affinity maturation but intact germinal center formation in mice expressing antisense RNA to DNA polymerase zeta. *J Immunol* 167, 327-335.
- Dickerson, S. K., Market, E., Besmer, E., and Papavasiliou, F. N. (2003). AID mediates hypermutation by deaminating single stranded DNA. *J Exp Med* 197, 1291-1296.
- Durandy, A., Revy, P., and Fischer, A. (2003). Hyper-immunoglobulin-M syndromes caused by an intrinsic B cell defect. *Curr Opin Allergy Clin Immunol* 3, 421-425.
- Durdik, J., Gerstein, R. M., Rath, S., Robbins, P. F., Nisonoff, A., and Selsing, E. (1989). Isotype switching by a microinjected mu immunoglobulin heavy chain gene in transgenic mice. *Proc Natl Acad Sci U S A* 86, 2346-2350.
- Ehrenstein, M. R., and Neuberger, M. S. (1999). Deficiency in Msh2 affects the efficiency and local sequence specificity of immunoglobulin class-switch recombination: parallels with somatic hypermutation. *Embo J* 18, 3484-3490.
- Ehrenstein, M. R., Rada, C., Jones, A. M., Milstein, C., and Neuberger, M. S. (2001). Switch junction sequences in PMS2-deficient mice reveal a microhomology-mediated mechanism of Ig class switch recombination. *Proc Natl Acad Sci U S A* 98, 14553-14558.
- Endoh, M., Zhu, W., Hasegawa, J., Watanabe, H., Kim, D. K., Aida, M., Inukai, N., Narita, T., Yamada, T., Furuya, A., *et al.* (2004). Human Spt6 stimulates transcription elongation by RNA polymerase II in vitro. *Mol Cell Biol* 24, 3324-3336.

- Etzerodt, M., Mikkelsen, T., Pedersen, F. S., Kjeldgaard, N. O., and Jorgensen, P. (1984). The nucleotide sequence of the Akv murine leukemia virus genome. *Virology* 134, 196-207.
- Fagarasan, S., Muramatsu, M., Suzuki, K., Nagaoka, H., Hiai, H., and Honjo, T. (2002). Critical roles of activation-induced cytidine deaminase in the homeostasis of gut flora. *Science* 298, 1424-1427.
- Faili, A., Aoufouchi, S., Flatter, E., Gueranger, Q., Reynaud, C. A., and Weill, J. C. (2002). Induction of somatic hypermutation in immunoglobulin genes is dependent on DNA polymerase iota. *Nature* 419, 944-947.
- Fayette, J., Dubois, B., Vandenabeele, S., Bridon, J. M., Vanbervliet, B., Durand, I., Banchereau, J., Caux, C., and Briere, F. (1997). Human dendritic cells skew isotype switching of CD40-activated naive B cells towards IgA1 and IgA2. *J Exp Med* 185, 1909-1918.
- Gearhart, P. J., and Bogenhagen, D. F. (1983). Clusters of point mutations are found exclusively around rearranged antibody variable genes. *Proc Natl Acad Sci U S A* 80, 3439-3443.
- Gearhart, P. J., Johnson, N. D., Douglas, R., and Hood, L. (1981). IgG antibodies to phosphorylcholine exhibit more diversity than their IgM counterparts. *Nature* 291, 29-34.
- Gellert, M. (1992). Molecular analysis of V(D)J recombination. *Annu Rev Genet* 26, 425-446.
- Gerondakis, S. (1990). Structure and expression of murine germ-line immunoglobulin epsilon heavy chain transcripts induced by interleukin 4. *Proc Natl Acad Sci U S A* 87, 1581-1585.
- Gilman, A. G., Simon, M. I., Bourne, H. R., Harris, B. A., Long, R., Ross, E. M., Stull, J. T., Taussig, R., Arkin, A. P., Cobb, M. H., *et al.* (2002). Overview of the Alliance for Cellular Signaling. *Nature* 420, 703-706.
- Goding, J. W., Scott, D. W., and Layton, J. E. (1977). Genetics, cellular expression and function of IgD and IgM receptors. *Immunol Rev* 37, 152-186.
- Goodman, M. F., and Tippin, B. (2000). The expanding polymerase universe. *Nat Rev Mol Cell Biol* 1, 101-109.
- Gordon, M. S., Kanegai, C. M., Doerr, J. R., and Wall, R. (2003). Somatic hypermutation of the B cell receptor genes B29 (Igbeta, CD79b) and mb1 (Igalpha, CD79a). *Proc Natl Acad Sci U S A* 100, 4126-4131.
- Gottschalk, A. R., and Quintans, J. (1995). Apoptosis in B lymphocytes: the WEHI-231 perspective. *Immunol Cell Biol* 73, 8-16.

- Green, N. S., Rabinowitz, J. L., Zhu, M., Kobrin, B. J., and Scharff, M. D. (1995). Immunoglobulin variable region hypermutation in hybrids derived from a pre-B- and a myeloma cell line. *Proc Natl Acad Sci U S A* *92*, 6304-6308.
- Gutman, G. A., Warner, N. L., and Harris, A. W. (1981). Immunoglobulin production by murine B-lymphoma cells. *Clin Immunol Immunopathol* *18*, 230-244.
- Haggerty, H. G., Wechsler, R. J., Lentz, V. M., and Monroe, J. G. (1993). Endogenous expression of delta on the surface of WEHI-231. Characterization of its expression and signaling properties. *J Immunol* *151*, 4681-4693.
- Harada, Y., Muramatsu, M., Shibata, T., Honjo, T., and Kuroda, K. (2003). Unmutated immunoglobulin M can protect mice from death by influenza virus infection. *J Exp Med* *197*, 1779-1785.
- Harris, R. S., Bishop, K. N., Sheehy, A. M., Craig, H. M., Petersen-Mahrt, S. K., Watt, I. N., Neuberger, M. S., and Malim, M. H. (2003a). DNA deamination mediates innate immunity to retroviral infection. *Cell* *113*, 803-809.
- Harris, R. S., Petersen-Mahrt, S. K., and Neuberger, M. S. (2002). RNA editing enzyme APOBEC1 and some of its homologs can act as DNA mutators. *Mol Cell* *10*, 1247-1253.
- Harris, R. S., Sheehy, A. M., Craig, H. M., Malim, M. H., and Neuberger, M. S. (2003b). DNA deamination: not just a trigger for antibody diversification but also a mechanism for defense against retroviruses. *Nat Immunol* *4*, 641-643.
- Hartl, F. U. (1996). Molecular chaperones in cellular protein folding. *Nature* *381*, 571-579.
- Hendrickson, B. A., Conner, D. A., Ladd, D. J., Kendall, D., Casanova, J. E., Corthesy, B., Max, E. E., Neutra, M. R., Seidman, C. E., and Seidman, J. G. (1995). Altered hepatic transport of immunoglobulin A in mice lacking the J chain. *J Exp Med* *182*, 1905-1911.
- Hickman, S., and Kornfeld, S. (1978). Effect of tunicamycin on IgM, IgA, and IgG secretion by mouse plasmacytoma cells. *J Immunol* *121*, 990-996.
- Honjo, T., and Kataoka, T. (1978). Organization of immunoglobulin heavy chain genes and allelic deletion model. *Proc Natl Acad Sci U S A* *75*, 2140-2144.
- Honjo, T., Muramatsu, M., and Fagarasan, S. (2004). AID: how does it aid antibody diversity? *Immunity* *20*, 659-668.
- Hozumi, N., and Tonegawa, S. (1976). Evidence for somatic rearrangement of immunoglobulin genes coding for variable and constant regions. *Proc Natl Acad Sci U S A* *73*, 3628-3632.
- Imai, K., Slupphaug, G., Lee, W. I., Revy, P., Nonoyama, S., Catalan, N., Yel, L., Forveille, M., Kavli, B., Krokan, H. E., *et al.* (2003). Human uracil-DNA glycosylase

deficiency associated with profoundly impaired immunoglobulin class-switch recombination. *Nat Immunol* 4, 1023-1028.

Imai, K., Zhu, Y., Revy, P., Morio, T., Mizutani, S., Fischer, A., Nonoyama, S., and Durandy, A. (2005). Analysis of class switch recombination and somatic hypermutation in patients affected with autosomal dominant hyper-IgM syndrome type 2. *Clin Immunol* 115, 277-285.

Ito, S., Nagaoka, H., Shinkura, R., Begum, N., Muramatsu, M., Nakata, M., and Honjo, T. (2004). Activation-induced cytidine deaminase shuttles between nucleus and cytoplasm like apolipoprotein B mRNA editing catalytic polypeptide 1. *Proc Natl Acad Sci U S A* 101, 1975-1980.

Iwasato, T., Shimizu, A., Honjo, T., and Yamagishi, H. (1990). Circular DNA is excised by immunoglobulin class switch recombination. *Cell* 62, 143-149.

Jack, H. M., Berg, J., and Wabl, M. (1989). Translation affects immunoglobulin mRNA stability. *Eur J Immunol* 19, 843-847.

Jacob, J., Kelsoe, G., Rajewsky, K., and Weiss, U. (1991). Intracloonal generation of antibody mutants in germinal centres. *Nature* 354, 389-392.

Jakway, J. P., Usinger, W. R., Gold, M. R., Mishell, R. I., and DeFranco, A. L. (1986). Growth regulation of the B lymphoma cell line WEHI-231 by anti-immunoglobulin, lipopolysaccharide, and other bacterial products. *J Immunol* 137, 2225-2231.

Jensen, N. A., Jorgensen, P., Kjeldgaard, N. O., and Pedersen, F. S. (1986). Mammalian expression-and-transmission vector derived from Akv murine leukemia virus. *Gene* 41, 59-65.

Jung, S., Rajewsky, K., and Radbruch, A. (1993). Shutdown of class switch recombination by deletion of a switch region control element. *Science* 259, 984-987.

Kelley, W. L. (1999). Molecular chaperones: How J domains turn on Hsp70s. *Curr Biol* 9, R305-308.

Kenter, A. L. (2003). Class-switch recombination: after the dawn of AID. *Curr Opin Immunol* 15, 190-198.

Kim, N., Bozek, G., Lo, J. C., and Storb, U. (1999). Different mismatch repair deficiencies all have the same effects on somatic hypermutation: intact primary mechanism accompanied by secondary modifications. *J Exp Med* 190, 21-30.

Kingzette, M., Spieker-Polet, H., Yam, P. C., Zhai, S. K., and Knight, K. L. (1998). Trans-chromosomal recombination within the Ig heavy chain switch region in B lymphocytes. *Proc Natl Acad Sci U S A* 95, 11840-11845.

Kinoshita, K., and Honjo, T. (2001). Linking class-switch recombination with somatic hypermutation. *Nat Rev Mol Cell Biol* 2, 493-503.

- Kinoshita, K., Tashiro, J., Tomita, S., Lee, C. G., and Honjo, T. (1998). Target specificity of immunoglobulin class switch recombination is not determined by nucleotide sequences of S regions. *Immunity* 9, 849-858.
- Klasen, M., Spillmann, F. J., Lorens, J. B., and Wabl, M. (2005). Retroviral vectors to monitor somatic hypermutation. *J Immunol Methods* 300, 47-62.
- Knight, K. L., Kingzette, M., Crane, M. A., and Zhai, S. K. (1995). Transchromosomally derived Ig heavy chains. *J Immunol* 155, 684-691.
- Lander, E. S., Linton, L. M., Birren, B., Nusbaum, C., Zody, M. C., Baldwin, J., Devon, K., Dewar, K., Doyle, M., FitzHugh, W., *et al.* (2001). Initial sequencing and analysis of the human genome. *Nature* 409, 860-921.
- Lau, P. P., Zhu, H. J., Baldini, A., Charnsangavej, C., and Chan, L. (1994). Dimeric structure of a human apolipoprotein B mRNA editing protein and cloning and chromosomal localization of its gene. *Proc Natl Acad Sci U S A* 91, 8522-8526.
- Laufen, T., Mayer, M. P., Beisel, C., Klostermeier, D., Mogk, A., Reinstein, J., and Bukau, B. (1999). Mechanism of regulation of hsp70 chaperones by DnaJ cochaperones. *Proc Natl Acad Sci U S A* 96, 5452-5457.
- Lee, G. S., Brandt, V. L., and Roth, D. B. (2004). B cell development leads off with a base hit: dU:dG mismatches in class switching and hypermutation. *Mol Cell* 16, 505-508.
- Li, Z., Woo, C. J., Iglesias-Ussel, M. D., Ronai, D., and Scharff, M. D. (2004). The generation of antibody diversity through somatic hypermutation and class switch recombination. *Genes Dev* 18, 1-11.
- Li, Z., Woo, C. J., and Scharff, M. D. (2003). Mutations in AID and UNG extend the function of AID. *Nat Immunol* 4, 945-946.
- Limoli, C. L., Giedzinski, E., and Cleaver, J. E. (2005). Alternative recombination pathways in UV-irradiated XP variant cells. *Oncogene* 24, 3708-3714.
- Liu, Y. J., Malisan, F., de Bouteiller, O., Guret, C., Lebecque, S., Banchereau, J., Mills, F. C., Max, E. E., and Martinez-Valdez, H. (1996). Within germinal centers, isotype switching of immunoglobulin genes occurs after the onset of somatic mutation. *Immunity* 4, 241-250.
- Lorens, J. B., Jang, Y., Rossi, A. B., Payan, D. G., and Bogenberger, J. M. (2000). Optimization of regulated LTR-mediated expression. *Virology* 272, 7-15.
- Lorenz, M., Jung, S., and Radbruch, A. (1995). Switch transcripts in immunoglobulin class switching. *Science* 267, 1825-1828.
- Lovmand, S., Kjeldgaard, N. O., Jorgensen, P., and Pedersen, F. S. (1990). Enhancer functions in U3 of Akv virus: a role for cooperativity of a tandem repeat unit and its flanking DNA sequences. *J Virol* 64, 3185-3191.

- Lutzker, S., and Alt, F. W. (1988). Structure and expression of germ line immunoglobulin gamma 2b transcripts. *Mol Cell Biol* 8, 1849-1852.
- Ma, L., Wortis, H. H., and Kenter, A. L. (2002). Two new isotype-specific switching activities detected for Ig class switching. *J Immunol* 168, 2835-2846.
- MacLennan, I. C. (1994). Germinal centers. *Annu Rev Immunol* 12, 117-139.
- Maley, F., Trimble, R. B., Tarentino, A. L., and Plummer, T. H., Jr. (1989). Characterization of glycoproteins and their associated oligosaccharides through the use of endoglycosidases. *Anal Biochem* 180, 195-204.
- Manis, J. P., Dudley, D., Kaylor, L., and Alt, F. W. (2002a). IgH class switch recombination to IgG1 in DNA-PKcs-deficient B cells. *Immunity* 16, 607-617.
- Manis, J. P., Gu, Y., Lansford, R., Sonoda, E., Ferrini, R., Davidson, L., Rajewsky, K., and Alt, F. W. (1998). Ku70 is required for late B cell development and immunoglobulin heavy chain class switching. *J Exp Med* 187, 2081-2089.
- Manis, J. P., Tian, M., and Alt, F. W. (2002b). Mechanism and control of class-switch recombination. *Trends Immunol* 23, 31-39.
- Martin, A., Bardwell, P. D., Woo, C. J., Fan, M., Shulman, M. J., and Scharff, M. D. (2002). Activation-induced cytidine deaminase turns on somatic hypermutation in hybridomas. *Nature* 415, 802-806.
- Martin, A., Li, Z., Lin, D. P., Bardwell, P. D., Iglesias-Ussel, M. D., Edelmann, W., and Scharff, M. D. (2003). Msh2 ATPase activity is essential for somatic hypermutation at a-T basepairs and for efficient class switch recombination. *J Exp Med* 198, 1171-1178.
- Martin, A., and Scharff, M. D. (2002a). AID and mismatch repair in antibody diversification. *Nat Rev Immunol* 2, 605-614.
- Martin, A., and Scharff, M. D. (2002b). Somatic hypermutation of the AID transgene in B and non-B cells. *Proc Natl Acad Sci U S A* 99, 12304-12308.
- Matsuoka, M., Yoshida, K., Maeda, T., Usuda, S., and Sakano, H. (1990). Switch circular DNA formed in cytokine-treated mouse splenocytes: evidence for intramolecular DNA deletion in immunoglobulin class switching. *Cell* 62, 135-142.
- McCarthy, H., Wierda, W. G., Barron, L. L., Cromwell, C. C., Wang, J., Coombes, K. R., Rangel, R., Elenitoba-Johnson, K. S., Keating, M. J., and Abruzzo, L. V. (2003). High expression of activation-induced cytidine deaminase (AID) and splice variants is a distinctive feature of poor-prognosis chronic lymphocytic leukemia. *Blood* 101, 4903-4908.
- McCormack, W. T., Carlson, L. M., Tjoelker, L. W., and Thompson, C. B. (1989). Evolutionary comparison of the avian IgL locus: combinatorial diversity plays a role in

the generation of the antibody repertoire in some avian species. *Int Immunol* **1**, 332-341.

McDonald, J. P., Frank, E. G., Plosky, B. S., Rogozin, I. B., Masutani, C., Hanaoka, F., Woodgate, R., and Gearhart, P. J. (2003). 129-derived strains of mice are deficient in DNA polymerase iota and have normal immunoglobulin hypermutation. *J Exp Med* **198**, 635-643.

McKean, D., Huppi, K., Bell, M., Staudt, L., Gerhard, W., and Weigert, M. (1984). Generation of antibody diversity in the immune response of BALB/c mice to influenza virus hemagglutinin. *Proc Natl Acad Sci U S A* **81**, 3180-3184.

Meacham, G. C., Lu, Z., King, S., Sorscher, E., Tousson, A., and Cyr, D. M. (1999). The Hdj-2/Hsc70 chaperone pair facilitates early steps in CFTR biogenesis. *Embo J* **18**, 1492-1505.

Medema, R. H., de Laat, W. L., Martin, G. A., McCormick, F., and Bos, J. L. (1992). GTPase-activating protein SH2-SH3 domains induce gene expression in a Ras-dependent fashion. *Mol Cell Biol* **12**, 3425-3430.

Meyer, J., Jack, H. M., Ellis, N., and Wabl, M. (1986). High rate of somatic point mutation in vitro in and near the variable-region segment of an immunoglobulin heavy chain gene. *Proc Natl Acad Sci U S A* **83**, 6950-6953.

Mills, F. C., Brooker, J. S., and Camerini-Otero, R. D. (1990). Sequences of human immunoglobulin switch regions: implications for recombination and transcription. *Nucleic Acids Res* **18**, 7305-7316.

Minegishi, Y., Lavoie, A., Cunningham-Rundles, C., Bedard, P. M., Hebert, J., Cote, L., Dan, K., Sedlak, D., Buckley, R. H., Fischer, A., *et al.* (2000). Mutations in activation-induced cytidine deaminase in patients with hyper IgM syndrome. *Clin Immunol* **97**, 203-210.

Morrison, H. L., Dai, H. Y., Pedersen, F. S., and Lenz, J. (1991). Analysis of the significance of two single-base-pair differences in the SL3-3 and Akv virus long terminal repeats. *J Virol* **65**, 1019-1022.

Muramatsu, M., Kinoshita, K., Fagarasan, S., Yamada, S., Shinkai, Y., and Honjo, T. (2000). Class switch recombination and hypermutation require activation-induced cytidine deaminase (AID), a potential RNA editing enzyme. *Cell* **102**, 553-563.

Muramatsu, M., Sankaranand, V. S., Anant, S., Sugai, M., Kinoshita, K., Davidson, N. O., and Honjo, T. (1999). Specific expression of activation-induced cytidine deaminase (AID), a novel member of the RNA-editing deaminase family in germinal center B cells. *J Biol Chem* **274**, 18470-18476.

- Nambu, Y., Sugai, M., Gonda, H., Lee, C. G., Katakai, T., Agata, Y., Yokota, Y., and Shimizu, A. (2003). Transcription-coupled events associating with immunoglobulin switch region chromatin. *Science* *302*, 2137-2140.
- Navaratnam, N., Bhattacharya, S., Fujino, T., Patel, D., Jarmuz, A. L., and Scott, J. (1995). Evolutionary origins of apoB mRNA editing: catalysis by a cytidine deaminase that has acquired a novel RNA-binding motif at its active site. *Cell* *81*, 187-195.
- Neuberger, M. S., Harris, R. S., Di Noia, J., and Petersen-Mahrt, S. K. (2003). Immunity through DNA deamination. *Trends Biochem Sci* *28*, 305-312.
- Neuberger, M. S., and Milstein, C. (1995). Somatic hypermutation. *Curr Opin Immunol* *7*, 248-254.
- Nielsen, A. L., Pallisgaard, N., Pedersen, F. S., and Jorgensen, P. (1992). Murine helix-loop-helix transcriptional activator proteins binding to the E-box motif of the Akv murine leukemia virus enhancer identified by cDNA cloning. *Mol Cell Biol* *12*, 3449-3459.
- Nikaido, T., Yamawaki-Kataoka, Y., and Honjo, T. (1982). Nucleotide sequences of switch regions of immunoglobulin C epsilon and C gamma genes and their comparison. *J Biol Chem* *257*, 7322-7329.
- Niles, M. J., Matsuuchi, L., and Koshland, M. E. (1995). Polymer IgM assembly and secretion in lymphoid and nonlymphoid cell lines: evidence that J chain is required for pentamer IgM synthesis. *Proc Natl Acad Sci U S A* *92*, 2884-2888.
- Nilsen, H., Otterlei, M., Haug, T., Solum, K., Nagelhus, T. A., Skorpen, F., and Krokan, H. E. (1997). Nuclear and mitochondrial uracil-DNA glycosylases are generated by alternative splicing and transcription from different positions in the UNG gene. *Nucleic Acids Res* *25*, 750-755.
- Nishikawa, S., Sasaki, Y., Kina, T., Amagai, T., and Katsura, Y. (1986). A monoclonal antibody against Igh6-4 determinant. *Immunogenetics* *23*, 137-139.
- Noguchi, E., Shibasaki, M., Inudou, M., Kamioka, M., Yokouchi, Y., Yamakawa-Kobayashi, K., Hamaguchi, H., Matsui, A., and Arinami, T. (2001). Association between a new polymorphism in the activation-induced cytidine deaminase gene and atopic asthma and the regulation of total serum IgE levels. *J Allergy Clin Immunol* *108*, 382-386.
- Okazaki, I. M., Kinoshita, K., Muramatsu, M., Yoshikawa, K., and Honjo, T. (2002). The AID enzyme induces class switch recombination in fibroblasts. *Nature* *416*, 340-345.
- Oppezzo, P., Vuillier, F., Vasconcelos, Y., Dumas, G., Magnac, C., Payelle-Brogard, B., Pritsch, O., and Dighiero, G. (2003). Chronic lymphocytic leukemia B cells

expressing AID display dissociation between class switch recombination and somatic hypermutation. *Blood* *101*, 4029-4032.

Otterlei, M., Haug, T., Nagelhus, T. A., Slupphaug, G., Lindmo, T., and Krokan, H. E. (1998). Nuclear and mitochondrial splice forms of human uracil-DNA glycosylase contain a complex nuclear localisation signal and a strong classical mitochondrial localisation signal, respectively. *Nucleic Acids Res* *26*, 4611-4617.

Page, D. M., and DeFranco, A. L. (1988). Role of phosphoinositide-derived second messengers in mediating anti-IgM-induced growth arrest of WEHI-231 B lymphoma cells. *J Immunol* *140*, 3717-3726.

Page, D. M., and DeFranco, A. L. (1990). Antigen receptor-induced cell cycle arrest in WEHI-231 B lymphoma cells depends on the duration of signaling before the G1 phase restriction point. *Mol Cell Biol* *10*, 3003-3012.

Page, D. M., Gold, M. R., Fahey, K. A., Matsuuchi, L., and DeFranco, A. L. (1991). Mutational analysis of antigen receptor regulation of B lymphocyte growth. Evidence for involvement of the phosphoinositide signaling pathway. *J Biol Chem* *266*, 5563-5574.

Pascual, V., Liu, Y. J., Magalski, A., de Bouteiller, O., Banchereau, J., and Capra, J. D. (1994). Analysis of somatic mutation in five B cell subsets of human tonsil. *J Exp Med* *180*, 329-339.

Petersen, S., Casellas, R., Reina-San-Martin, B., Chen, H. T., Difilippantonio, M. J., Wilson, P. C., Hanitsch, L., Celeste, A., Muramatsu, M., Pilch, D. R., *et al.* (2001). AID is required to initiate Nbs1/gamma-H2AX focus formation and mutations at sites of class switching. *Nature* *414*, 660-665.

Petersen-Mahrt, S. K., Harris, R. S., and Neuberger, M. S. (2002). AID mutates *E. coli* suggesting a DNA deamination mechanism for antibody diversification. *Nature* *418*, 99-103.

Pham, P., Bransteitter, R., Petruska, J., and Goodman, M. F. (2003). Processive AID-catalysed cytosine deamination on single-stranded DNA simulates somatic hypermutation. *Nature* *424*, 103-107.

Phung, Q. H., Winter, D. B., Cranston, A., Tarone, R. E., Bohr, V. A., Fishel, R., and Gearhart, P. J. (1998). Increased hypermutation at G and C nucleotides in immunoglobulin variable genes from mice deficient in the MSH2 mismatch repair protein. *J Exp Med* *187*, 1745-1751.

Powell, L. M., Wallis, S. C., Pease, R. J., Edwards, Y. H., Knott, T. J., and Scott, J. (1987). A novel form of tissue-specific RNA processing produces apolipoprotein-B48 in intestine. *Cell* *50*, 831-840.

- Rada, C., Ehrenstein, M. R., Neuberger, M. S., and Milstein, C. (1998). Hot spot focusing of somatic hypermutation in MSH2-deficient mice suggests two stages of mutational targeting. *Immunity* *9*, 135-141.
- Rada, C., Jarvis, J. M., and Milstein, C. (2002a). AID-GFP chimeric protein increases hypermutation of Ig genes with no evidence of nuclear localization. *Proc Natl Acad Sci U S A* *99*, 7003-7008.
- Rada, C., and Milstein, C. (2001). The intrinsic hypermutability of antibody heavy and light chain genes decays exponentially. *Embo J* *20*, 4570-4576.
- Rada, C., Williams, G. T., Nilsen, H., Barnes, D. E., Lindahl, T., and Neuberger, M. S. (2002b). Immunoglobulin isotype switching is inhibited and somatic hypermutation perturbed in UNG-deficient mice. *Curr Biol* *12*, 1748-1755.
- Radbruch, A., Liesegang, B., and Rajewsky, K. (1980). Isolation of variants of mouse myeloma X63 that express changed immunoglobulin class. *Proc Natl Acad Sci U S A* *77*, 2909-2913.
- Rajewsky, K., Forster, I., and Cumano, A. (1987). Evolutionary and somatic selection of the antibody repertoire in the mouse. *Science* *238*, 1088-1094.
- Reaban, M. E., and Griffin, J. A. (1990). Induction of RNA-stabilized DNA conformers by transcription of an immunoglobulin switch region. *Nature* *348*, 342-344.
- Reaban, M. E., Lebowitz, J., and Griffin, J. A. (1994). Transcription induces the formation of a stable RNA-DNA hybrid in the immunoglobulin alpha switch region. *J Biol Chem* *269*, 21850-21857.
- Reina-San-Martin, B., Difilippantonio, S., Hanitsch, L., Masilamani, R. F., Nussenzweig, A., and Nussenzweig, M. C. (2003). H2AX is required for recombination between immunoglobulin switch regions but not for intra-switch region recombination or somatic hypermutation. *J Exp Med* *197*, 1767-1778.
- Revy, P., Muto, T., Levy, Y., Geissmann, F., Plebani, A., Sanal, O., Catalan, N., Forveille, M., Dufourcq-Labeolouse, R., Gennery, A., *et al.* (2000). Activation-induced cytidine deaminase (AID) deficiency causes the autosomal recessive form of the Hyper-IgM syndrome (HIGM2). *Cell* *102*, 565-575.
- Reynaud, C. A., Anquez, V., Grimal, H., and Weill, J. C. (1987). A hyperconversion mechanism generates the chicken light chain preimmune repertoire. *Cell* *48*, 379-388.
- Reynaud, C. A., Dahan, A., Anquez, V., and Weill, J. C. (1989). Somatic hyperconversion diversifies the single Vh gene of the chicken with a high incidence in the D region. *Cell* *59*, 171-183.

- Reynaud, S., Delpy, L., Fleury, L., Dougier, H. L., Sirac, C., and Cogne, M. (2005). Interallelic class switch recombination contributes significantly to class switching in mouse B cells. *J Immunol* *174*, 6176-6183.
- Rogozin, I. B., and Kolchanov, N. A. (1992). Somatic hypermutagenesis in immunoglobulin genes. II. Influence of neighbouring base sequences on mutagenesis. *Biochim Biophys Acta* *1171*, 11-18.
- Rolink, A., Melchers, F., and Andersson, J. (1996). The SCID but not the RAG-2 gene product is required for S mu-S epsilon heavy chain class switching. *Immunity* *5*, 319-330.
- Rudikoff, S., Pawlita, M., Pumphrey, J., and Heller, M. (1984). Somatic diversification of immunoglobulins. *Proc Natl Acad Sci U S A* *81*, 2162-2166.
- Rudolph, A. K., Burrows, P. D., and Wabl, M. R. (1981). Thirteen hybridomas secreting hapten-specific immunoglobulin E from mice with Iga or Igb heavy chain haplotype. *Eur J Immunol* *11*, 527-529.
- Sale, J. E., Calandrini, D. M., Takata, M., Takeda, S., and Neuberger, M. S. (2001). Ablation of XRCC2/3 transforms immunoglobulin V gene conversion into somatic hypermutation. *Nature* *412*, 921-926.
- Schatz, D. G., Oettinger, M. A., and Baltimore, D. (1989). The V(D)J recombination activating gene, RAG-1. *Cell* *59*, 1035-1048.
- Schrader, C. E., Edelmann, W., Kucherlapati, R., and Stavnezer, J. (1999). Reduced isotype switching in splenic B cells from mice deficient in mismatch repair enzymes. *J Exp Med* *190*, 323-330.
- Schrader, C. E., Vardo, J., and Stavnezer, J. (2002). Role for mismatch repair proteins Msh2, Mlh1, and Pms2 in immunoglobulin class switching shown by sequence analysis of recombination junctions. *J Exp Med* *195*, 367-373.
- Schreier, P. H., Quester, S., and Bothwell, A. (1986). Allotypic differences in murine mu genes. *Nucleic Acids Res* *14*, 2381-2389.
- Shanmugam, A., Shi, M. J., Yauch, L., Stavnezer, J., and Kenter, A. L. (2000). Evidence for class-specific factors in immunoglobulin isotype switching. *J Exp Med* *191*, 1365-1380.
- Shen, H. M., Peters, A., Baron, B., Zhu, X., and Storb, U. (1998). Mutation of BCL-6 gene in normal B cells by the process of somatic hypermutation of Ig genes. *Science* *280*, 1750-1752.
- Shimizu, A., Takahashi, N., Yaoita, Y., and Honjo, T. (1982). Organization of the constant-region gene family of the mouse immunoglobulin heavy chain. *Cell* *28*, 499-506.

- Shinkura, R., Ito, S., Begum, N. A., Nagaoka, H., Muramatsu, M., Kinoshita, K., Sakakibara, Y., Hijikata, H., and Honjo, T. (2004). Separate domains of AID are required for somatic hypermutation and class-switch recombination. *Nat Immunol* *5*, 707-712.
- Shinkura, R., Tian, M., Smith, M., Chua, K., Fujiwara, Y., and Alt, F. W. (2003). The influence of transcriptional orientation on endogenous switch region function. *Nat Immunol* *4*, 435-441.
- Sibley, C. H., and Wagner, R. A. (1981). Glycosylation is not required for membrane localization or secretion of IgM in a mouse B cell lymphoma. *J Immunol* *126*, 1868-1873.
- Siekevitz, M., Kocks, C., Rajewsky, K., and Dildrop, R. (1987). Analysis of somatic mutation and class switching in naive and memory B cells generating adoptive primary and secondary responses. *Cell* *48*, 757-770.
- Smith, G. C., and Jackson, S. P. (1999). The DNA-dependent protein kinase. *Genes Dev* *13*, 916-934.
- Snapper, C. M., Marcu, K. B., and Zelazowski, P. (1997). The immunoglobulin class switch: beyond "accessibility". *Immunity* *6*, 217-223.
- Spillmann, F. J., and Wabl, M. (2004). Endogenous expression of activation-induced cytidine deaminase in cell line WEHI-231. *J Immunol* *173*, 1858-1867.
- Stavnezer, J. (2000). Molecular processes that regulate class switching. *Curr Top Microbiol Immunol* *245*, 127-168.
- Stavnezer, J., Radcliffe, G., Lin, Y. C., Nietupski, J., Berggren, L., Sitia, R., and Severinson, E. (1988). Immunoglobulin heavy-chain switching may be directed by prior induction of transcripts from constant-region genes. *Proc Natl Acad Sci U S A* *85*, 7704-7708.
- Stavnezer-Nordgren, J., and Sirlin, S. (1986). Specificity of immunoglobulin heavy chain switch correlates with activity of germline heavy chain genes prior to switching. *Embo J* *5*, 95-102.
- Storb, U., Peters, A., Klotz, E., Kim, N., Shen, H. M., Kage, K., Rogerson, B., and Martin, T. E. (1998). Somatic hypermutation of immunoglobulin genes is linked to transcription. *Curr Top Microbiol Immunol* *229*, 11-19.
- Sun, J., and Butler, J. E. (1997). Sequence analysis of pig switch mu, C mu, and C mu m. *Immunogenetics* *46*, 452-460.
- Svejstrup, J. Q. (2003). Transcription. Histones face the FACT. *Science* *301*, 1053-1055.

- Swanson, M. S., Carlson, M., and Winston, F. (1990). SPT6, an essential gene that affects transcription in *Saccharomyces cerevisiae*, encodes a nuclear protein with an extremely acidic amino terminus. *Mol Cell Biol* 10, 4935-4941.
- Swanson, M. S., and Winston, F. (1992). SPT4, SPT5 and SPT6 interactions: effects on transcription and viability in *Saccharomyces cerevisiae*. *Genetics* 132, 325-336.
- Syken, J., De-Medina, T., and Munger, K. (1999). TID1, a human homolog of the *Drosophila* tumor suppressor I(2)tid, encodes two mitochondrial modulators of apoptosis with opposing functions. *Proc Natl Acad Sci U S A* 96, 8499-8504.
- Ta, V. T., Nagaoka, H., Catalan, N., Durandy, A., Fischer, A., Imai, K., Nonoyama, S., Tashiro, J., Ikegawa, M., Ito, S., *et al.* (2003). AID mutant analyses indicate requirement for class-switch-specific cofactors. *Nat Immunol* 4, 843-848.
- Teng, B., Burant, C. F., and Davidson, N. O. (1993). Molecular cloning of an apolipoprotein B messenger RNA editing protein. *Science* 260, 1816-1819.
- Tisch, R., Roifman, C. M., and Hozumi, N. (1988). Functional differences between immunoglobulins M and D expressed on the surface of an immature B-cell line. *Proc Natl Acad Sci U S A* 85, 6914-6918.
- Tonegawa, S. (1983). Somatic generation of antibody diversity. *Nature* 302, 575-581.
- Trentin, G. A., Yin, X., Tahir, S., Lhotak, S., Farhang-Fallah, J., Li, Y., and Rozakis-Adcock, M. (2001). A mouse homologue of the *Drosophila* tumor suppressor I(2)tid gene defines a novel Ras GTPase-activating protein (RasGAP)-binding protein. *J Biol Chem* 276, 13087-13095.
- Venter, J. C., Adams, M. D., Myers, E. W., Li, P. W., Mural, R. J., Sutton, G. G., Smith, H. O., Yandell, M., Evans, C. A., Holt, R. A., *et al.* (2001). The sequence of the human genome. *Science* 291, 1304-1351.
- Ventura, A., Meissner, A., Dillon, C. P., McManus, M., Sharp, P. A., Van Parijs, L., Jaenisch, R., and Jacks, T. (2004). Cre-lox-regulated conditional RNA interference from transgenes. *Proc Natl Acad Sci U S A* 101, 10380-10385.
- von Schwedler, U., Jack, H. M., and Wabl, M. (1990). Circular DNA is a product of the immunoglobulin class switch rearrangement. *Nature* 345, 452-456.
- Wabl, M., Burrows, P. D., von Gabain, A., and Steinberg, C. (1985). Hypermutation at the immunoglobulin heavy chain locus in a pre-B-cell line. *Proc Natl Acad Sci U S A* 82, 479-482.
- Wabl, M., Cascalho, M., and Steinberg, C. (1999). Hypermutation in antibody affinity maturation. *Curr Opin Immunol* 11, 186-189.
- Wabl, M., Jack, H. M., Meyer, J., Beck-Engeser, G., von Borstel, R. C., and Steinberg, C. M. (1987). Measurements of mutation rates in B lymphocytes. *Immunol Rev* 96, 91-107.

- Wang, C. L., Harper, R. A., and Wabl, M. (2004a). Genome-wide somatic hypermutation. *Proc Natl Acad Sci U S A* *101*, 7352-7356.
- Wang, C. L., Hodgson, J. G., Malek, T., Pedersen, F. S., and Wabl, M. (2004b). A murine leukemia virus with Cre-LoxP excisable coding sequences allowing superinfection, transgene delivery, and generation of host genomic deletions. *Retrovirology* *1*, 5.
- Wang, C. L., and Wabl, M. (2004). DNA acrobats of the Ig class switch. *J Immunol* *172*, 5815-5821.
- Wang, C. L., and Wabl, M. (2005a). Hypermutation rate normalized by chronological time. *J Immunol* *174*, 5650-5654.
- Wang, C. L., and Wabl, M. (2005b). Mutational activity in cell line WEHI-231. *Immunogenetics* *56*, 849-853.
- Wang, H., Kavanaugh, M. P., North, R. A., and Kabat, D. (1991). Cell-surface receptor for ecotropic murine retroviruses is a basic amino-acid transporter. *Nature* *352*, 729-731.
- Waterston, R. H., Lindblad-Toh, K., Birney, E., Rogers, J., Abril, J. F., Agarwal, P., Agarwala, R., Ainscough, R., Alexandersson, M., An, P., *et al.* (2002). Initial sequencing and comparative analysis of the mouse genome. *Nature* *420*, 520-562.
- Weigert, M. G., Cesari, I. M., Yonkovich, S. J., and Cohn, M. (1970). Variability in the lambda light chain sequences of mouse antibody. *Nature* *228*, 1045-1047.
- Weill, J. C., and Reynaud, C. A. (1992). Early B-cell development in chickens, sheep and rabbits. *Curr Opin Immunol* *4*, 177-180.
- White, K. D., Frank, M. B., Foundling, S., and Waxman, F. J. (1996). Effect of immunoglobulin variable region structure on C3b and C4b deposition. *Mol Immunol* *33*, 759-768.
- Winkler, M., aus Dem Siepen, T., and Stamminger, T. (2000). Functional interaction between pleiotropic transactivator pUL69 of human cytomegalovirus and the human homolog of yeast chromatin regulatory protein SPT6. *J Virol* *74*, 8053-8064.
- Winston, F. (1992). Analysis fo SPT genes: A genetic approach toward analysis of TFIID, histones, and other transcription factors of yeast. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY *In " Transcriptional Regulations"*, 1271-1293.
- Winston, F., and Carlson, M. (1992). Yeast SNF/SWI transcriptional activators and the SPT/SIN chromatin connection. *Trends Genet* *8*, 387-391.
- Woo, C. J., Martin, A., and Scharff, M. D. (2003). Induction of somatic hypermutation is associated with modifications in immunoglobulin variable region chromatin. *Immunity* *19*, 479-489.

Yoshikawa, K., Okazaki, I. M., Eto, T., Kinoshita, K., Muramatsu, M., Nagaoka, H., and Honjo, T. (2002). AID enzyme-induced hypermutation in an actively transcribed gene in fibroblasts. *Science* 296, 2033-2036.

Yu, K., Chedin, F., Hsieh, C. L., Wilson, T. E., and Lieber, M. R. (2003). R-loops at immunoglobulin class switch regions in the chromosomes of stimulated B cells. *Nat Immunol* 4, 442-451.

Zan, H., Komori, A., Li, Z., Cerutti, A., Schaffer, A., Flajnik, M. F., Diaz, M., and Casali, P. (2001). The translesion DNA polymerase zeta plays a major role in Ig and bcl-6 somatic hypermutation. *Immunity* 14, 643-653.

Zeng, X., Winter, D. B., Kasmer, C., Kraemer, K. H., Lehmann, A. R., and Gearhart, P. J. (2001). DNA polymerase eta is an A-T mutator in somatic hypermutation of immunoglobulin variable genes. *Nat Immunol* 2, 537-541.