

9 REFERENCES*

- Abdulmawjood, A. and Bülte, M. (2002):
Identification of ostrich meat by restriction fragment length polymorphism (RFLP) analysis of *cytochrome b* gene.
J Food Sci, 67 (5), 1688-1691.
- Aksoy, S.; Maudlin, I.; Dale, C.; Robinson, A.S. and O'Neill, S.L. (2001):
Prospects for control of African trypanosomiasis by tsetse vector manipulation.
Trends Parasitol, 17 (1), 29-34.
- Allen, M.; Engstrom, A.S.; Meyers, S.; Handt, O.; Saldeen, T.; von Haeseler, A.; Paabo, S. and Gyllensten, U. (1998):
Mitochondrial DNA sequencing of shed hairs and saliva on robbery caps: Sensitivity and matching probabilities.
J Forensic Sci, 43 (3), 453-464.
- Allmann, M.; Candrian, U.; Höfelein, C. and Lüthy, J. (1993):
Polymerase chain reaction (PCR): A possible alternative to immunochemical methods assuring safety and quality of food.
Z Lebensm Unters Forsch, 196, 248-251.
- Anderson, S.; Bankier, A.T.; Barrell, B.G.; de Bruijn, M.H.L.; Coulson, A.R.; Drouin, J.; Eperon, I.C.; Nierlich, D.P.; Roe, B.A.; Sanger, F.; Schreier, P.H.; Smith, A.J.H.; Staden, R. and Young, I.G. (1981):
Sequence and organization of the human mitochondrial genome.
Nature, 290, 457-465.
- Ansell, J.; Hu, J.T.; Gilbert, S.C.; Hamilton, K.A.; Hill, A.V. and Lindsay, S.W. (2000):
Improved method for distinguishing the human source of mosquito bloodmeals between close family members.
Trans R Soc Trop Med Hyg, 94 (5), 572-574.
- Bauer, B. and Wetzel, H. (1976):
A new membrane for feeding *Glossina morsitans* Westw. (Diptera, Glossinidae).
Bull Ent Res, 65, 563-565.
- Bauer, B. and Aigner, H. (1978):
In vitro maintenance of *Glossina palpalis palpalis* (Robineau-Desvoidy) (Diptera: Glossinidae).
Bull Ent Res, 68, 393-400.
- Bauer, B.; Amsler-Delafosse, S.; Clausen, P.-H.; Kabore, I. and Petrich-Bauer, J. (1995):
Successful application of deltamethrin pour on to cattle in a campaign against tsetse flies (*Glossina spp.*) in the pastoral zone of Samorogouan, Burkina Fasso.
Trop Med Parasitol, 46, 183-189.

References

- Baur, C.; Teifel-Greding, J. and Liebhardt, E. (1989):
Species identification by fragmented DNA.
Beitr Gerichtl Med, 47, 165-170.
- Beier, J.C.; Perkins, P.V.; Wirtz, R.A.; Koros, J.; Diggs, D.; Gargan II, T.P. and Koech, D.K. (1988):
Bloodmeal identification by direct enzyme-linked immunosorbent assay (ELISA), tested on *Anopheles* (Diptera: Culicidae) in Kenya.
J Med Entomol, 25 (1), 9-16.
- Bellagamba, F.; Moretti, V.M.; Comincini, S. and Valfré, F. (2001):
Identification of species in animal feedstuffs by polymerase chain reaction-restriction fragment length polymorphism analysis of mitochondrial DNA.
J Agric Food Chem, 49 (8), 3775-3781.
- Bender, K.; Schneider, P.M. and Rittner, C. (2000):
Application of mtDNA sequence analysis in forensic casework for the identification of human remains.
Forensic Sci Int, 113 (1-3), 103-107.
- Boakye, D.A.; Tang, J.; Truc, P.; Merriweather, A. and Unnasch, T.R. (1999):
Identification of bloodmeals in haematophagous Diptera by cytochrome B heteroduplex analysis.
Med Vet Entomol, 13, 282-287.
- Boore, J.L. (1999):
Animal mitochondrial genomes.
Nucleic Acids Res, 27 (8), 1767-1780.
- Boreham, P.F.L. (1976):
Sterilization of arthropod bloodmeals prior to bloodmeal identification.
Mosquito News, 36 (4), 454-457.
- Boreham, P.F.L. and Gill, G.S. (1973):
Serological identification of reptile feeds of *Glossina*.
Acta Trop, 30 (4), 354-365.
- Bossier, P. (1999):
Authentication of seafood products by DNA patterns.
J Food Sci, 64 (2), 189-193.
- Bottero, M.T.; Civera, T.; Anastasio, A.; Turi, R.M. and Rosati, S. (2002):
Identification of cow's milk in "buffalo" cheese by duplex polymerase chain reaction.
J Food Prot, 65 (2), 362-366.
- Branciari, R.; Nijman, I.J.; Plas, M.E.; Di Antonio, E. and Lenstra, J.A. (2000):
Species origin of milk in Italian mozzarella and Greek feta cheese.
J Food Prot, 63 (3), 408-411.

References

- Branicki, W; Kupiec, T. and Pawlowski, R. (2003):
Validation of cytochrome b sequence analysis as a method of species identification.
J Forensic Sci, 48 (1), 1-5.
- Buntjer, J.B.; Lenstra, J.A. and Haagsma, N. (1995):
Rapid species identification in meat by using satellite DNA probes.
Z Lebensm Unters Forsch, 201 (6), 577-582.
- Burger, J.; Schoon, R.; Zeike, B.; Hummel, S. and Herrmann, B. (2002):
Species determination using species-discriminating PCR-RFLP of ancient DNA from perhistoric skeletal remains.
Ancient Biomolecules, 4 (1), 19-23.
- Burkot, T.R.; Goodman, W.G. and DeFoliart, G.R. (1981):
Identification of Mosquito bloodmeals by enzyme-linked immunosorbent assay.
Am J Trop Med Hyg, 30 (6), 1336-1341.
- Calvo, J.H.; Rodellar, C.; Zaragoza, P. and Osta, R. (2002):
Beef- and bovine-derived material identification in processed and unprocessed food and feed by PCR amplification.
J Agric Food Chem, 50, 5262- 5264.
- Calvo, J.H.; Zaragoza, P. and Osta, R. (2001a):
Technical note: A quick and more sensitive method to identify pork in processed and unprocessed food by PCR amplification of a new specific DNA fragment.
J Anim Sci, 79, 2108-2112.
- Calvo, J.H.; Zaragoza, P. and Osta, R. (2001b):
Random amplified polymorphic DNA fingerprints for identification of species in poultry paâté.
Poult Sci, 80, 522-524.
- Carrera, E.; Garcia, T.; Céspedes, A.; González, I.; Fernández, A.; Hernández, P.E. and Martín, R. (1999):
Salmon and trout analysis by PCR-RFLP for identity authentication.
J Food Sci, 64 (3), 410-413.
- Castro, J.A.; Picornell, A. and Ramon, M. (1998):
Mitochondrial DNA: A tool for populational genetics studies.
Int Microbiol, 1, 327-332.
- Céspedes, A.; García, T.; Caarrera, E.; González, I.; Sanz, B.; Hernández, P.E. and Martín, R. (1998):
Identification of flatfish species using polymerase chain reaction (PCR) amplification and restriction analysis of the cytochrome b gene.
J Food Sci, 63 (2), 206-209.

References

- Chikuni, K.; Ozutsumi, K.; Koishikawa, T. and Kato, S. (1990):
Species identification of cooked meats by DNA hybridization assay.
Meat Sci, 27, 119-128.
- Chow-Shaffer, E.; Sina, B.; Hawley, W.A.; De Benedictis, J. and Scott, T.W. (2000):
Laboratory and field evolution of polymerase chain reaction-based forensic DNA profiling
for use in identification of human blood meal sources of *Aedes aegypti* (Diptera: Culicidae).
J Med Entomol, 37 (4), 492-502.
- Clausen, P.-H.; Adeyemi, I.; Bauer, B.; Breloer, M.; Salchow, F. and Staak, C. (1998):
Host preferences of tsetse (Diptera: Glossinidae) based on bloodmeal identifications.
Med Vet Entomol, 12, 169-180.
- Cocolin, L.; D'Agaro, E.; Manzano, M.; Lanari, D. and Comi, G. (2000):
Rapid PCR-RFLP method for the identification of marine fish fillets (Seabass, Seabream,
Umbrine and Dentex).
J Food Sci, 65 (8), 1315-1317.
- Colombo, F.; Viacava, R. and Giaretti, M. (2000):
Differentiation of the species ostrich (*Struthio camelus*) and emu (*Dromaius novaehollandiae*) by polymerase chain reaction using an ostrich-specific primer pair.
Meat Sci, 56 (1), 15-17.
- Cunningham, C.O. (2002):
Molecular diagnosis of fish and shellfish diseases: Present status and potential use in
disease control.
Aquaculture, 206 (1-2), 19-55.
- Deloach, J.R. and Taher, M. (1983):
Investigations on development of an artificial diet for in vitro rearing of *Glossina palpalis palpalis* (Diptera: Glossinidae).
J Econ Entomol, 76, 1112-1117.
- DeWoody, J.A.; Chesser, R.K. and Baker, R.J. (1999):
A translocated mitochondrial cytochrome b pseudogene in Voles (Rodentia: Microtus).
J Mol Evol, 48, 380-382.
- Dimo-Simonin, N.; Grange, F.; Taroni, F.; Brandt-Casadevall, C. and Mangin, P. (2000):
Forensic evolution of mtDNA in a population from south west Switzerland.
Int J Legal Med, 113, 89-97.
- Ebbehoj, K.F. and Thomsen, P.D. (1991a):
Species differentiation of heated meat products by DNA hybridization.
Meat Sci, 30, 221-234.
- Ebbehoj, K.F. and Thomsen, P.D. (1991b):
Differentiation of closely related species by DNA hybridization.
Meat Sci, 30, 359-366.

References

- Edrissian, G.H. and Hafizi, A. (1982):
Application of enzyme-linked immunosorbent assay (ELISA) to identification of *Anopheles* mosquito bloodmeals.
Trans R Soc Trop Med Hyg, 76 (1), 54-56.
- Elsen, P. Amoudi, M.A. and Leclercq, M. (1990):
First record of *Glossina fuscipes* Newstead, 1910 and *Glossina morsitans submorsitans* Newstead, 1910 in Southwestern Saudi Arabia.
Ann Soc Belg Med Trop, 70, 281-287.
- Emslie, V.W. and Steinberg, E.A. (1973):
The production in the bovine of antibody specific for closely related species and the use of such antisera in the identification of tsetse bloodmeals.
Ann Trop Med Parasitol, 67 (2), 213-217.
- Eposti, M.D.; De Vries, S.; Crimi, M.; Ghelli, A.; Patarnello, T. and Meyer, A. (1993):
Mitochondrial cytochrome b: Evolution and structure of the protein.
Biochim Biophys Acta, 1143 (3), 243-271.
- Ettore, R.; Vittorio, L. and Hoong, D.C. (1996):
Evolutionary genetics of the Suiformes as reconstructed using mtDNA sequencing.
J Mammalian Evol, 3 (2), 163-194.
- FAO (1982):
Training manual for tsetse control personnel. Vol.:1 Tsetse biology, systematic and distribution techniques (ed. by J.N. Pollock). Food and Agriculture Organization of the United Nations (FAO), Rome.
- FAO (1993):
Training manual for tsetse control personnel. Vol. 5: Insecticides for tsetse and trypanosomiasis control using attractive bait techniques. Food and Agriculture Organization (FAO), Rome.
- Farias, I.P.; Orti, G.; Sampaio, I.; Schneider, H. and Meyer, A. (2001):
The cytochrome b gene as a phylogenetic marker: The limits of resolution for analyzing relationships among Cichlid fishes.
J Mol Evol, 53, 89-103.
- Fernández, A.; García, T.; Asensio, L.; Rodríguez, M.Á.; González, I.; Céspedes, A.: Hernández, P.E. and Martín, R. (2000):
Identification of the Clam species *Ruditapes decussatus* (Grooved Carpet Shell), *Venerupis pullastra* (Pullet Carpet Shell) and *Ruditapes philippinarum* (Japanese Carpet Shell) by PCR-RFLP.
J Agric Food Chem, 48 (8), 3336-3341.
- Fernández, A.; García, T.; Asensio, L.; Rodríguez, M.Á.; González, I.; Hernández, P.E. and Martín, R. (2001):
PCR-RFLP analysis of the internal transcribed spacer (ITS) region for identification of 3 clam species.
J Food Sci, 66 (5), 657-661.

References

- Fleischer, R.C.; Perry, E.A.; Muralidharan, K.; Stevens, E.E. and Wemmer, C.M. (2001):
Phylogeography of the Asian elephant (*Elephas maximus*) based on mitochondrial DNA.
Evolution, 55 (9), 1882-1892.
- Ford, J. (1962):
Microclimates of tsetse fly resting sites in the Zambesi Valley, Southern Rhodesia.
ISCTR 9 th meeting Conakry, 165-170.
- Gibson, W.; Backhouse, T. and Griffiths, A. (2002):
The human serum resistance associated gene is ubiquitous and conserved in
Trypanosoma brucei rhodesiense throughout East Africa.
Infect Genet Evol, 1, 207-214.
- Gokool, S.; Curtis, C.F. and Smith, D.F. (1993):
Analysis of mosquito bloodmeals by DNA profiling.
Med Vet Entomol, 7, 208-215.
- Gomes, L.A.M.; Duarte, R.; Lima, D.C.; Diniz, B.S.; Serrão, M.L. and Labarthe, N. (2001):
Comparison between precipitin and ELISA tests in the bloodmeal detection of *Aedes aegypti* (Linnaeus) and *Aedes vexans* (Lutz) mosquitoes experimentally fed on feline, canine and human hosts.
Mem Inst Oswaldo Cruz, Rio de Janeiro, 96 (5), 693-695.
- Goossens, B.; Waits, L.P. and Taberlet, P. (1998):
Plucked hair samples as a source of DNA: Reliability of dinucleotide microsatellite genotyping.
Mol Ecol, 7, 1237-1241.
- Greenwood, A.D. and Pääbo, S. (1999):
Nuclear insertion sequences of mitochondrial DNA predominate in hair but not in blood of elephants.
Mol Ecol, 8, 133-137.
- Guzow-Krzeminska, B.; Gorniak, M. and Wegrzyn, G. (2001):
Molecular determination keys: Construction of keys for species identification based on restriction fragment length polymorphism.
Int Arch Biosci, 1057-1067.
- Härlid, A.; Janke, A. and Arnason, U. (1997):
The mtDNA sequence of the ostrich and the divergence between paleognathous and neognathous birds.
Mol Biol Evol, 14 (7), 754-761.
- Herman, B.L. (2001):
Determination of the animal origin of raw food by species-specific PCR.
J Dairy Res, 68 (3), 429-436.

References

- Hershfield, B.; Chader, G. and Aguirre, G. (1994):
A polymerase chain reaction-based method for the identification of DNA samples from common vertebrate species.
Electrophoresis, 15 (7), 880-884.
- Hiendleder, S.; Lewalski, H.; Wassmuth, R. and Janke, A. (1998):
The complete mitochondrial DNA sequence of the domestic sheep (*Ovis aries*) and comparison with other major ovine haplotype.
J Mol Evol, 47, 441-448.
- Higuchi, R.; von Beroldingen, C.H.; Sensabaugh, G.F. and Erlich, H.A. (1988):
DNA typing from single hairs.
Nature, 332, 543-546.
- Higuchi, R. (1989):
Simple and rapid preparation of samples for PCR.
In: Ehrlich (Ed.), PCR Technology, Stockton Press, New York, 31-38.
- Hold, G.L.; Russell, V.J.; Pryde, S.E.; Rehbein, H.; Quinteiro, J.; Vidal, R.; Rey-Mendez, M.; Sotelo, C.G.; Pérez-Martin, R.I.; Santos, A.T. and Rosa, C. (2001):
Development of a DNA-based method aimed at identifying the fish species present in food products.
J Agric Food Chem, 49 (3), 1175-1179.
- Hsieh, H.M.; Chiang, H.L.; Tsai, L.C.; Lai, S.Y.; Huang, N.E.; Linacre, A. and Lee, J.C. (2001):
Cytochrome b gene for species identification of the conservation animals.
Forensic Sci Int, 122, (1), 7-18.
- Hühne, J.; Pfeiffer, H.; Waterkamp, K. and Brinkmann, B. (1999):
Mitochondrial DNA in human hair shafts - existence of intra-individual differences?
Int J Legal Med, 112, 172-175.
- Ishida, N.; Hasegawa, T.; Oyunsuren, T. and Mukoyama, H. (1996):
PCR-RFLP analysis of the cytochrome b gene in horse mitochondrial DNA.
Anim Genet, 27 (5), 359-363.
- Janke, A. and Arnason, U. (1997):
The complete mitochondrial genome of *Alligator mississippiensis* and the separation between recent Archosauria (Birds and Crocodiles).
Mol Biol Evol, 14, (12), 1266-1272.
- Katsumi, A.; Isao, M.; Isao, I.; Akio, K.; Kojiro, N.; Takuya, K. and Motoaki, K. (1997):
Phylogenetic relationship of caprini estimated by cytochrome b gene sequence analysis.
Anim Sci Technol, 68 (2), 148-155.

References

- Kengne, P.; Trung, H.D.; Baimai, V.; Coosemans, M. and Manguin, S. (2001):
A multiplex PCR-based method derived from random amplified polymorphic DNA (RAPD) markers for the identification of species of the *Anopheles minimus* group in Southeast Asia.
Insect Mol Biol, 10 (5), 427-435.
- Kirstein, F. and Gray, J.S. (1996):
A molecular marker for the identification of the zoonotic reservoirs of lyme borreliosis by analysis of blood meal in its European vector *Ixodes ricinus*.
Appl Environ Microbiol, Nov., 4060-4065.
- Kocher, T.D.; Thomas, W.K.; Meyer, A.; Edwards, S.V.; Pääbo, S.; Villablanca, F.X. and Wilson, A.C. (1989):
Dynamics of mitochondrial DNA evolution in animals: Amplification and sequencing with conserved primers.
Proc Natl Acad Sci USA, 86, 6196-6200.
- Leak, S.G.A. (1999):
Tsetse biology and ecology: Their role in the epidemiology and control of trypanosomosis.
CABI publishing, UK, pp.9, 81, 119.
- Lebedeva, I.A.; Kulikov, E.E.; Ivanova, N.V. and Poltaraus, A.B. (2000):
Mitochondrial DNA sequencing of human hair shafts stored for long time.
Sud Med Ekspert, 43 (3), 9-14.
- Lenstra, J.A.; Buntjer, J.B. and Janssen, F.W. (2001):
On the origin of meat - DNA techniques for species identification in meat products.
Vet Sci Tomorrow, 2, 1-13.
- Lin, Y.S; Poh, Y.P; Lin, S.M. and Tzeng, C.S. (2002):
Molecular techniques to identify freshwater eels: RFLP analyses of PCR-amplified DNA fragments and allele-specific PCR from mitochondrial DNA.
Zoological Studies, 41 (4), 421-430.
- Linville, J.G. and Wells, J.D. (2002):
Surface sterilization of a maggot using bleach does not interfere with mitochondrial DNA analysis of crop contents.
J Forensic Sci, 47 (5), 1-5.
- Lockley, A.K. and Bardsley, R.G. (2000):
Novel method for the discrimination of tuna (*Thunnus thynnus*) and bonito (*Sarda sarda*) DNA.
J Agric Food Chem, 48 (10), 4463-4468.
- Lombardi, S. and Esposito, F. (1983):
Enzyme-linked immunosorbent assay (ELISA) for the identification of mosquito bloodmeals.
Parasitologia, 25 (1), 49-56.

References

- Lovette, I.J.; Bermingham, E.; Rohwer, S. and Wood, C. (1999):
Mitochondrial restriction fragment length polymorphism (RFLP) and sequence variation among closely related avian species and the genetic characterization of hybrid *Dendroica* warblers.
Mol Ecol, 8, 1431-1441.
- Luis, C.; Bastos-Silveira, C.; Cothran, E.G. and Oom, M.M. (2002):
Variation in the mitochondrial control region sequence between the two maternal lines of the Sorraia horse breed.
Genet Mol Biol, 25 (3), 309-311.
- Lutz, S.; Weisser, H.J.; Heizmann, J. and Pollak, S. (1996):
MtDNA as a tool for identification of human remains. Identification using mtDNA.
Int J Legal Med, 109 (4), 205-209.
- Mackie, I.M.; Pryde, S.E.; Gonzales-Sotelo, C.; Medina, I.; Peréz-Martín, R.; Quinteiro, J.; Rey-Mendez, M. and Rehbein, H. (1999):
Challenges in the identification of species of canned fish.
Trends Food Sci Technol, 10, 9-14.
- Masri, S.; Rast, H.; Ripley, T.; James, D.; Green, M.; Jia, X. and Devlin, R.H. (2002):
Detection of genetically modified coho salmon using polymerase chain reaction (PCR) amplification.
J Agric Food Chem, 50 (11), 3161-3164.
- Matsunaga, T.; Chikuni, K.; Tanabe, R.; Muroya, S.; Shibata, K.; Yamada, J. and Shinmura, Y. (1999):
A quick and simple method for the identification of meat species and meat products by PCR assay.
Meat Sci, 51, 143-148.
- Maudet, C. and Taberlet, P. (2001):
Detection of cows' milk in goats' cheeses inferred from mitochondrial DNA polymorphism.
J Dairy Res, 68 (2), 229-235.
- McDowell, J. and Graves, J. (2002):
Nuclear and mitochondrial DNA markers for specific identification of istiophorid and xiphiid billfishes.
Fish Bull, 100 (3), 537-544.
- Mehlitz, D. Brinkmann, U. and Haller, L. (1982):
Epidemiological studies on the animal reservoir of gambiense sleeping sickness.
Tropenmed Parasitol, 32, 129-133.
- Meyer, R. and Candrian, U. (1996):
PCR-based DNA analysis for the identification and characterization of food components.
Lebensmittel-Wissenschaft und-Technologie/FST, 29 (1-2), 1-9.

References

- Meyer, R.; Candrian, U. and Lüthy, J. (1994):
Detection of pork in heated meat products by polymerase chain reaction (PCR).
J AOAC Int, 77 (3), 617-622.
- Meyer, R.; Höfelein, C.; Lüthy, J. and Candrian, U. (1995):
Polymerase chain reaction-restriction fragment length polymorphism analysis: A simple method for species identification in food.
J AOAC Int, 78 (6), 1542-1551.
- Mihok, S.; Otieno, L.H. and Darji, N. (1990):
Population genetics of *Trypanosoma brucei* and the epidemiology of human sleeping sickness in the Lambwe Valley, Kenya.
Parasitology, 100, 219-233.
- Millar, C.D.; Taylor, G.A.; Moore, P.J.; Halverson, J.L. and Lambert, D.M. (2000):
A novel restriction fragment length polymorphism for petrels or tube-nosed seabirds.
Mol Ecol, 9, 1915-1917.
- Mindell, D.P.; Sorenson, M.D. and Dimcheff, D.E. (1998):
Multiple independent origins of mitochondrial gene order in birds.
Proc Natl Acad Sci USA, 95, 10693-10697.
- Mohamed-Ahmed, M.M. and Odulaja, A. (1997):
Deil activity patterns and host preferences of *Glossina fuscipes fuscipes* (Diptera: Glossinidae) along the shores of Lake Victoria, Kenya.
Bull Ent Res, 87, 179-186.
- Montiel-Sosa, J.F.; Ruiz-Pesini, E.; Montoya, J.; Roncalés, P.; López-Pérez, M.J. and Pérez-Martos, A. (2000):
Direct and highly species-specific detection of pork meat and fat in meat products by PCR amplification of mitochondrial DNA.
J Agric Food Chem, 48 (7), 2829-2832.
- Mukabana, W.R.; Takken, W. and Knols, B.G.J. (2002):
Analysis of arthropod bloodmeals using molecular genetic markers.
Trends Parasitol, 18 (11), 505-509.
- Mullis, K.B. and Faloona, F.A. (1987):
Specific synthesis of DNA in vitro via polymerase-catalyzed chain reaction.
Methods Enzymol, 155, 335-350.
- Murray M. and Dexter, T.M. (1988):
Anaemia in bovine african trypanosomiasis.
Acta Trop, 45, 389-432.
- Murray, B.W.; McClymont, R.A. and Strobeck, C. (1995):
Forensic identification of ungulate species using restriction digests of PCR-amplified mitochondrial DNA.
J Forensic Sci, 40 (6), 943-951.

References

- Murray, M. and Gray, A.R. (1984):
The current situation on animal trypanosomiasis in Africa.
Prev Vet Med, 2, 23-30.
- Newton, C.R. and Graham, A. (1997):
PCR Second Edition, Springer-Verlag New York, pp. 1, 113, 115.
- Ngo, K.A. and Kramer, L.D. (2003):
Identification of mosquito bloodmeals using polymerase chain reaction (PCR) with order-specific primers.
J Med Entomol, 40 (2), 215-222.
- Ngumbi, P.M.; Lawyer, P.G.; Johnson, R.N.; Kiilu, G. and Asiago, C. (1992):
Identification of phlebotomine sand fly bloodmeals from Baringo District, Kenya, by direct enzyme-linked immunosorbent assay (ELISA).
Med Vet Entomol, 6, 385-388.
- Nguyen, T.T.T.; Murphy, N.P. and Austin, C.M. (2002):
Amplification of multiple copies of mitochondrial cytochrome b gene fragments in the Australian freshwater crayfish, *Cherax destructor* Clark (Parastacidae: Decapoda).
Int Soc Anim Genet, 33, 304-308.
- Okoth, J.O. and Kapaata, R. (1988):
The hosts of *Glossina fuscipes fuscipes* (Newstead) in Busoga, Uganda, and epidemiological implications for trypanosomiasis.
Ann Trop Med Parasitol, 82 (5), 517-518.
- Pääbo, S (1989):
Ancient DNA: Extraction, characterization, molecular cloning and enzymatic amplification.
Proc Natl Acad Sci, USA, 86, 1939-1943.
- Parson, W.; Pegoraro, K.; Niederstätter, H.; Föger, M. and Steinlechner, M. (2000):
Species identification by means of the cytochrome b gene.
Int J Legal Med, 114 (1-2), 23-28.
- Partis, L.; Croan, D.; Guo, Z.; Clark, R.; Coldham, T. and Murby, J. (2000):
Evaluation of a DNA fingerprinting method for determining the species origin of meats.
Meat Sci, 54 (4), 369-376.
- Pfeiffer, H.; Hühne, J.; Ortmann, C. and Waterkamp, K. (1999):
Mitochondrial DNA typing from human axillary, pubic and head hair shafts- success rates and sequence comparisons.
Int J Legal Med, 112, 287-290.
- Pierpaoli, M.; Riga, F.; Trocchi, V. and Randi, E. (1999):
Species distinction and evolutionary relationships of the Italian hare (*Lepus corsicanus*) as described by mitochondrial DNA sequencing.
Mol Ecol, 8 (11), 1805-1817.

References

- Prychitko, T.M. and Moore, W.S. (2000): Comparative evolution of the mitochondrial cytochrome b gene and nuclear β -fibrinogen intron7 in woodpeckers. *Mol Biol Evol*, 17, 1101-1111.
- Quinteiro, J.; Vidal, R.; Izquierdo, M.; Sotelo, C.G.; Chapela, M.J.; Pérez-Martín, R.I.; Rehbein, H.; Hold, G.L.; Russell, V.J.; Pryde, S.E.; Rosa, C.; Santos, A.T. and Rey-Méndez, M. (2001): Identification of Hake species (*Merluccius* Genus) using sequencing and PCR-RFLP analysis of mitochondrial DNA control region sequences. *J Agric Food Chem*, 49 (11), 5108-5114.
- Radostits, O.M.; Gay, C.C.; Blood, D.C. and Hinchcliff, K.W. (2000): A textbook of the diseases of cattle, sheep, pigs, goats and horses. WB Saunders, London, 9th ed. pp.1329-1337.
- Rajapaksha, W.R.A.K.J.S.; Thilakaratne, I.D.S.I.P.; Chandrasiri, A.D.N. and Niroshan, T.D. (2002): Development of PCR assay for differentiation of some important wild animal meat of Sri Lanka. *J Vet Med B*, 49, 322-324.
- Rego, I.; Martínez, A.; González-Tizón, A.; Vieites, J.; Leira, F. and Méndez, J. (2002): PCR technique for identification of mussel species. *J Agric Food Chem*, 50 (7), 1780-1784.
- Rephogle, J.; Lord, W.D.; Budowle, B.; Meinking, T.L. and Taplin, D. (1994): Identification of host DNA by amplified fragment length polymorphism analysis: Preliminary analysis of human crab louse (*Anoplura: Pediculidae*) excreta. *J Med Entomol*, 31 (5), 686-690.
- Ron, M.; Verner, N.; Feldmesser, E.; Hochman, D.; Band, M. and Shani, M. (1996): Amplification of the conserved cytochrome-b locus as a versatile internal control for PCR analysis in animals. *Biotechniques*, 20 (4), 604-608.
- Rurangirwa, F.R.; Minja, S.H.; Musoke, A.J.; Nantulya, V.M.; Grootenhuis, J.G. and Moloo, S.K. (1986): Production and evaluation of specific antisera against sera of various vertebrate species for identification of bloodmeals of *Glossina morsitans*. *Acta Trop*, 43 (4), 379-389.
- Russell, V.J.; Hold, G.L.; Pryde, S.E.; Rehbein, H.; Quinteiro, J.; Rey-Mendez, M.; Sotelo, C.G.; Pérez-Martin, R.I.; Santos, A.T. and Rosa, C. (2000): Use of restriction fragment length polymorphism to distinguish between salmon species. *J Agric Food Chem*, 48 (6), 2184-2188.

References

- Saiki, R.K.; Gelfand, D.H.; Stoffel, S.; Scharf, S.J.; Higuchi, R.; Horn, G.T.; Mullis, K.B. and Erlich, H.A. (1988):
Primer-directed enzymatic amplification of DNA with a thermostable DNA polymerase.
Science, 239, 487-491.
- Savelkoul, P.H.M.; Aarts, H.J.M.; DeHaas, J.; Dijkshoorn, L.; Duim, B.; Otsen, M.; Rademaker, J.L.W.; Schouls, L. and Lenstra, J.A. (1999):
Amplified-fragment length polymorphism analysis: The state of an art.
J Clin Microbiol, 37 (10), 3083-3091.
- Schneider, P.M.; Seo, Y. and Rittner, C. (1999):
Forensic mtDNA hair analysis excludes a dog from having caused a traffic accident.
Int J Legal Med, 112, 315-316.
- Sebastio, P.; Zanelli, P. and Neri, T.M. (2001):
Identification of Anchovy (*Engraulis encrasicholus* L.) and Gilt Sardine (*Sardinella aurita*) by polymerase chain reaction, sequence of their mitochondrial cytochrome b gene, and restriction analysis of polymerase chain reaction products in semipreserves.
J Agric Food Chem, 49, 1194-1199.
- Service, M.W.; Voller, A. and Bidwell, D.E. (1986):
The enzyme-linked immunosorbent assay (ELISA) test for identification of bloodmeals of haematophagous insects.
Bull Ent Res, 76, 321-330.
- Sotelo, C.G.; Calo-Mata, P.; Chapela, M.J.; Pérez-Martín, R.I.; Rehbein, H.; Hold, G.L.; Russell, V.J.; Pryde, S.; Quinteiro, J.; Izquierdo, M.; Rey-Méndez, M.; Rosa, C. and Santos, A.T. (2001):
Identification of flatfish (Pleuronectiforme) species using DNA-based techniques.
J Agric Food Chem, 49 (10), 4562-4569.
- Späth, J. (2000):
Feeding patterns of three sympatric tsetse species (*Glossina spp.*) (Diptera: Glossinidae) in the preforest zone of Côte d'Ivoire.
Acta Trop, 75 (1), 109-118.
- Staak, C.; Allmang, B.; Kämpe, U. and Mehlitz, D. (1981):
The complement fixation test for the species identification of bloodmeals from tsetse flies.
Tropenmed Parasit, 32, 97-98.
- Staak, C.; Kämpe, U. and Korkowski, G. (1986):
Species identification of bloodmeals from tsetse flies (Glossinidae): Results 1979-1985.
Trop Med Parasit, 37, 59-60.
- Stich, A.; Abel, P.M. And Krishna, S. (2002):
Human African trypanosomosis.
BMJ, 325, 203-206.

References

- Stuart, C. and Stuart, T. (1997):
The larger mammals of Africa.
1st Ed., Kyodo printing Co. (Singapore) Pte Ltd. pp. 16-223.
- Sullivan, K.M.; Alliston-Greiner, R.; Archampong, F.I.A.; Piercy, R.; Tully, G.; Gill, P. and Liroy-Davies, C. (2002):
A single difference in mtDNA control region sequence observed between hair shaft and reference samples from a single donor.
<http://www.promega.com/geneticidproc/ussymp7proc/0721.html>
- Sun, Y.L. and Lin, C.S. (2003):
Establishment and application of a fluorescent polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) method for identifying porcine, caprine and bovine meats.
J Agric Food Chem, 51 (7), 1771-1776.
- Tartaglia, M.; Saulle, E.; Pestalozza, S.; Morelli, L.; Antonucci, G. and Battaglia, P.A. (1998):
Detection of bovine mitochondrial DNA in ruminant feeds: A molecular approach test for the presence of bovine-derived materials.
J Food Prot, 61 (5), 513-518.
- Tempelis, C.H. (1975):
Host-feeding patterns of mosquitoes, with a review of advances in analysis of blood-meals by serology.
J Med Entomol, 11 (6), 635-653.
- Tobolewski, J.; Kaliszewski, M.J.; Colwell, R.K. and Oliver, J.R. (1992):
Detection and identification of mammalian DNA from the gut of museum specimens of ticks.
J Med Entomol, 29 (6), 1049-1051.
- Torr, S.J.; Wilson, P.J.; Schofield, S.; Mangwiyo, T.N.C.; Akber, S. and White, B.N. (2001):
Application of DNA markers to identify the individual-specific hosts of tsetse feeding on cattle.
Med Vet Entomol, 15, 78-86.
- Ursing, B.M. and Arnason, U. (1998a):
The complete mitochondrial DNA sequence of the pig (*Sus scrofa*).
J Mol Evol, 47, 302-306.
- Ursing, B.M. and Arnason, U. (1998b):
Analyses of mitochondrial genomes strongly support a hippopotamus-whale clade.
Proc R Soc Lond B, 265, 2251-2255.
- Vigilant, L. (1999):
An evaluation of techniques for the extraction and amplification of DNA from naturally shed hairs.
Biol Chem, 380, 1329-1331.

References

- Wang, H.Y.; Tsai, M.P.; Tu, M.C. and Lee, S.C. (2000):
Universal primers for amplification of the complete mitochondrial 12S rRNA gene in
vertebrates.
Zoological Studies, 39 (1), 61-66.
- Weitz, B. (1960):
Feeding habits of bloodsucking arthropods.
Exp Parasitol, 9, 63-82.
- Weitz, B. (1963):
The feeding habits of *Glossina*.
Bull Wld Hlth Org, 28, 711-729.
- Welburn, S.C. and Maudlin, I. (1999):
Tsetse-trypanosome interactions: Rites of passage.
Parasitology Today, 15 (10), 399-403.
- Welburn, S.C.; Févre, E.M.; Coleman, P.G.; Odiit, M. and Maudlin, I. (2001):
Sleeping sickness: A tale of two diseases.
Trends Parasitol, 17 (1), 19-24.
- Wells, J.D.; Intronà, F.Jr.; Di Vella, G.; Campobasso, C.P.; Hayes, J. and Sperling, F.A.H.
(2001):
Human and insect mitochondrial DNA analysis from maggots.
J Forensic Sci, 46 (3), 685-687.
- WHO (1988):
Vector control series, tsetse flies, training and information guide.
World Health Organization (WHO) vector biology and control division, Geneva.
- Wintero, A.K.; Thomsen, P.D. and Davies, W. (1990):
A comparison of DNA-hybridization, immunodiffusion, countercurrent immunoelectrophoresis and isoelectric focusing for detecting the admixture of pork to beef.
Meat Sci, 27, 75-85.
- Wolf, C.; Hübner, P. and Lüthy, J. (1999 a):
Differentiation of sturgeon species by PCR-RFLP.
Food Res Int, 32, 699-705.
- Wolf, C.; Rentsch, J. and Hübner, P. (1999 b):
PCR-RFLP analysis of mitochondrial DNA: A reliable method for species identification.
J Agric Food Chem, 47 (4), 1350-1355.
- Yoon, J.M. and Kim, G.W. (2001):
Randomly amplified polymorphic DNA-polymerase chain reaction analysis of two
different populations of cultured Korean catfish *Silurus asotus*.
J Biosci, 26 (5), 641-647.

References

- Yoshii, T.; Tamura, K. and Ishiyama, I. (1992):
Presence of a PCR inhibitor in hairs.
Nippon Hoigaku Zasshi, 46 (5), 313-316.
- Yoshii, T.; Tamura, K.; Taniguchi, T.; Akiyama, K. and Ishiyama, I. (1993):
Water-soluble eumelanin as a PCR-inhibitor and a simple method for its removal.
Nippon Hoigaku Zasshi, 47 (4), 323-329.
- Zehner, R.; Zimmermann, S. and Mebs, D. (1998):
RFLP and sequence analysis of the cytochrome b gene of selected animals and man:
Methodology and forensic application.
Int J Legal Med, 111, 323-327.

* The magazines were shortened after:

List of journals indexed in Index Medicus 2002, National Library of Medicine, U.S.
Department of Health and Human Services, Bethesda, Maryland, 2002.