

Literaturverzeichnis

- Accili, E. A., Kiehn, J., Yang, Q., Wang, Z., Brown, A. M., und Wible, B. A. (1997). Separable Kvbeta subunit domains alter expression and gating of potassium channels. **J Biol Chem**, 272(41):25824-31.
- Adachi, S., Uchida, S., Ito, H., Hata, M., Hiroe, M., Marumo, F., und Sasaki, S. (1994). Two isoforms of a chloride channel predominantly expressed in thick ascending limb of Henle's loop and collecting ducts of rat kidney. **J Biol Chem**, 269(26):17677-17683.
- Adrian, R. H. und Marshall, M. W. (1976). Action potentials reconstructed in normal and myotonic muscle fibres. **J Physiol (Lond)**, 258(1):125-43.
- al Awqati, Q. (1995). Chloride channels of intracellular organelles. **Curr Opin Cell Biol**, 7(4):504-8.
- Amling, M., Neff, L., Priemel, M., Schilling, A. F., Rueger, J. M., und Baron, R. (2000). Progressive increase in bone mass and development of odontomas in aging osteopetrotic c-src-deficient mice. **Bone**, 27(5):603-10.
- Anderson, M. P., Gregory, R. J., Thompson, S., Souza, D. W., Paul, S., Mulligan, R. C., Smith, A. E., und Welsh, M. J. (1991). Demonstration that CFTR is a chloride channel by alteration of its anion selectivity. **Science**, 253(5016):202-5.
- Ashcroft, F. M. (2000). **Ion Channels and Diseases**. Academic Press.
- Baron, R., Neff, L., Brown, W., Courtoy, P. J., Louvard, D., und Farquhar, M. G. (1988). Polarized secretion of lysosomal enzymes: co-distribution of cation-independent mannose-6-phosphate receptors and lysosomal enzymes along the osteoclast exocytic pathway. **J Cell Biol**, 106(6):1863-72.
- Baron, R., Neff, L., Louvard, D., und Courtoy, P. J. (1985). Cell-mediated extracellular acidification and bone resorption: evidence for a low pH in resorbing lacunae and localization of a 100-kD lysosomal membrane protein at the osteoclast ruffled border. **J Cell Biol**, 101(6):2210-22.
- Bensaoula, T., Shibuya, H., Katz, M. L., Smith, J. E., Johnson, G. S., John, S. K., und Milam, A. H. (2000). Histopathologic and immunocytochemical analysis of the retina and ocular tissues in Batten disease. **Ophthalmology**, 107(9):1746-53.
- Berkelaar, M., Clarke, D. B., Wang, Y. C., Bray, G. M., und Aguayo, A. J. (1994). Axotomy results in delayed death and apoptosis of retinal ganglion cells in adult rats. **J Neurosci**, 14(7):4368-74.
- Betz, H., Kuhse, J., Schmieden, V., Laube, B., Kirsch, J., und Harvey, R. J. (1999). Structure and functions of inhibitory and excitatory glycine receptors. **Ann N Y Acad Sci**, 868:667-76.
- Birnboim, H. C. und Doly, J. (1979). A rapid alkaline extraction procedure for screening recombinant plasmid DNA. **Nucleic Acids Res**, 7(6):1513-23.
- Blair, H. C., Teitelbaum, S. L., Tan, H. L., Koziol, C. M., und Schlesinger, P. H. (1991). Passive chloride permeability charge coupled to H(+)-ATPase of avian osteoclast ruffled membrane. **Am J Physiol**, 260(6 Pt 1):C1315-24.

- Boyce, B. F., Yoneda, T., Lowe, C., Soriano, P., und Mundy, G. R. (1992). Requirement of pp60c-src expression for osteoclasts to form ruffled borders and resorb bone in mice. *J Clin Invest*, 90(4):1622-7.
- Brandt, S. (1997). **Klonierung und Charakterisierung neuer Mitglieder der ClC-Chloridkanal-Familie.** PhD thesis, Christian-Albrechts-Universität.
- Brandt, S. und Jentsch, T. J. (1995). ClC-6 and ClC-7 are two novel broadly expressed members of the ClC chloride channel family. *FEBS Lett*, 377(1):15-20.
- Bösl, M., Stein, V., Hübner, C., Zdebik, A., Jordt, S., Mukhopadhyay, A., Davidoff, M., Holstein, A., und Jentsch, T. (2001). Male germ cells and photoreceptors, both depending on close cell-cell interactions, degenerate upon ClC-2 Cl-channel disruption. **EMBO (im Druck).**
- Buckwalter, M. S., Cook, S. A., Davisson, M. T., White, W. F., und Camper, S. A. (1994). A frameshift mutation in the mouse alpha 1 glycine receptor gene (Gla1) results in progressive neurological symptoms and juvenile death. *Hum Mol Genet*, 3(11):2025-30.
- Bumsted, K. M. und Barnstable, C. J. (2000). Dorsal retinal pigment epithelium differentiates as neural retina in the microphthalmia (mi/mi) mouse. *Invest Ophthalmol Vis Sci*, 41(3):903-8.
- Buyse, G., Trouet, D., Voets, T., Missiaen, L., Droogmans, G., Nilius, B., und Eggermont, J. (1998). Evidence for the intracellular location of chloride channel (ClC)-type proteins: co-localization of ClC-6a and ClC-6c with the sarco/endoplasmic-reticulum Ca²⁺ pump SERCA2b. *Biochem J*, 330(Pt 2):1015-1021.
- Chuang, J. Z., Milner, T. A., Zhu, M., und Sung, C. H. (1999). A 29 kDa intracellular chloride channel p64H1 is associated with large dense-core vesicles in rat hippocampal neurons. *J Neurosci*, 19(8):2919-28.
- Counillon, L. und Pouyssegur, J. (2000). The expanding family of eucaryotic Na⁽⁺⁾/H⁽⁺⁾ exchangers. *J Biol Chem*, 275(1):1-4.
- Cully, D. F., Vassilatis, D. K., Liu, K. K., Pares, P. S., Van der Ploeg, L. H., Schaeffer, J. M., und Arena, J. P. (1994). Cloning of an avermectin-sensitive glutamate-gated chloride channel from *Caenorhabditis elegans*. *Nature*, 371(6499):707-11.
- Davis-Kaplan, S. R., Askwith, C. C., Bengtzen, A. C., Radisky, D., und Kaplan, J. (1998). Chloride is an allosteric effector of copper assembly for the yeast multicopper oxidase Fet3p: an unexpected role for intracellular chloride channels. *Proc Natl Acad Sci U S A*, 95(23):13641-5.
- D'Cruz, P. M., Yasumura, D., Weir, J., Matthes, M. T., Abderrahim, H., LaVail, M. M., und Vollrath, D. (2000). Mutation of the receptor tyrosine kinase gene *Mertk* in the retinal dystrophic RCS rat. *Hum Mol Genet*, 9(4):645-51.
- Deng, C. und Capecchi, M. R. (1992). Reexamination of gene targeting frequency as a function of the extent of homology between the targeting vector and the target locus. *Mol Cell Biol*, 12(8):3365-71.
- Dougall, W. C., Glaccum, M., Charrier, K., Rohrbach, K., Brasel, K., De Smedt, T., Daro, E., Smith, J., Tometsko, M. E., Maliszewski, C. R., Armstrong, A., Shen, V., Bain, S., Cosman, D., Anderson, D., Morrissey, P. J., Peshon, J. J., und Schuh, J. (1999). RANK is essential for osteoclast and lymph node development. *Genes Dev*, 13(18):2412-24.
- Ducy, P., Schinke, T., und Karsenty, G. (2000). The osteoblast: a sophisticated fibroblast under central surveillance. *Science*, 289(5484):1501-4.
- Dulhunty, A., Gage, P., Curtis, S., Chelvanayagam, G., und Board, P. (2001). The Glutathione Transferase Structural Family Includes a Nuclear Chloride Channel and a Ryanodine

- Receptor Calcium Release Channel Modulator. **J. Biol. Chem.**, 276(5):3319-3323.
- Edwards, J. C. (1999). A novel p64-related Cl⁻ channel: subcellular distribution and nephron segment-specific expression. **Am J Physiol**, 276(3 Pt 2):F398-408.
- Elster, A. D., Theros, E. G., Key, L. L., und Chen, M. Y. (1992). Cranial imaging in autosomal recessive osteopetrosis. Part II. Skull base and brain. **Radiology**, 183(1):137-44.
- Fahlke, C. (2000). Molecular mechanisms of ion conduction in ClC-type chloride channels: lessons from disease-causing mutations. **Kidney Int**, 57(3):780-6.
- Felix, R., Cecchini, M. G., Hofstetter, W., Elford, P. R., Stutzer, A., und Fleisch, H. (1990). Impairment of macrophage colony-stimulating factor production and lack of resident bone marrow macrophages in the osteopetrotic op/op mouse. **J Bone Miner Res**, 5(7):781-9.
- Finnemann, S. C., Bonilha, V. L., Marmorstein, A. D., und Rodriguez-Boulan, E. (1997). Phagocytosis of rod outer segments by retinal pigment epithelial cells requires alpha(v)beta5 integrin for binding but not for internalization. **Proc Natl Acad Sci U S A**, 94(24):12932-7.
- Forgac, M. (1998). Structure, function and regulation of the vacuolar (H⁺)-ATPases. **FEBS Lett**, 440(3):258-63.
- Frattini, A., Orchard, P. J., Sobacchi, C., Giliani, S., Abinun, M., Mattsson, J. P., Keeling, D. J., Andersson, A. K., Wallbrandt, P., Zecca, L., Notarangelo, L. D., Vezzoni, P., und Villa, A. (2000). Defects in TCIRG1 subunit of the vacuolar proton pump are responsible for a subset of human autosomal recessive osteopetrosis. **Nat Genet**, 25(3):343-6.
- Friedrich, T., Breiderhoff, T., und Jentsch, T. J. (1999). Mutational analysis demonstrates that ClC-4 and ClC-5 directly mediate plasma membrane currents. **J Biol Chem**, 274(2):896-902.
- Fritsch, G. und Hitzig, E. (1870). Über die elektrische Erregbarkeit des Großhirns. **Arch.Anat.Physiol.Wiss.med.**, 37:300-332.
- Gerritsen, E. J., Vossen, J. M., van Loo, I. H., Hermans, J., Helfrich, M. H., Griscelli, C., und Fischer, A. (1994). Autosomal recessive osteopetrosis: variability of findings at diagnosis and during the natural course. **Pediatrics**, 93(2):247-253.
- Günther, W., Lüchow, A., Cluzeaud, F., Vandewalle, A., und Jentsch, T. J. (1998). ClC-5, the chloride channel mutated in Dent's disease, colocalizes with the proton pump in endocytotically active kidney cells. **Proc Natl Acad Sci U S A**, 95(14):8075-80.
- Goebel, H. H. (1984). Morphology of the gangliosidoses. **Neuropediatrics**, 15 Suppl(4):97-106.
- Goebel, H. H. (1992). Neuronal ceroid lipofuscinoses: the current status. **Brain Dev**, 14(4):203-11.
- Goebel, H. H., Schochet, S. S., Jaynes, M., und Gutmann, L. (1998). Ultrastructure of the retina in adult neuronal ceroid lipofuscinosis. **Acta Anat**, 162(2-3):127-32. Using Smart Source Parsing.
- Golabek, A. A., Kida, E., Walus, M., Kaczmarek, W., Michalewski, M., und Wisniewski, K. E. (2000). CLN3 protein regulates lysosomal pH and alters intracellular processing of Alzheimer's amyloid-beta protein precursor and cathepsin D in human cells. **Mol Genet Metab**, 70(3):203-13.
- Greene, J. R., Brown, N. H., DiDomenico, B. J., Kaplan, J., und Eide, D. J. (1993). The GEF1 gene of *Saccharomyces cerevisiae* encodes an integral membrane protein; mutations in which have effects on respiration and iron-limited growth. **Mol Gen Genet**, 241(5-6):542-53.
- Gründer, S., Thiemann, A., Pusch, M., und Jentsch, T. J. (1992). Regions involved in the

- opening of CIC-2 chloride channel by voltage and cell volume. **Nature**, 360(6406):759-762.
- Hall, N. A., Lake, B. D., Dewji, N. N., und Patrick, A. D. (1991). Lysosomal storage of subunit c of mitochondrial ATP synthase in Batten's disease (ceroid-lipofuscinosis). **Biochem J**, 275(Pt 1)(12):269-72.
- Haskell, R. E., Carr, C. J., Pearce, D. A., Bennett, M. J., und Davidson, B. L. (2000). Batten disease: evaluation of CLN3 mutations on protein localization and function. **Hum Mol Genet**, 9(5):735-44.
- Hayman, A. R., Jones, S. J., Boyde, A., Foster, D., Colledge, W. H., Carlton, M. B., Evans, M. J., und Cox, T. M. (1996). Mice lacking tartrate-resistant acid phosphatase (Acp 5) have disrupted endochondral ossification and mild osteopetrosis. **Development**, 122(10):3151-62.
- Heaney, C., Shalev, H., Elbedour, K., Carmi, R., Staack, J. B., Sheffield, V. C., und Beier, D. R. (1998). Human autosomal recessive osteopetrosis maps to 11q13, a position predicted by comparative mapping of the murine osteosclerosis (oc) mutation. **Hum Mol Genet**, 7(9):1407-10.
- Hechenberger, M., Schwappach, B., Fischer, W. N., Frommer, W. B., Jentsch, T. J., und Steinmeyer, K. (1996). A family of putative chloride channels from Arabidopsis and functional complementation of a yeast strain with a CLC gene disruption. **J Biol Chem**, 271(52):33632-33638.
- Heiss, N. S. und Poustka, A. (1997). Genomic structure of a novel chloride channel gene, CLIC2, in Xq28. **Genomics**, 45(1):224-8.
- Hodgkin, A., Huxley, A., und Katz, B. (1949). Ionic currents underlying activity in the giant axon of the squid. **Arch Sci Physiol**, 3:129-50.
- Hollmann, M., Hartley, M., und Heinemann, S. (1991). Ca²⁺ permeability of KA-AMPA-gated glutamate receptor channels depends on subunit composition. **Science**, 252(5007):851-3.
- Hryciw, D. H., Rychkov, G. Y., Hughes, B. P., und Bretag, A. H. (1998). Relevance of the D13 region to the function of the skeletal muscle chloride channel, CLC-1. **J Biol Chem**, 273(8):4304-7.
- Hummler, E., Barker, P., Talbot, C., Wang, Q., Verdumo, C., Grubb, B., Gatzky, J., Burnier, M., Horisberger, J. D., Beermann, F., Boucher, R., und Rossier, B. C. (1997). A mouse model for the renal salt-wasting syndrome pseudohypoaldosteronism. **Proc Natl Acad Sci U S A**, 94(21):11710-5.
- Ismailov, I., Awayda, M. S., Jovov, B., Berdiev, B. K., Fuller, C. M., Dedman, J. R., Kaetzel, M., und Benos, D. J. (1996). Regulation of epithelial sodium channels by the cystic fibrosis transmembrane conductance regulator. **J Biol Chem**, 271(9):4725-32.
- Jaissner, F. und Beggah, A. T. (1999). The non-gastric H⁺-K⁺-ATPases: molecular and functional properties. **Am J Physiol**, 276(6 Pt 2):F812-24.
- Jarvela, I., Lehtovirta, M., Tikkanen, R., Kyttala, A., und Jalanko, A. (1999). Defective intracellular transport of CLN3 is the molecular basis of Batten disease (JNCL). **Hum Mol Genet**, 8(6):1091-8.
- Jentsch, T. J., Friedrich, T., Schriever, A., und Yamada, H. (1999). The CLC chloride channel family. **Pflugers Arch**, 437(6):783-795.
- Jentsch, T. J., Steinmeyer, K., und Schwarz, G. (1990). Primary structure of Torpedo marmorata chloride channel isolated by expression cloning in Xenopus oocytes. **Nature**, 348(6301):510-514.
- Joyner, A. (2000). **Gene Targeting. A Practical Approach**. IRL Press at Oxford University Press, 2nd edition.

-
- Kamb, A., Iverson, L. E., und Tanouye, M. A. (1987). Molecular characterization of Shaker, a *Drosophila* gene that encodes a potassium channel. **Cell**, 50(3):405-13.
- Karet, F. E., Finberg, K. E., Nelson, R. D., Nayir, A., Mocan, H., Sanjad, S. A., Rodriguez-Soriano, J., Santos, F., Cremers, C. W., Di Pietro, A., Hoffbrand, B. I., Winiarski, J., Bakaloglu, A., Ozen, S., Dusunsel, R., Goodyer, P., Hulton, S. A., Wu, D. K., Skvorak, A. B., Morton, C. C., Cunningham, M. J., Jha, V., und Lifton, R. P. (1999). Mutations in the gene encoding B1 subunit of H⁺-ATPase cause renal tubular acidosis with sensorineural deafness. **Nat Genet**, 21(1):84-90.
- Keith, C. G. (1968). Retinal atrophy in osteopetrosis. **Arch Ophthalmol**, 79(3):234-41.
- Ketchum, K. A., Joiner, W. J., Sellers, A. J., Kaczmarek, L. K., und Goldstein, S. A. (1995). A new family of outwardly rectifying potassium channel proteins with two pore domains in tandem. **Nature**, 376(6542):690-5.
- Kieferle, S., Fong, P., Bens, M., Vandewalle, A., und Jentsch, T. J. (1994). Two highly homologous members of the ClC chloride channel family in both rat and human kidney. **Proc Natl Acad Sci U S A**, 91(15):6943-6947.
- Koike, M., Nakanishi, H., Saftig, P., Ezaki, J., Ishihara, K., Ohsawa, Y., Schulz-Schaeffer, W., Watanabe, T., Waguri, S., Kametaka, S., Shibata, M., Yamamoto, K., Kominami, E., Peters, C., von Figura, K., und Uchiyama, Y. (2000). Cathepsin D deficiency induces lysosomal storage with ceroid lipofuscin in mouse CNS neurons. **J Neurosci**, 20(18):6898-906.
- Kong, Y. Y., Yoshida, H., Sarosi, I., Tan, H. L., Timms, E., Capparelli, C., Morony, S., Oliveira-dos Santos, A. J., Van, G., Itie, A., Khoo, W., Wakeham, A., Dunstan, C. R., Lacey, D. L., Mak, T. W., Boyle, W. J., und Penninger, J. M. (1999). OPGL is a key regulator of osteoclastogenesis, lymphocyte development and lymph-node organogenesis. **Nature**, 397(6717):315-23.
- Kornak, U. (1996). Aufklärung der genomischen Struktur des Gens für den Chloridkanal mClC-7 und Konstruktion zweier Targeting-Vektoren zum Ausschalten dieses Gens in embryonalen Stammzellen der Maus. **Diplomarbeit, Fachbereich Chemie, Freie Universität Berlin**.
- Kornak, U., Bosl, M. R., und Kubisch, C. (1999). Complete genomic structure of the CLCN6 and CLCN7 putative chloride channel genes(1). **Biochim Biophys Acta**, 1447(1):100-6.
- Kornak, U., Schulz, A., Friedrich, W., Uhlhaas, S., Kremens, B., Voit, T., Hasan, C., Bode, U., Jentsch, T. J., und Kubisch, C. (2000). Mutations in the $\alpha 3$ subunit of the vacuolar H⁽⁺⁾-ATPase cause infantile malignant osteopetrosis. **Hum Mol Genet**, 9(13):2059-63.
- Laemmli, U. K. (1970). Cleavage of structural proteins during the assembly of the head of bacteriophage T4. **Nature**, 227(259):680-5.
- Landry, D., Sullivan, S., Nicolaides, M., Redhead, C., Edelman, A., Field, M., al Awqati, Q., und Edwards, J. (1993). Molecular cloning and characterization of p64, a chloride channel protein from kidney microsomes. **J Biol Chem**, 268(20):14948-55.
- Lazner, F., Gowen, M., Pavasovic, D., und Kola, I. (1999). Osteopetrosis and osteoporosis: two sides of the same coin. **Hum Mol Genet**, 8(10):1839-46.
- Lehmann-Horn, F. und Jurkat-Rott, K. (1999). Voltage-gated ion channels and hereditary disease. **Physiol Rev**, 79(4):1317-72.
- Li, Y. P., Chen, W., Liang, Y., Li, E., und Stashenko, P. (1999). Atp6i-deficient mice exhibit severe osteopetrosis due to loss of osteoclast-mediated extracellular acidification. **Nat Genet**, 23(4):447-51.

- Li, Y. P., Chen, W., and Stashenko, P. (1996). Molecular cloning and characterization of a putative novel human osteoclast-specific 116-kDa vacuolar proton pump subunit. **Biochem Biophys Res Commun**, 218(3):813-21.
- Lloyd, S. E., Pearce, S. H., Fisher, S. E., Steinmeyer, K., Schwappach, B., Scheinman, S. J., Harding, B., Bolino, A., Devoto, M., Goodyer, P., Rigden, S. P., Wrong, O., Jentsch, T. J., Craig, I. W., and Thakker, R. V. (1996). A common molecular basis for three inherited kidney stone diseases. **Nature**, 379(6564):445-449.
- Lomaga, M. A., Yeh, W. C., Sarosi, I., Duncan, G. S., Furlonger, C., Ho, A., Morony, S., Capparelli, C., Van, G., Kaufman, S., van der Heiden, A., Itie, A., Wakeham, A., Khoo, W., Sasaki, T., Cao, Z., Penninger, J. M., Paige, C. J., Lacey, D. L., Dunstan, C. R., Boyle, W. J., Goeddel, D. V., and Mak, T. W. (1999). TRAF6 deficiency results in osteopetrosis and defective interleukin-1, CD40, and LPS signaling. **Genes Dev**, 13(8):1015-24.
- Ludewig, U., Pusch, M., and Jentsch, T. J. (1996). Two physically distinct pores in the dimeric ClC-0 chloride channel. **Nature**, 383(6598):340-343.
- Lukacs, G. L., Chang, X. B., Bear, C., Kartner, N., Mohamed, A., Riordan, J. R., and Grinstein, S. (1993). The delta F508 mutation decreases the stability of cystic fibrosis transmembrane conductance regulator in the plasma membrane. Determination of functional half-lives on transfected cells. **J Biol Chem**, 268(29):21592-8.
- Luzio, J. P., Rous, B. A., Bright, N. A., Pryor, P. R., Mullock, B. M., and Piper, R. C. (2000). Lysosome-endosome fusion and lysosome biogenesis. **J Cell Sci**, 113(Pt 9)(5):1515-24.
- Mansour, S. L., Thomas, K. R., Deng, C. X., and Capecchi, M. R. (1990). Introduction of a lacZ reporter gene into the mouse int-2 locus by homologous recombination. **Proc Natl Acad Sci U S A**, 87(19):7688-92.
- Marshansky, V. und Vinay, P. (1996). Proton gradient formation in early endosomes from proximal tubules. **Biochim Biophys Acta**, 1284(2):171-80.
- Matsumura, Y., Uchida, S., Kondo, Y., Miyazaki, H., Ko, S. B., Hayama, A., Morimoto, T., Liu, W., Arisawa, M., Sasaki, S., und Marumo, F. (1999). Overt nephrogenic diabetes insipidus in mice lacking the CLC-K1 chloride channel. **Nat Genet**, 21(1):95-98.
- Mattsson, J. P., Skyman, C., Palokangas, H., Vaananen, K. H., und Keeling, D. J. (1997). Characterization and cellular distribution of the osteoclast ruffled membrane vacuolar H⁺-ATPase B-subunit using isoform-specific antibodies. **J Bone Miner Res**, 12(5):753-60.
- McHugh, K. P., Hodivala-Dilke, K., Zheng, M. H., Namba, N., Lam, J., Novack, D., Feng, X., Ross, F. P., Hynes, R. O., und Teitelbaum, S. L. (2000). Mice lacking beta3 integrins are osteosclerotic because of dysfunctional osteoclasts. **J Clin Invest**, 105(4):433-40.
- Midell, J., Maduke, M., Miller, C., und Grigorieff, N. (2001). Projection structure of a ClC-type chloride channel at 6.5 Å resolution. **Nature**, 409:219-223.
- Miller, C. (1982). Open-state substructure of single chloride channels from Torpedo electroplax. **Philos Trans R Soc Lond B Biol Sci**, 299(1097):401-11.
- Moore, K. (1995). Insight into the microphthalmia gene. **Trends Genet**, 11(11):442-448.
- Mullock, B. M., Perez, J. H., Kuwana, T., Gray, S. R., und Luzio, J. P. (1994). Lysosomes can fuse with a late endosomal compartment in a cell-free system from rat liver. **J Cell Biol**, 126(5):1173-82.
- Nakamura, I., Takahashi, N., Udagawa, N., Moriyama, Y., Kurokawa, T., Jimi, E., Sasaki, T., und Suda, T. (1997). Lack of vacuolar proton ATPase association with the cytoskeleton in osteoclasts of osteosclerotic (oc/oc) mice. **FEBS Lett**, 401(2-3):207-12.

-
- Neher, E. und Sakmann, B. (1976). Single-channel currents recorded from membrane of denervated frog muscle fibres. **Nature**, 260(5554):799-802.
- Nishizawa, T., Nagao, T., Iwatsubo, T., Forte, J. G., und Urushidani, T. (2000). Molecular cloning and characterization of a novel chloride intracellular channel-related protein, parchorin, expressed in water-secreting cells. **J Biol Chem**, 275(15):11164-73.
- Noda, M., Shimizu, S., Tanabe, T., Takai, T., Kayano, T., Ikeda, T., Takahashi, H., Nakayama, H., Kanaoka, Y., Minamino, N., und et al. (1984). Primary structure of Electrophorus electricus sodium channel deduced from cDNA sequence. **Nature**, 312(5990):121-7.
- Obermüller, N., Gretz, N., Kriz, W., Reilly, R. F., und Witzgall, R. (1998). The swelling-activated chloride channel CIC-2, the chloride channel CIC-3, and CIC-5, a chloride channel mutated in kidney stone disease, are expressed in distinct subpopulations of renal epithelial cells. **J Clin Invest**, 101(3):635-42.
- Ohkuma, S., Moriyama, Y., und Takano, T. (1982). Identification and characterization of a proton pump on lysosomes by fluorescein-isothiocyanate-dextran fluorescence. **Proc Natl Acad Sci U S A**, 79(9):2758-62.
- Palmer, S., Perry, J., und Ashworth, A. (1995). A contravention of Ohno's law in mice [see comments]. **Nat Genet**, 10(4):472-6.
- Palokangas, H., Mulari, M., und Väänänen, H. K. (1997). Endocytic pathway from the basal plasma membrane to the ruffled border membrane in bone-resorbing osteoclasts. **J Cell Sci**, 110(Pt 15):1767-80.
- Papazian, D. M., Timpe, L. C., Jan, Y. N., und Jan, L. Y. (1991). Alteration of voltage-dependence of Shaker potassium channel by mutations in the S4 sequence. **Nature**, 349(6307):305-10.
- Perzov, N., Nelson, H., und Nelson, N. (2000). Altered distribution of the yeast plasma membrane H⁺-ATPase as a feature of vacuolar H⁺-ATPase null mutants. **J Biol Chem**, 275(51):40088-95.
- Piwon, N., Gunther, W., Schwake, M., Bosl, M. R., und Jentsch, T. J. (2000). CIC-5 Cl⁻-channel disruption impairs endocytosis in a mouse model for Dent's disease [In Process Citation]. **Nature**, 408(6810):369-73.
- Ponting, C. P. (1997). CBS domains in CIC chloride channels implicated in myotonia and nephrolithiasis (kidney stones). **J Mol Med**, 75(3):160-3.
- Presley, J. F., Mayor, S., McGraw, T. E., Dunn, K. W., und Maxfield, F. R. (1997). Bafilomycin A1 treatment retards transferrin receptor recycling more than bulk membrane recycling. **J Biol Chem**, 272(21):13929-36.
- Pusch, M., Jordt, S. E., Stein, V., und Jentsch, T. J. (1999). Chloride dependence of hyperpolarization-activated chloride channel gates. **J Physiol (Lond)**, 515(Pt 2):341-53.
- Pusch, M., Ludewig, U., Rehfeldt, A., und Jentsch, T. J. (1995). Gating of the voltage-dependent chloride channel CIC-0 by the permeant anion. **Nature**, 373(6514):527-531.
- Qian, Z., Okuhara, D., Abe, M. K., und Rosner, M. R. (1999). Molecular cloning and characterization of a mitogen-activated protein kinase-associated intracellular chloride channel. **J Biol Chem**, 274(3):1621-7.
- Rabow, L. E., Russek, S. J., und Farb, D. H. (1995). From ion currents to genomic analysis: recent advances in GABAA receptor research. **Synapse**, 21(3):189-274.
- Rajendra, S., Lynch, J. W., Pierce, K. D., French, C. R., Barry, P. H., und Schofield, P. R. (1994). Startle disease mutations reduce the agonist sensitivity of the human inhibitory glycine receptor. **J Biol Chem**, 269(29):18739-42.

- Ranganathan, R., Cannon, S. C., und Horvitz, H. R. (2000). MOD-1 is a serotonin-gated chloride channel that modulates locomotory behaviour in *C. elegans* [In Process Citation]. **Nature**, 408(6811):470-5.
- Riordan, J. R., Rommens, J. M., Kerem, B., Alon, N., Rozmahel, R., Grzelczak, Z., Zielenski, J., Lok, S., Plavsic, N., Chou, J. L., und et al. (1989). Identification of the cystic fibrosis gene: cloning and characterization of complementary DNA [published erratum appears in *Science* 1989 Sep 29;245(4925):1437]. **Science**, 245(4922):1066-73.
- Romey, G., Attali, B., Chouabe, C., Abitbol, I., Guillemare, E., Barhanin, J., und Lazdunski, M. (1997). Molecular mechanism and functional significance of the MinK control of the KvLQT1 channel activity. **J Biol Chem**, 272(27):16713-6.
- Rothman, J. H., Yamashiro, C. T., Raymond, C. K., Kane, P. M., und Stevens, T. H. (1989). Acidification of the lysosome-like vacuole and the vacuolar H⁺-ATPase are deficient in two yeast mutants that fail to sort vacuolar proteins. **J Cell Biol**, 109(1):93-100.
- Saftig, P., Hetman, M., Schmahl, W., Weber, K., Heine, L., Mossmann, H., Koster, A., Hess, B., Evers, M., von Figura, K., und et al. (1995). Mice deficient for the lysosomal proteinase cathepsin D exhibit progressive atrophy of the intestinal mucosa and profound destruction of lymphoid cells. **Embo J**, 14(15):3599-608.
- Saftig, P., Hunziker, E., Wehmeyer, O., Jones, S., Boyde, A., Rommerskirch, W., Moritz, J. D., Schu, P., und von Figura, K. (1998). Impaired osteoclastic bone resorption leads to osteopetrosis in cathepsin-K-deficient mice. **Proc Natl Acad Sci U S A**, 95(23):13453-8.
- Salo, J., Lehenkari, P., Mulari, M., Metsikko, K., und Vaananen, H. K. (1997). Removal of osteoclast bone resorption products by transcytosis [see comments]. **Science**, 276(5310):270-3.
- Sambrook, J., Fritsch, E., und Maniatis, T. (1989). **Molecular Cloning: A Laboratory Manual**. Cold Spring Harbour Press, 2. edition.
- Sanger, F., Nicklen, S., und Coulson, A. R. (1977). DNA sequencing with chain-terminating inhibitors. **Proc Natl Acad Sci U S A**, 74(12):5463-7.
- Schlesinger, P. H., Blair, H. C., Teitelbaum, S. L., und Edwards, J. C. (1997). Characterization of the osteoclast ruffled border chloride channel and its role in bone resorption. **J Biol Chem**, 272(30):18636-43.
- Schmidt-Rose, T. und Jentsch, T. J. (1997a). Reconstitution of functional voltage-gated chloride channels from complementary fragments of CLC-1. **J Biol Chem**, 272(33):20515-21.
- Schmidt-Rose, T. und Jentsch, T. J. (1997b). Transmembrane topology of a CLC chloride channel. **Proc Natl Acad Sci U S A**, 94(14):7633-7638.
- Schriever, A. M., Friedrich, T., Pusch, M., und Jentsch, T. J. (1999). CLC chloride channels in *Caenorhabditis elegans*. **J Biol Chem**, 274(48):34238-44.
- Schwappach, B., Stobrawa, S., Hechenberger, M., Steinmeyer, K., und Jentsch, T. J. (1998). Golgi localization and functionally important domains in the NH₂ and COOH terminus of the yeast CLC putative chloride channel Gef1p. **J Biol Chem**, 273(24):15110-8.
- Schwartzberg, P. L., Xing, L., Hoffmann, O., Lowell, C. A., Garrett, L., Boyce, B. F., und Varmus, H. E. (1997). Rescue of osteoclast function by transgenic expression of kinase-deficient Src in src^{-/-} mutant mice. **Genes Dev**, 11(21):2835-44.
- Scimeca, J. C., Franchi, A., Trojani, C., Parrinello, H., Grosgeorge, J., Robert, C., Jailion, O., Poirier, C., Gaudray, P., und Carle, G. F. (2000). The gene encoding the

-
- mouse homologue of the human osteoclast-specific 116-kDa V-ATPase subunit bears a deletion in osteosclerotic (oc/oc) mutants. **Bone**, 26(3):207-13.
- Seeliger, M., Ruther, K., Apfelstedt-Sylla, E., Schlote, W., Wohlrab, M., und Zrenner, E. (1997). [Juvenile neuronal ceroid lipofuscinosis (Batten-Mayou) disease. Ophthalmologic diagnosis and findings]. **Ophthalmologie**, 94(8):557-62.
- Simon, D. B., Bindra, R. S., Mansfield, T. A., Nelson-Williams, C., Mendonca, E., Stone, R., Schurman, S., Nayir, A., Alpay, H., Bakaloglu, A., Rodriguez-Soriano, J., Morales, J. M., Sanjad, S. A., Taylor, C. M., Pilz, D., Brem, A., Trachtman, H., Griswold, W., Richard, G. A., John, E., und Lifton, R. P. (1997). Mutations in the chloride channel gene, CLCNKB, cause Bartter's syndrome type III. **Nat Genet**, 17(2):171-178.
- Simon, D. B., Karet, F. E., Rodriguez-Soriano, J., Hamdan, J. H., DiPietro, A., Trachtman, H., Sanjad, S. A., und Lifton, R. P. (1996). Genetic heterogeneity of Bartter's syndrome revealed by mutations in the K⁺ channel, ROMK. **Nat Genet**, 14(2):152-6.
- Simpson, E., Linder, C., Sargent, E., Davisson, M., Mobraaten, L., und Sharp, J. (1997). Genetic variation among 129 substrains and its importance for targeted mutagenesis in mice. **NatGenet**, 16(1):19-27.
- Sly, W. S., Hewett-Emmett, D., Whyte, M. P., Yu, Y. S., und Tashian, R. E. (1983). Carbonic anhydrase II deficiency identified as the primary defect in the autosomal recessive syndrome of osteopetrosis with renal tubular acidosis and cerebral calcification. **Proc Natl Acad Sci U S A**, 80(9):2752-6.
- Smith, A. N., Skaug, J., Choate, K. A., Nayir, A., Bakaloglu, A., Ozen, S., Hulton, S. A., Sanjad, S. A., Al-Sabban, E. A., Lifton, R. P., Scherer, S. W., und Karet, F. E. (2000). Mutations in ATP6N1B, encoding a new kidney vacuolar proton pump 116-kD subunit, cause recessive distal renal tubular acidosis with preserved hearing. **Nat Genet**, 26(1):71-5.
- Snyders, D. J. (1999). Structure and function of cardiac potassium channels. **Cardiovasc Res**, 42(2):377-90.
- Soriano, P., Montgomery, C., Geske, R., und Bradley, A. (1991). Targeted disruption of the c-src proto-oncogene leads to osteopetrosis in mice. **Cell**, 64(4):693-702.
- Steinmeyer, K., Klocke, R., Ortland, C., Grone-meier, M., Jockusch, H., Gründer, S., und Jentsch, T. J. (1991a). Inactivation of muscle chloride channel by transposon insertion in myotonic mice. **Nature**, 354(6351):304-308.
- Steinmeyer, K., Ortland, C., und Jentsch, T. J. (1991b). Primary structure and functional expression of a developmentally regulated skeletal muscle chloride channel. **Nature**, 354(6351):301-4.
- Stobrawa, S. M., Breiderhoff, T., Takamori, S., Engel, D., Schweizer, M., Zdebik, A. A., Bösl, M. R., Ruether, K., Jahn, H., Draguhn, A., und Jentsch, T. (2001). Disruption of ClC-3, a chloride channel expressed on synaptic vesicles, leads to a loss of the hippocampus. **Neuron**, 29 (in press).
- Szewczyk, A. (1998). The intracellular potassium and chloride channels: properties, pharmacology and function (review). **Mol Membr Biol**, 15(2):49-58.
- Teitelbaum, S. L. (2000). Bone resorption by osteoclasts. **Science**, 289(5484):1504-8.
- Thomas, K. R. und Capecchi, M. R. (1987). Site-directed mutagenesis by gene targeting in mouse embryo-derived stem cells. **Cell**, 51(3):503-12.
- Tiffée, J. C., Xing, L., Nilsson, S., und Boyce, B. F. (1999). Dental abnormalities associated with failure of tooth eruption in src knockout and op/op mice. **Calcif Tissue Int**, 65(1):53-8.

- Tjelle, T. E., Brech, A., Juvet, L. K., Griffiths, G., und Berg, T. (1996). Isolation and characterization of early endosomes, late endosomes and terminal lysosomes: their role in protein degradation. *J Cell Sci*, 109(Pt 12)(16):2905-14.
- Tondravi, M. M., McKercher, S. R., Anderson, K., Erdmann, J. M., Quiroz, M., Maki, R., und Teitelbaum, S. L. (1997). Osteopetrosis in mice lacking haematopoietic transcription factor PU.1. *Nature*, 386(6620):81-4.
- Toyomura, T., Oka, T., Yamaguchi, C., Wada, Y., und Futai, M. (2000). Three subunit α isoforms of mouse vacuolar H⁽⁺⁾-ATPase. Preferential expression of the $\alpha 3$ isoform during osteoclast differentiation [In Process Citation]. *J Biol Chem*, 275(12):8760-5.
- Tulk, B. M. und Edwards, J. C. (1998). NCC27, a homolog of intracellular Cl⁻ channel p64, is expressed in brush border of renal proximal tubule. *Am J Physiol*, 274(6 Pt 2):F1140-9.
- Tyynela, J., Palmer, D. N., Baumann, M., und Haltia, M. (1993). Storage of saposins A and D in infantile neuronal ceroid-lipofuscinosis. *FEBS Lett*, 330(1):8-12.
- Uchida, S., Sasaki, S., Furukawa, T., Hiraoka, M., Imai, T., Hirata, Y., und Marumo, F. (1993). Molecular cloning of a chloride channel that is regulated by dehydration and expressed predominantly in kidney medulla [published erratum appears in *J Biol Chem* 1994 Jul 22;269(29):19192]. *J Biol Chem*, 268(6):3821-3824.
- Udagawa, N., Sasaki, T., Akatsu, T., Takahashi, N., Tanaka, S., Tamura, T., Tanaka, H., und Suda, T. (1992). Lack of bone resorption in osteosclerotic (oc/oc) mice is due to a defect in osteoclast progenitors rather than the local microenvironment provided by osteoblastic cells. *Biochem Biophys Res Commun*, 184(1):67-72.
- Udagawa, N., Takahashi, N., Akatsu, T., Sasaki, T., Yamaguchi, A., Kodama, H., Martin, T. J., und Suda, T. (1989). The bone marrow-derived stromal cell lines MC3T3-G2/PA6 and ST2 support osteoclast-like cell differentiation in cocultures with mouse spleen cells. *Endocrinology*, 125(4):1805-13.
- Udy, G. B., Parkes, B. D., und Wells, D. N. (1997). ES cell cycle rates affect gene targeting frequencies. *Exp Cell Res*, 231(2):296-301.
- Valenzuela, S. M., Martin, D. K., Por, S. B., Robbins, J. M., Warton, K., Bootcov, M. R., Schofield, P. R., Campbell, T. J., und Breit, S. N. (1997). Molecular cloning and expression of a chloride ion channel of cell nuclei. *J Biol Chem*, 272(19):12575-82.
- van Weert, A. W., Dunn, K. W., Gueze, H. J., Maxfield, F. R., und Stoorvogel, W. (1995). Transport from late endosomes to lysosomes, but not sorting of integral membrane proteins in endosomes, depends on the vacuolar proton pump. *J Cell Biol*, 130(4):821-34.
- Väänänen, H. K., Zhao, H., Mulari, M., und Halleen, J. M. (2000). The cell biology of osteoclast function. *J Cell Sci*, 113(Pt 3):377-381.
- Waldegger, S. und Jentsch, T. J. (2000). Functional and structural analysis of ClC-K chloride channels involved in renal disease. *J Biol Chem*, 275(32):24527-33.
- Wang, Z. Q., Ovitt, C., Grigoriadis, A. E., Mohle-Steinlein, U., Ruther, U., und Wagner, E. F. (1992). Bone and haematopoietic defects in mice lacking *c-fos*. *Nature*, 360(6406):741-5.
- Weinreich, F. und Jentsch, T. J. (2000). Pores formed by single subunits in mixed dimers of different ClC chloride channels. *J Biol Chem*.
- Whitaker, J. N. und Rhodes, R. H. (1983). The distribution of cathepsin D in rat tissues determined by immunocytochemistry. *Am J Anat*, 166(4):417-28.
- Wollnik, B., Kubisch, C., Steinmeyer, K., und Pusch, M. (1997). Identification of functionally important regions of the muscular

chloride channel CIC-1 by analysis of recessive and dominant myotonic mutations. **Hum Mol Genet**, 6(5):805-11.