



Ideas and perspectives: Biogeochemistry – some key foci for the future

Thomas S. Bianchi¹, Madhur Anand², Chris T. Bauch³, Donald E. Canfield⁴, Luc De Meester^{5,6,7}, Katja Fennel⁸, Peter M. Groffman^{9,13}, Michael L. Pace¹⁰, Mak Saito¹¹, and Myrna J. Simpson¹²

¹Dept. of Geological Science, University of Florida, Gainesville, FL, USA

²School of Environmental Sciences, University of Guelph, Guelph, Ontario, Canada

³University of Waterloo, Department of Applied Mathematics, Waterloo, Canada

⁴Nordcee, University of Southern Denmark, Odense, Denmark

⁵Dept. of Biology, University of Leuven, Leuven, Belgium

⁶Leibniz Institut für Gewässerökologie und Binnenfischerei (IGB), Berlin, Germany

⁷Institute of Biology, Freie Universität Berlin, Berlin, Germany

⁸Dept. of Oceanography, Dalhousie University, Halifax, Nova Scotia, Canada

⁹City University of New York Advanced Science Research Center at the Graduate Center, New York, NY, USA

¹⁰Dept. of Environmental Sciences, University of Virginia, Charlottesville, VA, USA

¹¹Marine Chemistry and Geochemistry, Woods Hole Oceanographic Institution, Woods Hole, MA, USA

¹²Dept. of Physical and Environmental Sciences, University of Toronto, Toronto, Canada

¹³Cary Institute of Ecosystem Studies, Millbrook, NY, USA

Correspondence: Thomas S. Bianchi (tbianchi@ufl.edu)

Received: 24 October 2020 – Discussion started: 4 November 2020

Revised: 30 March 2021 – Accepted: 1 April 2021 – Published: 19 May 2021

Abstract. Biogeochemistry has an important role to play in many environmental issues of current concern related to global change and air, water, and soil quality. However, reliable predictions and tangible implementation of solutions, offered by biogeochemistry, will need further integration of disciplines. Here, we refocus on how further developing and strengthening ties between biology, geology, chemistry, and social sciences will advance biogeochemistry through (1) better incorporation of mechanisms, including contemporary evolutionary adaptation, to predict changing biogeochemical cycles, and (2) implementing new and developing insights from social sciences to better understand how sustainable and equitable responses by society are achieved. The challenges for biogeochemists in the 21st century are formidable and will require both the capacity to respond fast to pressing issues (e.g., catastrophic weather events and pandemics) and intense collaboration with government officials, the public, and internationally funded programs. Keys to success will be the degree to which biogeochemistry can make biogeochemical knowledge more available to policy makers

and educators about predicting future changes in the biosphere, on timescales from seasons to centuries, in response to climate change and other anthropogenic impacts. Biogeochemistry also has a place in facilitating sustainable and equitable responses by society.

1 Introduction

Biogeochemistry was one of the first truly inter- or multi-disciplinary sciences (Bianchi, 2020; Gorham, 1991; Schlesinger, 1991; Vernadsky et al., 1926) and the field continues to expand in multiple directions at an amazing pace; from small scales via interactions with microbiology and “omics” approaches (genomics, transcriptomics, proteomics, and metabolomics) (Fig. 1) to large scales as a component of Earth system sciences (Steffen et al., 2020).

A recent review of biogeochemistry by Bianchi (2020), reflects on these more nascent linkages in molecular biology and their historical and disparate connections. Countries




Figure 1. The interdisciplinary field of biogeochemistry has advanced with expanded approaches that involve the integration of core biogeochemical areas with other disciplines needed to understand a rapidly changing Earth system and meet the needs for sustainability.

around the world scramble with new health and sociopolitical challenges ranging from global climate change to loss of biodiversity, changing ecosystems, and global pandemics (e.g., coronavirus disease 2019 (COVID-19)) – often linked with the growing human population. Addressing these challenges requires new cross-disciplinary approaches by scientists – coupled with a better science-educated public that are more involved with decision-making on sustainability issues (e.g., Derouin, 2020). Some may argue these are “old” questions/challenges previously debated in the academic circles of Earth science, ecology, ecosystem science, biogeochemistry, and the like. However, we contend that humanity now sits at a *Silent Spring* moment with the attention of the global public – in large part due to COVID-19 and recent catastrophic weather events. Recognition of earlier “moments” of environmental crisis led to the formation of organizations like the World Climate Research Programme (WCRP), International Geosphere-Biosphere Programme (IGBP), and Intergovernmental Panel on Climate Change (IPCC), to name a few. Kress et al. (2020) posit that

Now is the time to use the full power of science through cooperative efforts among initiatives such as BIOSCAN, the Global Virome Project (GVP), and Earth BioGenome Project (EBPP) to advance our understanding of the complex web of interactions that span the domains of life.

So, how can biogeochemists assist in linking human population dynamics, range expansion effects on element cycling,

organismal adaptation, and contaminant cycling, to name a few, with such global efforts? It has long been recognized that better integration of environmental sciences and social sciences is needed in seeking a viable sustainability for the future. For example, the introduction of *translational ecology* (Schlesinger, 2010) and continued emphasis by the National Science Foundation (NSF) and their supported working groups, such as the National Socio-Environmental Synthesis Center (SESYNC <https://www.sesync.org/>, last access: 2012), clearly reflect new solutions to the problem.

Are biogeochemists adequately unified in addressing some of these key global issues in the 21st century? Most biogeochemists would agree that better links are needed with Earth system models, including better links between biogeochemical cycling, organismal traits and their changes, and environmental modeling. This is a major challenge requiring connections between cellular and organismal level systems biology with observational and modeling studies of global biogeochemical cycles. Synergies between detailed process-level understanding through local or regional studies, and the ability to upscale and detect global change through global-scale observations, have already contributed strongly to progress in our field, advancing beyond some of its previously conceived shortfalls (Cutter, 2005; Likens, 2004). Nevertheless, biogeochemists, amongst others, have called for more improvements (Groffman et al., 2017) in the accessibility and sharing of complex data (Saito et al., 2020a; Tanhua et al., 2019; Villar et al., 2018), the integration of observations and predictive

models (Fennel et al., 2019), and the incorporation of societal factors (e.g., damming, nutrient management) in model projections (Seitzinger et al., 2010). Here, we call for better incorporation of mechanistic knowledge from omics studies (Urban et al., 2016; Coles et al., 2017) and a stronger integration of modern and past ecological and evolutionary dynamics with biogeochemistry. Climate-driven range expansion of organisms, including immigration–emigration patterns by humans, is expected to enhance zoonotic diseases (both viral and bacterial) (e.g., Han et al., 2015; Allen et al., 2017) and threaten global food supplies (via rise in soil pathogens) (e.g., Delgado-Baquerizo et al., 2020). These changes are likely to be coupled with broader shifts in community-level interactions, organismal adaptive change (Scheffers et al., 2016), and associated changes in biogeochemical cycling rates and fluxes (e.g., nutrient, contaminants, redox conditions) (e.g., Bianchi et al., 2021). In this perspective, we provide some insights on why these linkages, between biogeochemistry, evolutionary biology, and social sciences, are what we believe to be some key foci for biogeochemists in the coming decades. Thus, we argue for (1) better integration of adaptive evolutionary change, coupled with range expansion, and biogeochemical cycles and (2) continued integration of social sciences, focusing on the human–natural system – in the context of sustainability and biogeochemistry.

2 Eco-evolutionary dynamics and biogeochemistry

There has been a long-standing interest in the co-evolution of life and biogeochemical cycles on Earth, as chemical conditions of this planet have been strongly influenced by evolving biochemical capabilities of life (Canfield et al., 2007; Lenton et al., 2014; Saito et al., 2003). Earth's life support system is inextricably tied to biogeochemical cycling and prokaryote evolution (Falkowski et al., 2008). Research has shown that understanding such co-evolutionary patterns in prokaryotes is key in developing environmental engineering solutions for future sustainability in the Anthropocene (Newman and Banfield, 2002). Also, metazoans have long been recognized as important “engineers” of Earth's elemental cycles (Darwin, 1881). Moreover, a better understanding of organismal–biogeochemical interactions in the fossil record can help predict future linkages between climate change, organismal adaptation, range expansion, and biogeochemical cycles (Bianchi et al., 2021). For example, range expansion and/or contraction of marine benthic communities may be linked to climate change (Buatois et al., 2020). Furthermore, evolutionary radiations in marine environments reflect changing feeding guilds and bioturbation activities (Mángano and Buatois, 2014), which show important biogeochemical feedbacks (e.g., redox changes in sediments) (Boyle et al., 2014; Buatois et al., 2020). We argue that more structured collaboration (via joint workshops and meetings) between ecological, biogeochemical, evolutionary, and pa-

leo-environmental scientists is needed in the 21st century to better utilize the fossil record for such questions.

Recently, evidence has shown that significant evolutionary trait change can occur over timescales of just a few generations, and the rapidly changing environmental context at local, regional, and global scales during the Anthropocene leads to strong selection pressures on populations to adapt (Bell and Collins, 2008; Hutchins et al., 2015; Kuebbing et al., 2018; Seibel and Deutsch, 2020). In fact, there is now ample evidence for rapid evolution, where rapid refers to contemporary evolution or evolution in ecological time. Multiple review papers and books on this topic have emerged (Bell and Collins, 2008; Fussmann et al., 2007; Hutchins et al., 2019; Palumbi, 2002; Schoener, 2011), as well as a monograph on eco-evolutionary dynamics (Hendry and Green, 2017). In brief, from microbes to plankton, insects to plants, fishes, and birds, there are now hundreds of studies showing significant evolutionary change in trait values over short time spans – just a few tens of generations. Interestingly, this contemporary evolution impacts ecosystem functioning and elemental cycling dynamics (Bassar et al., 2010; Declerck et al., 2015). In one case, the evolution of zooplankton within a single growth season has been shown to shape the typical seasonal dynamics of phyto- and zooplankton in lakes (Schaffner et al., 2019). In another example, evolution in body size in salmon, through its effects on salmon consumption by bears, impacts nutrient transfer from aquatic to terrestrial systems (Carlson et al., 2011). Many additional surprising pathways and mechanisms, inclusive of ecological aspects like behavior, remain to be discovered. For example, fear of predation by spiders can alter the elemental composition of grasshoppers, resulting in changes in production and nutrient cycles in ecosystems (Hawlena et al., 2012). Theory indicates that rapid evolutionary trait change can also influence the occurrence and (recovery) trajectory of ecosystem regime shifts (Dakos et al., 2019). Integrating ecological and evolutionary responses is needed to make reliable predictions of how ecosystems respond to climate change (Matthews et al., 2011) and how this impacts biogeochemical cycles. While the evolutionary biologists are well aware of such changes, better linkages with ecosystem ecologists and biogeochemists are needed to examine how these adaptive changes in taxa and community composition impact biogeochemical cycles – and how this gets integrated in future IPCC reports.

Rapid evolutionary changes are gaining attention because they can influence ecological responses, including responses to global change where new steep/novel gradients in “critical zones” continue to develop in the Anthropocene – in both terrestrial and aquatic systems (e.g., Bianchi and Morrison, 2018; Chorover et al., 2007). This implies that our analyses of ecological responses and their biogeochemical implications should not assume that trait values of species are fixed in time and in equilibrium with biogeochemical rates and fluxes. Depending on the taxon and the selection pressure,

traits can significantly change, and these changes have been shown to influence ecosystem processes such as consumption, production, respiration, and nutrient cycles. Given the importance of microbes for biogeochemical cycles, this notion becomes even more important, because microbial generation times are short and can evolve significantly different trait values in a matter of a few days or weeks. For example, Lawrence et al. (2012) showed that when competing bacterial strains were forced to grow together, they changed their physiology so much that they became partially dependent on each other and reached higher densities than when initially grown in the presence of one other. Yet, these observations of rapid evolution must be reconciled with the fact that the evolution of novel biochemical pathways and their impacts on biogeochemical cycles have occurred rarely (David and Alm, 2011). Eco-evodynamics are generally underexplored relative to biogeochemistry; thus, we consider this topic as a research frontier where integration among disciplines will lead to significant advances in the 21st century.

In recent decades, the ability to directly track genes and gene functions of organisms in nature, especially in microbes, has greatly contributed to understanding their interactions with biogeochemical cycles (Martiny et al., 2006; Rusch et al., 2010). Measurements of microbial transcripts and proteins in natural environments has allowed direct observation of cellular functions as adaptive responses to the environment (Bergauer et al., 2018; Gifford et al., 2011). These functional systems include biogeochemically relevant enzymes, transporters, storage molecules, and regulatory systems, and the quantitation of enzymes can be used to generate omics-based potential biogeochemical rates (Saito et al., 2020b). This provides mechanistic information about the underpinnings of biological controls on biogeochemistry and allows direct quantification of rate changes along different pathways. There are, however, many knowledge gaps to fill: roughly half of all genes have unknown functions, the systems biology controlling gene regulation is poorly characterized (Held et al., 2019), and we know little about how the different biochemical pathways relate to resilience at the ecosystem level. Forging connections between the genetic and biochemical underpinnings to the production of metabolites that contribute to carbon and other element cycling is primed for discovery (Soule et al., 2015). Omics studies will also help reveal how the microbiomes of plants, invertebrates, and vertebrates, who make up the predator–prey and decomposition food webs, influence biogeochemical cycles (Macke et al., 2017).

3 Embracing the social dimension

With growing needs to understand how human population growth will interact with ecosystem evolution, in the face of climate change and organismal range expansion (including humans), biogeochemistry should continue to explore new

dimensions of sustainability throughout the research process – including linkages with the social sciences for a holistic understanding of the human–environment system (Fig. 2). These ideas build on earlier notable efforts such as translational ecology (Schlesinger, 2010) and are similar to the idea of translational medicine, an interdisciplinary area of research that aims to improve human health by accelerating the application of novel research discoveries to improving patient outcomes. Other fields, such as economics, are also warming to mission-driven research (Crain et al., 2014; Mazzucato, 2018). We argue that there is a large gap between discoveries in biogeochemistry and their application to improving ecosystem health. For example, biogeochemistry research may present many possible solutions to managing the global carbon budget – from planting a trillion trees, to carbon taxes and trading, to direct air carbon capture – but their adoption (or not) by various levels of society is not generally studied scientifically, which could very much hamper the development of solutions. As human range expansion continues to escalate with climate-change-driven food security issues in many countries, particularly developing nations (Carney and Krause, 2020), integrating human behavior at several scales is essential. In some cases, a lack of understanding of social processes can lead to unexpected societal “pushback”, rendering scientific knowledge less impactful. In 2018, the “yellow vest” protests in France emerged in response to a new carbon tax and greatly hindered progress toward carbon management goals in that country. Protestors agreed that climate change is an issue but were not willing to accept socially unjust solutions. Top-down approaches like the Paris agreement, with its ongoing political challenges and limited efficacy in combating climate, face severe challenges. Therefore scientists are rushing to study how to harness social forces in a polycentric manner in order to tackle global-scale sustainability challenges. Put succinctly, without involving individuals outside of the field in all stages of the research process, biogeochemistry research that seeks to advance sustainability through policy or behavior change risks answering questions that decision-makers are not asking, or proposing solutions that populations will not adopt. The window of opportunity to protect many of the natural systems we have the privilege of studying is rapidly disappearing. And, the primary barriers to adoption of many sustainability solutions are often political and social limitations, not lack of scientific knowledge or availability of technology. Increasing public awareness of how basic science is linked with environmental problems through early education will be key in reducing these limitations.

Does the need for socially conscious policy-driven research mean that basic science inquiry in biogeochemistry is dead? Are we not allowed to wonder about the origin of the Earth and the basic processes that underlie the cycling of energy, water, and nutrients across the surface of the Earth? We argue that there is a false dichotomy between biogeochemistry and a translational bio-geo-socio-chemistry. In-




Figure 2. “Translational biogeochemistry” would encompass the two-way feedbacks between human and biogeochemical systems. Understanding human societal pathways will be critical in discerning and mitigating future climate patterns and their biogeochemical consequences.

deed, our goals are consistent with Vernadsky’s original vision, as he stated “understanding our planet the way it is.” The difference is that human influence on the Earth system is now so pervasive that our challenge has moved from integration of biology, geology, and chemistry to inclusion of social sciences where evidence now exists for how organismal range expansion and/or contraction of, say, marine benthic communities respond to climate change (Buatois et al., 2020). Indeed, by including the social sciences, biogeochemistry can help predict future changes in the biosphere in response to climate change and other anthropogenic pressures, while helping to facilitate sustainable and equitable responses by society. Thus, biogeochemical knowledge comes closer to policy makers. Nevertheless, our knowledge of biogeochemical cycles remains relatively limited compared to other core sciences (biology, chemistry, physics, geology), and thus basic research will be key in understanding and laying the foundation for good policy development – relating to global change science. While the core scientific disciplines of physics, chemistry, geology, and biology have long been essential in the development in part of NASA exploration, pharmaceutical and engineering materials, petrochemical processes, and medicinal and agricultural biochemistry, as well as biogeochemistry, is uniquely poised (see

Fig. 1) to serve the public as a key core science in addressing climate change issues.

Biogeochemistry is already an inherent component of sustainability and Earth system sciences, which are addressing the overarching challenge of how global change pressures the habitability of the planet and the ability to sustainably use its resources to feed and supply the world population and economy. A pivotal issue is how organismal, environmental, and societal processes cause feedbacks that affect biogeochemical cycles and global change (Seitzinger et al., 2010). A “translational” biogeochemistry would be a natural pathway of research on transformational human–environment processes because (1) both sustainability science and biogeochemistry are systems science approaches, and (2) collaboration between biogeochemists and social scientists could address topical key questions at a scale that is both holistic with respect to social–climate interactions and suitably detailed in addressing biogeochemical issues (Fig. 2). A holistic human–environment systems approach to applied biogeochemistry that accounts for social feedback might help winnow down policy recommendations to those that are both effective and likely to be adopted. Calls for greater integration between social and natural sciences have been made for years. The barriers slowing this integration down are not

only due to differences in methodology and perspectives. We speculate that lack of data on the interactions between social and natural systems has also contributed to this problem. However, we suggest that the dawn of digital social data has created a vast amount of essentially free observational data on social systems and their relation to natural systems that could help address this limitation and should be taken advantage of.

Key questions include the following. Do changes in biogeochemical pathways, associated with specific climate change drivers (e.g., droughts/flooding events versus ocean acidification), influence risk perception and social behavior differently than other broader global change issues (e.g., greenhouse gas (GHG) emission reduction versus dietary change)? If so, how can the biogeochemical knowledge we provide better “activate” the social pathways required to support mitigation behavior? And, how will feedbacks between different biogeochemical systems hinder or accelerate these social pathways? An example of such an integrative approach is coupled social–climate modeling (Bury et al., 2019), in which sub-models are developed for both social dynamics and climate dynamics, and the two sub-models are then coupled together. As a result of this, socio-economic pathways become a prediction of theoretical models and thus become the subject of scientific study themselves, instead of being assumptions that are simply input into climate models. A human–environment perspective would change not only how we think about the natural world, but also how we design our research. A sustainability science approach to research may include stakeholders and policy experts at all stages in the research process.

4 Summary

The regional and global importance of biogeochemical processes for the homeostasis of Earth’s life support system necessitates accelerating research to achieve the goal of a sustainable global society. Starting from an awareness of the field’s history, new developments, and key limitations of current approaches, we aimed to develop a perspective on how biogeochemistry can better serve society.

The challenges and opportunities of 21st century biogeochemists are formidable and will require intense collaboration with government officials, the public, internationally funded programs, and other fields in the social sciences. A key to success will be the degree to which biogeochemistry succeeds in making biogeochemical knowledge more available to policy makers and educators, predicting future changes in the biosphere in response to climate change (and other anthropogenic impacts on timescales from seasons to centuries), and facilitating sustainable and equitable responses by society. Biogeochemistry can have an important role in bringing about a sustainable future. But, there are several impediments to fully realizing this role, including

the need for further integration across disciplines and spatial scales, the intrinsic challenges of combining increased breadth with mechanistic depth, and the need to strengthen connections to society. While biogeochemistry has made major achievements in the past century describing Earth’s global and regional biogeochemical cycles for the first time, we recognize that the field has acquired new societal responsibilities, in particular uncovering how humans are rapidly changing biogeochemical cycles, assessing the impact of these changes on biological communities and feedbacks on society, and effectively communicating this information to policy makers and society at large.

We call for more focused cross-disciplinary workshops and meetings in an already complex mixture of interdisciplinary sciences that allows for biogeochemistry (both paleo and modern) to utilize its novel origin and evolution (Bianchi, 2020) and to develop a better way forward in planning for climate change. Perhaps a global congress of biogeochemical and climate change is needed to move this ahead.

Data availability. No data sets were used in this article.

Author contributions. All authors contributed equally to the conceptualization and writing of this document. TSB was supported in part by the Beverly Thompson Endowed Chair in Geological Sciences; MJS acknowledges support from a Natural Sciences and Engineering Research Council of Canada Tier 1 Canada Research Chair in Integrative Molecular Biogeochemistry.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. We would like to thank William Schlesinger and a second anonymous reviewer for their constructive comments.

Financial support. TSB was supported in part by the Beverly Thompson Endowed Chair in Geological Sciences; MJS acknowledges support from the Natural Sciences and Engineering Research Council via a Tier 1 Canada Research Chair in Integrative Molecular Biogeochemistry.

Review statement. This paper was edited by Steven Bouillon and reviewed by William Schlesinger and one anonymous referee.

References

- Allen, T., Murray, K. A., Zambrana-Torrel, C., Morse, S. S., Rondinini, C., Di Marco, M., Breit, N., Olival, K. J., and Daszak, P.: Global hotspots and correlates of emerging zoonotic dis-

- eases. *Nat. Comm.*, 8, 1124, <https://doi.org/10.1038/s41467-017-00923-8>, 2017.
- Bassar, R. D., Marshall, M. C., López-Sepulcre, A., Zandonà, E., Auer, S. K., Travis, J., Pringle, C. M., Flecker, A. S., Thomas, S. A., Fraser, D. F., and Reznick, D. N.: Local adaptation in Trinidadian guppies alters ecosystem processes, *P. Natl. Acad. Sci. USA*, 107, 3616–3621, <https://doi.org/10.1073/pnas.0908023107>, 2010.
- Bell, G. and Collins, S.: Adaptation, extinction and global change, *Evol. Appl.*, 1, 3–16, <https://doi.org/10.1111/j.1752-4571.2007.00011.x>, 2008.
- Bergauer, K., Fernandez-Guerra, A., Garcia, J. A. L., Sprenger, R. R., Stepanauskas, R., Pachiadaki, M. G., Jensen, O. N., and Herndl, G. J.: Organic matter processing by microbial communities throughout the Atlantic water column as revealed by metaproteomics, *P. Natl. Acad. Sci. USA*, 115, 400–408, <https://doi.org/10.1073/pnas.1708779115>, 2018.
- Bianchi, T. S.: The evolution of biogeochemistry, revisited, *Biogeochemistry*, 13, 199–239, <https://doi.org/10.1007/s10533-020-00708-0>, 2020.
- Bianchi, T. S., Aller, R. C., Atwood, T., Buatois, L. A., Levin, L. A., Levinton, J. S., Middelburg, J. J., Morrison, E. S., Regnier, P., Shields, M. R., Snelgrove, P. V. R., Sotka, E. E., and Stanley, R. R. E.: What Global Biogeochemical Consequences Will Marine Animal-Sediment Interactions Have During Climate Change? *Elementa: Science of the Anthropocene*, <https://doi.org/10.1525/elementa.2020.00180>, 2021.
- Bianchi, T. S. and Morrison, E.: Human activities create corridors of change in aquatic zones, *EOS*, 99, 13–15, <https://doi.org/10.1029/2018EO104743>, 2018.
- Boyle, R. A., Dahl, T. W., Dale, A. W., Shields-Zhou, G. A., Zhu, M., Brasier, M. D., Canfield, D. E., and Lenton, T. M.: Stabilization of the coupled oxygen and phosphorus cycles by the evolution of bioturbation, *Nat. Geosci.*, 7, 671–676, <https://doi.org/10.1038/ngeo2213>, 2014.
- Buatois, L. A., Mangano, M. G., Monter, N. J., and Zhou, K.: Quantifying ecospace utilization and ecosystem engineering during the early Phanerozoic – The role of bioturbation and bioerosion, *Sci. Adv.*, 6, eabb0618 <https://doi.org/10.1126/sciadv.abb0618>, 2020.
- Bury, T. M., Bauch, C. T., and Anand, M.: Charting pathways to climate change mitigation in a coupled socio-climate model, *PLoS Comput. Biol.*, 15, e1007000, <https://doi.org/10.1371/journal.pcbi.1007000>, 2019.
- Canfield, D. E., Poulton, S. W., and Narbonne, G. M.: Late-Neoproterozoic deep-ocean oxygenation and the rise of animal life, *Science*, 315, 92–95, <https://doi.org/10.1126/science.1135013>, 2007.
- Carlson, S. M., Quinn, T. P., and Hendry, A. P.: Eco-evolutionary dynamics in Pacific salmon, *Heredity*, 106, 438–447, <https://doi.org/10.1038/hdy.2010.163>, 2011.
- Carney, M. A. and Krause, K. C.: Immigration/migration and healthy publics: the threat of food insecurity, *Palgrave Communications*, 6, 1–12, 2020.
- Chorover, J., Kretzschmar, R., Garcia-Pichel, F., and Sparks, D. L.: Soil biogeochemical processes within the critical zone, *Elements*, 3, 321–326, 2007.
- Coles, V. J., Stukel, M. R., Brooks, M. T., Burd, A., Crump, B. C., Moran, M. A., Paul, J. H., Satinsky, B. M., Yager, P. L., Zielinski, B. L., and Hood, R. R.: Ocean biogeochemistry modeled with emergent trait-based genomics, *Science*, 358, 1149–1154, <https://doi.org/10.1126/science.aan5712>, 2017.
- Collins, S. and Bell, G.: Evolution of natural algal populations at elevated CO₂, *Ecol. Lett.*, 9, 129–135, <https://doi.org/10.1111/j.1461-0248.2005.00854.x>, 2006.
- Crain, R., Cooper, C., and Dickinson, J. L.: Citizen science: a tool for integrating studies of human and natural systems, *Annu. Rev. Environ. Res.*, 39, 641–665, <https://doi.org/10.1146/annurev-environ-030713-154609>, 2014.
- Cutter, G. A.: Biogeochemistry: now and into the future, *Palaeogeogr., Palaeoclimatol., Palaeoecol.*, 219, 191–198, <https://doi.org/10.1016/j.palaeo.2004.10.021>, 2005.
- Dakos, V., Matthews, B., Hendry, A. P., Levine, J., Loeuille, N., Norberg, J., Nosil, P., Scheffer, M., and De Meester, L.: Ecosystem tipping points in an evolving world, *Nat. Ecol. Evol.*, 3, 355–362, <https://doi.org/10.1038/s41559-019-0797-2>, 2019.
- Darwin, C.: The formation of vegetable mould through the action of worms with some observations on their habits, John Murray, London, available at: <https://hdl.handle.net/2027/hvd.32044092526755>, 1881.
- David, L. A. and Alm, E. J.: Rapid evolutionary innovation during an Archaeal genetic expansion, *Nature*, 469, 93–96, 2011.
- Declerck, S. A. J., Malo, A. R., Diehl, S., Waasdorp, D., Lemmen, K. D., Proios, K., and Papakosta, S.: Rapid adaptation of herbivore consumers to nutrient limitation: eco-evolutionary feedbacks to population demography and resource control, *Ecol. Lett.*, 18, 553–562, <https://doi.org/10.1111/ele.12436>, 2015.
- Delgado-Baquerizo, M., Guerra, C. A., Cano-Díaz, C., Egidi, E., Wang, J., Eisenhauer, N., Singh, B. K., and Maestre, F. T.: The proportion of soil-borne pathogens increases with warming at the global scale, *Nat. Clim. Chang.*, 10, 550–554, <https://doi.org/10.1038/s41558-020-0759-3>, 2020.
- Derouin, S.: Geoscientists help map the pandemic, *Eos*, 101, <https://doi.org/10.1029/2020EO143538>, 2020.
- Falkowski, P. G., Fenchel, T., and Delong, E. F.: The microbial engines that drive Earth's biogeochemical cycles, *Science* 320, 1034–1039, <https://doi.org/10.1126/science.1153213>, 2008.
- Fennel, K., Gehlen, M., Brasseur, P., Brown, C.W., Ciavatta, S., Cossarini, G., Crise, A., Edwards, C.A., Ford, D., Friedrichs, M.A., Gregoire, M., Jones, E., Kim, H., Lamouroux, J., Murtagudde, R., Perruche, C., and the GODAE Ocean View: Marine Ecosystem Analysis and Prediction Task Team Advancing marine biogeochemical and ecosystem reanalyses and forecasts as tools for monitoring and managing ecosystem health, *Front. Mar. Sci.*, 6, <https://doi.org/10.3389/fmars.2019.00089>, 2019.
- Fussman, G. F., Loreau, M., and Abrams, P. A.: Eco-evolutionary dynamics of communities and ecosystems, *Func. Ecol.*, 21, 465–477, <https://doi.org/10.1111/j.1365-2435.2007.01275.x>, 2007.
- Gifford, S. M., Sharma, S., Rinta-Kanto, J. M., and Moran, M. A.: Quantitative analysis of a deeply sequenced marine microbial metatranscriptome, *The ISME J.*, 5, 461–472, <https://doi.org/10.1038/ismej.2010.14>, 2011.
- Gorham, E.: Biogeochemistry: Its origins and development, *Biogeochemistry*, 13, 199–239, <https://doi.org/10.1007/BF00002942>, 1991.
- Groffman, P. M., Cadenasso, M. L., Cavender-Bares, J., Childers, D. L., Grimm, N. B., Morgan Grove, J., Hobbie, S. E., Hutrya, L. R., Darrel Jenerette, G., McPhearson, T., Pataki, D. E., Pickett,

- S. T. S., Pouyat, R. V., Rosi-Marshall, E., and Ruddell, B. L.: Moving towards a new urban systems science, *Ecosystems*, 20, 38–43, <https://doi.org/10.1007/s10021-016-0053-4>, 2017.
- Han, B. A., Schmidt, J. P., Bowden, S. E., and Drake, J. M.: Rodent reservoirs of future zoonotic diseases, *P. Natl. Acad. Sci. USA*, 112, 7039–7044, <https://doi.org/10.1073/pnas.1501598112>, 2015.
- Hawlena, D., Strickland, M. S., Bradford, M. A., and Schmitz, O. J.: Fear of predation slows plant-litter decomposition, *Science*, 336, 1434–1438, <https://doi.org/10.1126/science.1220097>, 2012.
- Held, N. A., McIlvin, M. R., Moran, D. M., Laub, M. T., and Saito, M. A.: Unique patterns and biogeochemical relevance of two-component sensing in marine bacteria, *MSystems*, 4, e00317-18, <https://doi.org/10.1128/mSystems.00317-18>, 2019.
- Hendry, A. P. and Green, D. M.: Eco-Evolutionary Dynamics in Cold Blood, *Ichthyol., Herpetol.*, 105, 441–450, <https://doi.org/10.1643/OT-17-631>, 2017.
- Hutchins, D. A., Jansson, J. K., Remais, J. V., Rich, V. I., Singh, B. K., and Trivedi, P.: Climate change microbiology – problems and perspectives, *Nat. Rev. Microbiol.*, 17, 391–396, <https://doi.org/10.1038/s41579-019-0178-5>, 2019.
- Hutchins, D. A., Walworth, N. G., Webb, E. A., Saito, M. A., Moran, D., McIlvin, M. R., Gale, J., and Fu, F. X.: Irreversibly increased nitrogen fixation in *Trichodesmium* experimentally adapted to elevated carbon dioxide, *Nat. Comm.*, 6, 1–7, <https://doi.org/10.1038/ncomms9155>, 2015.
- Kress, W. B., Mazet, J. A. K., and Hebert, P. D. N.: Intercepting pandemics through genomics, *P. Natl. Acad. Sci. USA*, 117, 13852–13855, <https://doi.org/10.1073/pnas.2009508117>, 2020.
- Kuebbing, S. E., Reimer, A. P., Rosenthal, S. A., Feinberg, G., Leiserowitz, A., Lau, J. A., and Bradford, M. A.: Long-term research in ecology and evolution: a survey of challenges and opportunities, *Ecol. Monogr.*, 88, 245–258, <https://doi.org/10.1002/ecm.1289>, 2018.
- Lawrence, D., Fiegna, F., Behrends, V., Bundy, J. G., Phillimore, A. B., Bell, T., and Barraclough, T. G.: Species interactions alter evolutionary responses to a novel environment, *PLoS Biol.*, 10, e1001330, <https://doi.org/10.1371/journal.pbio.1001330>, 2012.
- Lenton, T. M., Boyle, R. A., Poulton, S. W., Shields-Zhou, G. A., and Butterfield, N. J.: Co-evolution of eukaryotes and ocean oxygenation in the Neoproterozoic era, *Nat. Geosci.*, 7, 257–265, <https://doi.org/10.1038/ngeo2108>, 2014.
- Likens, G. E.: Biogeochemistry: some opportunities and challenges for the future, *Wat. Air Soil Pollut. Focus*, 4, 5–24, <https://doi.org/10.1023/B:WAFO.0000028341.75842.08>, 2004.
- Macke, E., Tasiemski, A., Massol, F., Callens, M., and DeCaestecker, E.: Life history and eco-evolutionary dynamics in light of the gut microbiota, *Oikos*, 126, 508–531, <https://doi.org/10.1111/oik.03900>, 2017.
- Mángano, M. G. and Buatois, L. A.: Decoupling of body-plan diversification and ecological structuring during the Ediacaran-Cambrian transition: Evolutionary and geobiological feedbacks, *P. Roy. Soc. A-Biol. Sci.*, 281, 20140038, <https://doi.org/10.1098/rspb.2014.0038>, 2014.
- Martiny, A. C., Coleman, M. L., and Chisholm, S. W.: Phosphate acquisition genes in *Prochlorococcus* ecotypes: evidence for genome-wide adaptation, *P. Natl. Acad. Sci. USA*, 103, 12552–12557, <https://doi.org/10.1073/pnas.0601301103>, 2006.
- Matthews, B., Narwani, A., Hausch, S., Nonaka, E., Peter, H., Yamamichi, M., Sullam, K. E., Bird, K. C., Thomas, M. K., Hanley, T. C., and Turner, C. B.: Towards an integration of evolutionary biology and ecosystem science, *Ecol. Lett.*, 14, 690–701, <https://doi.org/10.1111/j.1461-0248.2011.01627.x>, 2011.
- Mazzucato, M.: Mission-oriented innovation policies: challenges and opportunities, *Industrial and Corporate Change*, 27, 803–815, <https://doi.org/10.1093/icc/dty034>, 2018.
- Newman, D. K. and J. F.: Banfield. Geomicrobiology: how molecular-scale interactions underpin biogeochemical systems, *Science*, 10, 1071–1077, <https://doi.org/10.1126/science.1010716>, 2002.
- Palumbi, S. R.: The Evolution Explosion: How Humans Cause Rapid Evolutionary Change, W. W. Norton & Company, ISBN 9780393323382, 2002.
- Rusch, D. B., Martiny, A. C., Dupont, C. L., Halpern, A. L., and Venter, J. C.: Characterization of *Prochlorococcus* clades from iron-depleted oceanic regions, *P. Natl. Acad. Sci. USA*, 107, 16184–16189, <https://doi.org/10.1073/pnas.1009513107>, 2010.
- Saito, M. A., Saunders, J. K., Chagnon, M., Gaylord, D., Shepherd, A., Held, N. A., Dupont, C., Symmonds, N., York, A., Charron, M., and Kinkade, D.: Development of an ocean protein portal for interactive discovery and education, *Journal of Proteome Research*, <https://doi.org/10.1101/2020.05.29.124388>, 2020a.
- Saito, M. A., McIlvin, M. R., Moran, D. M., Santoro, A. E., Dupont, C. L., Rafters, P. A., Saunders, J. K., Kaul, D., Lamberg, C. H., Westley, M., Valois, F., and Waterbury, J. B.: Abundant nitrite-oxidizing metalloenzymes in the mesopelagic zone of the tropical Pacific Ocean, *Nat. Geosci.*, 13, 355–362, <https://doi.org/10.1038/s41561-020-0565-6>, 2020b.
- Saito, M. A., Sigman, D. M., and Morel, F. M.: The bioinorganic chemistry of the ancient ocean: the co-evolution of cyanobacterial metal requirements and biogeochemical cycles at the Archean-Proterozoic boundary, *Inorg. Chim. Acta*, 356, 308–318, [https://doi.org/10.1016/S0020-1693\(03\)00442-0](https://doi.org/10.1016/S0020-1693(03)00442-0), 2003.
- Schaffner, L. R., Govaert, L., De Meester, L., Ellner, S. P., Fairchild, E., Miner, B. E., Rudstam, L. G., Spaak, P., and Hairston, N. G.: Consumer-resource dynamics is an eco-evolutionary process in a natural plankton community, *Nat. Ecol. Evol.*, 3, 1351–1358, <https://doi.org/10.1038/s41559-019-0960-9>, 2019.
- Scheffers, B. R., De Meester, L., Bridge, T. C. L., Hoffmann, A. A., Pandolfi, J. M., Corlett, R. T., Butchart, S. H. M., Pearce-Kelly, P., Kovacs, K. M., Dudgeon, D., Pacifici, M., Rondinini, C., Foden, W. B., Martin, T. G., Mora, C., Bickford, D., and Watson, J. E. M.: The broad footprint of climate change from genes to biomes to people, *Science*, 354, aaf7671, <https://doi.org/10.1126/science.aaf7671>, 2016.
- Schlesinger, W. H.: Biogeochemistry: An Analysis of Global Change, Academic Press, ISBN 9780323137843, 1991.
- Schlesinger, W. H.: Translational Ecology, *Science*, 609, <https://doi.org/10.1126/science.1195624>, 2010.
- Schoener, T. W.: The newest synthesis: understanding the interplay of evolutionary and ecological dynamics, *Science*, 331, 426–429, <https://doi.org/10.1126/science.1193954>, 2011.
- Seibel, B. A. and Deutsch, C.: Oxygen supply capacity in animals evolves to meet maximum demand at the current oxygen partial pressure regardless of size or temperature, *J. Exp. Biol.*, <https://doi.org/10.1242/jeb.210492>, 2020.

- Seitzinger, S. P., Mayorga, E., Bouwman, A. F., Kroeze, C., Beusen, A. H. W., Billen, G., Van Drecht, G., Dumont, E., Fekete, B. M., Garnier, J., and Harrison J. A.: Global river nutrient export: A scenario analysis of past and future trends, *Global Biogeochem. Cy.*, 24, GB0A08, <https://doi.org/10.1029/2009GB003587>, 2010.
- Soule, M. C. K., Longnecker, K., Johnson, W. M., and Kujawinski, E. B.: Environmental metabolomics: Analytical strategies, *Mar. Chem.*, 177, 374–387, <https://doi.org/10.1016/j.marchem.2015.06.029>, 2015.
- Steffen, W., Richardson, K., Rockström, J., Schellnhuber, H. J., Dube, O. P., Dutreuil, S., Lenton, T. M., and Lubchenco, L.: The emergence and evolution of Earth System Science, *Nat. Rev. Earth Environ.*, 1, 54–63, <https://doi.org/10.1038/s43017-019-0005-6>, 2020.
- Tanhua, T., Pouliquen, S., Hausman, J., O'Brien, K., Bricher, P., de Bruin, T., Buck, J. J. H., Burger, E. F., Carval, T., Casey, K. S., Diggs, S., Giorgetti, A., Glaves, H., Harscoat, V., Kinkade, D., Muelbert, J. H., Novellino, A., Pfeil, B., Pulsifer, P. L., Van de Putte, A., Robinson, E., Schaap, D., Smirnov, A., Smith, N., Snowden, D., Spears, T., Stall, S., Tacoma, M., Thijsse, P., Tronstad, S., Vandenberghe, T., Wengren, M., Lyborn, L., and Zhao, Z.: Ocean FAIR Data Services, *Front. Mar. Sci.*, 6, 440, <https://doi.org/10.3389/fmars.2019.00440>, 2019.
- Urban, M., Bocedi, G., Hendry, A. P., Mihoub, J. B., Pe'er, G., Singer, A., Bridle, J. R., Crozier, L. G., De Meester, L., Godsoe, W., Gonzalez, A., Hellmann, J. J., Holt, R. D., Huth, A., Johst, K., Krug, C. B., Leadley, P. W., Palmer, S. C. F., Pantel, J. H., Schmitz, A., Zollner, P. A., and Travis, J. M. J.: Improving the forecast for biodiversity under climate change, *Science*, 353, aad8466, <https://doi.org/10.1126/science.aad8466>, 2016.
- Vernadsky, V. I.: *Biosfera*. Leningrad, Seen in abridged English translation, Synergetic Press, Oracle, Arizona (1986), also available in a complete French translation, *La Biosphere*, by the author, Alcan, Paris (1929) (1945), <https://doi.org/10.1017/S0376892900036584>, 1926.
- Villar, E., Vannier, T., Vernet, C., Lescot, M., Cuenca, M., Alexandre, A., Bachelerie, P., Rosnet, T., Pelletier, E., Sunagawa, S., and Hingamp, P.: The Ocean Gene Atlas: exploring the biogeography of plankton genes online, *Nucl. Acids Res.*, 46, 289–295 <https://doi.org/10.1093/nar/gky376>, 2018.