

7 REFERENCES

- Abele, R. and Tampe, R. (1999) Function of the transport complex TAP in cellular immune recognition. *Biochim Biophys Acta*, **1461**, 405-419.
- Ahn, K., Angulo, A., Ghazal, P., Peterson, P.A., Yang, Y. and Fruh, K. (1996a) Human cytomegalovirus inhibits antigen presentation by a sequential multistep process. *Proc Natl Acad Sci U S A*, **93**, 10990-10995.
- Ahn, K., Gruhler, A., Galocha, B., Jones, T.R., Wiertz, E.J., Ploegh, H.L., Peterson, P.A., Yang, Y. and Fruh, K. (1997) The ER-luminal domain of the HCMV glycoprotein US6 inhibits peptide translocation by TAP. *Immunity*, **6**, 613-621.
- Ahn, K., Meyer, T.H., Uebel, S., Sempe, P., Djaballah, H., Yang, Y., Peterson, P.A., Fruh, K. and Tampe, R. (1996b) Molecular mechanism and species specificity of TAP inhibition by herpes simplex virus ICP47. *Embo J*, **15**, 3247-3255.
- Alimonti, J., Zhang, Q.J., Gabathuler, R., Reid, G., Chen, S.S. and Jefferies, W.A. (2000) TAP expression provides a general method for improving the recognition of malignant cells in vivo. *Nat Biotechnol*, **18**, 515-520.
- Allikmets, R., Singh, N., Sun, H., Shroyer, N.F., Hutchinson, A., Chidambaram, A., Gerrard, B., Baird, L., Stauffer, D., Peiffer, A., Rattner, A., Smallwood, P., Li, Y., Anderson, K.L., Lewis, R.A., Nathans, J., Leppert, M., Dean, M. and Lupski, J.R. (1997) A photoreceptor cell-specific ATP-binding transporter gene (ABCR) is mutated in recessive Stargardt macular dystrophy. *Nat Genet*, **15**, 236-246.
- Ambagala, A.P., Gopinath, R.S. and Srikuamran, S. (2003) Inhibition of TAP by pseudorabies virus is independent of its vhs activity. *Virus Res*, **96**, 37-48.
- Ambagala, A.P., Gopinath, R.S. and Srikuamran, S. (2004) Peptide transport activity of the transporter associated with antigen processing (TAP) is inhibited by an early protein of equine herpesvirus-1. *J Gen Virol*, **85**, 349-353.
- Ambagala, A.P., Hinkley, S. and Srikuamran, S. (2000) An early pseudorabies virus protein down-regulates porcine MHC class I expression by inhibition of transporter associated with antigen processing (TAP). *J Immunol*, **164**, 93-99.
- Antoniou, A.N., Ford, S., Pilley, E.S., Blake, N. and Powis, S.J. (2002) Interactions formed by individually expressed TAP1 and TAP2 polypeptide subunits. *Immunology*, **106**, 182-189.

- Arora, S., Lapinski, P.E. and Raghavan, M. (2001) Use of chimeric proteins to investigate the role of transporter associated with antigen processing (TAP) structural domains in peptide binding and translocation. *Proc Natl Acad Sci U S A*, **98**, 7241-7246.
- Atalay, R., Zimmermann, A., Wagner, M., Borst, E., Benz, C., Messerle, M. and Hengel, H. (2002) Identification and expression of human cytomegalovirus transcription units coding for two distinct Fc gamma receptor homologs. *J Virol*, **76**, 8596-8608.
- Azarian, S.M. and Travis, G.H. (1997) The photoreceptor rim protein is an ABC transporter encoded by the gene for recessive Stargardt's disease (ABCR). *FEBS Lett*, **409**, 247-252.
- Bahram, S., Arnold, D., Bresnahan, M., Strominger, J.L. and Spies, T. (1991) Two putative subunits of a peptide pump encoded in the human major histocompatibility complex class II region. *Proc Natl Acad Sci U S A*, **88**, 10094-10098.
- Banks, T.A., Jenkins, F.J., Kanangat, S., Nair, S., Dasgupta, S., Foster, C.M. and Rouse, B.T. (1994) Vaccination with the immediate-early protein ICP47 of herpes simplex virus-type 1 (HSV-1) induces virus-specific lymphoproliferation, but fails to protect against lethal challenge. *Virology*, **200**, 236-245.
- Barry, M. and Bleackley, R.C. (2002) Cytotoxic T lymphocytes: all roads lead to death. *Nat Rev Immunol*, **2**, 401-409.
- Bauer, D. and Tampe, R. (2002) Herpes viral proteins blocking the transporter associated with antigen processing TAP--from genes to function and structure. *Curr Top Microbiol Immunol*, **269**, 87-99.
- Begley, G.S., Horvath, A.R., Taylor, J.C. and Higgins, C.F. (2005) Cytoplasmic domains of the transporter associated with antigen processing and P-glycoprotein interact with subunits of the proteasome. *Mol Immunol*, **42**, 137-141.
- Beismann-Driemeyer, S. and Tampe, R. (2004) Function of the antigen transport complex TAP in cellular immunity. *Angew Chem Int Ed Engl*, **43**, 4014-4031.
- Beninga, J., Rock, K.L. and Goldberg, A.L. (1998) Interferon-gamma can stimulate post-proteasomal trimming of the N terminus of an antigenic peptide by inducing leucine aminopeptidase. *J Biol Chem*, **273**, 18734-18742.
- Bennett, E.M., Bennink, J.R., Yewdell, J.W. and Brodsky, F.M. (1999) Cutting edge: adenovirus E19 has two mechanisms for affecting class I MHC expression. *J Immunol*, **162**, 5049-5052.
- Benz, C. and Hengel, H. (2000) MHC class I-subversive gene functions of cytomegalovirus and their regulation by interferons—an intricate balance. *Virus Genes*, **21**, 39-47.

REFERENCES

- Blom, D., Hirsch, C., Stern, P., Tortorella, D. and Ploegh, H.L. (2004) A glycosylated type I membrane protein becomes cytosolic when peptide: N-glycanase is compromised. *Embo J*, **23**, 650-658.
- Boname, J.M., de Lima, B.D., Lehner, P.J. and Stevenson, P.G. (2004) Viral degradation of the MHC class I peptide loading complex. *Immunity*, **20**, 305-317.
- Boname, J.M. and Stevenson, P.G. (2001) MHC class I ubiquitination by a viral PHD/LAP finger protein. *Immunity*, **15**, 627-636.
- Burgert, H.G. and Kvist, S. (1987) The E3/19K protein of adenovirus type 2 binds to the domains of histocompatibility antigens required for CTL recognition. *Embo J*, **6**, 2019-2026.
- Cerundolo, V., Kelly, A., Elliott, T., Trowsdale, J. and Townsend, A. (1995) Genes encoded in the major histocompatibility complex affecting the generation of peptides for TAP transport. *Eur J Immunol*, **25**, 554-562.
- Chang, G. (2003) Structure of MsbA from Vibrio cholera: a multidrug resistance ABC transporter homolog in a closed conformation. *J Mol Biol*, **330**, 419-430.
- Chang, G. and Roth, C.B. (2001) Structure of MsbA from *E. coli*: a homolog of the multidrug resistance ATP binding cassette (ABC) transporters. *Science*, **293**, 1793-1800.
- Chee, M.S., Bankier, A.T., Beck, S., Bohni, R., Brown, C.M., Cerny, R., Horsnell, T., Hutchison, C.A., 3rd, Kouzarides, T., Martignetti, J.A. and et al. (1990) Analysis of the protein-coding content of the sequence of human cytomegalovirus strain AD169. *Curr Top Microbiol Immunol*, **154**, 125-169.
- Chen, M., Abele, R. and Tampe, R. (2004) Functional non-equivalence of ATP-binding cassette signature motifs in the transporter associated with antigen processing (TAP). *J Biol Chem*, **279**, 46073-46081.
- Collins, F.S. (1992) Cystic fibrosis: molecular biology and therapeutic implications. *Science*, **256**, 774-779.
- Craiu, A., Akopian, T., Goldberg, A. and Rock, K.L. (1997) Two distinct proteolytic processes in the generation of a major histocompatibility complex class I-presented peptide. *Proc Natl Acad Sci U S A*, **94**, 10850-10855.
- de la Salle, H., Hanau, D., Fricker, D., Urlacher, A., Kelly, A., Salamero, J., Powis, S.H., Donato, L., Bausinger, H., Laforet, M. and et al. (1994) Homozygous human TAP peptide transporter mutation in HLA class I deficiency. *Science*, **265**, 237-241.

- Degen, E., Cohen-Doyle, M.F. and Williams, D.B. (1992) Efficient dissociation of the p88 chaperone from major histocompatibility complex class I molecules requires both beta 2-microglobulin and peptide. *J Exp Med*, **175**, 1653-1661.
- Derfuss, T. and Meinl, E. (2002) Herpesviral proteins regulating apoptosis. *Curr Top Microbiol Immunol*, **269**, 257-272.
- Diederichs, K., Diez, J., Greller, G., Muller, C., Breed, J., Schnell, C., Vonrhein, C., Boos, W. and Welte, W. (2000) Crystal structure of MalK, the ATPase subunit of the trehalose/maltose ABC transporter of the archaeon *Thermococcus litoralis*. *Embo J*, **19**, 5951-5961.
- Driscoll, J., Brown, M.G., Finley, D. and Monaco, J.J. (1993) MHC-linked LMP gene products specifically alter peptidase activities of the proteasome. *Nature*, **365**, 262-264.
- Emmerich, N.P., Nussbaum, A.K., Stevanovic, S., Priemer, M., Toes, R.E., Rammensee, H.G. and Schild, H. (2000) The human 26 S and 20 S proteasomes generate overlapping but different sets of peptide fragments from a model protein substrate. *J Biol Chem*, **275**, 21140-21148.
- Fowler, K.B., Stagno, S., Pass, R.F., Britt, W.J., Boll, T.J. and Alford, C.A. (1992) The outcome of congenital cytomegalovirus infection in relation to maternal antibody status. *N Engl J Med*, **326**, 663-667.
- Fruh, K., Ahn, K., Djaballah, H., Sempe, P., van Endert, P.M., Tampe, R., Peterson, P.A. and Yang, Y. (1995) A viral inhibitor of peptide transporters for antigen presentation. *Nature*, **375**, 415-418.
- Fu, Y., Yan, G., Shi, L. and Faustman, D. (1998) Antigen processing and autoimmunity. Evaluation of mRNA abundance and function of HLA-linked genes. *Ann N Y Acad Sci*, **842**, 138-155.
- Galocha, B., Hill, A., Barnett, B.C., Dolan, A., Raimondi, A., Cook, R.F., Brunner, J., McGeoch, D.J. and Ploegh, H.L. (1997) The active site of ICP47, a herpes simplex virus-encoded inhibitor of the major histocompatibility complex (MHC)-encoded peptide transporter associated with antigen processing (TAP), maps to the NH₂-terminal 35 residues. *J Exp Med*, **185**, 1565-1572.
- Gao, M., Cui, H.R., Loe, D.W., Grant, C.E., Almquist, K.C., Cole, S.P. and Deeley, R.G. (2000) Comparison of the functional characteristics of the nucleotide binding domains of multidrug resistance protein 1. *J Biol Chem*, **275**, 13098-13108.

REFERENCES

- Gaudet, R. and Wiley, D.C. (2001) Structure of the ABC ATPase domain of human TAP1, the transporter associated with antigen processing. *Embo J*, **20**, 4964-4972.
- Gewurz, B.E., Gaudet, R., Tortorella, D., Wang, E.W., Ploegh, H.L. and Wiley, D.C. (2001) Antigen presentation subverted: Structure of the human cytomegalovirus protein US2 bound to the class I molecule HLA-A2. *Proc Natl Acad Sci U S A*, **98**, 6794-6799.
- Gorbulev, S., Abele, R. and Tampe, R. (2001) Allosteric crosstalk between peptide-binding, transport, and ATP hydrolysis of the ABC transporter TAP. *Proc Natl Acad Sci U S A*, **98**, 3732-3737.
- Groettrup, M., Khan, S., Schwarz, K. and Schmidtke, G. (2001) Interferon-gamma inducible exchanges of 20S proteasome active site subunits: why? *Biochimie*, **83**, 367-372.
- Gruhler, A., Peterson, P.A. and Fruh, K. (2000) Human cytomegalovirus immediate early glycoprotein US3 retains MHC class I molecules by transient association. *Traffic*, **1**, 318-325.
- Halenius, A., Momburg, F., Bauer, D., Lobigs, M. and Hengel, H. (2005) Physical and functional interactions of the cytomegalovirus US6 glycoprotein with the transporter associated with antigen processing (TAP). *revised*.
- Hamprecht, K., Maschmann, J., Vochem, M., Dietz, K., Speer, C.P. and Jahn, G. (2001) Epidemiology of transmission of cytomegalovirus from mother to preterm infant by breastfeeding. *Lancet*, **357**, 513-518.
- Hegde, N.R., Tomazin, R.A., Wisner, T.W., Dunn, C., Boname, J.M., Lewinsohn, D.M. and Johnson, D.C. (2002) Inhibition of HLA-DR assembly, transport, and loading by human cytomegalovirus glycoprotein US3: a novel mechanism for evading major histocompatibility complex class II antigen presentation. *J Virol*, **76**, 10929-10941.
- Hengel, H., Flohr, T., Hammerling, G.J., Koszinowski, U.H. and Momburg, F. (1996) Human cytomegalovirus inhibits peptide translocation into the endoplasmic reticulum for MHC class I assembly. *J Gen Virol*, **77 (Pt 9)**, 2287-2296.
- Hengel, H., Koopmann, J.O., Flohr, T., Muranyi, W., Goulmy, E., Hammerling, G.J., Koszinowski, U.H. and Momburg, F. (1997) A viral ER-resident glycoprotein inactivates the MHC-encoded peptide transporter. *Immunity*, **6**, 623-632.
- Hewitt, E.W., Gupta, S.S. and Lehner, P.J. (2001) The human cytomegalovirus gene product US6 inhibits ATP binding by TAP. *Embo J*, **20**, 387-396.
- Higgins, C.F. (2001) ABC transporters: physiology, structure and mechanism--an overview. *Res Microbiol*, **152**, 205-210.

- Hill, A., Jugovic, P., York, I., Russ, G., Bennink, J., Yewdell, J., Ploegh, H. and Johnson, D. (1995) Herpes simplex virus turns off the TAP to evade host immunity. *Nature*, **375**, 411-415.
- Howarth, M., Williams, A., Tolstrup, A.B. and Elliott, T. (2004) Tapasin enhances MHC class I peptide presentation according to peptide half-life. *Proc Natl Acad Sci U S A*, **101**, 11737-11742.
- Hung, L.-W., Wang, I.X., Nikaido, K., Liu, P.-Q., Ames, G.F.-L. and Kim, S.-H. (1998) Crystal structure of the ATP-binding subunit of an ABC transporter. *Nature*, **396**, 703-707.
- Johnson, D.C. and Hegde, N.R. (2002) Inhibition of the MHC class II antigen presentation pathway by human cytomegalovirus. *Curr Top Microbiol Immunol*, **269**, 101-115.
- Jones, T.R. and Muzithras, V.P. (1991) Fine mapping of transcripts expressed from the US6 gene family of human cytomegalovirus strain AD169. *J Virol*, **65**, 2024-2036.
- Jones, T.R., Wiertz, E.J., Sun, L., Fish, K.N., Nelson, J.A. and Ploegh, H.L. (1996) Human cytomegalovirus US3 impairs transport and maturation of major histocompatibility complex class I heavy chains. *Proc Natl Acad Sci U S A*, **93**, 11327-11333.
- Jugovic, P., Hill, A.M., Tomazin, R., Ploegh, H. and Johnson, D.C. (1998) Inhibition of Major Histocompatibility Complex Class I Antigen Presentation in Pig and Primate Cells by Herpes Simplex Virus Type 1 and 2 ICP47. *J. Virol.*, **72**, 5076-5084.
- Karttunen, J.T., Lehner, P.J., Gupta, S.S., Hewitt, E.W. and Cresswell, P. (2001) Distinct functions and cooperative interaction of the subunits of the transporter associated with antigen processing (TAP). *Proc Natl Acad Sci U S A*, **98**, 7431-7436.
- Kikkert, M., Hassink, G., Barel, M., Hirsch, C., van der Wal, F.J. and Wiertz, E. (2001) Ubiquitination is essential for human cytomegalovirus US11-mediated dislocation of MHC class I molecules from the endoplasmic reticulum to the cytosol. *Biochem J*, **358**, 369-377.
- Kisselev, A.F., Akopian, T.N., Woo, K.M. and Goldberg, A.L. (1999) The sizes of peptides generated from protein by mammalian 26 and 20 S proteasomes. Implications for understanding the degradative mechanism and antigen presentation. *J Biol Chem*, **274**, 3363-3371.
- Klar, D. and Hammerling, G.J. (1989) Induction of assembly of MHC class I heavy chains with beta 2microglobulin by interferon-gamma. *Embo J*, **8**, 475-481.

- Kleijmeer, M.J., Kelly, A., Geuze, H.J., Slot, J.W., Townsend, A. and Trowsdale, J. (1992) Location of MHC-encoded transporters in the endoplasmic reticulum and cis-Golgi. *Nature*, **357**, 342-344.
- Koch, J., Guntrum, R., Heintke, S., Kyritsis, C. and Tampe, R. (2003) Functional dissection of the transmembrane domains of the transporter associated with antigen processing (TAP). *J Biol Chem*.
- Koppers-Lalic, D., Reits, E.A.J., Ressing, M.E., Lipinska, A.D., Abele, R., Koch, J., Rezende, M.M., Admiraal, P., van Leeuwen, D., Bienkowska-Szewczyk, K., Mettenleiter, T.C., Rijsewijk, F.A.M., Tampe, R., Neefjes, J. and Wiertz, E.J.H.J. (2005) Varicelloviruses avoid T cell recognition by UL49.5-mediated inactivation of the transporter associated with antigen processing. *PNAS*, 0501463102.
- Koppers-Lalic, D., Rychlowski, M., van Leeuwen, D., Rijsewijk, F.A., Ressing, M.E., Neefjes, J.J., Bienkowska-Szewczyk, K. and Wiertz, E.J. (2003) Bovine herpesvirus 1 interferes with TAP-dependent peptide transport and intracellular trafficking of MHC class I molecules in human cells. *Arch Virol*, **148**, 2023-2037.
- Kotenko, S.V., Saccani, S., Izotova, L.S., Mirochnitchenko, O.V. and Pestka, S. (2000) Human cytomegalovirus harbors its own unique IL-10 homolog (cmvIL-10). *Proc Natl Acad Sci U S A*, **97**, 1695-1700.
- Kunisawa, J. and Shastri, N. (2003) The group II chaperonin TRiC protects proteolytic intermediates from degradation in the MHC class I antigen processing pathway. *Mol Cell*, **12**, 565-576.
- Kyritsis, C., Gorbulev, S., Hutschenreiter, S., Pawlitschko, K., Abele, R. and Tampe, R. (2001) Molecular mechanism and structural aspects of transporter associated with antigen processing inhibition by the cytomegalovirus protein US6. *J Biol Chem*, **276**, 48031-48039.
- Lacaille, V.G. and Androlewicz, M.J. (1998) Herpes simplex virus inhibitor ICP47 destabilizes the transporter associated with antigen processing (TAP) heterodimer. *J Biol Chem*, **273**, 17386-17390.
- Laemmli, U.K. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*, **227**, 680-685.
- Lapinski, P.E., Neubig, R.R. and Raghavan, M. (2001) Walker A lysine mutations of TAP1 and TAP2 interfere with peptide translocation but not peptide binding. *J Biol Chem*, **276**, 7526-7533.

REFERENCES

- Lee, S., Park, B. and Ahn, K. (2003) Determinant for endoplasmic reticulum retention in the luminal domain of the human cytomegalovirus US3 glycoprotein. *J Virol*, **77**, 2147-2156.
- Lehner, P.J., Karttunen, J.T., Wilkinson, G.W. and Cresswell, P. (1997) The human cytomegalovirus US6 glycoprotein inhibits transporter associated with antigen processing-dependent peptide translocation. *Proc Natl Acad Sci U S A*, **94**, 6904-6909.
- Lehner, P.J., Surman, M.J. and Cresswell, P. (1998) Soluble tapasin restores MHC class I expression and function in the tapasin-negative cell line .220. *Immunity*, **8**, 221-231.
- Leib, D.A. (2002) Counteraction of interferon-induced antiviral responses by herpes simplex viruses. *Curr Top Microbiol Immunol*, **269**, 171-185.
- Lilley, B.N. and Ploegh, H.L. (2004) A membrane protein required for dislocation of misfolded proteins from the ER. *Nature*, **429**, 834-840.
- Livneh, E., Benveniste, M., Prywes, R., Felder, S., Kam, Z. and Schlessinger, J. (1986) Large deletions in the cytoplasmic kinase domain of the epidermal growth factor receptor do not affect its lateral mobility. *J Cell Biol*, **103**, 327-331.
- Lobigs, M. and Mullbacher, A. (1995) Recombinant expression of vaccinia virus-encoded TAP1 and TAP2 promotes MHC class I-restricted antigen presentation in a Syrian hamster cell line. *Immunol Cell Biol*, **73**, 181-184.
- Locher, K.P., Lee, A.T. and Rees, D.C. (2002) The *E. coli* BtuCD structure: a framework for ABC transporter architecture and mechanism. *Science*, **296**, 1091-1098.
- Meyer, T.H., van Endert, P.M., Uebel, S., Ehring, B. and Tampe, R. (1994) Functional expression and purification of the ABC transporter complex associated with antigen processing (TAP) in insect cells. *FEBS Lett*, **351**, 443-447.
- Misaghi, S., Sun, Z.Y., Stern, P., Gaudet, R., Wagner, G. and Ploegh, H. (2004) Structural and functional analysis of human cytomegalovirus US3 protein. *J Virol*, **78**, 413-423.
- Momburg, F. and Hengel, H. (2002) Corking the bottleneck: the transporter associated with antigen processing as a target for immune subversion by viruses. *Curr Top Microbiol Immunol*, **269**, 57-74.
- Momburg, F., Ortiz-Navarrete, V., Neefjes, J., Goulmy, E., van de Wal, Y., Spits, H., Powis, S.J., Butcher, G.W., Howard, J.C., Walden, P. and et al. (1992) Proteasome subunits encoded by the major histocompatibility complex are not essential for antigen presentation. *Nature*, **360**, 174-177.

- Mosser, J., Douar, A.M., Sarde, C.O., Kioschis, P., Feil, R., Moser, H., Poustka, A.M., Mandel, J.L. and Aubourg, P. (1993) Putative X-linked adrenoleukodystrophy gene shares unexpected homology with ABC transporters. *Nature*, **361**, 726-730.
- Mullis, K.B. and Faloona, F.A. (1987) Specific synthesis of DNA in vitro via a polymerase-catalyzed chain reaction. *Methods Enzymol*, **155**, 335-350.
- Murphy, E., Rigoutsos, I., Shibuya, T. and Shenk, T.E. (2003) Reevaluation of human cytomegalovirus coding potential. *Proc Natl Acad Sci U S A*, **100**, 13585-13590.
- Neefjes, J.J., Momburg, F. and Hammerling, G.J. (1993) Selective and ATP dependent translocation of peptides by the MHC-encoded transporter. *Science*, **261**, 769-771.
- Neumann, L., Kraas, W., Uebel, S., Jung, G. and Tampe, R. (1997) The active domain of the herpes simplex virus protein ICP47: a potent inhibitor of the transporter associated with antigen processing. *J Mol Biol*, **272**, 484-492.
- Nijenhuis, M. and Hammerling, G.J. (1996) Multiple regions of the transporter associated with antigen processing (TAP) contribute to its peptide binding site. *J Immunol*, **157**, 5467-5477.
- Nijenhuis, M., Schmitt, S., Armandola, E.A., Obst, R., Brunner, J. and Hammerling, G.J. (1996) Identification of a contact region for peptide on the TAP1 chain of the transporter associated with antigen processing. *J Immunol*, **156**, 2186-2195.
- Nossner, E. and Parham, P. (1995) Species-specific differences in chaperone interaction of human and mouse major histocompatibility complex class I molecules. *J Exp Med*, **181**, 327-337.
- Ortmann, B., Copeman, J., Lehner, P.J., Sadasivan, B., Herberg, J.A., Grandea, A.G., Riddell, S.R., Tampe, R., Spies, T., Trowsdale, J. and Cresswell, P. (1997) A critical role for tapasin in the assembly and function of multimeric MHC class I-TAP complexes. *Science*, **277**, 1306-1309.
- Parham, P., Barnstable, C.J. and Bodmer, W.F. (1979) Use of a monoclonal antibody (W6/32) in structural studies of HLA-A,B,C, antigens. *J Immunol*, **123**, 342-349.
- Park, B., Kim, Y., Shin, J., Lee, S., Cho, K., Fruh, K. and Ahn, K. (2004) Human cytomegalovirus inhibits tapasin-dependent peptide loading and optimization of the MHC class I peptide cargo for immune evasion. *Immunity*, **20**, 71-85.
- Peckham, C.S. (1991) Cytomegalovirus infection: congenital and neonatal disease. *Scand J Infect Dis Suppl*, **80**, 82-87.
- Reddehase, M.J. (2002) Antigens and immuno-evasins: opponents in cytomegalovirus immune surveillance. *Nat Rev Immunol*, **2**, 831-844.

- Reits, E.A., Vos, J.C., Gromme, M. and Neefjes, J. (2000) The major substrates for TAP in vivo are derived from newly synthesized proteins. *Nature*, **404**, 774-778.
- Ressing, M.E., Keating, S.E., van Leeuwen, D., Koppers-Lalic, D., Pappworth, I.Y., Wiertz, E.J. and Rowe, M. (2005) Impaired Transporter Associated with Antigen Processing-Dependent Peptide Transport during Productive EBV Infection. *J Immunol*, **174**, 6829-6838.
- Russ, G., Esquivel, F., Yewdell, J.W., Cresswell, P., Spies, T. and Bennink, J.R. (1995) Assembly, intracellular localization, and nucleotide binding properties of the human peptide transporters TAP1 and TAP2 expressed by recombinant vaccinia viruses. *J Biol Chem*, **270**, 21312-21318.
- Sadasivan, B., Lehner, P.J., Ortmann, B., Spies, T. and Cresswell, P. (1996) Roles for calreticulin and a novel glycoprotein, tapasin, in the interaction of MHC class I molecules with TAP. *Immunity*, **5**, 103-114.
- Sanger, F., Nicklen, S. and Coulson, A.R. (1977) DNA sequencing with chain-terminating inhibitors. *Proc Natl Acad Sci U S A*, **74**, 5463-5467.
- Saric, T., Beninga, J., Graef, C.I., Akopian, T.N., Rock, K.L. and Goldberg, A.L. (2001) Major histocompatibility complex class I-presented antigenic peptides are degraded in cytosolic extracts primarily by thimet oligopeptidase. *J Biol Chem*, **276**, 36474-36481.
- Saric, T., Chang, S.C., Hattori, A., York, I.A., Markant, S., Rock, K.L., Tsujimoto, M. and Goldberg, A.L. (2002) An IFN-gamma-induced aminopeptidase in the ER, ERAP1, trims precursors to MHC class I-presented peptides. *Nat Immunol*, **3**, 1169-1176.
- Saurin, W., Hofnung, M. and Dassa, E. (1999) Getting in or out: early segregation between importers and exporters in the evolution of ATP-binding cassette (ABC) transporters. *J Mol Evol*, **48**, 22-41.
- Saveanu, L., Daniel, S. and van Endert, P.M. (2001) Distinct functions of the ATP binding cassettes of transporters associated with antigen processing: a mutational analysis of Walker A and B sequences. *J Biol Chem*, **276**, 22107-22113.
- Schneider, E. and Hunke, S. (1998) ATP-binding-cassette (ABC) transport systems: functional and structural aspects of the ATP-hydrolyzing subunits/domains. *FEMS Microbiol Rev*, **22**, 1-20.
- Schwartz, D.C. and Hochstrasser, M. (2003) A superfamily of protein tags: ubiquitin, SUMO and related modifiers. *Trends Biochem Sci*, **28**, 321-328.
- Seliger, B., Maeurer, M.J. and Ferrone, S. (1997) TAP off--tumors on. *Immunol Today*, **18**, 292-299.

- Senior, A.E., al-Shawi, M.K. and Urbatsch, I.L. (1995) The catalytic cycle of P-glycoprotein. *FEBS Lett*, **377**, 285-289.
- Serwold, T., Gonzalez, F., Kim, J., Jacob, R. and Shastri, N. (2002) ERAAP customizes peptides for MHC class I molecules in the endoplasmic reticulum. *Nature*, **419**, 480-483.
- Shamu, C.E., Flierman, D., Ploegh, H.L., Rapoport, T.A. and Chau, V. (2001) Polyubiquitination is required for US11-dependent movement of MHC class I heavy chain from endoplasmic reticulum into cytosol. *Mol Biol Cell*, **12**, 2546-2555.
- Shamu, C.E., Story, C.M., Rapoport, T.A. and Ploegh, H.L. (1999) The pathway of US11-dependent degradation of MHC class I heavy chains involves a ubiquitin-conjugated intermediate. *J Cell Biol*, **147**, 45-58.
- Shepherd, J.C., Schumacher, T.N., Ashton-Rickardt, P.G., Imaeda, S., Ploegh, H.L., Janeway, C.A., Jr. and Tonegawa, S. (1993) TAP1-dependent peptide translocation in vitro is ATP dependent and peptide selective. *Cell*, **74**, 577-584.
- Solheim, J.C., Harris, M.R., Kindle, C.S. and Hansen, T.H. (1997) Prominence of beta 2-microglobulin, class I heavy chain conformation, and tapasin in the interactions of class I heavy chain with calreticulin and the transporter associated with antigen processing. *J Immunol*, **158**, 2236-2241.
- Spies, T., Bresnahan, M., Bahram, S., Arnold, D., Blanck, G., Mellins, E., Pious, D. and DeMars, R. (1990) A gene in the human major histocompatibility complex class II region controlling the class I antigen presentation pathway. *Nature*, **348**, 744-747.
- Srivastava, P. (2002) Interaction of heat shock proteins with peptides and antigen presenting cells: chaperoning of the innate and adaptive immune responses. *Annu Rev Immunol*, **20**, 395-425.
- Stoltze, L., Dick, T.P., Deeg, M., Pommerl, B., Rammensee, H.G. and Schild, H. (1998) Generation of the vesicular stomatitis virus nucleoprotein cytotoxic T lymphocyte epitope requires proteasome-dependent and -independent proteolytic activities. *Eur J Immunol*, **28**, 4029-4036.
- Tan, P., Kropshofer, H., Mandelboim, O., Bulbuc, N., Hammerling, G.J. and Momburg, F. (2002) Recruitment of MHC class I molecules by tapasin into the transporter associated with antigen processing-associated complex is essential for optimal peptide loading. *J Immunol*, **168**, 1950-1960.
- Teisserenc, H., Schmitt, W., Blake, N., Dunbar, R., Gadola, S., Gross, W.L., Exley, A. and Cerundolo, V. (1997) A case of primary immunodeficiency due to a defect of the

- major histocompatibility gene complex class I processing and presentation pathway. *Immunol Lett*, **57**, 183-187.
- Thompson, C.L., Hooda-Dhingra, U. and Condit, R.C. (1989) Fine structure mapping of five temperature-sensitive mutants in the 22- and 147-kilodalton subunits of vaccinia virus DNA-dependent RNA polymerase. *J Virol*, **63**, 705-713.
- Tirosh, B., Furman, M.H., Tortorella, D. and Ploegh, H.L. (2003) Protein unfolding is not a prerequisite for endoplasmic reticulum-to-cytosol dislocation. *J Biol Chem*, **278**, 6664-6672.
- Tomazin, R., Hill, A.B., Jugovic, P., York, I., van Endert, P., Ploegh, H.L., Andrews, D.W. and Johnson, D.C. (1996) Stable binding of the herpes simplex virus ICP47 protein to the peptide binding site of TAP. *Embo J*, **15**, 3256-3266.
- Tomazin, R., van Schoot, N.E., Goldsmith, K., Jugovic, P., Sempe, P., Fruh, K. and Johnson, D.C. (1998) Herpes simplex virus type 2 ICP47 inhibits human TAP but not mouse TAP. *J Virol*, **72**, 2560-2563.
- Uebel, S., Kraas, W., Kienle, S., Wiesmuller, K.H., Jung, G. and Tampe, R. (1997) Recognition principle of the TAP transporter disclosed by combinatorial peptide libraries. *Proc Natl Acad Sci U S A*, **94**, 8976-8981.
- Uebel, S., Meyer, T.H., Kraas, W., Kienle, S., Jung, G., Wiesmuller, K.H. and Tampe, R. (1995) Requirements for peptide binding to the human transporter associated with antigen processing revealed by peptide scans and complex peptide libraries. *J Biol Chem*, **270**, 18512-18516.
- Van den Berg, B., Clemons, W.M., Jr., Collinson, I., Modis, Y., Hartmann, E., Harrison, S.C. and Rapoport, T.A. (2004) X-ray structure of a protein-conducting channel. *Nature*, **427**, 36-44.
- van Endert, P.M., Saveanu, L., Hewitt, E.W. and Lehner, P. (2002) Powering the peptide pump: TAP crosstalk with energetic nucleotides. *Trends Biochem Sci*, **27**, 454-461.
- van Endert, P.M., Tampe, R., Meyer, T.H., Tisch, R., Bach, J.F. and McDevitt, H.O. (1994) A sequential model for peptide binding and transport by the transporters associated with antigen processing. *Immunity*, **1**, 491-500.
- Vos, J.C., Reits, E.A., Wojcik-Jacobs, E. and Neefjes, J. (2000) Head-head/tail-tail relative orientation of the pore-forming domains of the heterodimeric ABC transporter TAP. *Curr Biol*, **10**, 1-7.

REFERENCES

- Vos, J.C., Spee, P., Momburg, F. and Neefjes, J. (1999) Membrane topology and dimerization of the two subunits of the transporter associated with antigen processing reveal a three-domain structure. *J Immunol*, **163**, 6679-6685.
- Wiertz, E.J., Jones, T.R., Sun, L., Bogyo, M., Geuze, H.J. and Ploegh, H.L. (1996a) The human cytomegalovirus US11 gene product dislocates MHC class I heavy chains from the endoplasmic reticulum to the cytosol. *Cell*, **84**, 769-779.
- Wiertz, E.J., Tortorella, D., Bogyo, M., Yu, J., Mothes, W., Jones, T.R., Rapoport, T.A. and Ploegh, H.L. (1996b) Sec61-mediated transfer of a membrane protein from the endoplasmic reticulum to the proteasome for destruction. *Nature*, **384**, 432-438.
- Ye, Y., Shibata, Y., Yun, C., Ron, D. and Rapoport, T.A. (2004) A membrane protein complex mediates retro-translocation from the ER lumen into the cytosol. *Nature*, **429**, 841-847.
- Yewdell, J.W., Anton, L.C. and Bennink, J.R. (1996) Defective ribosomal products (DRiPs): a major source of antigenic peptides for MHC class I molecules? *J Immunol*, **157**, 1823-1826.
- Yin, Y., Manoury, B. and Fahraeus, R. (2003) Self-inhibition of synthesis and antigen presentation by Epstein-Barr virus-encoded EBNA1. *Science*, **301**, 1371-1374.
- York, I.A., Roop, C., Andrews, D.W., Riddell, S.R., Graham, F.L. and Johnson, D.C. (1994) A cytosolic herpes simplex virus protein inhibits antigen presentation to CD8+ T lymphocytes. *Cell*, **77**, 525-535.
- Yuan, Y.R., Blecker, S., Martsinkevich, O., Millen, L., Thomas, P.J. and Hunt, J.F. (2001) The crystal structure of the MJ0796 ATP-binding cassette. Implications for the structural consequences of ATP hydrolysis in the active site of an ABC transporter. *J Biol Chem*, **276**, 32313-32321.
- Zeidler, R., Eissner, G., Meissner, P., Uebel, S., Tampe, R., Lazis, S. and Hammerschmidt, W. (1997) Downregulation of TAP1 in B lymphocytes by cellular and Epstein-Barr virus-encoded interleukin-10. *Blood*, **90**, 2390-2397.
- Zimmermann, A., Trilling, M., Wagner, M., Wilborn, M., Bubic, I., Jonjic, S., Koszinowski, U. and Hengel, H. (2005) A cytomegaloviral protein reveals a dual role for STAT2 in IFN-{gamma} signaling and antiviral responses. *J Exp Med*, **201**, 1543-1553.