

Larval Development, the Origin of the Coelom and the Phylogenetic Relationships of the Phoronida

Dissertation
zur Erlangung des Doktorgrades
am Fachbereich Biologie, Chemie und Pharmazie
der Freien Universität Berlin

vorgelegt von
Peter Grobe
November 2007



Phoronopsis harmeri (Phoronida)

1. Gutachter: Prof. Dr. Thomas Bartolomaeus (Freie Universität Berlin)
2. Gutachter: PD Dr. Andreas Schmidt-Rhaesa (Universität Hamburg)

Datum der Disputation: 21.12.2007

Contents

General Introduction	5
Coelomogenesis in young larvae of <i>Phoronis ovalis</i> and <i>P. muelleri</i>	
Introduction	13
Material and Methods	15
Terminology	16
Results	
Coelomogenesis in <i>P. ovalis</i>	17
Overview	17
Gastrulation	20
Young larva	22
Late larva	27
Nephridia in the larva of <i>P. ovalis</i>	30
Origin of the coelom in <i>P. muelleri</i>	33
Overview	33
Gastrula stage	34
Young larva after gastrulation	36
Two lobed larva	37
Discussion	
Origin of the coelom in Phoronida	38
Homology of <i>P. ovalis</i> larva with the actinotroch larva	39
The condition in the ground pattern of Phoronida	40
Literature	42

Contents

Morphological and molecular affinities of Phoronida and Brachiopoda	
Introduction	45
Material and Methods	
Examined taxa	46
Molecular data	47
Morphological data	48
Phylogenetic analysis	49
Results	
Morphological data	52
Combined data	53
Sequence data	55
Phoroniformea	57
Character congruence	57
Character descriptions	58
Discussion	
Groundpattern of the Phoronida	72
Evolution of characters within Phoronida	75
The Phoroniformea hypothesis	76
Literature	77
Summary	85
Zusammenfassung	87
Appendix	89