

## 6 Literatur

- Aboud, M., Segal, S., et al. (1992). "Effect of temperature on the expression of major histocompatibility complex class-I antigens." *Immunological Investigations* **21**(3): 219-229.
- Altmeyer, A., Maki, R.G., et al. (1996). "Tumor-specific cell surface expression of the-KDEL containing, endoplasmic reticular heat shock protein gp96." *Int J Cancer* **69**(4): 340-9.
- Arnold, D., Faath, S., et al. (1995). "Cross-priming of minor histocompatibility antigen-specific cytotoxic T cells upon immunization with the heat shock protein gp96." *J Exp Med* **182**(3): 885-9.
- Arnold, D., Wahl, C., et al. (1997). "Influences of transporter associated with antigen processing (TAP) on the repertoire of peptides associated with the endoplasmic reticulum-resident stress protein gp96." *J Exp Med* **186**(3): 461-6.
- Asea, A., Kraeft, S.K., et al. (2000). "HSP70 stimulates cytokine production through a CD14-dependant pathway, demonstrating its dual role as a chaperone and cytokine." *Nat Med* **6**(4): 435-42.
- Avery, M., Nathanson, S.D., et al. (1992). "Lymph flow from murine footpad tumors before and after sublethal hyperthermia." *Radiation Research* **132**(1): 50-3.
- Batalia, M.A., Kirksey, T.J., et al. (2000). "Class I MHC is stabilized against thermal denaturation by physiological concentrations of NaCl." *Biochemistry* **39**(30): 9030-8.
- Becker, N. and Wahrendorf, J.r. (1998). *Krebsatlas der Bundesrepublik Deutschland 1981-1990*. Heidelberg, Springer Verlag.
- Bercovich, B., Stancovski, I., et al. (1997). "Ubiquitin-dependent degradation of certain protein substrates in vitro requires the molecular chaperone Hsc70." *J Biol Chem* **272**(14): 9002-10.
- Bidri, M., Royer, B., et al. (1999). "Inhibition of mouse mast cell proliferation and proinflammatory mediator release by benzodiazepines." *Immunopharmacology* **43**(1): 75-86.
- Binder, R.J., Han, D.K., et al. (2000). "CD91: a receptor for heat shock protein gp96." *Nat Immunol* **1**(2): 151-5.

- Blachere, N.E., Li, Z., et al. (1997). "Heat shock protein-peptide complexes, reconstituted in vitro, elicit peptide-specific cytotoxic T lymphocyte response and tumor immunity." *Journal of Experimental Immunology* **186**(8): 1315-1322.
- Bloom, M.B., Perry-Lalley, D., et al. (1997). "Identification of tyrosinase-related protein 2 as a tumor rejection antigen for the B16 melanoma." *J Exp Med* **185**(3): 453-9.
- Boehm, U., Klamp, T., et al. (1997). "Cellular responses to interferon-gamma." *Annu Rev Immunol* **15**: 749-95.
- Botzler, C., Li, G., et al. (1998). "Definition of extracellular localized epitopes of Hsp70 involved in an NK immune response." *Cell Stress Chaperones* **3**(1): 6-11.
- Breloer, M., Fleischer, B., et al. (1999). "In vivo and in vitro activation of T cells after administration of Ag-negative heat shock proteins." *J Immunol* **162**(6): 3141-7.
- Breloer, M., Marti, T., et al. (1998). "Isolation of processed, H-2K<sup>b</sup>-binding ovalbumin-derived peptides associated with the stress proteins HSP70 and GP96." *European Journal of Immunology* **28**: 1016-1021.
- Brichard, V., Van Pel, A., et al. (1993). "The tyrosinase gene codes for an antigen recognized by autologous cytolytic T lymphocytes on HLA-A2 melanomas." *J Exp Med* **178**(2): 489-95.
- Busch, W. (1868). "Über den Einfluß, welche heftigere Erysepelen zuweilen auf organisierte Neubildungen ausüben." *Berliner Klinische Wochenschrift* **3**: 28-30.
- Chen, D. and Androlewicz, M.J. (2001). "Heat shock protein 70 moderately enhances peptide binding and transport by the transporter associated with antigen processing." *Immunol Lett* **75**(2): 143-8.
- Chen, W., Syldath, U., et al. (1999). "Human 60-kDa heat-shock protein: a danger signal to the innate immune system." *J Immunol* **162**(6): 3212-9.
- Chouchane, L., Ahmed, S.B., et al. (1997). "Polymorphism in the tumor necrosis factor-alpha promotor region and in the heat shock protein 70 genes associated with malignant tumors." *Cancer* **80**(8): 1489-96.
- Dasgupta, J.D., Watkins, S., et al. (1988). "Receptor-like nature of class I HLA: endocytosis via coated pits." *J Immunol* **141**(8): 2577-80.

- DeMatos, P., Abdel-Wahab, Z., et al. (1998). "Pulsing of dendritic cells with cell lysates from either B16 melanoma or MCA-106 fibrosarcoma yields equally effective vaccines against B16 tumors in mice." *J Surg Oncol* **68**(2): 79-91.
- Dietzel, F. (1983). "Basic principles in hyperthermic tumor therapy." *Recent Results Cancer Res* **86**: 177-90.
- Dressel, R., Elsner, L., et al. (2000). "Heat shock protein 70 is able to prevent heat shock-induced resistance of target cells to CTL." *J Immunol* **164**(5): 2362-71.
- Dyall, R., Bowne, W.B., et al. (1998). "Heteroclitic immunization induces tumor immunity." *J Exp Med* **188**(9): 1553-61.
- Ehrlich, R., Smorodinsky, N., et al. (1984). "B16 melanoma development, NK activity cytostasis and natural antibodies in 3 and 12 month old mice." *Br J Cancer* **49**(6): 769-77.
- Eisenthal, A., Lafreniere, R., et al. (1987). "Effect of anti-B16 melanoma monoclonal antibody on established murine B16 melanoma liver metastases." *Cancer Res* **47**(11): 2771-6.
- Eisenthal, A. and McIntosh, J.K. (1990). "Effect of recombinant human tumor necrosis factor alpha on the induction of antibody-dependent cellular cytotoxicity in the treatment of established B16 melanoma liver nodules." *Cancer Immunol Immunother* **31**(4): 243-9.
- Feng, H., Zeng, Y., et al. (2001). "Stressed apoptotic tumor cells express heat shock proteins and elicit tumor-specific immunity." *Blood* **97**(11): 3505-12.
- Ferrarini, M., Heltai, S., et al. (1992). "Unusual expression and localization of heat-shock proteins in human tumor cells." *Int J Cancer* **51**(4): 613-9.
- Fidler, I.J. (1975). "Biological behavior of malignant melanoma cells correlated to their survival in vivo." *Cancer Res* **35**(1): 218-24.
- Field, S.B. (1988). "The concept of thermal dose." *Recent Results Cancer Research* **107**: 1-6.
- Fisher, E.A., Zhou, M., et al. (1997). "The degradation of apolipoprotein B100 is mediated by the ubiquitin-proteasome pathway and involves heat shock protein 70." *J Biol Chem* **272**(33): 20427-34.
- Fishman, D., Irena, B., et al. (2001). "The role of MHC class I glycoproteins in the regulation of induction of cell death in immunocytes by malignant melanoma cells." *Proc Natl Acad Sci U S A* **98**(4): 1740-4.

- Gabathuler, R., Reid, G., et al. (1994). "Comparison of cell lines deficient in antigen presentation reveals a functional role for TAP-1 alone in antigen processing." *J Exp Med* **180**(4): 1415-25.
- Garbe, C. and Krasagakis, K. (1993). "Effects of interferons and cytokines on melanoma cells." *J Invest Dermatol* **100**(2 Suppl): 239S-244S.
- Garrido, C., Gurbuxani, S., et al. (2001). "Heat shock proteins: endogenous modulators of apoptotic cell death." *Biochem Biophys Res Commun* **286**(3): 433-42.
- Geertsen, R., Boni, R., et al. (2002). "Loss of single HLA class I allospecificities in melanoma cells due to selective genomic abbreviations." *Int J Cancer* **99**(1): 82-7.
- Geertsen, R.C., Hofbauer, G.F., et al. (1998). "Higher frequency of selective losses of HLA-A and -B allospecificities in metastasis than in primary melanoma lesions." *J Invest Dermatol* **111**(3): 497-502.
- Gough, M.J., Melcher, A.A., et al. (2001). "Macrophages orchestrate the immune response to tumor cell death." *Cancer Res* **61**(19): 7240-7.
- Green, D.M., Samulski, T.V., et al. (1990). "Effects of hyperthermia and irradiation on the growth of mouse melanoma tumors following immunotherapy." *Journal Of Surgical Oncology* **44**(4): 256-9.
- Gromme, M., Uytdehaag, F.G., et al. (1999). "Recycling MHC class I molecules and endosomal peptide loading." *Proc Natl Acad Sci U S A* **96**(18): 10326-31.
- Halaban, R., Cheng, E., et al. (1997). "Aberrant retention of tyrosinase in the endoplasmic reticulum mediates accelerated degradation of the enzyme and contributes to the dedifferentiated phenotype of amelanotic melanoma cells." *Proc Natl Acad Sci U S A* **94**(12): 6210-5.
- Harris, A.L. (2002). "Hypoxia--a key regulatory factor in tumour growth." *Nat Rev Cancer* **2**(1): 38-47.
- Hazenberg, M.D., Verschuren, M.C., et al. (2001). "T cell receptor excision circles as markers for recent thymic emigrants: basic aspects, technical approach, and guidelines for interpretation." *J Mol Med* **79**(11): 631-40.
- Hobohm, U. (2001). "Fever and cancer in perspective." *Cancer Immunol Immunother* **50**(8): 391-6.

- Hochman, J.H., Jiang, H., et al. (1991). "Endocytosis and dissociation of class I MHC molecules labeled with fluorescent beta-2 microglobulin." *J Immunol* **146**(6): 1862-7.
- Hosken, N.A. and Bevan, M.J. (1992). "An endogenous antigenic peptide bypasses the class I antigen presentation defect in RMA-S." *J Exp Med* **175**(3): 719-29.
- Hosono, M.N., Hosono, M., et al. (1994). "Effect of hyperthermia on tumor uptake of radiolabeled anti-neural cell adhesion molecule antibody in small-cell lung cancer xenografts." *Journal Of Nuclear Medicine* **35**(3): 504-9.
- Iezzi, G., Rivolta, L., et al. (1997). "The immunogenicity of experimental tumors is strongly biased by the expression of dominant viral cytotoxic T-lymphocyte epitopes." *Cancer Res* **57**(13): 2564-8.
- Imamura, M., Seki, T., et al. (1998). "Change of temperature in mouse tumour tissue immersed in a water bath at 44 degrees C." *Oncol Rep* **5**(5): 1061-4.
- Imani, F. and Soloski, M.J. (1991). "Heat shock proteins can regulate expression of the Tla region-encoded class Ib molecule Qa-1." *Proc Natl Acad Sci U S A* **88**(23): 10475-9.
- Ishii, T., Udon, H., et al. (1999). "Isolation of MHC class I restricted tumor antigen peptide and its precursors associated with heat shock proteins hsp70, hsp90 and gp96." *The Journal of Immunology* **162**: 1303-1309.
- Jiang, H. and Chess, L. (2000). "The specific regulation of immune responses by CD8+ T cells restricted by the MHC class Ib molecule, Qa-1." *Annu Rev Immunol* **18**: 185-216.
- Johnsen, A., France, J., et al. (1998). "Down-regulation of the transporter for antigen presentation, proteasome subunits, and class I major histocompatibility complex in tumor cell lines." *Cancer Res* **58**(16): 3660-7.
- Jordan, A., Scholz, R., et al. (1997). "Effects of magnetic fluid hyperthermia (MFH) on C3H mammary carcinoma in vivo." *International Journal Of Hyperthermia* **13**(6): 587-605.
- Kelsall, S.R., Le Fur, N., et al. (1997). "Qualitative and quantitative catalog of tyrosinase alternative transcripts in normal murine skin melanocytes as a basis for detecting melanoma-specific changes." *Biochem Biophys Res Commun* **236**(1): 173-7.
- Kim, H.T., Nelson, E.L., et al. (1995). "Gamma delta T cell recognition of tumor Ig peptide." *J Immunol* **154**(4): 1614-23.

- Kölmel, K., Vehmeyer, K., et al. (1991). "Treatment of advanced malignant melanoma by a pyrogenic bacterial lysate. A pilot study." *Onkologie* **14**: 411.
- Kölmel, K.F., Pfahlberg, A., et al. (1999). "Infections and melanoma risk: results of a multicentre EORTC case-control study. European Organization for Research and Treatment of Cancer." *Melanoma Res* **9**(5): 511-9.
- Kroeger, P.E., Sarge, K.D., et al. (1993). "Mouse heat shock transcription factors 1 and 2 prefer a trimeric binding site but interact differently with the HSP70 heat shock element." *Mol Cell Biol* **13**(6): 3370-83.
- Kurz, A., Sessler, D.I., et al. (1995). "Midazolam minimally impairs thermoregulatory control." *Anesth Analg* **81**(2): 393-8.
- Laad, A.D., Thomas, M.L., et al. (1999). "Human gamma delta T cells recognize heat shock protein-60 on oral tumor cells." *Int J Cancer* **80**(5): 709-14.
- Lage, H., Jordan, A., et al. (2000). "Thermosensitivity of multidrug-resistant human gastric and pancreatic carcinoma cells." *Int J Hyperthermia* **16**(4): 291-303.
- Lammert, E., Arnold, D., et al. (1997). "The endoplasmic reticulum-resident stress protein gp96 binds peptides translocated by TAP." *Eur J Immunol* **27**(4): 923-7.
- Leibovici, J., Klorin, G., et al. (1991). "Differential sensitivity to hyperthermia of the F1 and F10 B16 melanoma variants." *Int J Exp Pathol* **72**(2): 139-50.
- Linderoth, N.A., Popowicz, A., et al. (2000). "Identification of the peptide-binding site in the heat shock chaperone/tumor rejection antigen gp96 (Grp94)." *J Biol Chem* **275**(8): 5472-7.
- Liu, Y., Lillehei, K., et al. (2001). "Overcoming MDR by ultrasound-induced hyperthermia and P-glycoprotein modulation." *Biochem Biophys Res Commun* **289**(1): 62-8.
- Ljunggren, H.G. and Karre, K. (1985). "Host resistance directed selectively against H-2-deficient lymphoma variants. Analysis of the mechanism." *J Exp Med* **162**(6): 1745-59.
- Lu, X., Michaud, C., et al. (2001). "Heat shock protein-90 and the catalytic activities of the 20 S proteasome (multicatalytic proteinase complex)." *Arch Biochem Biophys* **387**(1): 163-71.
- Ma, D. and Ariyan, S. (2000). "The use of isolated limb perfusion to manage recurrent malignant melanoma." *Clin Plast Surg* **27**(3): 441-50, ix.

- Martinez-Esparza, M., Jimenez-Cervantes, C., et al. (1997). "Transforming growth factor-beta1 inhibits basal melanogenesis in B16/F10 mouse melanoma cells by increasing the rate of degradation of tyrosinase and tyrosinase-related protein-1." *J Biol Chem* **272**(7): 3967-72.
- Martinez-Esparza, M., Jimenez-Cervantes, C., et al. (1998). "Mechanisms of melanogenesis inhibition by tumor necrosis factor-alpha in B16/F10 mouse melanoma cells." *Eur J Biochem* **255**(1): 139-46.
- Mathew, A., Mathur, S.K., et al. (2001). "Stress-specific activation and repression of heat shock factors 1 and 2." *Mol Cell Biol* **21**(21): 7163-71.
- Matsumoto, Y., Suzuki, N., et al. (1997). "A possible mechanism for hyperthermic radiosensitization mediated through hyperthermic lability of Ku subunits in DNA-dependent protein kinase." *Biochem Biophys Res Commun* **234**(3): 568-72.
- Mehdi, S.Q. and Hahn, G.M. (1990). "Effects of hyperthermic temperatures and the synthesis of heat-shock proteins on the lateral diffusion of H-2Kk." *Int J Hyperthermia* **6**(3): 553-61.
- Melcher, A., Todryk, S., et al. (1998). "Tumor immunogenicity is determined by the mechanism of cell death via induction of heat shock protein expression." *Nature Medicine* **4**(5): 581-587.
- Mestiri, S., Bouaouina, N., et al. (2001). "Genetic variation in the tumor necrosis factor-alpha promoter region and in the stress protein hsp70-2: susceptibility and prognostic implications in breast carcinoma." *Cancer* **91**(4): 672-8.
- Mosse, C.A., Meadows, L., et al. (1998). "The class I antigen-processing pathway for the membrane protein tyrosinase involves translation in the endoplasmic reticulum and processing in the cytosol." *J Exp Med* **187**(1): 37-48.
- Multhoff, G. (1997). "Heat shock protein 72 (HSP72), a hyperthermia-inducible immunogenic determinant on leukemic K562 and Ewing's Sarcoma cells." *International Journal of Hyperthermia* **13**(1): 39-48.
- Multhoff, G., Botzler, C., et al. (1997). "Heat shock protein 72 on tumor cells: a recognition structure for natural killer cells." *J Immunol* **158**(9): 4341-50.
- Multhoff, G., Mizzen, L., et al. (1999). "Heat shock protein 70 (Hsp70) stimulates proliferation and cytolytic activity of natural killer cells." *Exp Hematol* **27**(11): 1627-36.

- Nashan, D. and Luger, T.A. (2001). "Zytokine: Stand und Perspektiven in der Behandlung von Hauttumoren." *Hautarzt* **52**(8): 691-6.
- Nathanson, S.D., Haas, G.P., et al. (1987). "Regional lymph node and pulmonary metastases after local hyperthermia of melanomas in C57BL/6 mice." *Int J Radiat Oncol Biol Phys* **13**(2): 243-9.
- Nathanson, S.D., Nelson, L., et al. (1991). "Development of lymph node and pulmonary metastases after local irradiation and hyperthermia of footpad melanomas." *Clin Exp Metastasis* **9**(4): 377-92.
- Obrig, T.G., Culp, W.J., et al. (1971). "The mechanism by which cycloheximide and related glutarimide antibiotics inhibit peptide synthesis on reticulocyte ribosomes." *J Biol Chem* **246**(1): 174-81.
- Ohtsubo, T., Igawa, H., et al. (2001). "Acidic environment modifies heat- or radiation-induced apoptosis in human maxillary cancer cells." *Int J Radiat Oncol Biol Phys* **49**(5): 1391-8.
- Ohtsubo, T., Park, H.J., et al. (2000). "Effect of acidic environment and p53 on apoptosis induction by hyperthermia." *Int J Hyperthermia* **16**(6): 481-91.
- Okazaki, K., Davis, D.D., et al. (1987). "T cell receptor beta gene sequences in the circular DNA of thymocyte nuclei: direct evidence for intramolecular DNA deletion in V-D-J joining." *Cell* **49**(4): 477-85.
- Oliveira Filho, R.S., Bevilacqua, R.G., et al. (1997). "Hyperthermia increases the metastatic potential of murine melanoma." *Brazilian Journal Of Medical And Biological Research* **30**(8): 941-5.
- Ostrand-Rosenberg, S. and Kruisbeek, A.M. (2000). Animal Models for Tumor Immunology. Current Protocols in Immunology, John Wiley & Sons. **Supplement 39:** 20.0.1-20.0.8.
- Overwijk, W.W. and Restifo, N.P. (2000). B16 s a mouse model for human melanoma. Current Protocols in Immunology. J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach and W. Strober, John Wiley & Sons, Inc.: 20.1.1-29.
- Overwijk, W.W., Tsung, A., et al. (1998). "gp100/pmel 17 is a murine tumor rejection antigen: induction of "self"-reactive, tumoricidal T cells using high-affinity, altered peptide ligand." *J Exp Med* **188**(2): 277-86.

- Owen, B.A. and Pease, L.R. (2001). "Thermal stability of MHC class I-beta 2-microglobulin peptide complexes in the endoplasmic reticulum is determined by the peptide occupancy of the transporter associated with antigen processing complex." *J Immunol* **166**(3): 1740-7.
- Panjwani, N.N., Popova, L., et al. (2002). "Heat shock proteins gp96 and hsp70 activate the release of nitric oxide by APCs." *J Immunol* **168**(6): 2997-3003.
- Peck, J.W. and Gibbs, F.A.J. (1983). "Capillary blood flow in murine tumors, feet, and intestines during localized hyperthermia." *Radiation Research* **96**(1): 65-81.
- Peter, I., Mezzacasa, A., et al. (2001). "Comparative analysis of immunocritical melanoma markers in the mouse melanoma cell lines B16, K1735 and S91-M3." *Melanoma Res* **11**(1): 21-30.
- Rammensee, H., Bachmann, J., et al. (1999). "SYFPEITHI: database for MHC ligands and peptide motifs." *Immunogenetics* **50**(3-4): 213-9.
- Reich, Z., Altman, J.D., et al. (1997). "Stability of empty and peptide-loaded class II major histocompatibility complex molecules at neutral and endosomal pH: comparison to class I proteins." *Proc Natl Acad Sci U S A* **94**(6): 2495-500.
- Rippmann, F., Taylor, W.R., et al. (1991). "A hypothetical model for the peptide binding domain of hsp70 based on the peptide binding domain of HLA." *Embo J* **10**(5): 1053-9.
- Robert, J., Menoret, A., et al. (1999). "Cell surface expression of the endoplasmic reticular heat shock protein gp96 is phylogenetically conserved." *J Immunol* **163**(8): 4133-9.
- Schaadt, J., Crowley, R., et al. (2002). "Isolated limb perfusion: a literature review." *J Extra Corpor Technol* **34**(2): 130-43.
- Seliger, B., Harders, C., et al. (1998). "Down-regulation of the MHC class I antigen-processing machinery after oncogenic transformation of murine fibroblasts." *Eur J Immunol* **28**(1): 122-33.
- Seliger, B., Wollscheid, U., et al. (2001). "Characterization of the major histocompatibility complex class I deficiencies in B16 melanoma cells." *Cancer Res* **61**(3): 1095-9.
- Sempowski, G.D., Gooding, M.E., et al. (2002). "T cell receptor excision circle assessment of thymopoiesis in aging mice." *Mol Immunol* **38**(11): 841-8.

- Shrayer, D., Kouttab, N., et al. (1993). "Generation of cytotoxic antibodies to the B16 murine melanoma using a formalinized vaccine." *Int J Cancer* **53**(4): 696-702.
- Singh-Jasuja, H., Scherer, H.U., et al. (2000). "The heat shock protein gp96 induces maturation of dendritic cells and down-regulation of its receptor." *Eur J Immunol* **30**(8): 2211-5.
- Srivastava, P.K., Udon, H., et al. (1994). "Heat shock proteins transfer peptides during antigen processing and CTL priming." *Immunogenetics* **39**(2): 93-8.
- Stein, U., Rau, B., et al. (1999). "Hyperthermia for treatment of rectal cancer: evaluation for induction of multidrug resistance gene (mdr1) expression." *Int J Cancer* **80**(1): 5-12.
- Stewart, J.R., Gibbs, F.A.J., et al. (1983). "Change in the in vivo hyperthermic response resulting from the metabolic effects of temporary vascular occlusion." *International Journal Of Radiation Oncology, Biology, Physics* **9**(2): 197-201.
- Stojkovic, R. and Radacic, M. (2002). "Cell killing of melanoma B16 in vivo by hyperthermia and cytotoxins." *Int J Hyperthermia* **18**(1): 62-71.
- Stubbs, M., McSheehy, P.M., et al. (2000). "Causes and consequences of tumour acidity and implications for treatment." *Mol Med Today* **6**(1): 15-9.
- Suto, R. and Srivastava, P.K. (1995). "A mechanism for the specific immunogenicity of heat shock protein-chaperoned peptides." *Science* **269**: 1585-1588.
- Takahashi, I., Emi, Y., et al. (2002). "Clinical application of hyperthermia combined with anticancer drugs for the treatment of solid tumors." *Surgery* **131**(1 Suppl): S78-84.
- The Jackson Laboratory, B.H., USA Mouse Genome Informatics, The Jackson Laboratory, Bar Harbor, USA. **2001**.
- Thomas, M.L., Samant, U.C., et al. (2000). "gammadelta T cells lyse autologous and allogenic oesophageal tumours: involvement of heat-shock proteins in the tumour cell lysis." *Cancer Immunol Immunother* **48**(11): 653-9.
- Todryk, S., Melcher, A.A., et al. (1999). "Heat shock protein 70 induced during tumor cell killing induces Th1 cytokines and targets immature dendritic cell precursors to enhance antigen uptake." *J Immunol* **163**(3): 1398-408.

- Todryk, S.M., Melcher, A.A., et al. (2000). "Heat shock proteins refine the danger theory." *Immunology* **99**(3): 334-7.
- Tsukamoto, K., Hirata, S., et al. (2000). "Detection of circulating melanoma cells by RT-PCR amplification of three different melanocyte-specific mRNAs in a mouse model." *Pigment Cell Res* **13**(3): 185-9.
- Udono, H. and Srivastava, P.K. (1993). "Heat shock protein 70-associated peptides elicit specific cancer immunity." *J Exp Med* **178**(4): 1391-6.
- Vance, R.E., Kraft, J.R., et al. (1998). "Mouse CD94/NKG2A is a natural killer cell receptor for the nonclassical major histocompatibility complex (MHC) class I molecule Qa-1(b)." *J Exp Med* **188**(10): 1841-8.
- Vega, M.A. and Strominger, J.L. (1989). "Constitutive endocytosis of HLA class I antigens requires a specific portion of the intracytoplasmic tail that shares structural features with other endocytosed molecules." *Proc Natl Acad Sci U S A* **86**(8): 2688-92.
- Wahl, A., Brustad, T., et al. (1979). "Effect of hyperthermia alone and in combination with <sup>60</sup>Co radiation on the growth of B16 melanoma in mice." *Acta Radiol Oncol Radiat Phys Biol* **18**(4): 343-56.
- Wassenberg, J.J., Dezfulian, C., et al. (1999). "Receptor mediated and fluid phase pathways for internalization of the ER Hsp90 chaperone GRP94 in murine macrophages." *J Cell Sci* **112**(Pt 13): 2167-75.
- Weber, J.S., Salinger, A., et al. (2000). "Optimal N-ethyl-N-nitrosourea (ENU) doses for inbred mouse strains." *Genesis* **26**(4): 230-3.
- Wei, Y., Zhao, X., et al. (1996). "Induction of autologous tumor killing by heat treatment of fresh human tumor cells: involvement of gamma delta T cells and heat shock protein 70." *Cancer Res* **56**(5): 1104-10.
- Welch, W.J. and Mizzen, L.A. (1988). "Characterization of the thermotolerant cell. II. Effects on the intracellular distribution of heat-shock protein 70, intermediate filaments, and small nuclear ribonucleoprotein complexes." *J Cell Biol* **106**(4): 1117-30.
- Zolzer, F. and Streffer, C. (1999). "Radiation and/or hyperthermia sensitivity of human melanoma cells grown for several days in media with reduced pH." *Strahlenther Onkol* **175**(7): 325-32.