

# Lewis Superacidic Tellurenyl Cation-Induced Electrophilic Activation of an Inert Carborane

Martin Hejda,<sup>\*[a, b]</sup> Daniel Duvinage,<sup>[a]</sup> Enno Lork,<sup>[a]</sup> Antonín Lyčka,<sup>[c]</sup> Zdeněk Černošek,<sup>[b]</sup> Jan Macháček,<sup>[d]</sup> Sergey Makarov,<sup>[e]</sup> Sergey Ketkov,<sup>\*[e]</sup> Stefan Mebs,<sup>\*[f]</sup> Libor Dostál,<sup>[b]</sup> and Jens Beckmann<sup>\*[a]</sup>

**Abstract:** The aryltellurenyl cation  $[2-(t\text{BuNCH})\text{C}_6\text{H}_4\text{Te}]^+$ , a Lewis super acid, and the weakly coordinating carborane anion  $[\text{CB}_{11}\text{H}_{12}]^-$ , an extremely weak Brønsted acid ( $\text{p}K_{\text{a}} = 131.0$  in MeCN), form an isolable ion pair complex  $[2-(t\text{BuNCH})\text{C}_6\text{H}_4\text{Te}][\text{CB}_{11}\text{H}_{12}]$ , in which the Brønsted acidity ( $\text{p}K_{\text{a}} = 7.4$  in MeCN) of the formally hydridic B–H bonds is dramatically increased by more than 120 orders of magnitude. The electrophilic activation of B–H bonds in the carborane moiety gives rise to a proton transfer from boron to nitrogen at slightly elevated temperatures, as rationalized by the isolation of a mixture of the zwitterionic isomers 12- and 7-[2-(tBuN{H}CH)C<sub>6</sub>H<sub>4</sub>Te(CB<sub>11</sub>H<sub>11</sub>)] in ratios ranging from 62:38 to 80:20.


Main-group elements mediating bond activation of small molecules and catalytic transformations have attracted considerable attention in recent years.<sup>[1]</sup> That holds particularly for the cooperative reactivity of Lewis acids and Lewis bases that are restricted to form energetically favourable donor–acceptor complexes. Amongst those, the most prominent are arguably the frustrated Lewis pairs (FLPs), in which bulky substituents prevent the formation of stable (and unreactive) donor–acceptor bonds. Besides FLPs, there is a growing number of regular, yet reactive Lewis pairs that are capable of activating small molecules.<sup>[2]</sup>

The aryltellurenyl cation,  $[2-(t\text{BuNCH})\text{C}_6\text{H}_4\text{Te}]^+$  (I), containing an imino donor functionality can be regarded as an intramolecular regular N→Te Lewis pair, however, the intramolecularly coordinating 2-*tert*-butyl-iminomethylphenyl group compensates the electron deficiency at the tellurium atom only insufficiently (Figure 1).<sup>[3]</sup> Despite the N-donor coordination and the aromatic character of the five-membered C<sub>3</sub>NTe ring, I is a highly electrophilic Lewis superacid that gives rise to ion pairs even with weakly coordinating anions (WCAs).<sup>[3]</sup> These ion pairs  $[2-(t\text{BuNCH})\text{C}_6\text{H}_4\text{Te}][\text{X}]$  ( $[\text{X}]^- = [\text{O}_3\text{SCF}_3]^-$ ,  $[\text{SbF}_6]^-$  and  $[\text{Al}\{\text{OC}(\text{CF}_3)_3\}_4]^-$ ) show significant Te⋯O and Te⋯F interactions in the solid state, whereas in nonpolar solvents the electrolytic dissociation remains incomplete. In an effort to obtain an essentially isolated  $[2-(t\text{BuNCH})\text{C}_6\text{H}_4\text{Te}]^+$  (I) cation, we turned our attention to an alternative WCA, namely, the *closo*-carborane anion  $[\text{CB}_{11}\text{H}_{12}]^-$ , which is known as robust entity with a very low reactivity.<sup>[4]</sup> However, the salt metathesis reaction of 2-(*t*BuNCH)C<sub>6</sub>H<sub>4</sub>TeCl (II)<sup>[5]</sup> with Ag[CB<sub>11</sub>H<sub>12</sub>]<sup>[6]</sup> provided again a contact ion pair  $[2-(t\text{BuNCH})\text{C}_6\text{H}_4\text{Te}][\text{CB}_{11}\text{H}_{12}]$  (1), which was isolated as yellow crystals in 99% yield (Scheme 1, Figure 2).<sup>[7]</sup> In the solid state, the dative N→Te bond (2.088(2) Å) of 1 is shorter than that of (tBuNCH)C<sub>6</sub>H<sub>4</sub>Te[O<sub>3</sub>SCF<sub>3</sub>] (2.113(1) Å), but longer than those of (tBuNCH)C<sub>6</sub>H<sub>4</sub>Te[SbF<sub>6</sub>] (2.076(2) Å) and


- [a] Dr. M. Hejda, D. Duvinage, E. Lork, Prof. Dr. J. Beckmann  
Institut für Anorganische Chemie und Kristallographie  
Universität Bremen  
Leobener Straße 7, 28359 Bremen (Germany)  
E-mail: j.beckmann@uni-bremen.de
- [b] Dr. M. Hejda, Prof. Dr. Z. Černošek, Prof. Dr. L. Dostál  
Department of General and Inorganic Chemistry  
University of Pardubice  
Studentská 573, 532 10 Pardubice (Czech Republic)  
E-mail: martin.hejda@upce.cz
- [c] Prof. Dr. A. Lyčka  
Faculty of Science  
University of Hradec Králové  
Rokitanského 62, 500 03 Hradec Králové 3 (Czech Republic)
- [d] Dr. J. Macháček  
Institute of Inorganic Chemistry, Czech Academy of Sciences  
250 68 Řež near Prague (Czech Republic)
- [e] Dr. S. Makarov, Prof. Dr. S. Ketkov  
G. A. Razuvaev Institute of Organometallic Chemistry RAS  
49 Tropinin St., 603950 Nizhny Novgorod (Russian Federation)  
E-mail: sketkov@iomc.ras.ru
- [f] Dr. S. Mebs  
Institut für Experimentalphysik, Freie Universität Berlin  
Arnimallee 14, 14195 Berlin (Germany)  
E-mail: stebs@chemie.fu-berlin.de

Supporting information for this article is available on the WWW under <https://doi.org/10.1002/chem.202103181>

© 2021 The Authors. Chemistry - A European Journal published by Wiley-VCH GmbH. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.




**Figure 1.** Resonance formula representations and electrostatic potential (ESP) of the Lewis superacid  $[2-(t\text{BuNCH})\text{C}_6\text{H}_4\text{Te}]^+$  (I).<sup>[3]</sup>



**Scheme 1.** Reaction of 2-(*t*BuNCH) $\text{C}_6\text{H}_4\text{TeCl}$  (II) with  $\text{Ag}[\text{CB}_{11}\text{H}_{12}]$  and further transformations.

(*t*BuNCH) $\text{C}_6\text{H}_4\text{Te}[\text{Al}\{\text{OC}(\text{CF}_3)_3\}_4]$  (2.051(4) Å). The cation and anion of **1** are associated by a prominent Te...H contact (2.550(1) Å) with the B–H functionality in the B12 position.<sup>[8]</sup> As further reactivity studies showed, **1** is in fact only metastable and susceptible to further transformations at slightly elevated temperatures, both in  $\text{CH}_2\text{Cl}_2$  and the solid state. Gentle heating of **1** in inert solvents produced a mixture of two isomeric dinuclear donor-acceptor complexes [2-(*t*BuNCH) $\text{C}_6\text{H}_4\text{Te}\cdot\text{D}][\text{CB}_{11}\text{H}_{12}]$  (**2a**, D = 12-[2-(*t*BuN{H}CH) $\text{C}_6\text{H}_4\text{Te}][\text{CB}_{11}\text{H}_{11}]$  (**5a**) and **2b**, D = 7-[2-(*t*BuN{H}CH) $\text{C}_6\text{H}_4\text{Te}][\text{CB}_{11}\text{H}_{11}]$  (**5b**) in ratio 80:20,<sup>[9]</sup> which were isolated as mixed orange crystals in 49% yield (Scheme 1, Figure 2).<sup>[7,10]</sup> In the solid-state, the Te→Te bond of **2a(2b)** (3.034(1) Å) is substantially longer than that of [MesTe(TeMes<sub>2</sub>)]( $\text{O}_3\text{SCF}_3$ ) (2.808(1) Å)<sup>[11]</sup> or [MesTe(TeMes<sub>2</sub>)]( $\text{SbF}_6$ ) (2.765(1) Å)<sup>[12]</sup> due to the additional N-donation. The dative N→Te bond of **2a(2b)** (2.228(1) Å) is substantially longer than in **1**. As a result, the corresponding Te→Te vibration mode in Raman spectra of **2a(2b)** (122  $\text{cm}^{-1}$ ) is considerably shifted bathochromically in comparison to ([MesTe(TeMes<sub>2</sub>)]( $\text{SbF}_6$ ))

(143  $\text{cm}^{-1}$ ) indicating a rather weak Te–Te bond.<sup>[12]</sup> Although the donor–acceptor complex **2a(2b)** is virtually insoluble in  $\text{CH}_2\text{Cl}_2$  and aromatic solvents, it very easily dissolves in THF via dissociation of the Te→Te bond (Figures S6–17 in the Supporting Information). Consequently, the formation of 1 equiv. of [2-(*t*BuNCH) $\text{C}_6\text{H}_4\text{Te}\cdot\text{THF}][\text{CB}_{11}\text{H}_{12}]$  (**3**) is observed along with releasing of (1–*x*) equiv. of 12-[2-(*t*BuN{H}CH) $\text{C}_6\text{H}_4\text{Te}][\text{CB}_{11}\text{H}_{11}]$  (**5a**) and *x* equiv. of 7-[2-(*t*BuN{H}CH) $\text{C}_6\text{H}_4\text{Te}][\text{CB}_{11}\text{H}_{11}]$  (**5b**) from the mixture of **2a** and **2b** in molar ratio (1–*x*):*x* (Scheme 1).<sup>[13]</sup> In the presence of the donor molecules THF and DMAP, the reaction of 2-(*t*BuNCH) $\text{C}_6\text{H}_4\text{TeCl}$  with  $\text{Ag}[\text{CB}_{11}\text{H}_{12}]$  directly afforded the mononuclear donor-acceptor complexes [2-(*t*BuNCH) $\text{C}_6\text{H}_4\text{Te}\cdot\text{D}][\text{CB}_{11}\text{H}_{12}]$  **3** (D = THF) and **4** (D = DMAP) as yellowish crystals in 95% and 83% yield (Scheme 1, Figure 2).<sup>[7,14]</sup> It is worth noting that heating of the DMAP complex **4** in THF did not provide any hint of further transformation as the Lewis acidity is attenuated in comparison to **1**.<sup>[15]</sup> On the other hand, prolonged heating of the THF complex **3** in THF quantitatively provided mixtures of the isomers **5a** and **5b** (Scheme 1).



**Figure 2.** Molecular structures of **1**, **2a(2b)**, **3**, **4** and **5a(5b)** showing 50% probability ellipsoids and the atomic numbering scheme. For **2a(2b)**, **3** and **4**, the  $[\text{CB}_{11}\text{H}_{12}]^-$  anion is omitted for clarity.


Thermodynamic measurements based on  $^1\text{H}$  NMR integration within transformation of **3** upon heating at various temperatures in  $[\text{D}_8]\text{THF}$  proved its first-order kinetics (Figures S28 and 29) with the following activation parameters:  $\Delta G_{298}^\ddagger = 118.4 \text{ kJ mol}^{-1}$ ;  $\Delta H^\ddagger = 114.9 \text{ kJ mol}^{-1}$ ;  $\Delta S^\ddagger = -11.8 \text{ J mol}^{-1} \text{ K}^{-1}$ . The most straightforward way providing the quantitative yield of the zwitterionic species **5a** and **5b** (in a ratio of 62:38) was heating of complex **3** in the solid state at  $140^\circ\text{C}$  for 2.5 h, which effectively removes the THF (details in the Supporting Information). The formation of **5a** and **5b** (donors **D** in case of **2a(2b)**) may be rationalized by hydrogen transfer from the B–H functions in the B12 and B7 positions<sup>[9]</sup> of the *closo*-carborane anion  $[\text{CB}_{11}\text{H}_{12}]^-$  to the lone pair of the N atom in **1**, upon which a Te–B12(B7) bond is formed. This hydrogen transfer is facilitated by the cooperative reactivity of the intramolecular regular N→Te Lewis pair. While the Lewis acidic Te site reverses the formal polarity of the B–H bond from hydridic to protic, the Lewis basic N-atom serves as a final proton acceptor within its gradual transfer from the Te atom to the N atom (for the DFT based mechanism, see below). Thus, the Lewis super acid  $[\text{2-(tBuNCH)C}_6\text{H}_4\text{Te}]^+$  (**I**) has activated the *closo*-carborane ion  $[\text{CB}_{11}\text{H}_{12}]^-$ , which serves as a proton donor. In fact, such a bond arrangement gives a unique insight into a long unresolved issue of electrophilic activation of *closo*- $[\text{CB}_{11}\text{H}_{12}]^-$  (see below).<sup>[4]</sup> This process is accompanied not only by a significant change in color from light-yellow (**3**) to dark red, but also by a dramatic change in the  $\delta(^{125}\text{Te})$  values<sup>[15]</sup> reflecting the transition of the cationic  $\text{Te}^{\text{II}}$  site to neutral in the zwitterionic **5a(5b)** by the Te–B12(B7) bond formation. Consequently, the boron atom at

the B12 position of the free  $[\text{CB}_{11}\text{H}_{12}]^-$  anion of **3** resonating in  $^{11}\text{B}$  NMR at  $-7.3 \text{ ppm}$  is high-field shifted upon formation of Te–B12 bond in compound **5a** to  $-11.5 \text{ ppm}$ . While the unsubstituted B12 atom in case of **5b** containing Te–B7 bond resonates slightly more low-field at  $-5.8 \text{ ppm}$ , the signal of boron atom at the B7 position is shifted to high field from  $\delta(^{11}\text{B}) = -13.7 \text{ ppm}$  in **3** to  $\delta(^{11}\text{B}) = -17.3 \text{ ppm}$ . Interestingly, both Te–B12 and Te–B7 atoms in **5a** and **5b** are in  $^{11}\text{B}$  NMR significantly low-field shifted ( $\Delta\delta \sim 7 \text{ ppm}$ ) in comparison to analogously monoiodo substituted derivatives of *closo*-carbadodecaborate.<sup>[16]</sup> In the solid state **5a(5b)** features a weak intramolecular hydrogen bond of the type N–H...Te (N–H:  $0.860(2) \text{ \AA}$ , H...Te ( $2.568(1) \text{ \AA}$ , N...Te:  $3.320(1) \text{ \AA}$ ), in which Te serves as unprecedented hydrogen bond acceptor. This bonding situation is a result of the **5a(5b)** formation mechanism (see below) enforcing the *E* configuration of the protonated imino  $\text{CH}=\text{NH}^+$  moiety for both products, which is manifested by observation of typical  $^3J(^1\text{H},^1\text{H})$  values (acquired in  $\text{CD}_2\text{Cl}_2$ ) of 17.3 and 17.4 Hz, respectively. Although the intramolecular hydrogen bond of the type N–H...Te was found in the solid state, the coupling constant  $^1J(^{15}\text{N},^1\text{H}) = 85.3 \text{ Hz}$  for both **5a** and **5b** in the solution spectra lies in range for protonated imines.<sup>[17]</sup> Similarly, despite the fact that both **5a** and **5b** contain such bonding interaction, FTIR is showing no shift of the N–H bond  $\nu$  vibration in the solid state sample (**5a**:  $3244 \text{ cm}^{-1}$  **5b**:  $3295 \text{ cm}^{-1}$ ) in comparison to adducts **2a** and **2b** ( $3244 \text{ cm}^{-1}$  and  $3297 \text{ cm}^{-1}$ ) having no N–H...Te interaction. Finally, deprotonation of **5a(5b)** was achieved upon addition of triethylamine, which afforded a mixture of  $[\text{Et}_3\text{NH}][12\text{-}\{2\text{-(tBuNCH)C}_6\text{H}_4\text{Te}\}]$

CB<sub>11</sub>H<sub>11</sub>] (**6a**) and [Et<sub>3</sub>NH][12-{2-(*t*BuNCH)C<sub>6</sub>H<sub>4</sub>Te}CB<sub>11</sub>H<sub>11</sub>] (**6b**) in the same molar ratio as given by parent compounds **5a**(**5b**), which was isolated as orange oil in quantitative yield (Scheme 1). The deprotonation was shown to have a significant effect on the shielding of the nitrogen atom, as the signal with value of  $\delta(^{15}\text{N}) = -172.1$  ppm acquired in [D<sub>8</sub>]THF for **5a**(**5b**) is shifted to low field for **6a** and **6b** up to  $-35.4$  and  $-32.8$  ppm, respectively. Despite the assessable lone pair of the N atom in the **6a**(**6b**) for coordination of Te atom, we can conclude that in this case no N→Te interaction is present, as  $\delta(^{15}\text{N})$  for these compounds approaches the value of  $\delta(^{15}\text{N})$  for the unsubstituted parent Schiff base, namely (*t*BuNCH)C<sub>6</sub>H<sub>5</sub> ( $-24.3$  ppm; [D<sub>8</sub>]THF).

In an effort to shed light on the activation of the *closo*-carborane anion by the Lewis superacidic aryltellurenyl cation, two contact ion pairs, namely, **1** resembling the B12–H...Te connectivity found in the solid state, and the by  $7.6$  kJ mol<sup>-1</sup> less stable isomer **1'** featuring a B7–H...Te connectivity, were fully optimized in the gas phase (Figure 3a). The calculated B...Te distances of **1** (2.858 Å) and **1'** (2.867 Å) are significantly shorter than the experimentally observed distance of **1** (3.410(2) Å). As a result of the contact to adjacent Te atom, the B–H bond lengths of **1** (1.234) and **1'** (1.229) are substantially longer than those of the free *closo*-carborane anion (1.194 and 1.193 Å). The B–H stretching upon Te coordination is accompanied by a dramatic increase of the Brønsted acidity. As expected for a weakly coordinating anion, the calculated p*K*<sub>a</sub>

value<sup>[18]</sup> of the free *closo*-carborane (131.0) in MeCN is extremely high, rendering it an extremely weak acid. The p*K*<sub>a</sub> values of **1** (7.4) and **1'** (7.2) are dramatically smaller by more than 120 units. These p*K*<sub>a</sub> values compare well with that of HCl (7.9) in MeCN.<sup>[18]</sup> Thus, the Lewis superacidic aryltellurenyl cation, [2-(*t*BuNCH)C<sub>6</sub>H<sub>4</sub>Te]<sup>+</sup> (**1**) induces an electrophilic activation of the 12- or 7-B–H bond in the *closo*-carborane ion<sup>[4]</sup> which seems to be instrumental for the bond activation and the proton transfer to the N atoms of **5a** and **5b**. The proton transfer from **1'** to **5a** is associated with an energy gain of  $193.3$  kJ mol<sup>-1</sup> and most likely proceeds via a concerted wagging motion involving the transition state **TS** with double triangular arrangement of N, Te and B12–H, which accounts for an activation barrier of  $118.1$  kJ mol<sup>-1</sup> (Figure 3a). Such a value of activation barrier is in great agreement with the experimental one ( $\Delta G_{298}^{\ddagger} = 118.4$  kJ mol<sup>-1</sup>) as discussed above. The AIM bond topology of **TS** reveals a curved Te–H(B) bond path, indicating the onset of Te–B bond formation (Figure 3b).<sup>[19–21]</sup> With an electron density (ED,  $\rho(r)$ ) of  $0.59$  eÅ<sup>-3</sup> and considerably negative total energy over ED ratio ( $H/\rho(r)$ ) of  $-0.38$  a.u., covalent bonding aspects of the Te–H contact in the **TS** are much higher than in **1** ( $\rho(r) = 0.34$ ,  $H/\rho(r) = -0.21$  a.u.) and **5a** ( $\rho(r) = 0.23$ ,  $H/\rho(r) = -0.14$  a.u.). This is supported by the NCI, which shows a ring-shaped and red-colored NCI basin enclosing the Te–H and Te–B bonding axes, in contrast to the disc-shaped and blue-colored NCI basins in **1** and **5** (Figure 3d). In accordance with a H(Te)B arrangement, the ED within the corresponding ELI–D basin is



**Figure 3.** a) Calculated gas-phase structures, relative energies and p*K*<sub>a</sub> values of **1**, **1'**, **TS**, and **5a**. b) The atoms-in-molecules (AIM) topology, c) iso-surface representation of the electron localizability indicator (ELI–D) and d) the noncovalent interaction (NCI) index of the **TS**.



distributed over the atoms as follows: H=53%, Te=26%, and B=22%, compared to the bisynaptic contact mode in **1** (H=16%, Te=4%, B=80%). In **5a**, only the N (77%) and now protic H (23%) atoms contributions are relevant, supporting a coordinative Te...H bonding mode (Figure 3c).

In summary, the aryltellurenyl cation [2-(tBuNCH)<sub>2</sub>C<sub>6</sub>H<sub>4</sub>Te]<sup>+</sup> (**1**), and the weakly coordinating *closo*-carborane anion [CB<sub>11</sub>H<sub>12</sub>]<sup>-</sup> give rise to a metastable contact ion pair [2-(tBuNCH)<sub>2</sub>C<sub>6</sub>H<sub>4</sub>Te][CB<sub>11</sub>H<sub>12</sub>]<sup>-</sup> (**1**), in which the B–H bond in the 12- or 7-position of *closo*-[CB<sub>11</sub>H<sub>12</sub>]<sup>-</sup> is activated by the proximity of a Lewis superacidic cation, the Brønsted acidity of the *closo*-carborane is extremely increased, and this triggers proton transfer and the formation of 12-[2-(tBuN{H}CH)<sub>2</sub>C<sub>6</sub>H<sub>4</sub>Te]CB<sub>11</sub>H<sub>11</sub> (**5a**) and 7-[2-(tBuN{H}CH)<sub>2</sub>C<sub>6</sub>H<sub>4</sub>Te]CB<sub>11</sub>H<sub>11</sub> (**5b**) in ratios ranging from 62:38 to 80:20.<sup>[9]</sup> The contact ion pair **1** can be regarded as a snapshot of the first step of the electrophilic substitution of *closo*-[CB<sub>11</sub>H<sub>12</sub>]<sup>-</sup>, which is poorly understood in general.<sup>[4]</sup> Our proposed mechanism is in full agreement with the theoretically calculated mechanism of the methylation of *closo*-[CB<sub>11</sub>H<sub>12</sub>]<sup>-</sup><sup>[22]</sup> involving the formation of three-center bonded intermediate [MeBH]<sup>+</sup>. We are currently investigating the utility of **1** for the activation of small molecules.

## Acknowledgements

The Deutsche Forschungsgemeinschaft (DFG) is gratefully acknowledged for financial support. The pK<sub>a</sub> values were calculated within the frames of Project 19-03-00755 supported by the Russian Foundation for Basic Research (RFBR). The authors L.D. and M.H. thank the Czech Science Foundation (grant no. 18-10222 S). A.L. thanks the University of Hradec Králové for a support. Open Access funding enabled and organized by Projekt DEAL.

## Conflict of Interest

The authors declare no conflict of interest.

**Keywords:** bond activation · boron · carboranes · Lewis superacids · tellurium

- [1] a) P. P. Power, *Nature* **2010**, *463*, 171; b) C. Weetman, S. Inoue, *ChemCatChem* **2018**, *10*, 4213–4228.  
[2] a) D. W. Stephan, G. Erker, *Angew. Chem. Int. Ed.* **2010**, *49*, 46–76; b) D. W. Stephan *J. Am. Chem. Soc.* **2015**, *137*, 10018–10032; c) D. W. Stephan, *Acc. Chem. Res.* **2015**, *48*, 306–316; d) D. W. Stephan, G. Erker, *Angew. Chem. Int. Ed.* **2015**, *54*, 6400–6441.

- [3] M. Hejda, D. Duvinage, E. Lork, A. Lyčka, S. Mebs, L. Dostál, J. Beckmann, *Organometallics* **2020**, *39*, 1202–1212.  
[4] a) S. Körbe, P. J. Schreiber, J. Michl, *Chem. Rev.* **2006**, *106*, 5208–5249; b) C. Douvris, J. Michl, *Chem. Rev.* **2013**, *113*, PR179–PR233.  
[5] M. Hejda, E. Lork, S. Mebs, L. Dostál, J. Beckmann, *Eur. J. Inorg. Chem.* **2017**, 3435–3445.  
[6] K. Shelly, D. C. Finster, Y. J. Lee, W. R. Scheidt, C. A. Reed, *J. Am. Chem. Soc.* **1985**, *107*, 5955–5959.  
[7] Deposition Number 2092933 (**1**), 2092934 (**2a/2b**), 2092935 (**3**), 2092936 (**4**) and 2092937 (**5a/5b**) contain the supplementary crystallographic data for this paper. These data are provided free of charge by the joint Cambridge Crystallographic Data Centre and Fachinformationszentrum Karlsruhe Access Structures service. The isomerism of **2a/2b** and **5a/5b** is reflected by statistical disorder of B and C sites in the icosahedral carborane structure.  
[8] B12 refers to the *antipodal* position, whereas B7 is one of the five positions in the lower pentagonal belt.  
[9] Mutual molar ratio of **2a/2b** and **5a/5b**, respectively, is very dependent on the reaction conditions, for example, the temperature.  
[10] The <sup>125</sup>Te MAS NMR spectrum of **2a/2b** shows three signals at δ<sub>iso</sub> = 991, 802 and 374 ppm. See the Supporting Information, Figure S15.  
[11] J. Beckmann, J. Bolsinger, A. Duthie, P. Finke, E. Lork, C. Lütke, O. Mallow, S. Mebs, *Inorg. Chem.* **2012**, *51*, 12395–12406.  
[12] J. Jeske, W.-W. du Mont, F. Ruthe, P. G. Jones, L. M. Mercuri, P. Deplano, *Eur. J. Inorg. Chem.* **2000**, 1591–1599.  
[13] Obtained values of δ(<sup>125</sup>Te) for **5a** (94.0 ppm), **5b** (58.4 ppm) and **3** (1749.7 ppm) acquired by solution <sup>125</sup>Te NMR in [D<sub>8</sub>]THF significantly differ from values of **2a(2b)** given in ref. [10] and thus serves as another proof of Te→Te bond dissociation after dissolving **2a(2b)** in THF.  
[14] The donor-acceptor Te–E bonds of **2** (E=Te, 3.034(1) Å), **3** (E=O, 2.403(1) Å), **4** (E=N, 2.293(2) Å) are significantly longer than the sum of covalent radii (2.76, 2.04 and 0.09 respectively). B. Cordero, V. Gómez, A. E. Platero-Prats, M. Revés, J. Echeverría, E. Ceemades, F. Barragán, S. Alvarez, *Dalton Trans.* **2008**, 2832–2838.  
[15] Despite extremely long acquisition times (several days), no <sup>125</sup>Te NMR signal was obtained for **1** in CD<sub>2</sub>Cl<sub>2</sub>. It is known that the <sup>125</sup>Te NMR chemical shift of **1** dramatically depends on the WCA (see ref. [3] for details). The donor acceptor complexes **3** and **4** resonate at 1749.7 and 1446.9 ppm, respectively, in [D<sub>8</sub>]THF, whereas the isomeric mixture of **5a** and **5b** shows two signals at 94.0 and 58.4 ppm in [D<sub>8</sub>]THF.  
[16] D. Hnyk, J. Holub, A. Růžicka, Z. Padělková, M. Bühl, *Struct. Chem.* **2013**, *24*, 927–932.  
[17] G. Binsch, J. B. Lambert, B. W. Roberts, J. D. Roberts, *J. Am. Chem. Soc.* **1964**, *86*, 5564–5570.  
[18] The pK<sub>a</sub> values were calculated as described previously: a) R. Kather, E. Rychagova, P. Sanz Camacho, S. E. Ashbrook, J. D. Woollins, L. Robben, E. Lork, S. Ketkov, J. Beckmann *Chem. Commun.* **2016**, *52*, 10992–10995; b) M. Olaru, M. F. Hesse, E. Rychagova, S. Ketkov, S. Mebs, J. Beckmann, *Angew. Chem. Int. Ed.* **2017**, *56*, 16490–16494; *Angew. Chem.* **2017**, *129*, 16713–16717. See Supporting Information for further details.  
[19] R. W. F. Bader, *Atoms in Molecules: A Quantum Theory*, Cambridge University Press Oxford, **1991**.  
[20] E. R. Johnson, S. Keinan, P. Mori-Sanchez, J. Contreras-García, A. J. Cohen, W. Yang, *J. Am. Chem. Soc.* **2010**, *132*, 6498–6506.  
[21] a) M. Kohout, *Int. J. Quantum Chem.* **2004**, *97*, 651–658; b) M. Kohout, F. R. Wagner, Y. Grin, *Theor. Chem. Acc.* **2008**, *119*, 413–420.  
[22] J. Kaleta, A. Akdag, R. Crespo, M.-C. Piqueras, J. Michl, *ChemPlusChem* **2013**, *78*, 1174–1183.

Manuscript received: September 2, 2021

Accepted manuscript online: September 8, 2021

Version of record online: September 28, 2021