

References

- Akopian, A.N., V. Souslova, S. England, K. Okuse, N. Ogata, J. Ure, A. Smith, B.J. Kerr, S.B. McMahon, S. Boyce, R. Hill, L.C. Stanfa, A.H. Dickenson, and J.N. Wood. 1999. The tetrodotoxin-resistant sodium channel SNS has a specialized function in pain pathways. *Nat Neurosci.* 2:541-8.
- Babes, A., D. Zorzon, and G. Reid. 2006. A novel type of cold-sensitive neuron in rat dorsal root ganglia with rapid adaptation to cooling stimuli. *Eur J Neurosci.* 24:691-8.
- Bardin, L., J. Lavarenne, and A. Eschalier. 2000. Serotonin receptor subtypes involved in the spinal antinociceptive effect of 5-HT in rats. *Pain.* 86:11-8.
- Baumann, T.K., D.A. Simone, C.N. Shain, and R.H. LaMotte. 1991. Neurogenic hyperalgesia: the search for the primary cutaneous afferent fibers that contribute to capsaicin-induced pain and hyperalgesia. *J Neurophysiol.* 66:212-27.
- Bautista, D.M., S.E. Jordt, T. Nikai, P.R. Tsuruda, A.J. Read, J. Poblete, E.N. Yamoah, A.I. Basbaum, and D. Julius. 2006. TRPA1 mediates the inflammatory actions of environmental irritants and proalgesic agents. *Cell.* 124:1269-82.
- Benham, C.D., M.J. Gunthorpe, and J.B. Davis. 2003. TRPV channels as temperature sensors. *Cell Calcium.* 33:479-87.
- Benn, S.C., M. Costigan, S. Tate, M. Fitzgerald, and C.J. Woolf. 2001. Developmental expression of the TTX-resistant voltage-gated sodium channels Nav1.8 (SNS) and Nav1.9 (SNS2) in primary sensory neurons. *J Neurosci.* 21:6077-85.
- Bennett, D.L., S. Averill, D.O. Clary, J.V. Priestley, and S.B. McMahon. 1996. Postnatal changes in the expression of the trkB high-affinity NGF receptor in primary sensory neurons. *Eur J Neurosci.* 8:2204-8.
- Bennett, D.L., M. Koltzenburg, J.V. Priestley, D.L. Shelton, and S.B. McMahon. 1998. Endogenous nerve growth factor regulates the sensitivity of nociceptors in the adult rat. *Eur J Neurosci.* 10:1282-91.
- Berman, D.M., T.M. Wilkie, and A.G. Gilman. 1996. GAIP and RGS4 are GTPase-activating proteins for the Gi subfamily of G protein alpha subunits. *Cell.* 86:445-52.
- Besmer, P. 1991. The kit ligand encoded at the murine Steel locus: a pleiotropic growth and differentiation factor. *Curr Opin Cell Biol.* 3:939-46.
- Bessou, P., and E.R. Perl. 1969. Response of cutaneous sensory units with unmyelinated fibers to noxious stimuli. *J Neurophysiol.* 32:1025-43.
- Birder, L.A., Y. Nakamura, S. Kiss, M.L. Nealen, S. Barrick, A.J. Kanai, E. Wang, G. Ruiz, W.C. De Groat, G. Apodaca, S. Watkins, and M.J. Caterina. 2002. Altered

urinary bladder function in mice lacking the vanilloid receptor TRPV1. *Nat Neurosci.* 5:856-60.

Bonnington, J.K., and P.A. McNaughton. 2003. Signalling pathways involved in the sensitisation of mouse nociceptive neurones by nerve growth factor. *J Physiol.* 551:433-46.

Burgess, P.R., and E.R. Perl. 1967. Myelinated afferent fibres responding specifically to noxious stimulation of the skin. *J Physiol.* 190:541-62.

Burgess, P.R., D. Petit, and R.M. Warren. 1968. Receptor types in cat hairy skin supplied by myelinated fibers. *J Neurophysiol.* 31:833-48.

Cain, D.M., S.G. Khasabov, and D.A. Simone. 2001. Response properties of mechanoreceptors and nociceptors in mouse glabrous skin: an in vivo study. *J Neurophysiol.* 85:1561-74.

Campbell, J.N., R.A. Meyer, and R.H. LaMotte. 1979. Sensitization of myelinated nociceptive afferents that innervate monkey hand. *J Neurophysiol.* 42:1669-79.

Caterina, M.J., and D. Julius. 1999. Sense and specificity: a molecular identity for nociceptors. *Curr Opin Neurobiol.* 9:525-30.

Caterina, M.J., A. Leffler, A.B. Malmberg, W.J. Martin, J. Trafton, K.R. Petersen-Zeitz, M. Koltzenburg, A.I. Basbaum, and D. Julius. 2000. Impaired nociception and pain sensation in mice lacking the capsaicin receptor. *Science.* 288:306-13.

Caterina, M.J., T.A. Rosen, M. Tominaga, A.J. Brake, and D. Julius. 1999. A capsaicin-receptor homologue with a high threshold for noxious heat. *Nature.* 398:436-41.

Caterina, M.J., M.A. Schumacher, M. Tominaga, T.A. Rosen, J.D. Levine, and D. Julius. 1997. The capsaicin receptor: a heat-activated ion channel in the pain pathway. *Nature.* 389:816-24.

Cesare, P., and P. McNaughton. 1996. A novel heat-activated current in nociceptive neurons and its sensitization by bradykinin. *Proc Natl Acad Sci U S A.* 93:15435-9.

Chabot, B., D.A. Stephenson, V.M. Chapman, P. Besmer, and A. Bernstein. 1988. The proto-oncogene c-kit encoding a transmembrane tyrosine kinase receptor maps to the mouse W locus. *Nature.* 335:88-9.

Chen, C.L., D.C. Broom, Y. Liu, J.C. de Nooij, Z. Li, C. Cen, O.A. Samad, T.M. Jessell, C.J. Woolf, and Q. Ma. 2006. Runx1 determines nociceptive sensory neuron phenotype and is required for thermal and neuropathic pain. *Neuron.* 49:365-77.

Cheong, K.H., D. Zacchetti, E.E. Schneeberger, and K. Simons. 1999. VIP17/MAL, a lipid raft-associated protein, is involved in apical transport in MDCK cells. *Proc Natl Acad Sci U S A.* 96:6241-8.

Chung, M.K., H. Lee, A. Mizuno, M. Suzuki, and M.J. Caterina. 2004. TRPV3 and TRPV4 mediate warmth-evoked currents in primary mouse keratinocytes. *J Biol Chem.* 279:21569-75.

- Costigan, M., T.A. Samad, A. Allchorne, C. Lanoue, S. Tate, and C.J. Woolf. 2003. High basal expression and injury-induced down regulation of two regulator of G-protein signaling transcripts, RGS3 and RGS4 in primary sensory neurons. *Mol Cell Neurosci.* 24:106-16.
- Crowley, C., S.D. Spencer, M.C. Nishimura, K.S. Chen, S. Pitts-Meek, M.P. Armanini, L.H. Ling, S.B. McMahon, D.L. Shelton, A.D. Levinson, and et al. 1994. Mice lacking nerve growth factor display perinatal loss of sensory and sympathetic neurons yet develop basal forebrain cholinergic neurons. *Cell.* 76:1001-11.
- Dalwadi, H., B. Wei, M. Schrage, K. Spicher, T.T. Su, L. Birnbaumer, D.J. Rawlings, and J. Braun. 2003. B cell developmental requirement for the G alpha i2 gene. *J Immunol.* 170:1707-15.
- Davies, A.M. 1994. The role of neurotrophins in the developing nervous system. *J Neurobiol.* 25:1334-48.
- Davis, B.M., G.R. Lewin, L.M. Mendell, M.E. Jones, and K.M. Albers. 1993. Altered expression of nerve growth factor in the skin of transgenic mice leads to changes in response to mechanical stimuli. *Neuroscience.* 56:789-92.
- Davis, J.B., J. Gray, M.J. Gunthorpe, J.P. Hatcher, P.T. Davey, P. Overend, M.H. Harries, J. Latcham, C. Clapham, K. Atkinson, S.A. Hughes, K. Rance, E. Grau, A.J. Harper, P.L. Pugh, D.C. Rogers, S. Bingham, A. Randall, and S.A. Sheardown. 2000. Vanilloid receptor-1 is essential for inflammatory thermal hyperalgesia. *Nature.* 405:183-7.
- Dechant, G. 2001. Molecular interactions between neurotrophin receptors. *Cell Tissue Res.* 305:229-38.
- Deckwerth, T.L., J.L. Elliott, C.M. Knudson, E.M. Johnson, Jr., W.D. Snider, and S.J. Korsmeyer. 1996. BAX is required for neuronal death after trophic factor deprivation and during development. *Neuron.* 17:401-11.
- Dhaka, A., A.N. Murray, J. Mathur, T.J. Earley, M.J. Petrus, and A. Patapoutian. 2007. TRPM8 is required for cold sensation in mice. *Neuron.* 54:371-8.
- Dittert, I., J. Benedikt, L. Vyklicky, K. Zimmermann, P.W. Reeh, and V. Vlachova. 2006. Improved superfusion technique for rapid cooling or heating of cultured cells under patch-clamp conditions. *J Neurosci Methods.* 151:178-85.
- Djouhri, L., and S.N. Lawson. 2004. Abeta-fiber nociceptive primary afferent neurons: a review of incidence and properties in relation to other afferent A-fiber neurons in mammals. *Brain Res Brain Res Rev.* 46:131-45.
- Dray, A. 1995. Inflammatory mediators of pain. *Br J Anaesth.* 75:125-31.
- Drew, L.J., D.K. Rohrer, M.P. Price, K.E. Blaver, D.A. Cockayne, P. Cesare, and J.N. Wood. 2004. Acid-sensing ion channels ASIC2 and ASIC3 do not contribute to mechanically activated currents in mammalian sensory neurones. *J Physiol.* 556:691-710.
- Drusco, A., N. Zanesi, C. Roldo, F. Trapasso, J.L. Farber, L.Y. Fong, and C.M. Croce. 2005. Knockout mice reveal a tumor suppressor function for Testin. *Proc Natl Acad Sci U S A.* 102:10947-51.

- Dubreuil, A.S., H. Boukhaddaoui, G. Desmadryl, C. Martinez-Salgado, R. Moshourab, G.R. Lewin, P. Carroll, J. Valmier, and F. Scamps. 2004. Role of T-type calcium current in identified d-hair mechanoreceptor neurons studied in vitro. *J Neurosci.* 24:8480-4.
- Dubreuil, P., R. Rottapel, A.D. Reith, L. Forrester, and A. Bernstein. 1990. The mouse W/c-kit locus. A mammalian gene that controls the development of three distinct cell lineages. *Ann N Y Acad Sci.* 599:58-65.
- Elgström, K., M. Bjursten, R. Willen, P.W. Bland, and E.H. Hornquist. 2007. Aberrant T-cell ontogeny and defective thymocyte and colonic T-cell chemotactic migration in colitis-prone Galphai2-deficient mice. *Immunology.* 122:199-209.
- Elitt, C.M., S.L. McIlwrath, J.J. Lawson, S.A. Malin, D.C. Molliver, P.K. Cornuet, H.R. Koerber, B.M. Davis, and K.M. Albers. 2006. Artemin overexpression in skin enhances expression of TRPV1 and TRPA1 in cutaneous sensory neurons and leads to behavioral sensitivity to heat and cold. *J Neurosci.* 26:8578-87.
- Freeman, W., and A.J. Morton. 2004. Regional and progressive changes in brain expression of complexin II in a mouse transgenic for the Huntington's disease mutation. *Brain Res Bull.* 63:45-55.
- Galoyan, S.M., J.C. Petruska, and L.M. Mendell. 2003. Mechanisms of sensitization of the response of single dorsal root ganglion cells from adult rat to noxious heat. *Eur J Neurosci.* 18:535-41.
- Gong, Z., W. Son, Y.D. Chung, J. Kim, D.W. Shin, C.A. McClung, Y. Lee, H.W. Lee, D.J. Chang, B.K. Kaang, H. Cho, U. Oh, J. Hirsh, M.J. Kernan, and C. Kim. 2004. Two interdependent TRPV channel subunits, inactive and Nanchung, mediate hearing in *Drosophila*. *J Neurosci.* 24:9059-66.
- Greene, L.A., and E.M. Shooter. 1980. The nerve growth factor: biochemistry, synthesis, and mechanism of action. *Annu Rev Neurosci.* 3:353-402.
- Grillet, N., A. Pattyn, C. Contet, B.L. Kieffer, C. Goridis, and J.F. Brunet. 2005. Generation and characterization of Rgs4 mutant mice. *Mol Cell Biol.* 25:4221-8.
- Handwerker, H.O., S. Kilo, and P.W. Reeh. 1991. Unresponsive afferent nerve fibres in the sural nerve of the rat. *J Physiol.* 435:229-42.
- Hardy, J.D., H.G. Wolff, and H. Goodell. 1952. Pricking pain threshold in different body areas. *Proc Soc Exp Biol Med.* 80:425-7.
- Hayashi, S., T. Kunisada, M. Ogawa, K. Yamaguchi, and S. Nishikawa. 1991. Exon skipping by mutation of an authentic splice site of c-kit gene in W/W mouse. *Nucleic Acids Res.* 19:1267-71.
- Hepler, J.R., D.M. Berman, A.G. Gilman, and T. Kozasa. 1997. RGS4 and GAIP are GTPase-activating proteins for Gq alpha and block activation of phospholipase C beta by gamma-thio-GTP-Gq alpha. *Proc Natl Acad Sci U S A.* 94:428-32.
- Hirata, T., T. Kasugai, E. Morii, S. Hirota, S. Nomura, H. Fujisawa, and Y. Kitamura. 1995. Characterization of c-kit-positive neurons in the dorsal root ganglion of mouse. *Brain Res Dev Brain Res.* 85:201-11.

- Hirata, T., E. Morii, M. Morimoto, T. Kasugai, T. Tsujimura, S. Hirota, Y. Kanakura, S. Nomura, and Y. Kitamura. 1993. Stem cell factor induces outgrowth of c-kit-positive neurites and supports the survival of c-kit-positive neurons in dorsal root ganglia of mouse embryos. *Development*. 119:49-56.
- Hoger, U., and A.S. French. 1999. Temperature sensitivity of transduction and action potential conduction in a spider mechanoreceptor. *Pflugers Arch.* 438:837-42.
- Horch, K.W., D. Whitehorn, and P.R. Burgess. 1974. Impulse generation in type I cutaneous mechanoreceptors. *J Neurophysiol.* 37:267-81.
- Hornquist, C.E., X. Lu, P.M. Rogers-Fani, U. Rudolph, S. Shappell, L. Birnbaumer, and G.R. Harriman. 1997. G(alpha)i2-deficient mice with colitis exhibit a local increase in memory CD4+ T cells and proinflammatory Th1-type cytokines. *J Immunol.* 158:1068-77.
- Hu, J., and G.R. Lewin. 2006. Mechanosensitive currents in the neurites of cultured mouse sensory neurones. *J Physiol.* 577:815-28.
- Hu, J., N. Milenkovic, and G.R. Lewin. 2006. The high threshold mechanotransducer: a status report. *Pain*. 120:3-7.
- Huang, E.J., and L.F. Reichardt. 2003. Trk receptors: roles in neuronal signal transduction. *Annu Rev Biochem.* 72:609-42.
- Hulsebosch, C.E., R.E. Coggeshall, and J.R. Perez-Polo. 1987. Persistence of anti-NGF induced dorsal root axons: possible penetration into the mammalian spinal cord. *Brain Res.* 411:267-74.
- Ikoma, A., M. Steinhoff, S. Stander, G. Yosipovitch, and M. Schmelz. 2006. The neurobiology of itch. *Nat Rev Neurosci.* 7:535-47.
- Jameson, H.L., and K.A. Lillycrop. 2001. Nerve growth factor induces the expression of the LIM homeodomain transcription factor Isl-1 with the kinetics of an immediate early gene in adult rat dorsal root ganglion. *Neurosci Lett.* 309:130-4.
- Kahn-Kirby, A.H., and C.I. Bargmann. 2005. TRP Channels in *C. elegans*. *Annu Rev Physiol.*
- Kahn-Kirby, A.H., J.L. Dantzker, A.J. Apicella, W.R. Schafer, J. Browse, C.I. Bargmann, and J.L. Watts. 2004. Specific Polyunsaturated Fatty Acids Drive TRPV-Dependent Sensory Signaling In Vivo. *Cell.* 119:889-900.
- Kashiba, H., Y. Uchida, and E. Senba. 2001. Difference in binding by isolectin B4 to trkA and c-ret mRNA-expressing neurons in rat sensory ganglia. *Brain Res Mol Brain Res.* 95:18-26.
- Katoh, Y., and M. Katoh. 2004. Identification and characterization of CDC50A, CDC50B and CDC50C genes in silico. *Oncol Rep.* 12:939-43.
- Kendall, G., H. Crankson, E. Ensor, D.M. Lublin, and D.S. Latchman. 1996. Activation of the gene encoding decay accelerating factor following nerve growth factor treatment of sensory neurons is mediated by promoter sequences within 206 bases of the transcriptional start site. *J Neurosci Res.* 45:96-103.

- Keshet, E., S.D. Lyman, D.E. Williams, D.M. Anderson, N.A. Jenkins, N.G. Copeland, and L.F. Parada. 1991. Embryonic RNA expression patterns of the c-kit receptor and its cognate ligand suggest multiple functional roles in mouse development. *Embo J.* 10:2425-35.
- Kessler, F., C. Habelt, B. Averbeck, P.W. Reeh, and M. Kress. 1999. Heat-induced release of CGRP from isolated rat skin and effects of bradykinin and the protein kinase C activator PMA. *Pain*. 83:289-95.
- Koerber, H.R., Mendell, L. M. 1992. Functional heterogeneity of dorsal root ganglion cells. In *Sensory Neurons: Diversity, Development and Plasticity*. New York: Oxford Univ. Press. p. 77-96.
- Koltzenburg, M., C.L. Stucky, and G.R. Lewin. 1997. Receptive properties of mouse sensory neurons innervating hairy skin. *J Neurophysiol.* 78:1841-50.
- Kothapalli, R., S.J. Yoder, S. Mane, and T.P. Loughran, Jr. 2002. Microarray results: how accurate are they? *BMC Bioinformatics*. 3:22.
- Kwan, K.Y., A.J. Allchorne, M.A. Vollrath, A.P. Christensen, D.S. Zhang, C.J. Woolf, and D.P. Corey. 2006. TRPA1 contributes to cold, mechanical, and chemical nociception but is not essential for hair-cell transduction. *Neuron*. 50:277-89.
- Lawson, S.N. 2002. Phenotype and function of somatic primary afferent nociceptive neurones with C-, Adelta- or Aalpha/beta-fibres. *Exp Physiol.* 87:239-44.
- Levi-Montalcini, R., and P.U. Angeletti. 1968. Nerve growth factor. *Physiol Rev.* 48:534-69.
- Lewin, G.R., and L.M. Mendell. 1994. Regulation of cutaneous C-fiber heat nociceptors by nerve growth factor in the developing rat. *J Neurophysiol.* 71:941-9.
- Lewin, G.R., and R. Moshourab. 2004. Mechanosensation and pain. *J Neurobiol.* 61:30-44.
- Lewin, G.R., A.M. Ritter, and L.M. Mendell. 1992. On the role of nerve growth factor in the development of myelinated nociceptors. *J Neurosci.* 12:1896-905.
- Lewin, G.R., A. Rueff, and L.M. Mendell. 1994. Peripheral and central mechanisms of NGF-induced hyperalgesia. *Eur J Neurosci.* 6:1903-12.
- Li, W., Z. Feng, P.W. Sternberg, and X.Z. Xu. 2006. A *C. elegans* stretch receptor neuron revealed by a mechanosensitive TRP channel homologue. *Nature*. 440:684-7.
- Little, C.C. 1915. The Inheritance of Black-Eyed White Spotting in Mice. *The American Naturalist*. 49:727-740.
- Lublin, D.M., and J.P. Atkinson. 1989. Decay-accelerating factor: biochemistry, molecular biology, and function. *Annu Rev Immunol.* 7:35-58.
- Lumpkin, E.A., and D.M. Bautista. 2005. Feeling the pressure in mammalian somatosensation. *Curr Opin Neurobiol.* 15:382-8.

- Lumpkin, E.A., and M.J. Caterina. 2007. Mechanisms of sensory transduction in the skin. *Nature*. 445:858-65.
- Luo, W., S.R. Wickramasinghe, J.M. Savitt, J.W. Griffin, T.M. Dawson, and D.D. Ginty. 2007. A hierarchical NGF signaling cascade controls Ret-dependent and Ret-independent events during development of nonpeptidergic DRG neurons. *Neuron*. 54:739-54.
- Ma, Q.P. 2001. Vanilloid receptor homologue, VRL1, is expressed by both A- and C-fiber sensory neurons. *Neuroreport*. 12:3693-5.
- Maingret, F., I. Lauritzen, A.J. Patel, C. Heurteaux, R. Reyes, F. Lesage, M. Lazdunski, and E. Honore. 2000. TREK-1 is a heat-activated background K(+) channel. *Embo J*. 19:2483-91.
- Marchand, F., M. Perretti, and S.B. McMahon. 2005. Role of the immune system in chronic pain. *Nat Rev Neurosci*. 6:521-32.
- Maricq, A.V., A.S. Peterson, A.J. Brake, R.M. Myers, and D. Julius. 1991. Primary structure and functional expression of the 5HT3 receptor, a serotonin-gated ion channel. *Science*. 254:432-7.
- Matsui, Y., K.M. Zsebo, and B.L. Hogan. 1990. Embryonic expression of a haematopoietic growth factor encoded by the SI locus and the ligand for c-kit. *Nature*. 347:667-9.
- McCleskey, E.W., and M.S. Gold. 1999. Ion channels of nociception. *Annu Rev Physiol*. 61:835-56.
- Melzack, R., and P.D. Wall. 1965. Pain mechanisms: a new theory. *Science*. 150:971-9.
- Milenkovic, N., C. Frahm, M. Gassmann, C. Griffel, B. Erdmann, C. Birchmeier, G.R. Lewin, and A.N. Garratt. 2007. Nociceptive tuning by Stem Cell Factor/c-Kit signaling *Neuron*.
- Millan, J., and M.A. Alonso. 1998. MAL, a novel integral membrane protein of human T lymphocytes, associates with glycosylphosphatidylinositol-anchored proteins and Src-like tyrosine kinases. *Eur J Immunol*. 28:3675-84.
- Miwa, T., N. Okada, and H. Okada. 2000. Alternative exon usage in the 3' region of a single gene generates glycosylphosphatidylinositol-anchored and transmembrane forms of rat decay-accelerating factor. *Immunogenetics*. 51:129-37.
- Modregger, J., B. Ritter, B. Witter, M. Paulsson, and M. Plomann. 2000. All three PACSIN isoforms bind to endocytic proteins and inhibit endocytosis. *J Cell Sci*. 113 Pt 24:4511-21.
- Mogil, J.S., F. Miermeister, F. Seifert, K. Strasburg, K. Zimmermann, H. Reinold, J.S. Austin, N. Bernardini, E.J. Chesler, H.A. Hofmann, C. Hordo, K. Messlinger, K.V. Nemmani, A.L. Rankin, J. Ritchie, A. Siegling, S.B. Smith, S. Sotocinal, A. Vater, S.G. Lehto, S. Klussmann, R. Quirion, M. Michaelis, M. Devor, and P.W. Reeh. 2005. Variable sensitivity to noxious heat is mediated by differential expression of the CGRP gene. *Proc Natl Acad Sci U S A*. 102:12938-43.

- Molliver, D.C., M.J. Radeke, S.C. Feinstein, and W.D. Snider. 1995. Presence or absence of TrkA protein distinguishes subsets of small sensory neurons with unique cytochemical characteristics and dorsal horn projections. *J Comp Neurol.* 361:404-16.
- Molliver, D.C., and W.D. Snider. 1997. Nerve growth factor receptor TrkA is down-regulated during postnatal development by a subset of dorsal root ganglion neurons. *J Comp Neurol.* 381:428-38.
- Molliver, D.C., D.E. Wright, M.L. Leitner, A.S. Parsadanian, K. Doster, D. Wen, Q. Yan, and W.D. Snider. 1997. IB4-binding DRG neurons switch from NGF to GDNF dependence in early postnatal life. *Neuron.* 19:849-61.
- Moqrich, A., S.W. Hwang, T.J. Earley, M.J. Petrus, A.N. Murray, K.S. Spencer, M. Andahazy, G.M. Story, and A. Patapoutian. 2005. Impaired thermosensation in mice lacking TRPV3, a heat and camphor sensor in the skin. *Science.* 307:1468-72.
- Motro, B., D. van der Kooy, J. Rossant, A. Reith, and A. Bernstein. 1991. Contiguous patterns of c-kit and steel expression: analysis of mutations at the W and SI loci. *Development.* 113:1207-21.
- Mu, X., I. Silos-Santiago, S.L. Carroll, and W.D. Snider. 1993. Neurotrophin receptor genes are expressed in distinct patterns in developing dorsal root ganglia. *J Neurosci.* 13:4029-41.
- Negri, L., R. Lattanzi, E. Giannini, M. Colucci, F. Margheriti, P. Melchiorri, V. Vellani, H. Tian, M. De Felice, and F. Porreca. 2006. Impaired nociception and inflammatory pain sensation in mice lacking the prokineticin receptor PKR1: focus on interaction between PKR1 and the capsaicin receptor TRPV1 in pain behavior. *J Neurosci.* 26:6716-27.
- Nocka, K., J. Buck, E. Levi, and P. Besmer. 1990. Candidate ligand for the c-kit transmembrane kinase receptor: KL, a fibroblast derived growth factor stimulates mast cells and erythroid progenitors. *Embo J.* 9:3287-94.
- Nykjaer, A., R. Lee, K.K. Teng, P. Jansen, P. Madsen, M.S. Nielsen, C. Jacobsen, M. Kliemann, E. Schwarz, T.E. Willnow, B.L. Hempstead, and C.M. Petersen. 2004. Sortilin is essential for proNGF-induced neuronal cell death. *Nature.* 427:843-8.
- O'Hagan, R., M. Chalfie, and M.B. Goodman. 2005. The MEC-4 DEG/ENaC channel of *Caenorhabditis elegans* touch receptor neurons transduces mechanical signals. *Nat Neurosci.* 8:43-50.
- Okajima, T., S. Fukumoto, H. Ito, M. Kiso, Y. Hirabayashi, T. Urano, and K. Furukawa. 1999. Molecular cloning of brain-specific GD1alpha synthase (ST6GalNAc V) containing CAG/Glutamine repeats. *J Biol Chem.* 274:30557-62.
- Olausson, H., Y. Lamarre, H. Backlund, C. Morin, B.G. Wallin, G. Starck, S. Ekholm, I. Strigo, K. Worsley, A.B. Vallbo, and M.C. Bushnell. 2002. Unmyelinated tactile afferents signal touch and project to insular cortex. *Nat Neurosci.* 5:900-4.

- Orozco, O.E., L. Walus, D.W. Sah, R.B. Pepinsky, and M. Sanicola. 2001. GFRalpha3 is expressed predominantly in nociceptive sensory neurons. *Eur J Neurosci.* 13:2177-82.
- Orr-Utreger, A., A. Avivi, Y. Zimmer, D. Givol, Y. Yarden, and P. Lonai. 1990. Developmental expression of c-kit, a proto-oncogene encoded by the W locus. *Development.* 109:911-23.
- Patel, T.D., A. Jackman, F.L. Rice, J. Kucera, and W.D. Snider. 2000. Development of sensory neurons in the absence of NGF/TrkA signaling in vivo. *Neuron.* 25:345-57.
- Peier, A.M., A.J. Reeve, D.A. Andersson, A. Moqrich, T.J. Earley, A.C. Hergarden, G.M. Story, S. Colley, J.B. Hogenesch, P. McIntyre, S. Bevan, and A. Patapoutian. 2002. A heat-sensitive TRP channel expressed in keratinocytes. *Science.* 296:2046-9.
- Perez-Otano, I., R. Lujan, S.J. Tavalin, M. Plomann, J. Modregger, X.B. Liu, E.G. Jones, S.F. Heinemann, D.C. Lo, and M.D. Ehlers. 2006. Endocytosis and synaptic removal of NR3A-containing NMDA receptors by PACSIN1/syndapin1. *Nat Neurosci.* 9:611-21.
- Pezet, S., and S.B. McMahon. 2006. Neurotrophins: mediators and modulators of pain. *Annu Rev Neurosci.* 29:507-38.
- Plomann, M., R. Lange, G. Vopper, H. Cremer, U.A. Heinlein, S. Scheff, S.A. Baldwin, M. Leitges, M. Cramer, M. Paulsson, and D. Barthels. 1998. PACSIN, a brain protein that is upregulated upon differentiation into neuronal cells. *Eur J Biochem.* 256:201-11.
- Price, M.P., G.R. Lewin, S.L. McIlwrath, C. Cheng, J. Xie, P.A. Heppenstall, C.L. Stucky, A.G. Mannsfeldt, T.J. Brennan, H.A. Drummond, J. Qiao, C.J. Benson, D.E. Tarr, R.F. Hrstka, B. Yang, R.A. Williamson, and M.J. Welsh. 2000. The mammalian sodium channel BNC1 is required for normal touch sensation. *Nature.* 407:1007-11.
- Rasband, M.N., E.W. Park, T.W. Vanderah, J. Lai, F. Porreca, and J.S. Trimmer. 2001. Distinct potassium channels on pain-sensing neurons. *Proc Natl Acad Sci U S A.* 98:13373-8.
- Reeh, P.W. 1986. Sensory receptors in mammalian skin in an in vitro preparation. *Neurosci Lett.* 66:141-6.
- Reim, K., M. Mansour, F. Varoqueaux, H.T. McMahon, T.C. Sudhof, N. Brose, and C. Rosenmund. 2001. Complexins regulate a late step in Ca²⁺-dependent neurotransmitter release. *Cell.* 104:71-81.
- Reith, A.D., R. Rottapel, E. Giddens, C. Brady, L. Forrester, and A. Bernstein. 1990. W mutant mice with mild or severe developmental defects contain distinct point mutations in the kinase domain of the c-kit receptor. *Genes Dev.* 4:390-400.
- Richardson, B.P. 1990. Serotonin and nociception. *Ann N Y Acad Sci.* 600:511-9; discussion 519-20.

- Ringkamp, M., Y.B. Peng, G. Wu, T.V. Hartke, J.N. Campbell, and R.A. Meyer. 2001. Capsaicin responses in heat-sensitive and heat-insensitive A-fiber nociceptors. *J Neurosci.* 21:4460-8.
- Ritter, A.M., G.R. Lewin, N.E. Kremer, and L.M. Mendell. 1991. Requirement for nerve growth factor in the development of myelinated nociceptors *in vivo*. *Nature*. 350:500-2.
- Ritter, A.M., G.R. Lewin, and L.M. Mendell. 1993. Regulation of myelinated nociceptor function by nerve growth factor in neonatal and adult rats. *Brain Res Bull*. 30:245-9.
- Rogaeva, E., Y. Meng, J.H. Lee, Y. Gu, T. Kawarai, F. Zou, T. Katayama, C.T. Baldwin, R. Cheng, H. Hasegawa, F. Chen, N. Shibata, K.L. Lunetta, R. Pardossi-Piquard, C. Bohm, Y. Wakutani, L.A. Cupples, K.T. Cuenco, R.C. Green, L. Pinessi, I. Rainero, S. Sorbi, A. Bruni, R. Duara, R.P. Friedland, R. Inzelberg, W. Hampe, H. Bujo, Y.Q. Song, O.M. Andersen, T.E. Willnow, N. Graff-Radford, R.C. Petersen, D. Dickson, S.D. Der, P.E. Fraser, G. Schmitt-Ulms, S. Younkin, R. Mayeux, L.A. Farrer, and P. St George-Hyslop. 2007. The neuronal sortilin-related receptor SORL1 is genetically associated with Alzheimer disease. *Nat Genet.* 39:168-77.
- Ruberti, F., S. Capsoni, A. Comparini, E. Di Daniel, J. Franzot, S. Gonfoni, G. Rossi, N. Berardi, and A. Cattaneo. 2000. Phenotypic knockout of nerve growth factor in adult transgenic mice reveals severe deficits in basal forebrain cholinergic neurons, cell death in the spleen, and skeletal muscle dystrophy. *J Neurosci.* 20:2589-601.
- Rudolph, U., M.J. Finegold, S.S. Rich, G.R. Harriman, Y. Srinivasan, P. Brabet, A. Bradley, and L. Birnbaumer. 1995. Gi2 alpha protein deficiency: a model of inflammatory bowel disease. *J Clin Immunol.* 15:101S-105S.
- Salmon, A.M., I. Damaj, S. Sekine, M.R. Picciotto, L. Marubio, and J.P. Changeux. 1999. Modulation of morphine analgesia in alphaCGRP mutant mice. *Neuroreport.* 10:849-54.
- Sasagawa, K., Y. Matsudo, M. Kang, L. Fujimura, Y. Iitsuka, S. Okada, T. Ochiai, T. Tokuhisa, and M. Hatano. 2002. Identification of Nd1, a novel murine kelch family protein, involved in stabilization of actin filaments. *J Biol Chem.* 277:44140-6.
- Schmelz, M., G. Osiander, J. Blunk, M. Ringkamp, P.W. Reeh, and H.O. Handwerker. 1997. Intracutaneous injections of platelets cause acute pain and protracted hyperalgesia. *Neurosci Lett.* 226:171-4.
- Schmidt, R., M. Schmelz, C. Forster, M. Ringkamp, E. Torebjork, and H. Handwerker. 1995. Novel classes of responsive and unresponsive C nociceptors in human skin. *J Neurosci.* 15:333-41.
- Sherrington, C.S. 1906. The integrative action of the nervous system. In C Scribner and Sons, New York.
- Shin, J.B., C. Martinez-Salgado, P.A. Heppenstall, and G.R. Lewin. 2003. A T-type calcium channel required for normal function of a mammalian mechanoreceptor. *Nat Neurosci.* 6:724-30.

- Shu, X., and L.M. Mendell. 2001. Acute sensitization by NGF of the response of small-diameter sensory neurons to capsaicin. *J Neurophysiol.* 86:2931-8.
- Sinclair, D.C. 1955. Cutaneous sensation and the doctrine of specific energy. *Brain.* 78:584-614.
- Smeyne, R.J., R. Klein, A. Schnapp, L.K. Long, S. Bryant, A. Lewin, S.A. Lira, and M. Barbacid. 1994. Severe sensory and sympathetic neuropathies in mice carrying a disrupted Trk/NGF receptor gene. *Nature.* 368:246-9.
- Snider, W.D. 1994. Functions of the neurotrophins during nervous system development: what the knockouts are teaching us. *Cell.* 77:627-38.
- Snider, W.D., and S.B. McMahon. 1998. Tackling pain at the source: new ideas about nociceptors. *Neuron.* 20:629-32.
- So, H.C., R.Y. Chen, E.Y. Chen, E.F. Cheung, T. Li, and P.C. Sham. 2007. An association study of RGS4 polymorphisms with clinical phenotypes of schizophrenia in a Chinese population. *Am J Med Genet B Neuropsychiatr Genet.*
- Stein, A.T., C.A. Ufret-Vincenty, L. Hua, L.F. Santana, and S.E. Gordon. 2006. Phosphoinositide 3-kinase binds to TRPV1 and mediates NGF-stimulated TRPV1 trafficking to the plasma membrane. *J Gen Physiol.* 128:509-22.
- Stucky, C.L., M. Koltzenburg, M. Schneider, M.G. Engle, K.M. Albers, and B.M. Davis. 1999. Overexpression of nerve growth factor in skin selectively affects the survival and functional properties of nociceptors. *J Neurosci.* 19:8509-16.
- Stucky, C.L., and G.R. Lewin. 1999. Isolectin B(4)-positive and -negative nociceptors are functionally distinct. *J Neurosci.* 19:6497-505.
- Stucky, C.L., J. Rossi, M.S. Airaksinen, and G.R. Lewin. 2002. GFR alpha2/neurturin signalling regulates noxious heat transduction in isolectin B4-binding mouse sensory neurons. *J Physiol.* 545:43-50.
- Suzuki, M., A. Mizuno, K. Kodaira, and M. Imai. 2003. Impaired pressure sensation in mice lacking TRPV4. *J Biol Chem.* 278:22664-8.
- Taiwo, Y.O., and J.D. Levine. 1992. Serotonin is a directly-acting hyperalgesic agent in the rat. *Neuroscience.* 48:485-90.
- Tominaga, M., M.J. Caterina, A.B. Malmberg, T.A. Rosen, H. Gilbert, K. Skinner, B.E. Raumann, A.I. Basbaum, and D. Julius. 1998. The cloned capsaicin receptor integrates multiple pain-producing stimuli. *Neuron.* 21:531-43.
- Tracey, W.D., Jr., R.I. Wilson, G. Laurent, and S. Benzer. 2003. *painless*, a Drosophila gene essential for nociception. *Cell.* 113:261-73.
- Trevisani, M., D. Smart, M.J. Gunthorpe, M. Tognetto, M. Barbieri, B. Campi, S. Amadesi, J. Gray, J.C. Jerman, S.J. Brough, D. Owen, G.D. Smith, A.D. Randall, S. Harrison, A. Bianchi, J.B. Davis, and P. Geppetti. 2002. Ethanol elicits and potentiates nociceptor responses via the vanilloid receptor-1. *Nat Neurosci.* 5:546-51.

- Voets, T., G. Droogmans, U. Wissenbach, A. Janssens, V. Flockerzi, and B. Nilius. 2004. The principle of temperature-dependent gating in cold- and heat-sensitive TRP channels. *Nature*. 430:748-54.
- Vollrath, M.A., K.Y. Kwan, and D.P. Corey. 2007. The micromachinery of mechanotransduction in hair cells. *Annu Rev Neurosci*. 30:339-65.
- Wade, R., and L. Kedes. 1989. Developmental regulation of contractile protein genes. *Annu Rev Physiol*. 51:179-88.
- Waldmann, R., G. Champigny, F. Bassilana, C. Heurteaux, and M. Lazdunski. 1997. A proton-gated cation channel involved in acid-sensing. *Nature*. 386:173-7.
- Wang, P.H., W.L. Lee, C.M. Juang, Y.H. Yang, W.H. Lo, C.R. Lai, S.L. Hsieh, and C.C. Yuan. 2005. Altered mRNA expressions of sialyltransferases in ovarian cancers. *Gynecol Oncol*. 99:631-9.
- Waskow, C., S. Paul, C. Haller, M. Gassmann, and H.R. Rodewald. 2002. Viable c-Kit(W/W) mutants reveal pivotal role for c-kit in the maintenance of lymphopoiesis. *Immunity*. 17:277-88.
- Way, J.C., and M. Chalfie. 1988. *mec-3*, a homeobox-containing gene that specifies differentiation of the touch receptor neurons in *C. elegans*. *Cell*. 54:5-16.
- Westergaard, U.B., E.S. Sorensen, G. Hermey, M.S. Nielsen, A. Nykjaer, K. Kirkegaard, C. Jacobsen, J. Gliemann, P. Madsen, and C.M. Petersen. 2004. Functional organization of the sortilin Vps10p domain. *J Biol Chem*. 279:50221-9.
- Wetzel, C., J. Hu, D. Riethmacher, A. Benckendorff, L. Harder, A. Eilers, R. Moshourab, A. Kozlenkov, D. Labuz, O. Caspani, B. Erdmann, H. Machelska, P.A. Heppenstall, and G.R. Lewin. 2007. A stomatin-domain protein essential for touch sensation in the mouse. *Nature*. 445:206-9.
- Woodbury, C.J., A.M. Ritter, and H.R. Koerber. 2001. Central anatomy of individual rapidly adapting low-threshold mechanoreceptors innervating the "hairy" skin of newborn mice: early maturation of hair follicle afferents. *J Comp Neurol*. 436:304-23.
- Woodbury, C.J., M. Zwick, S. Wang, J.J. Lawson, M.J. Caterina, M. Koltzenburg, K.M. Albers, H.R. Koerber, and B.M. Davis. 2004. Nociceptors lacking TRPV1 and TRPV2 have normal heat responses. *J Neurosci*. 24:6410-5.
- Woolf, C.J. 1996. Phenotypic modification of primary sensory neurons: the role of nerve growth factor in the production of persistent pain. *Philos Trans R Soc Lond B Biol Sci*. 351:441-8.
- Wright, D.E., and W.D. Snider. 1995. Neurotrophin receptor mRNA expression defines distinct populations of neurons in rat dorsal root ganglia. *J Comp Neurol*. 351:329-38.
- Zeitz, K.P., N. Guy, A.B. Malmberg, S. Dirajlal, W.J. Martin, L. Sun, D.W. Bonhaus, C.L. Stucky, D. Julius, and A.I. Basbaum. 2002. The 5-HT3 subtype of serotonin receptor contributes to nociceptive processing via a novel subset of myelinated and unmyelinated nociceptors. *J Neurosci*. 22:1010-9.

- Zimmermann, K., A. Leffler, A. Babes, C.M. Cendan, R.W. Carr, J. Kobayashi, C. Nau, J.N. Wood, and P.W. Reeh. 2007. Sensory neuron sodium channel Nav1.8 is essential for pain at low temperatures. *Nature*. 447:855-8.
- Zimmermann, K., A. Leffler, M.M. Fischer, K. Messlinger, C. Nau, and P.W. Reeh. 2005. The TRPV1/2/3 activator 2-aminoethoxydiphenyl borate sensitizes native nociceptive neurons to heat in wildtype but not TRPV1 deficient mice. *Neuroscience*. 135:1277-84.
- Zirlinger, M., G. Kreiman, and D.J. Anderson. 2001. Amygdala-enriched genes identified by microarray technology are restricted to specific amygdaloid subnuclei. *Proc Natl Acad Sci U S A*. 98:5270-5.

