

4. Literaturverzeichnis

Apcher GS, Maitland J, Dawson S, Sheppard P, Mayer RJ (2004) The alpha4 and alpha7 subunits and assembly of the 20S proteasome. *FEBS Lett.* 569(1-3):211-6.

Aki M, Shimbara N, Takashina M, Akiyama K, Kagawa S, Tamura T, Tanahashi N, Yoshimura T, Tanaka K, Ichihara A (1994) Interferon-gamma induces different subunit organizations and functional diversity of proteasomes. *J Biochem (Tokyo)* 115:257-269

Bienkowska JR, Hartman H, Smith TF. (2003) A search method for homologs of small proteins. Ubiquitin-like proteins in prokaryotic cells? *Protein Eng.* 16(12):897-904.

Bochtler M, Ditzel L, Groll M, Hartmann C, Huber R (1999) The proteasome. *Annu Rev Biophys Biomol Struct*, 28, 295-317.

Bochtler M, Hartmann C, Song HK, Bourenkov GP, Bartunik HD, Huber R (2000) The structures of HslU and the ATP-dependent protease HslU-HslV. *Nature* 403, 800-805.

Branninigan JA, Dodson G, Duggleby HJ, Moody PC, Smith JL, Tomchick, DR, Murzin AG (1995) A protein catalytic framework with an N-terminal nucleophile is capable of self-activation. *Nature* 378:414-419

Braun BC, Glickman M, Kraft R, Dahlmann B, Kloetzel PM, Finley D, Schmidt M. (1999) The base of the proteasome regulatory particle exhibits chaperone-like activity. *Nat Cell Biol.* 1(4):221-6.

Brotz-Oesterhelt H, Beyer D, Kroll HP, Endermann R, Ladel C, Schroeder W, Hinzen B, Raddatz S, Paulsen H, Henninger K, Bandow JE, Sahl HG, Labischinski H. (2005) Dysregulation of bacterial proteolytic machinery by a new class of antibiotics. *Nat Med.* 11(10):1082-7.

Bukau, B. (1997) The heat shock response in *Escherichia coli*. In Gething, M.J. (ed.), *Guidebook to molecular chaperones and protein-folding catalysts*. Oxford University Press, Oxford, UK, pp. 525-528.

Burri L, Höckendorff J, Boehm U, Klamp T, Dohmen JR, Lévy F (2000) Identification and characterization of a mammalian protein interacting with 20S proteasome precursors. *Proc Natl Acad Sci USA* 97:10348-10353

Bush KT, Goldberg AL, Nigam SK (1997) Proteasome inhibition leads to a heat-shock response, induction of endoplasmic reticulum chaperones, and thermotolerance. *J Biol Chem* 272:9086-92

Cagney G, Uetz P, Fields S (2001) Two-hybrid analysis of the *Saccharomyces cerevisiae* 26S proteasome. *Physiol Genomics*, 7, 27-34.

Chen P, Hochstrasser M (1996) Autocatalytic subunit processing couples active site formation in the 20S proteasome to completion of assembly. *Cell* 86:961-972

Chen W, Norbury CC, Cho Y, Yewdell JW, Bennink JR (2001) Immunoproteasomes shape immunodominance hierarchies of antiviral CD8(+) T cells at the levels of T cell repertoire and presentation of viral antigens. *J Exp Med* 193, 1319-26.

Ciechanover A (2005). Proteolysis: from the lysosome to ubiquitin and the proteasome. *Nat Rev Mol Cell Biol* 6, 79-87.

Dahlmann B, Kopp F, Kuehn L, Reinauer H, Schwenen M (1986) Studies on the multicatalytic proteinase from rat skeletal muscle. *Biomed Biochim Acta*, 45, 1493-1501.

Derré I, Rappoport G, Msadek T (1999) CtsR, a novel regulator of stress and heat shock response, controls *clp* and molecular chaperone gene expression in Gram-positive bacteria. *Mol Microbiol.* 31, 117-131.

Ehrlich R. Modulation of antigen processing and presentation by persistent virus infections and in tumors. (1997) *Hum Immunol.* 54(2):104-16.

Eleuteri AM, Kohanski RA, Cardozo C, Orlowski M (1997) Bovine spleen multicatalytic proteinase complex (proteasome). Replacement of X, Y, and Z subunits by LMP7, LMP2, and MECL1 and changes in properties and specificity. *J Biol Chem*, 272, 11824-11831.

Fleming JA, Lightcap ES, Sadis S, Thoroddsen V, Bulawa CE, Blackman RK (2002) Complementary whole-genome technologies reveal the cellular response to proteasome inhibition by PS-341. *Proc Natl Acad Sci U S A* 99:1461-1466

Frees D, Chastanet A, Qazi S, Sorensen K, Hill P, Msadek T, Ingmer H. (2004) Clp ATPases are required for stress tolerance, intracellular replication and biofilm formation in *Staphylococcus aureus*. *Mol Microbiol.* 54(5):1445-62.

Frees D, Sorensen K, Ingmer H. (2005) Global virulence regulation in *Staphylococcus aureus*: pinpointing the roles of ClpP and ClpX in the *sar/agr* regulatory network. *Infect Immun.* 73(12):8100-8.

Frentzel S, Pesold-Hurt B, Seelig A, Kloetzel PM (1994) 20 S proteasomes are assembled via distinct precursor complexes. Processing of LMP2 and LMP7 proproteins takes place in 13-16 S preproteasome complexes. *J Mol Biol* 236:975-981

Fricke B, Heink S, Kloetzel PM, Krüger E (2006) POMP facilitates 20S Proteasome Formation at the Endoplasmic Reticulum. *EMBO reports*, submitted

Fruh K, Yang Y, Arnold D, Chambers J, Wu L, Waters JB, Spies T, Peterson PA. (1992) Alternative exon usage and processing of the major histocompatibility complex-encoded proteasome subunits. *J Biol Chem* 267:22131-40.

Gallo RC (2002) Human retroviruses after 20 years: a perspective from the past and prospects for their future control. *Immunol Rev* 185, 236-65.

Gerards WL, Enzlin J, Haner M, Hendriks IL, Aebi U, Bloemendal H, Boelens W (1997) The human alpha-type proteasomal subunit HsC8 forms a double ringlike structure, but does not assemble into proteasome-like particles with the beta-type subunits HsDelta or HsBPROS26. *J Biol Chem.* 272:10080-10086

Gerards WLH, deJong WW, Bloemendal H, Boelens W (1998) The human proteasomal subunit HsC8 induces ring formation of other β -type subunits. *J Mol Biol* 275:113-121

Gerth U, Krüger E, Derré I, Msadek T, Hecker M (1998) Stress induction of the *Bacillus subtilis* clpP gene encoding the proteolytic component of the Clp protease and involvement of ClpP and ClpX in stress tolerance. *Mol. Microbiol.* 28, 787-802.

Gottesman S (1996) Proteases and their targets in *Escherichia coli*. *Annu. Rev. Genet.* 30, 465-506.

Gottesman S, Roche E, Zhou YN, Sauer RT (1998) The ClpXP and ClpAP proteases degrade proteins with carboxy-terminal peptide tails added by the SsrA tagging-system. *Genes and Dev.* 12, 1338-1347.

Gottesman S, Wickner S, Maurizi MR (1997) Protein quality control: triage by chaperones and proteases. *Genes and Dev.* 11, 815-823.

Griffin TA, Nandi D, Cruz M, Fehling HJ, Kaer LV, Monaco JJ, Colbert RA (1998) Immunoproteasome assembly: cooperative incorporation of interferon gamma (IFN- γ)-inducible subunits. *J Exp Med* 187:97-104

Griffin TA, Slack JP, McCluskey TS, Monaco JJ, Colbert RA (2000) Identification of proteasemblin, a mammalian homologue of the yeast protein, Ump1p, that is required for normal proteasome assembly *Mol Cell Biol Res Comm* 3:212-217

Groll M, Brandstetter H, Bartunik H, Bourenkow G, Huber R. (2003) Investigations on the maturation and regulation of archaeobacterial proteasomes. *J Mol Biol.* 327(1):75-83.

Groll M, Bajorek M, Kohler A, Moroder L, Rubin DM, Huber R, Glickman MH, Finley D (2000) A gated channel into the proteasome core particle. *Nat Struct Biol*, 7, 1062-1067.

Groll M, Brandstetter H, Bartunik H, Bourenkow G, Huber R (2003) Investigations on the maturation and regulation of archaeobacterial proteasomes. *J Mol Biol*, 327, 75-83.

Groll M, Ditzel L, Lowe J, Stock D, Bochtler M, Bartunik HD, Huber R (1997) Structure of 20S proteasome from yeast at 2.4 Å resolution. *Nature*, 386, 463-471.

Heinemeyer W, Fischer M, Krimmer T, Stachon U, Wolf DH (1997) The active sites of the eukaryotic 20S proteasome and their involvement in subunit precursor processing. *J Biol Chem* 272:25200-25209

Heink S, Fricke B, Ludwig D, Kloetzel PM, Krüger E (2006) Tumor Cell Lines Expressing the Proteasome Subunit Isoform LMP7E1 Exhibit Immunoproteasome Deficiency. *Cancer Res.* 66 (2):649-52.

Heink S, Ludwig D, Kloetzel PM, Krüger E (2005) Interferon- γ -induced immune-adaptation of the proteasome system is an accelerated and transient response. *Proc Natl Acad Sci U S A.* 102 (26): 9241-246.

Hirano Y, Hendil KB, Yashiroda H, Iemura S, Nagane R, Hioki Y, Natsume T, Tanaka K, Murata S. (2005) A heterodimeric complex that promotes the assembly of mammalian 20S proteasomes. *Nature*. 437(7063):1381-5.

Huang X, Seifert U, Salzmann U, Henklein P, Preissner R, Henke W, Sijts AJ, Kloetzel PM, Dubiel W (2002) The RTP site shared by the HIV-1 Tat protein and the 11S regulator subunit alpha is crucial for their effects on proteasome function including antigen processing. *J Mol. Biol.* 323(4), 771-82.

Hughes AL (1997) Evolution of the proteasome components. *Immunogenetics*, 46, 82-92.

Hwang BJ, Park WJ, Chung CH, Goldberg AL (1987). *Escherichia coli* contains a soluble ATP-dependent protease (Ti) distinct from protease La. *Proc. Nat. Acad. Sci. USA* 84, 5550-5554.

Ibrahim YM, Kerr AR, Silva NA, Mitchell TJ. (2005) Contribution of the ATP-dependent protease ClpCP to the autolysis and virulence of *Streptococcus pneumoniae*. *Infect Immun.* 73(2):730-40.

Jäger S, Groll M, Huber R, Wolf DH, Heinemeyer W (1999) Proteasome beta-type subunits: unequal roles of propeptides in core particle maturation and a hierarchy of active site function. *J Mol Biol* 291:997-1013

Jesenberger, V Jentsch S (2002) Deadly encounter: ubiquitin meets apoptosis. *Nat Rev Mol Cell Biol*, 3, 112-121.

Kang MS, Lim BK, Seong IS, Seol JH, Tanahashi N, Tanaka K, Chung CH. (2001) The ATP-dependent CodWX (HslVU) protease in *Bacillus subtilis* is an N-terminal serine protease. *EMBO J.* 20(4):734-42.

Katayama-Fujimura Y, Gottesman S, Maurizi MR (1987) A multiple-component, ATP-dependent protease from *Escherichia coli*. *J. Biol. Chem.* 262, 4447-4485.

Kirstein J, Zuhlke D, Gerth U, Turgay K, Hecker M. (2005) A tyrosine kinase and its activator control the activity of the CtsR heat shock repressor in *B. subtilis*. *EMBO J.* 24(19):3435-45.

Kloetzel PM (2001) Antigen processing by the proteasome. *Nat Rev Mol Cell Biol* 2: 179-187.

Kloetzel PM (2004) Generation of major histocompatibility complex class I antigens: functional interplay between proteasomes and TPPII. *Nat Immunol* 5: 661-669.

Kobayashi M, Yamamoto M. Molecular mechanisms activating the Nrf2-Keap1 pathway of antioxidant gene regulation. (2005) *Antioxid Redox Signal.* 7(3-4):385-94.

Kreft J, Vazquez-Boland JA. (2001) Regulation of virulence genes in *Listeria*. *Int J Med Microbiol.* 291(2):145-57

Krüger E, Kloetzel PM, Enenkel C (2001) 20S proteasome biogenesis. *Biochimie* 83:289-294

Krüger E, Hecker M (1998) The first gene of the *clpC*-operon in *Bacillus subtilis*, *ctsR*, encodes a negative regulator of its own operon and other classIII heat shock genes. *J Bacteriol.* 180: 6681-6688.

Krüger E, Kuckelkorn U, Sijts A, Kloetzel PM (2003) The components of the proteasome system and their role in MHC class I antigen processing. *Rev Phys Biochem Pharmac.* 148:81-104.

Krüger E, Msadek T, Hecker M (1996) Alternate promoters direct stress induced transcription of the *Bacillus subtilis clpC* operon. *Mol. Microbiol.* 20: 713-723.

Krüger E, Völker U, Hecker M (1994) Stress induction of *clpC* in *Bacillus subtilis* and its involvement in stress tolerance. *J Bacteriol.* 176: 3360-3367.

Krüger E, Witt E, Ohlmeier S, Hanschke R, Hecker M (2000) The Clp proteases of *Bacillus subtilis* are directly involved in degradation of misfolded proteins. *J Bacteriol.* 182, 3259-3265.

Kwon HY, Ogunniyi AD, Choi MH, Pyo SN, Rhee DK, Paton JC. (2004) The ClpP protease of *Streptococcus pneumoniae* modulates virulence gene expression and protects against fatal pneumococcal challenge. *Infect Immun.* 72(10):5646-53.

Lehmann A, Janek K, Braun B, Kloetzel PM, Enenkel C (2002) 20 S proteasomes are imported as precursor complexes into the nucleus of yeast. *J Mol Biol* 317: 401-413.

Löwe J, Stock D, Jap B, Zwickl P, Baumeister W, Huber R. (1995) Crystal structure of the 20S proteasome from the archaeon *T. acidophilum* at 3.4 Å resolution. *Science.* 268(5210):533-9.

Mannhaupt G, Schnall R, Karpov V, Vetter I, Feldmann H (1999) Rpn4p acts as a transcription factor by binding to PACE, a nonamer box found upstream of 26S proteasomal and other genes in yeast. *FEBS Lett* 450:27-34

Maupin-Furlow JA, Wilson HL, Kaczowka SJ, Ou MS. (2000) Proteasomes in the archaea: from structure to function. *Front Biosci.* 5:D837-65.

Mayr J, Seemüller E, Müller SA, Engel A, Baumeister W (1998) Late events in the assembly of 20S proteasomes. *J Struct Biol* 124:179-188

Meiners S, Heyken D, Weller A, Ludwig A, Stangl K, Kloetzel PM, Krüger E (2003) Inhibition of Proteasome Activity Induces Concerted Expression of Proteasome Genes and de novo Formation of Mammalian Proteasomes. *J Biol Chem.* 278: 21517-25.

Meusser B, Hirsch C, Jarosch E Sommer T (2005) ERAD: the long road to destruction. *Nat Cell Biol.* 7: 766-772.

Michel A, Agerer F, Hauck CR, Herrmann M, Ullrich J, Hacker J, Ohlsen K. (2006) Global regulatory impact of ClpP protease of *Staphylococcus aureus* on regulons involved in virulence, oxidative stress response, autolysis, and DNA repair. *J Bacteriol.* 188(16):5783-96.

Missiakas D, Schwager F, Betton JM, Georgopolus C, Raina S (1996) Identification and characterization of HslV HslU (ClpQ ClpY) proteins involved in overall proteolysis of misfolded proteins in Escherichia coli. *EMBO J.* 15: 6899-6909.

Morel S, Levy F, Burllet-Schiltz O, Brasseur F, Probst-Kepper M, Peitrequin AL, Monsarrat B, Van Velthoven R, Cerottini JC, Boon T, Gairin JE, and Van den Eynde BJ (2000) Processing of some antigens by the standard proteasome but not by the immunoproteasome results in poor presentation by dendritic cells. *Immunity* 12: 107-117.

Morimoto RI (1998) Regulation of the heat shock transcription response: cross talk between a family of heat shock factors, molecular chaperones, and negative regulators. *Genes Dev.* 12: 3788-3796.

Mullapudi S, Pullan L, Bishop OT, Khalil H, Stoops JK, Beckmann R, Kloetzel PM, Krüger E, Penczek PA. (2004) Rearrangement of the 16S Precursor Subunits Is Essential for the Formation of the Active 20S Proteasome. *Biophys J.* 87(6):4098-105.

Nair S, Derre I, Msadek T, Gaillot O, Berche P (2000) CtsR controls class III heat shock gene expression in the human pathogen *Listeria monocytogens*. *Mol. Microbiol.* 35: 800-811.

Nandi D, Woodward E, Ginsburg DB, Monaco JJ (1997) Intermediates in the formation of mouse 20S proteasomes: implications for the assembly of precursor subunits. *EMBO J* 16:5363-5375

Nussbaum AK, Dick TP, Keilholz W, Schirle M, Stevanovic S, Dietz K, Heinemeyer W, Groll M, Wolf DH, Huber R, Rammensee HG, Schild H (1998) Cleavage motifs of the yeast 20S proteasome beta subunits deduced from digests of enolase 1. *Proc Natl Acad Sci U S A*, 95: 12504-12509.

Ogura T, Wilkinson AJ. (2001) AAA+ superfamily ATPases: common structure--diverse function. *Genes Cells.*6(7):575-97.

Orlowski M, Cardozo C, Michaud C (1993) Evidence for the presence of five distinct proteolytic components in the pituitary multicatalytic proteinase complex. Properties of two components cleaving bonds on the carboxyl side of branched chain and small neutral amino acids. *Biochemistry*, 32, 1563-1572.

Pak M, Hoskins JR, Singh SK, Maurizi MR, Wickner S (1999) Concurrent chaperone and protease activities of ClpAP and requirement for the N-terminal ClpA ATP binding site for chaperone activity. *J. Biol. Chem.* 265, 12536-12545.

Qian SB, Princiotta MF, Bennink JR, Yewdell JW. (2006) Characterization of rapidly degraded polypeptides in mammalian cells reveals a novel layer of nascent protein quality control. *J Biol Chem.* 281(1):392-400.

Qian SB, Reits E, Neefjes J, Deslich JM, Bennink JR, Yewdell JW. (2006) Tight linkage between translation and MHC class I peptide ligand generation implies specialized antigen processing for defective ribosomal products. *J Immunol.* 177(1):227-33.

Ramos PC, Höckendorff J, Johnson ES, Varshavsky A, Dohmen RJ (1998) Ump1p is required for proper maturation of the 20S proteasome and becomes its substrate upon completion of the assembly. *Cell* 92:489-499

Reits EA, Vos JC, Gromme M, Neeffjes J. (2000) The major substrates for TAP in vivo are derived from newly synthesized proteins. *Nature*. 404(6779):774-8.

Roccaro AM, Hideshima T, Richardson PG, Russo D, Ribatti D, Vacca A, Dammacco F, Anderson KC. (2006) Bortezomib as an antitumor agent. *Curr Pharm Biotechnol*. 7(6):441-8.
Rosen R, Ron EZ. (2002) Proteome analysis in the study of the bacterial heat-shock response. *Mass Spectrom Rev*. 21(4):244-65.

Rouquette C, de Chastellier C, Nair S, Berche P (1998) The ClpC ATPase of *Listeria monocytogenes* is a general stress protein required for virulence and promoting of early bacterial escape from the phagosome of macrophages. *Mol. Microbiol*. 27: 1235-1245.

Ruepp A, Eckerskorn C, Bogyo M, Baumeister W. (1998) Proteasome function is dispensable under normal but not under heat shock conditions in *Thermoplasma acidophilum*. *FEBS Lett*. 425(1):87-90.

Schultz ES, Chapiro J, Lurquin C, Claverol S, Burlet-Schiltz O, Warnier G, Russo V, Morel S, Levy F, Boon T, Van den Eynde BJ, and van der Bruggen P. (2002) The production of a new MAGE-3 peptide presented to cytolytic T lymphocytes by HLA-B40 requires the immunoproteasome. *J Exp Med* 195: 391-399.

Schmidt M, Schmidtke G, Kloetzel PM (1997) Structure and structure formation of the 20S proteasome. *Mol Biol Rep*, 24, 103-112.

Schmidtke G, Kraft R, Kostka S, Henklein P, Frömmel C, Löwe J, Huber R, Kloetzel PM, Schmidt M (1996) Analysis of mammalian 20S proteasome biogenesis: the maturation of beta-subunits is an ordered two-step mechanism involving autocatalysis. *EMBO J* 15: 6887-6898.

Schmidtke G, Schmidt M, Kloetzel PM (1997) Maturation of mammalian 20 S proteasome: purification and characterization of 13 S and 16 S proteasome precursor complexes. *J Mol Biol*: 268, 95-106.

Schubert U, Anton LC, Gibbs J, Norbury CC, Yewdell JW, Bennink JR (2000) Rapid degradation of a large fraction of newly synthesized proteins by proteasomes. *Nature* 404: 770-774.

Seeger M, Ferrell K, Frank R, Dubiel W (1997) HIV-1 tat inhibits the 20 S proteasome and its 11 S regulator-mediated activation. *J Biol Chem* 272: 8145-8.

Strehl B, Seifert U, Krüger E, Heink S, Kuckelkorn U, Kloetzel PM (2005) Interferon- γ , the functional plasticity of the ubiquitin-proteasome system, and MHC class I antigen processing. *Immunol Rev*. 207: 19-30.

Tanaka K, Ii K, Ichihara A, Waxman L, Goldberg AL (1986) A high molecular weight protease in the cytosol of rat liver. I. Purification, enzymological properties, and tissue distribution. *J Biol Chem* 261: 15197-15203.

- Toes RE, Nussbaum AK, Degermann S, Schirle M, Emmerich NP, Kraft M, Laplace C, Zwinderman A, Dick TP, Muller J, Schonfisch B, Schmid C, Fehling HJ, Stevanovic S, Rammensee HG, Schild H (2001) Discrete cleavage motifs of constitutive and immunoproteasomes revealed by quantitative analysis of cleavage products J Exp Med 194: 1-12.
- Tomoyasu T, Gamer J, Bukau B, Kanemori M, Mori H, Rutman AJ, Oppenheim AB, Yura T, Yamanaka K, Niki H, Hiraga S, Ogura T (1995) Escherichia coli FtsH is a membrane-bound, ATP-dependent protease which degrades the heat-shock transcription factor σ^{32} . EMBO J 14: 2551-2560.
- Unno M, Mizushima T, Morimoto Y, Tomisugi Y, Tanaka K, Yasuoka N, Tsukihara T (2002) The structure of the mammalian 20S proteasome at 2.75 Å resolution. Structure (Camb), 10: 609-618.
- Wang J, Hartling JA, Flanagan JM (1997) The structure of ClpP at 2.3 Å Resolution suggests a model for ATP-dependent proteolysis. Cell 91: 447-456.
- Ward DE, Shockley KR, Chang LS, Levy RD, Michel JK, Connors SB, Kelly RM. (2002) Proteolysis in hyperthermophilic microorganisms. Archaea. 1:63-74.
- Whitby FG, Master EI, Kramer L, Knowlton JR, Yao Y, Wang CC, Hill CP (2000) Structural basis for the activation of 20S proteasomes by 11S regulators. Nature 408:115-120
- Wickner S, Maurizi MR, Gottesman S (1999) Posttranslational quality control: folding, refolding, and degrading proteins. Science. 286(5446):1888-93.
- Wilk S and Orłowski M (1983) Evidence that pituitary cation-sensitive neutral endopeptidase is a multicatalytic protease complex. J Neurochem 40:842-849
- Witt S, Kwon YD, Sharon M, Felderer K, Beuttler M, Robinson CV, Baumeister W, Jap BK. (2006) Proteasome assembly triggers a switch required for active-site maturation. Structure. 14(7):1179-88.
- Witt E, Zantopf D, Schmidt M, Kraft R, Kloetzel PM, Krüger E (2000) Characterisation of the newly identified human Ump1 homologue POMP and analysis of LMP7 ($\beta 5i$) incorporation into 20S proteasomes. J Mol. Biol 301: 1-9
- Wojcik C, DeMartino GN (2003) Intracellular localization of proteasomes. Int J Biochem Cell Biol, 35: 579-589.
- Xie Y, Varshavsky A (2001) RPN4 is a ligand, substrate, and transcriptional regulator of the 26S proteasome: a negative feedback circuit. Proc Natl Acad Sci U S A 98:3056-3061
- Yewdell JW, Reits E, Neefjes J. (2003) Making sense of mass destruction: quantitating MHC class I antigen presentation. Nat Rev Immunol. 3(12):952-61.
- Yewdell JW (2005) Immunoproteasomes: Regulating the regulator. Proc Natl Acad Sci U S A. 102 (26): 9089-90.
- Yoo SJ, Shim YK, Seong IS, Seol JH, Kang MS, Chung CH. (1997) Mutagenesis of two N-terminal Thr and five Ser residues in HslV, the proteolytic component of the ATP-dependent HslVU protease. FEBS Lett. 21:412(1):57-60.