

9. References

- Arndt-Jovin-DJ; Udvardy-A; Garner-MM; Ritter-S; Jovin-TM (1993). Z-DNA Binding and Inhibition by GTP of *Drosophila* Topoisomerase II. *Biochemistry*. 32:4862-72
- Bartel-DP; Szostak-JW (1993): Selection of new Ribozymes from a Large Pool of Random Sequences. *Science*. 261: 1411-8
- Bartel-DP; Szostak-JW (1993). Study of RNA-protein recognition by in vitro selection. (In: *The RNA World*, Gesteland-RF; Atkins-JF editors; Cold Spring Harbor Laboratory Press) 248-268
- Bechert-T; Diekmann-S; Arndt-Jovin-DJ (1994). Human 170 kDa and 180 kDa topoisomerases II bind preferentially to curved and left-handed linear DNA. *J-Biomol-Struct-Dyn*. 12(3): 605-23
- Behe-M; Felsenfeld-G (1981). Effects of methylation on a synthetic polynucleotide: The B-Z transition in poly(dG-m⁵dC)-poly(dG-m⁵dC). *Proc-Natl-Acad-Sci-U-S-A*. 78; 1619-23
- Beutel-BA; Gold-L (1992). In vitro evolution of intrinsically bent DNA. *J-Mol-Biol*. 228(3): 803-12
- Biedenkamp-H; Borgmeyer-U; Sippel-AE; Klempnauer-KH (1988). Viral *myb* oncogene encodes a sequence-specific DNA-binding activity. *Nature* 335:835-7
- Blaho-JA; Wells-RD (1987). Left-handed Z-DNA binding by the recA protein of *Escherichia coli* [published erratum appears in *J Biol Chem* 1988 Aug 5;263(22):11015]. *J-Biol-Chem*. 262(13): 6082-8
- Blackwell-KT; Weintraub-H (1990). Differences and Similarities in DNA-Binding Preferences of MyoD and E2A Protein Complexes Revealed by Binding Site Selection. *Science*. 250:1104-1110
- Bronstein-IN; Semendjajew-KA (1987). *Taschenbuch der Mathematik*. Verlag Harri Deutsch, Thun und Frankfurt/Main. 662
- Burns-CM; Chu-H; Rueter-SM; Hutchinson-LK; Canton-H; Sanders-Bush-E; Emeson-RB (1997). Regulation of serotonin 2C receptor G-protein coupling by RNA editing. *Nature*. 387: 303-8
- Casey-JL; Gerin-JL (1995). Hepatitis D Virus RNA Editing: Specific Modification of Adenosine in the Antigenomic RNA. *J-Virology*. 69(12): 7593-600
- Cattaneo-R; Billeter-MA (1992). Mutations and A/I Hypermutations in Measles Virus Persistent Infections. *Curr-Top-Microbiol-Immunol*. 176: 63-74

- Chattopadhyaya-R; Ikuta-S; Grzeskowiak-K; Dickerson-RE (1988); Structure of a T4 hairpin loop on a Z-DNA stem and comparison with A-RNA and B-DNA loops. *Nature* 334:175-9
- Chawla-AK; Tomasz-M (1988). Interaction of the antitumor antibiotic Mitomycin C with Z-DNA. *J-Biomol-Struct-Dyn.* 6(3): 459-70
- Doerfler-W (1983). DNA methylation and gene activity. *Annu-Rev-Biochem.* 52; 93-124
- Eichman-BF; Schroth-GP; Basham-BE; Shing Ho-P (1999). The intrinsic structure and stability of out-of-alternation base pairs in Z-DNA. *Nucleic-Acid-Res.* 27(2): 543-50
- Ellington-AD; Szostak-JW (1990). In vitro selection of RNA molecules that bind specific ligands. *Nature* 346: 818-22
- Ellison-MJ; Feigon-J; Kelleher-RJ 3d; Wang-AH; Habener-JF; Rich-A (1986). An assessment of the Z-DNA forming potential of alternating dA-dT stretches in supercoiled plasmids. *Biochemistry.* 25(12): 3648-55
- Ellison-MJ; Kelleher-RJ 3d; Wang-AH; Habener-JF; Rich-A (1985). Sequence-dependent energetics of the B-Z transition in supercoiled DNA containing nonalternating purine-pyrimidine sequences. *Proc-Natl-Acad-Sci-U-S-A.* 82(24): 8320-4
- Esposito-F; Sinden-RR (1987). DNA supercoiling and eukaryotic gene expression. *Oxford Surv. Euk. Genes.* 5:1-50
- Fitzwater-T; Polisky-B (1996). A Selex Primer. *Meth. Enzymol.* 267: 275-301
- Freund-AM; Bichara-M; Fuchs-RPP (1989). Z-DNA-forming sequences are spontaneous deletion hot spots. *Proc-Natl-Acad-Sci-U-S-A.* 86: 7465-9
- Freund-AM; Bichara-M; Fuchs-RPP (1990). Z-Forming DNA sequences are deletion hot-spots. *Mutation and the Environment, Part A;* 1990; 41-9
- Glikin-GC; Jovin-TM; Arndt-Jovin-DJ (1991). Interactions of Drosophila DNA topoisomerase II with left-handed Z-DNA in supercoiled minicircles. *Nucleic-Acids-Res.* 19(25): 7139-44
- Griffith-JD (1976). Visualization of prokaryotic DNA in a regularly condensed chromatin-like fiber. *Proc-Natl-Acad-Sci-U-S-A.* 73: 563-7
- Greene-A (1990). Stratagies 3: 5-6
- Gruskin-EA; Rich-A (1993). B-DNA to Z-DNA Structural Transitions in the SV40 Enhancer: Stabilization of Z-DNA in Negatively Supercoiled DNA Minicircles. *Biochemistry* 32:2167-76

- Gut-SH; Bischoff-M; Hobi-R; Kuenzle-CC (1987). Z-DNA-binding proteins from bull testis. *Nucleic-Acids-Res.* 15(23): 9691-705
- Haniford-DB; Pulleyblank-DE (1983). Facile transition of poly[d(TG)-d(CA)] into a left-handed helix in physiological conditions. *Nature.* 302: 632-4
- Hayes-TE; Dixon-JE (1985). Z-DNA in the rat somatostatin gene. *J-Biol-Chem.* 260(13): 8145-56
- Herbert-AG; Rich-A (1993). A method to identify and characterize Z-DNA binding proteins using a linear oligodeoxynucleotide. *Nucleic-Acids-Res.* 21(11): 2669-72
- Herbert-AG; Spitzner-JR; Lowenhaupt-K; Rich-A (1993). Z-DNA binding protein from chicken blood nuclei. *Proc-Natl-Acad-Sci-U-S-A.* 90(8): 3339-42
- Herbert-A; Lowenhaupt-K; Spitzner-J; Rich-A (1995). Chicken double-stranded RNA adenosine deaminase has apparent specificity for Z-DNA. *Proc-Natl-Acad-Sci-U-S-A.* 92(16): 7550-4
- Herbert-A; Rich-A (1996). The Biology of Left-handed Z-DNA. *J-Biol-Chem.* 271(29): 11595-8
- Herbert-A; Schade-M; Lowenhaupt-K; Alfken-J; Schwartz-T; Shlyakhtenko-LS; Lyubchenko-YL; Rich-A (1998). The Za domain form human ADAR1 binds to the Z-DNA conformer of many different sequences. *Nucleic-Acids-Res.* 26(15): 3486-93
- Hill-RJ; Stollar-BD (1983). Dependence of Z-DNA antibody binding to polytene chromosomes on acid fixation and DNA torsional strain. *Nature.* 305(5932): 338-40
- Ho-PS; Ellison-MJ; Quigley-GJ; Rich-A (1986). A computer aided thermodynamic approach for predicting the formation of Z-DNA in naturally occurring sequences. *EMBO-J.* 5(10): 2737-44
- IUPAC-IUB Commission on Biochemical Nomenclature (1970). Abbreviations and Symbols for Nucleic Acids, Polynucleotides and their Constituents. Recommendations 1970. *Eur-J-Biochem.* 15: 203-8
- Jaworski-A; Hsieh-WT; Blaho-JA; Larson-JE; Wells-RD (1987). Left-handed DNA *in vivo*. *Science.* 238: 773-7
- Kim-J-M; DasSarma-S (1996). Isolation and Chromosomal Distribution of Natural Z-DNA-forming Sequences in *Halobacterium halobium*. *J-Biol-Chem.* 271(33): 19724-31
- Klysik-J; Stirdvant-SM; Wells-RD (1982). Left-handed DNA. *J-Biol-Chem.* 257(17): 10152-8

- Konopka-AK; Reiter-J; Jung-M; Zarling-DA; Jovin-TM (1985). Accordance of experimentally mapped or predicted Z-DNA sites with positions of selected alternating purine-pyrimidine tracts. *Nucleic-Acids-Res.* 13(5): 1683-1701
- Krishna-P; Kennedy-BP; Waisman-DM; van-de-Sande-JH; McGhee-JD (1990). Are many Z-DNA binding proteins actually phospholipid-binding proteins? *Proc-Natl-Acad-Sci-U-S-A.* 87(4): 1292-5
- Krishna-P; Morgan-AR; van-de-Sande-JH (1991). Interaction of recA protein with left-handed Z-DNA. *Biochem-J.* 275 (Pt 3): 711-9
- Lafer-EM; Moller-A; Nordheim-A; Stollar-BD; Rich-A (1981). Antibodies specific for left-handed Z-DNA. *Proc-Natl-Acad-Sci-U-S-A.* 78(6): 3546-50
- Lafer-EM; Valle-RP; Moller-A; Nordheim-A; Schur-PH; Rich-A; Stollar-BD (1983). Z-DNA-specific antibodies in human systemic lupus erythematosus. *J-Clin-Invest.* 71(2): 314-21
- Lafer-EM; Sousa-RJ; Rich-A (1988). Z-DNA-binding proteins in *Escherichia coli* purification, generation of monoclonal antibodies and gene isolation. *J-Mol-Biol.* 203(2): 511-6
- Landt-O; Grunert-HP; Hahn-U (1990). A general method for rapid site-directed mutagenesis using the polymerase chain reaction. *Gene.* 96(1): 125-8
- Levinson-G; Gutman-GA (1987). High frequencies of short frameshifts in poly-CA/TG tandem repeats borne by bacteriophage M13 in *Escherichia coli* K-12. *Nucleic-Acids-Res.* 15(13): 5323-5338
- Liu-LF; Wang-JC (1987). Supercoiling of the DNA template during transcription. *Proc-Natl-Acad-Sci-U-S-A.* 84: 7024-7
- McLean-MJ; Blaho-JA; Kilpatrick-MW; Wells-RD (1986). Consecutive A-T pairs can adopt a left-handed DNA structure. *Proc-Natl-Acad-Sci-U-S-A.* 83:5884-8
- Menzel-R; Gellert-M (1983). Regulation of the genes for *E.coli* DNA gyrase: Homeostatic control of DNA supercoiling. *Cell.* 34: 105-13
- Mercado-CM; Tomasz-M (1977). Circular dichroism of mitomycin-DNA complexes. Evidence for a conformational change in DNA. *Biochemistry.* 16: 2040-6
- Möller-A; Gabriels-JE; Lafer-EM; Nordheim-A; Rich-A, Stollar-BD (1982). Monoclonal antibodies recognize different parts of Z-DNA. *J-Biol-Chem.* 257(20): 12081-5
- Möller-A; Nordheim-A; Kozlowski-SA; Patel-DJ; Rich-A (1984). Bromination stabilizes poly(dG-dC) in the Z-DNA form under low-salt conditions. *Biochemistry.* 23(1): 54-62
- Müller-V; Takeya-M; Brendel-S; Wittig-B; Rich-A (1996). Z-DNA forming sites within the human β -globin gene cluster. *Proc-Natl-Acad-Sci-U-S-A.* 93: 780-4

- Murphy-KE; Stringer-JR (1986). RecA independent recombination of poly[d(GT)-d(CA)] in pBR322. *Nucleic-Acids-Res.* 14(18): 7325-40
- Nordheim-A; Pardue-ML; Lafer-EM; Moller-A; Stollar-BD; Rich-A (1981). Antibodies to left-handed Z-DNA bind to interband regions of *Drosophila* polytene chromosomes. *Nature.* 294(5840): 417-22
- Nordheim-A; Meese-K (1988). Topoisomer gel retardation: detection of anti-Z-DNA antibodies bound to Z-DNA within supercoiled DNA minicircles. *Nucleic-Acid-Res.* 16(1): 21-37
- Oliphant-AR; Brand-CJ; Struhl-K (1989). Defining the Sequence Specificity of DNA-Binding Proteins by Selecting Binding Sites from Random-Sequence Oligonucleotides: Analysis of Yeast GCN4 Protein. *Mol-Cell-Biol.* 9: 2444-9
- Patterson-JB; Samuel-CE (1995). Expression and Regulation by Interferon of a Double-Stranded-RNA-Specific Adenosine Deaminase from Human Cells: Evidence for Two Forms of the Deaminase. *Molecular-and-Cellular-Biology.* 15(10): 5376-88
- Peck-LJ; Nordheim-A; Rich-A; Wang-JC (1982). Flipping of cloned d(pCpG)_n-d(pGpC)_n DNA sequences from right- to left-handed helical structure by salt, Co(III), or negative supercoiling. *Proc-Natl-Acad-Sci-U-S-A.* 79; 4560-4
- Pohl-FM; Jovin-TM (1972). Salt-induced Co-operative Conformational Change of a Synthetic DNA: Equilibrium and Kinetic Studies with Poly(dG-dC). *J-Mol-Biol.* 67: 375-96
- Pollock-R; Treisman-R (1990). A sensitive method for the determination of protein-DNA binding specificities. *Nucleic-Acids-Research.* 18(21): 6197-204
- Rahmouni-AR, Wells-RD (1989). Stabilization of Z DNA *in vivo* by localized supercoiling. *Science.* 246:358-63
- Rich-A; Nordheim-A; Wang-AH (1984). The chemistry and biology of left-handed Z-DNA. *Annu-Rev-Biochem.* 53: 791-846
- Rich-A (1994). Speculation on the biological roles of left-handed Z-DNA. *Ann-N-Y-Acad-Sci.* 726: 1-16; Discussion 16-7
- Rohner-KJ; Hobi-R; Kuenzle-CC (1990). Z-DNA-binding proteins. Identification critically depends on the proper choice of ligands. *J-Biol-Chem.* 265(31): 19112-5
- Sarkar-G; Bolander-ME (1995). Semi exponentiell cycle sequencing. *Nucleic-Acids-Research.* 23(7): 1269-70
- Sastray-M; Fiala-R; Lipman-R; Tomasz-M; Patel-DJ (1995). Solution structure of the mono alkylated mitomycin C-DNA complex. *J-Mol-Biol.* 247: 338-59

- Scadden-AD; Smith-CW (1997). A ribonuclease specific for inosine-containing RNA: a potential role in antiviral defence ? EMBO J. 16(8): 2140-9
- Schroth-GP; Chou-PJ; Ho-PS (1992). Mapping Z-DNA in the human genome. Computer-aided mapping reveals a nonrandom distribution of potential Z-DNA-forming sequences in human genes. J-Biol-Chem. 267(17): 11846-55
- Shore-D; Langowski-J; Baldwin-RL (1981). DNA flexibility studied by covalent closure of short fragments into circles. Proc-Natl-Acad-Sci-U-S-A. 78(5): 4833-7
- Shore-D; Baldwin-RL (1983). Energetics of DNA Twisting. I. Relation between Twist and Cyclization Probability. J-Mol-Biol. 170: 957-81
- Sinden-RR; Kochel-TJ (1987). Reduced 4,5',8-trimethylpsoralen cross-linking of left-handed Z-DNA stabilized by DNA supercoiling. Biochemistry. 26(5): 1343-50
- Sinden-RR (1996). DNA Structure and Function (Academic Press). Page 211
- Smith-G (1992). Cloning in Fuse vectors; edition of February 10, 1992
- Sommer-B; Köhler-M; Sprengel-R; Seeburg-PH (1991). RNA editing in brain controls a determinant of ion flow in glutamate-gated channels. Cell. 67:11-9
- Takeuchi-H; Hanamura-N; Harada-I (1994). Structural specificity of peptides in Z-DNA formation and energetics of the peptide-induced B-Z transition of poly(dG-m5dC). J-Mol-Biol. 236(2): 610-7
- Thiesen-HJ; Bach-C (1990). Target Detection Assay (TDA): a versatile procedure to determine DNA binding sites as demonstrated on SP1 protein. Nucleic-Acids-Res. 18(11): 3203-3209.
- Tomasz-M; Barton-JK; Magliozzo-CC; Tucker-D; Lafer-EM; Stollar-BD (1983). Lack of Z-DNA conformation in mitomycin-modified polynucleotides having inverted circular dichroism. Proc-Natl-Acad-Sci-U-S-A. 80(10): 2874-8
- Tuerk-C; Gold-L (1990). Systematic Evolution of Ligands by Exponential Enrichment: RNA Ligands to Bacteriophage T4 DNA Polymerase. Science. 249: 505-10
- Tse-Dinh-YC; Beran-RK (1988). Multiple promoters for transcription of the *Escherichia coli* DNA topoisomerase I gene and their regulation by DNA supercoiling. J-Mol-Biol. 202: 735-42
- van de Sande-JH; McIntosh-LP; Jovin-TM (1982). Mn²⁺ and other transition metals at low concentrations induce the right-to-left helical transformation of poly[d(G-C)]. EMBO J. 1:777-82
- Votavova-H; Matlova-R; Sponar-J (1994). B-Z Transition of Synthetic Oligonucleotides Containing Non-Z Forming Elements. J-Biomol-Struct-Dyn. 12(1): 163-172

- Ulanov-BP; Matorina-TI; Pozdeev-PP (1992). Nitrogen Mustard Mechlorethamine fixes the Z-conformation in poly[d(G-C)]poly[d(G-C)] and plasmid DNA. Mol-Biol (Moskau); 26(3): 583-90
- Vahrson-W (1991). Computergestützte und experimentelle Untersuchungen zum Zusammenhang zwischen Transkription und dem Vorkommen von Z-DNA. Diploma Thesis FU-Berlin
- Wang-AHJ; Quigley-GJ; Kolpak-FJ; Crawford-JL; van Boom-JH; van der Marel-G; Rich-A (1979). Molecular structure of a left-handed double helical DNA fragment at atomic resolution. Nature. 282: 680-6
- Watson-JD; Crick-FHC (1953). Molecular structure of nucleic acids-A structure for deoxyribose nucleic acid. Nature. 171: 737-8
- Wilhelm-ML; Reinbolt-J; Gangloff-J; Dirheimer-G; Wilhelm-FX (1994). Transfer RNA binding protein in the nucleus of *Saccharomyces cerevisiae*. FEBS-Lett. 349(2): 260-4
- Wittig-B; Dorbic-T; Rich-A (1989). The level of Z-DNA in metabolically active, permeabilized mammalian cell nuclei is regulated by torsional strain. J-Cell-Biol. 108(3): 755-64
- Wittig-B; Dorbic-T; Rich-A (1991). Transcription is associated with Z-DNA formation in metabolically active permeabilized mammalian cell nuclei [published erratum appears in Proc-Natl-Acad-Sci-U-S-A. 88(15):6898] Proc-Natl-Acad-Sci-U-S-A. 88(6): 2259-63
- Wittig-B; Wölfl-S; Dorbic-T; Vahrson-W; Rich-A (1992). Transcription of human c-myc in permeabilized nuclei is associated with formation of Z-DNA in three discrete regions of the gene. EMBO-J. 11(12): 4653-63
- Wölfl-S; Vahrson-W, Herbert-A (1995) Analysis of Left-Handed Z-DNA in vivo (In DNA and Nucleoprotein Structure In Vivo; edited by Saluz-HP and Wiebauer-K R.G. Landes Company) : 137-58
- Wölfl-S; Martinez-C; Rich-A; Majzoub-JA (1996). Transcription of the human corticotropin-releasing hormone gene in NPLC cells is correlated with Z-DNA formation. Proc-Natl-Acad-Sci-U-S-A. 93: 3664-8
- Wu-HM; Dattagupta-N; Crothers-DM (1981). Solution structural studies of the A and Z forms of DNA. Proc-Natl-Acad-Sci-U-S-A. 78(11): 6808-11
- Zhang-S; Lockshin-C; Herbert-A; Winter-E; Rich-A (1992). Zuotin, a putative Z-DNA binding protein in *Saccharomyces cerevisiae*. EMBO-J. 11(10): 3787-96